

# Binding of calcium and magnesium to human cardiac troponin C

Received for publication, November 30, 2020, and in revised form, January 25, 2021. Published, Papers in Press, February 3, 2021, <https://doi.org/10.1016/j.jbc.2021.100350>

Kaveh Rayani<sup>1</sup>, Justin Seffernick<sup>2</sup>, Alison Yueh Li<sup>1,3</sup>, Jonathan P. Davis<sup>4</sup>, Anne Marie Spuches<sup>5</sup>, Filip Van Petegem<sup>3</sup>, R. John Solaro<sup>6</sup>, Steffen Lindert<sup>2</sup>, and Glen F. Tibbits<sup>1,7,8,\*</sup>

From the <sup>1</sup>Molecular Cardiac Physiology Group, Simon Fraser University, Burnaby, British Columbia, Canada; <sup>2</sup>Department of Chemistry and Biochemistry, Ohio State University, Columbus, Ohio, USA; <sup>3</sup>Department of Biochemistry and Molecular Biology, The University of British Columbia, Vancouver, British Columbia, Canada; <sup>4</sup>Department of Physiology and Cell Biology, The Ohio State University, Columbus, Ohio, USA; <sup>5</sup>Department of Chemistry, East Carolina University, 300 Science and Technology Building, Greenville, North Carolina, USA; <sup>6</sup>Department of Physiology and Biophysics and the Center for Cardiovascular Research, College of Medicine, University of Illinois at Chicago, Chicago, Illinois, USA; <sup>7</sup>Department of Molecular Biology and Biochemistry, Simon Fraser University, Burnaby, British Columbia, Canada; <sup>8</sup>Cardiac Group, BC Children's Hospital Research Institute, Vancouver, British Columbia, Canada

Edited by Enrique De La Cruz

Cardiac muscle thin filaments are composed of actin, tropomyosin, and troponin that change conformation in response to  $\text{Ca}^{2+}$  binding, triggering muscle contraction. Human cardiac troponin C (cTnC) is the  $\text{Ca}^{2+}$ -sensing component of the thin filament. It contains structural sites (III/IV) that bind both  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  and a regulatory site (II) that has been thought to bind only  $\text{Ca}^{2+}$ . Binding of  $\text{Ca}^{2+}$  at this site initiates a series of conformational changes that culminate in force production. However, the mechanisms that underpin the regulation of binding at site II remain unclear. Here, we have quantified the interaction between site II and  $\text{Ca}^{2+}/\text{Mg}^{2+}$  through isothermal titration calorimetry and thermodynamic integration simulations. Direct and competitive binding titrations with WT N-terminal cTnC and full-length cTnC indicate that physiologically relevant concentrations of both  $\text{Ca}^{2+}/\text{Mg}^{2+}$  interacted with the same locus. Moreover, the D67A/D73A N-terminal cTnC construct in which two coordinating residues within site II were removed was found to have significantly reduced affinity for both cations. In addition, 1 mM  $\text{Mg}^{2+}$  caused a 1.4-fold lower affinity for  $\text{Ca}^{2+}$ . These experiments strongly suggest that cytosolic-free  $\text{Mg}^{2+}$  occupies a significant population of the available site II. Interaction of  $\text{Mg}^{2+}$  with site II of cTnC likely has important functional consequences for the heart both at baseline as well as in diseased states that decrease or increase the availability of  $\text{Mg}^{2+}$ , such as secondary hyperparathyroidism or ischemia, respectively.

Cardiac troponin (cTn) is a heterotrimeric complex that includes components for  $\text{Ca}^{2+}$  binding (cardiac troponin C [cTnC]), inhibition of contraction (cardiac troponin I [cTnI]), and tropomyosin binding (cardiac troponin T [cTnT]) (1).  $\text{Ca}^{2+}$  binding to site II of cTnC is the precursor to a series of structural perturbations in the thin filament (TF) that

culminate in a force-generating reaction between the actin filament and myosin heads (1–5).

Human cTnC is a 161-amino acid protein composed of nine helices (N and A–H), which form four EF-hand or helix–loop–helix binding motifs (sites I–IV). Within these domains, residues in positions 1 (+x), 3 (+y), 5 (+z), 7 (–y), 9 (–x), and 12 (–z) contain oxygen atoms arranged in a pentagonal bipyramid allowing for coordination of metal cations (Figs. S1 and S2) (6–8). Skeletal muscle troponin TnC (sTnC) has four functional  $\text{Ca}^{2+}$ -binding motifs (9, 10). cTnC has a similar overall structure to sTnC but a slightly different primary sequence; the insertion of a valine at residue 28, along with the substitutions D29L and D31A, has rendered site I of cTnC non-receptive to  $\text{Ca}^{2+}$  binding (11, 12).

$\text{Ca}^{2+}$  binding to sites III and IV in the C domain of cTnC occurs with high affinity ( $\sim 10^7 \text{ M}^{-1}$ ) ( $\sim 10\times$  higher than the N domain) and a slow exchange rate ( $\sim 100\times$  slower than binding to the N domain) (13–15). Given the abundance of contractile filaments throughout cardiomyocytes, sites III/IV of cTnC buffer  $\sim 80\%$  of the 100 to 200  $\mu\text{M}$   $[\text{Ca}^{2+}]_{\text{in}}$  at resting concentrations of free  $\text{Ca}^{2+}$  ( $\sim 100 \text{ nM}$ ) (16). At resting free cytosolic  $\text{Ca}^{2+}$  concentrations, sites III and IV are usually saturated with  $\text{Ca}^{2+}$  (16).  $\text{Mg}^{2+}$  also binds at sites III and IV but with lower affinity ( $K_A \sim 10^4 \text{ M}^{-1}$ ) (4). However, the cytosolic concentration of  $\text{Mg}^{2+}$  allows this cation to compete with and reduce the binding of  $\text{Ca}^{2+}$  to the “structural” sites (17, 18). The binding of  $\text{Ca}^{2+}/\text{Mg}^{2+}$  to sites III and IV alters the structure of TnC and is a prerequisite for tethering to the rest of the TF (19, 20).

The C domain of cTnC is linked to the N domain by a linker region composed of a nine-turn  $\alpha$ -helix (21, 22). Within the N domain (N-cTnC),  $\text{Ca}^{2+}$  binds the low-affinity ( $\sim 10^{-5} \text{ M}$ ) site II such that this site is only partially occupied at diastolic-free  $\text{Ca}^{2+}$  concentrations ( $\sim 0.1 \mu\text{M}$ ) with very few sites being bound (23). The degree of occupancy is significantly higher at systolic-free  $\text{Ca}^{2+}$  concentrations ( $\sim 0.5$ – $1.2 \mu\text{M}$ ), which follow  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release (24).  $\text{Ca}^{2+}$  binding to site II provides

\* For correspondence: Glen F. Tibbits, [tibbits@sfu.ca](mailto:tibbits@sfu.ca).

## Ca and Mg binding to cTnC

the free energy to allow for exposure of a hydrophobic pocket, which is otherwise less favorable (25, 26). Helices B and C move away from helices N, A, and D to expose the hydrophobic cleft, with the short antiparallel  $\beta$ -sheet between EF-hands I and II acting as a hinge (27–29). Binding of the “switch peptide” of cTnI<sub>147–163</sub> to this pocket facilitates exposure of this hydrophobic region (22, 30, 31).

A persistent and underinvestigated question is the role of cellular Mg<sup>2+</sup> in the signaling of activation by cTnC. Of the total intracellular [Mg<sup>2+</sup>]<sub>i</sub> (~10 mM), the majority is bound to cellular components such as ATP with only ~0.5 to 1.0 mM being freely available in the cytosol (32, 33). In conditions with diminished Mg<sup>2+</sup> buffering capacity, such as ATP-depleted states, the free [Mg<sup>2+</sup>] can increase significantly (34, 35) prior to being extruded from the cell (36), but it could also compete with Ca<sup>2+</sup> for binding to cTnC.

Increase in Mg<sup>2+</sup>–ATP in both skeletal and cardiac tissues decreases the Ca<sup>2+</sup> sensitivity of skinned fibers (37–39). These systems contain the entirety of the TF. The binding of Ca<sup>2+</sup> normally induces a conformation change, which allows for exposure of the hydrophobic core of cTnC (40). It is unlikely that Mg<sup>2+</sup> would cause this structural change. Indeed, previous findings suggest that Mg<sup>2+</sup> competes for binding with Ca<sup>2+</sup> but does not itself cause the same conformational changes (20).

Further evidence has been obtained through fluorescence-based studies of isolated cTnC (41–46), the Tn complex (41, 43), and reconstituted fibers (43, 47), where Mg<sup>2+</sup> appears to decrease Ca<sup>2+</sup> sensitivity. In isolated TnC, the  $K_A$  of the low-affinity sites (III/IV) for Ca<sup>2+</sup> and Mg<sup>2+</sup> was measured to be on the order of 10<sup>6</sup> and 10<sup>2</sup> M<sup>-1</sup>, respectively (42).

Interaction of Ca<sup>2+</sup>/Mg<sup>2+</sup> with sites III/IV results in a large change in enthalpy ( $\Delta H$ ), in contrast to changes resulting from site I/II binding, which are small by comparison. Detection of heat changes associated with the interactions of metal ions and proteins is both challenging and a highly technique-dependent method, such that small changes may be deemed negligible (48–50). Experiments used to study this system decades ago were limited by the technology of the time. In contrast, modern isothermal titration calorimetry (ITC) is a sensitive method that can be used to define the thermodynamic parameters of binding without the use of labeling methods that could interfere. While there is a place for use of fluorophores to investigate cTnC function (51–53), even conservative substitutions (e.g., sTnC F29W) have been shown to modify Ca<sup>2+</sup>-binding properties (54). Modern ITC allows for the study of single-binding sites within isolated protein domains and can be used to detect heat changes as small as 0.1  $\mu$ cal (55–58).

We have used ITC to explore the binding of Mg<sup>2+</sup> and Ca<sup>2+</sup> to site II at the level of N-cTnC and full-length cTnC. Competitive binding to the N domain and mutations in the site II caused a reduction in apparent affinity, indicating interaction of both cations with the same locus in the protein. In addition, a double mutant removing two of the coordinating residues within the EF-hand of site II was used to investigate the impact on Ca<sup>2+</sup> and Mg<sup>2+</sup> binding to this site. In full-length cTnC, Mg<sup>2+</sup> competed with and reduced Ca<sup>2+</sup> binding to all three sites. These findings further corroborate and expand

upon what has been shown by a few laboratories, but the findings are largely ignored by most; the role of Mg<sup>2+</sup> in modulating the Ca<sup>2+</sup> sensitivity of force production in cardiomyocytes is one that merits further discussion (40, 51–53).

## Results

The ratio of the ligand to titrant in the single-binding site condition (as given by the stoichiometry—N) is a measure of the functional moles of protein and was approximately 1.00 in all the N-cTnC titrations (Table S2). Given the method of concentration determination, the number of binding sites, cooperativity, and the variable binding strength of each titrants, the N cannot be used in the same way for the full-length cTnC experiments (Table S3). Therefore, the values presented can be compared between conditions, but care should be taken when comparing these to other systems, such as the cTn complex or the reconstituted thin filaments (59, 60). Ease of manipulation of the N-cTnC/cTnC system contrasts with those that include the cTn/TF. Thus, the binding parameters measured here may not translate in absolute term when cTnC is incorporated into a more complex system; these limitations are further explored in the Discussion section.

### Ca<sup>2+</sup> and Mg<sup>2+</sup> binding to apo-state N-cTnC

The interaction of N-cTnC with either Ca<sup>2+</sup> or Mg<sup>2+</sup> was found to be associated with a positive  $\Delta H$ , so the interaction is driven by entropy (Fig. 1) consistent with previously published data (61–64). Given the characteristic of the heat signals observed in the site II-containing system of N-cTnC, the endothermic component of the multiple binding site system (full-length cTnC) can be attributed to binding at this site within the N domain.

The affinity of N-cTnC for Ca<sup>2+</sup> ( $K_d = 15.2 \pm 0.5 \mu$ M) was found to be significantly greater ( $p = 0.001$ ) and 42.9-fold different than for Mg<sup>2+</sup> ( $K_d = 652.8 \pm 28.4 \mu$ M) (Figs. 1 and 2; Table S2).


The  $\Delta H$  of the Ca<sup>2+</sup>–N-cTnC interaction ( $3.82 \pm 0.04 \text{ kcal} \cdot \text{mol}^{-1}$ ) was significantly greater ( $p < 0.0001$ ) than that with Mg<sup>2+</sup> ( $2.64 \pm 0.10 \text{ kcal} \cdot \text{mol}^{-1}$ ), indicating a greater enthalpic cost of binding for the Ca<sup>2+</sup> titration. Moreover, the entropic contribution for the Ca<sup>2+</sup> titrations ( $T \cdot \Delta S = 10.39 \pm 0.03 \text{ kcal} \cdot \text{mol}^{-1}$ ) was more favorable ( $p < 0.0001$ ) than the Mg<sup>2+</sup> titrations ( $T \cdot \Delta S = 6.99 \pm 0.07 \text{ kcal} \cdot \text{mol}^{-1}$ ) (Table S2).

As expected, the affinity of Ca<sup>2+</sup> binding to apo-state N-cTnC ( $K_d = 15.2 \pm 0.5 \mu$ M) was found to be greater than Mg<sup>2+</sup> binding at this site, or when compared with the preincubation experiments (Fig. 2 and Table S2).

### N-terminal cTnC

#### Mg<sup>2+</sup> binding to Ca<sup>2+</sup>-preincubated N-cTnC

To investigate Mg<sup>2+</sup> binding in the presence of Ca<sup>2+</sup>, apo-state N-cTnC was preincubated with three concentrations of Ca<sup>2+</sup> (0, 1, and 3 mM), then titrated with 20 mM Mg<sup>2+</sup> (Figs. 1 and 2 and Table S2). Moreover, the change in enthalpy in these conditions was lower with greater amounts of Ca<sup>2+</sup> preincubated. Titration of Mg<sup>2+</sup> into apo-state protein yielded a  $\Delta H = 2.64 \pm 0.10 \text{ kcal} \cdot \text{mol}^{-1}$ , lower than Ca<sup>2+</sup> into apoprotein, which liberated



**Figure 1. Representative isotherms of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  binding to apo-state and preincubated N-cTnC.** All titrations were carried out into 200  $\mu\text{M}$  WT human N-cTnC in 50 mM Hepes at pH 7.2, 150 mM KCl, and 2 mM EDTA. The *top row* shows the titration of 4 mM  $\text{Ca}^{2+}$  into apo-state N-cTnC, followed by the same titration into 1 and 3 mM  $\text{Mg}^{2+}$  preincubated N-cTnC. The *bottom row* shows the titration of 20  $\text{Mg}^{2+}$  into apo-state N-cTnC, followed by 1 and 3 mM  $\text{Ca}^{2+}$ -preincubated N-cTnC. In the last injections,  $\text{Mg}^{2+}$  in the  $\text{Ca}^{2+}$ -preincubation conditions, the heat changes approached the detection limits of the instrument. Thermograms were fit to a “single-binding site” model using Origin 7 MicroCal2000 ITC software package. N-cTnC, cardiac troponin C with N domain.

$3.82 \pm 0.04 \text{ kcal} \cdot \text{mol}^{-1}$ . Moreover, the  $K_d$  values were  $1870.0 \pm 171.5$  and  $2037.5 \pm 172.2 \mu\text{M}$  for the 1 and 3 mM  $\text{Ca}^{2+}$  conditions, respectively, indicating a trend of decreasing affinity with increasing concentrations of  $\text{Ca}^{2+}$  preincubated with the protein sample and a more than two orders of magnitude lower affinity compared with the  $\text{Ca}^{2+}$  into WT condition. The reduction in affinity,  $\Delta H$ , and lower  $\Delta S$  associated with higher  $\text{Ca}^{2+}$  preincubation suggests that both metal cations may be binding to the same EF-hand-binding motif in site II of N-cTnC.


#### *Ca<sup>2+</sup> binding to Mg<sup>2+</sup>-preincubated N-cTnC*

Apoprotein preincubated with  $\text{Mg}^{2+}$  was titrated with  $\text{Ca}^{2+}$  to assess the “apparent” affinity of the protein for  $\text{Ca}^{2+}$  when

the site might be occupied by the other divalent cation. As expected, increasing the  $\text{Mg}^{2+}$  concentration significantly ( $p < 0.0001$ ) reduced the  $\Delta H$  associated with binding from  $3.82 \pm 0.04 \text{ kcal} \cdot \text{mol}^{-1}$  in the apotitration to  $1.73 \pm 0.05 \text{ kcal} \cdot \text{mol}^{-1}$  in the 3 mM  $\text{Mg}^{2+}$  preincubated construct. The binding affinity changed from  $15.2 \pm 0.5 \mu\text{M}$  in the apo-state to  $48.9 \pm 2.8 \mu\text{M}$  in the 3 mM  $\text{Mg}^{2+}$ -preincubated condition. The  $\text{Ca}^{2+}$  affinity was lower when comparing the apo-N-cTnC binding condition with higher concentrations of preincubated  $\text{Mg}^{2+}$  (Fig. 2 and Table S2).

It is possible to directly obtain a measure of the binding of a secondary ion when a competition experiment is carried out. Figure 3 shows the binding of  $\text{Ca}^{2+}$  to N-cTnC in the apo-state

## Ca and Mg binding to cTnC



**Figure 2. Binding of Ca<sup>2+</sup> and Mg<sup>2+</sup> to apo-state and preincubated N-cTnC.** *Left panel*, the affinity of site II for Mg<sup>2+</sup> is compared in the apo-state and with Ca<sup>2+</sup> preincubation in human N-cTnC. *Right panel*, the affinity of site II for Ca<sup>2+</sup> is compared in the apo-state and with Mg<sup>2+</sup> preincubation in N-cTnC. Statistical differences were assessed through ANOVA followed by Tukey's post hoc test. Ca<sup>2+</sup> titrations were not significantly different. Titration of Mg<sup>2+</sup> into N-cTnC preincubated with 1 and 3 mM Ca<sup>2+</sup> was statistically indifferent, but both significantly differed from the apo-state titration ( $p < 0.0001$ ). N-cTnC, cardiac troponin C with N domain.

and following preincubation with 1 mM Mg<sup>2+</sup>. This figure also shows the binding of Mg<sup>2+</sup> to N-cTnC in the apo-state and following preincubation with 1 mM Ca<sup>2+</sup>. The values of the thermodynamic parameters obtained from this model for the apo-state titration were as follows:  $N = 1.00 \pm 0.015$  sites,  $K_A = 4.58 \pm 0.37 \times 10^4 \text{ M}^{-1}$ , and  $\Delta H = 4.11 \pm 0.08 \text{ kcal/mol}$ ; these values are comparable to those obtained from the single-binding site model (Table S2). The parameters for the preincubation condition were as follows:  $N = 1.18 \pm 0.02$  sites,  $K_A = 5.60 \pm 0.38 \times 10^4 \text{ M}^{-1}$ , and  $\Delta H = 4.15 \pm 0.07 \text{ kcal/mol}$  for Ca<sup>2+</sup>; Mg<sup>2+</sup> was estimated to bind with a  $K_A$  of  $1.54 \times 10^3 \text{ M}^{-1}$  and a  $\Delta H$  of 2.64 kcal/mol; these values fall within the range obtained using the single-binding site model (Table S2). Moreover, the values obtained for the binding of Mg<sup>2+</sup> to the apo-state N-cTnC and with preincubated Ca<sup>2+</sup> are comparable to a single-binding site model (Fig. 3 and Table S2). The  $N$ -value obtained when applying the competitive binding model to the titration of Mg<sup>2+</sup> into 1 mM Ca<sup>2+</sup> preincubated N-cTnC ( $4.48 \pm 0.06$ ) is thought to result from a weak binding interaction and indicates a less reliable fit under the experimental conditions (Fig. 3).

### Ca<sup>2+</sup> and Mg<sup>2+</sup> binding to apo-D67A/D73A N-cTnC

Point mutations (D67A and D73A) were made (Fig. S2), affecting two known Ca<sup>2+</sup> coordinating residues in site II. Binding of both divalent cations was reduced by these mutations, but the  $K_d$  was lower for Ca<sup>2+</sup> binding ( $180.3 \pm 16.2 \text{ μM}$ ) compared with Mg<sup>2+</sup> binding ( $1148.6 \pm 95.0 \text{ μM}$ ) (Figs. 4 and 5 and Table S2). The double mutant caused a 11.9-fold alteration in Ca<sup>2+</sup> binding, yet this difference was not found to be statistically significant ( $p = 0.88$ ); it also altered Mg<sup>2+</sup> binding 1.8-fold ( $p = 0.04$ ); this change in  $K_d$  supports the binding of Mg<sup>2+</sup> to the EF-hand of site II.

### Ca<sup>2+</sup>- and Mg<sup>2+</sup>-binding affinities from thermodynamic integration

Thermodynamic integration (TI) was performed to calculate absolute binding affinities (with change in Gibbs free energy


reported) for the ions in the following systems: Ca<sup>2+</sup> to WT N-cTnC, Ca<sup>2+</sup> to D67A/D73A N-cTnC, Mg<sup>2+</sup> to WT N-cTnC, and Mg<sup>2+</sup> to D67A/D73A N-cTnC (Table 1). The average calculated binding affinities over five independent runs were  $-6.9 \pm 1.3$ ,  $-4.5 \pm 2.4$ ,  $-0.6 \pm 2.8$ , and  $+0.4 \pm 2.3 \text{ kcal * mol}^{-1}$ , respectively. The TI-determined Ca<sup>2+</sup>-binding affinities were in good agreement with the ITC data. While the calculated absolute Mg<sup>2+</sup>-binding affinities were not in perfect agreement with the ITC data, they did show that Mg<sup>2+</sup> had a weaker binding affinity than Ca<sup>2+</sup> for all systems ( $-6.57$  to  $-4.38 \text{ kcal * mol}^{-1}$  and  $-6.9$  to  $-0.6 \text{ kcal * mol}^{-1}$  for ITC and TI, respectively, for WT system and  $-5.12$  to  $-4.02 \text{ kcal * mol}^{-1}$  and  $-4.5$  to  $+0.4 \text{ kcal * mol}^{-1}$  for ITC and TI, respectively, for D67A/D73A system). In addition, between the Mg<sup>2+</sup>-binding affinities, the binding affinity was consistently weaker for the D67A/D73A mutation. The  $\Delta\Delta G$  values comparing  $\Delta G$  between WT and D67A/D73A systems were similar for ITC and TI ( $0.36 \text{ kcal * mol}^{-1}$  and  $1.0 \text{ kcal * mol}^{-1}$ , respectively).

### Full-length cTnC

#### Ca<sup>2+</sup> binding to apo-state full-length cTnC

Binding of Ca<sup>2+</sup>/Mg<sup>2+</sup> to site II is characterized by an endothermic interaction as indicated by our titrations on the N-terminal domain in this and previous publications (63, 64). From this and the ITC work on full-length cTnC by others (65), we can deduce that the exothermic heat changes seen in Figure 6 result from interactions with site III/IV. The data (Fig. 6 and Table S3) show that Mg<sup>2+</sup> binds to apo-state full-length cTnC at two distinct sets of sites.

The binding of Ca<sup>2+</sup> to sites III/IV occurred with an apparent  $K_d$  of  $0.12 \pm 0.02 \text{ μM}$ , characterized by an exothermic component ( $\Delta H = -8.12 \pm 0.07 \text{ kcal * mol}^{-1}$ ) with a positive change in entropy ( $T * \Delta S = 1.24 \pm 0.07 \text{ kcal * mol}^{-1}$ ). In the same full-length construct, the  $K_d$  associated with binding of Ca<sup>2+</sup> to site II was  $22.7 \pm 0.5 \text{ μM}$ . This interaction had a positive  $\Delta H$  ( $3.71 \pm 0.06 \text{ kcal * mol}^{-1}$ ) and was entropically driven ( $T * \Delta S = 10.0 \pm 0.01 \text{ kcal * mol}^{-1}$ ) (Figs. 6 and 7 and Table S3).




**Figure 3. Competition model of Ca<sup>2+</sup> and Mg<sup>2+</sup> binding to apo-state and N-cTnC preincubated with 1 mM Mg<sup>2+</sup> or 1 mM Ca<sup>2+</sup>.** All titrations were carried out into 200  $\mu\text{M}$  WT human N-cTnC in 50 mM Hepes at pH 7.2, 150 mM KCl, and 2 mM EDTA. The *top two panels* show Ca<sup>2+</sup> titration, and the *bottom two panels* show Mg<sup>2+</sup> titrations. The *top left panel* shows the titration of 4 mM Ca<sup>2+</sup> into apo-state N-cTnC, and the *top right panel* illustrates the same titration with 1 mM Mg<sup>2+</sup>-preincubated N-cTnC. The *bottom left panel* shows the titration of 20 mM Mg<sup>2+</sup> into apo-state N-cTnC, and the *bottom right panel* illustrates the same titration with 1 mM Ca<sup>2+</sup>-preincubated N-cTnC. The thermograms were fit using a "competition" model using Origin 7 MicroCal2000 ITC software package, in which the concentration of the ion in the cell and an estimated  $K_A$  value is input. The values for thermodynamic parameters obtained are listed in the table described. Thermodynamic parameters without a reported error value were fixed in the model. N-cTnC, cardiac troponin C with N domain.

#### Mg<sup>2+</sup> binding to apo-state full-length cTnC

Mg<sup>2+</sup> binding to site II ( $K_d = 406.1 \pm 7.9 \mu\text{M}$ ) and sites III/IV ( $K_d = 16.7 \pm 0.7 \mu\text{M}$ ) was characterized by a positive  $\Delta H$  ( $0.091 \pm 0.001 \text{ kcal} \cdot \text{mol}^{-1}$ ) and negative  $\Delta H$  ( $-0.23 \pm 0.01 \text{ kcal} \cdot \text{mol}^{-1}$ ), respectively (Fig. 6 and Table S3). The difference in  $K_d$  values indicates that greater amounts of Mg<sup>2+</sup> binding occur at the C-terminal sites, in comparison to the N-terminal

site of cTnC. The interaction of Mg<sup>2+</sup> with sites III/IV is two orders of magnitude weaker ( $p < 0.0001$ ) than that seen for Ca<sup>2+</sup>. The interaction of Mg<sup>2+</sup> with site II and sites III/IV was both entropically favorable ( $T \cdot \Delta S = 4.71 \pm 0.01 \text{ kcal} \cdot \text{mol}^{-1}$  and  $T \cdot \Delta S = 6.28 \pm 0.03 \text{ kcal} \cdot \text{mol}^{-1}$ , respectively) and resulted in spontaneous interactions ( $\Delta G = -4.62 \pm 0.11 \text{ kcal} \cdot \text{mol}^{-1}$  and  $\Delta G = -6.51 \pm 0.31 \text{ kcal} \cdot \text{mol}^{-1}$ , respectively).



**Figure 4. Representative binding isotherms for binding of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  WT and D67A/D73A N-cTnC.** All titrations were carried out into 200  $\mu\text{M}$  human N-cTnC in 50 mM Hepes at pH 7.2, 150 mM KCl, and 2 mM EDTA. *Top row*, titration of 4 mM  $\text{Ca}^{2+}$  into apo-state N-cTnC, followed by the same titration into the D67A/D73A mutant. *Bottom row*, titrations of 20 mM  $\text{Mg}^{2+}$  into apo-state N-cTnC, followed by the same titration into the D67A/D73A mutant. Thermograms were fit to a “single-binding site” model using Origin 7 MicroCal2000 ITC software package. N-cTnC, cardiac troponin C with N domain.


#### $\text{Ca}^{2+}$ binding to $\text{Mg}^{2+}$ -preincubated full-length cTnC

At greater concentrations,  $\text{Mg}^{2+}$  occupied a greater proportion of binding sites and limited binding  $\text{Ca}^{2+}$  to cTnC at all sites (Figs. 6 and 7 and Table S3). Binding of  $\text{Ca}^{2+}$  to site II was reduced by preincubation with 1 and 3 mM  $\text{Mg}^{2+}$  as indicated by an increase in  $K_d$ ; however, these changes were not found to be statistically significant ( $p = 0.52$  and  $p = 0.14$ ), moreover  $\Delta H$  was lowered ( $p < 0.0001$  for both). Binding of  $\text{Ca}^{2+}$  to sites III/IV in the presence of 1 mM  $\text{Mg}^{2+}$  resulted in a  $K_d$  ( $0.14 \pm 0.01 \mu\text{M}$ ) that was not significantly different ( $p < 0.05$ ) than seen for the 3 mM  $\text{Mg}^{2+}$  preincubation ( $K_d = 0.08 \pm 0.01 \mu\text{M}$ ) (Fig. 6 and Table S3).

At sites III/IV, for the 1 mM  $\text{Mg}^{2+}$  preincubation, the interaction proceeded with favorable enthalpy ( $\Delta H = -6.87 \pm 0.09 \text{ kcal} \cdot \text{mol}^{-1}$ ) and entropy ( $T \cdot \Delta S = 2.50 \pm 0.10 \text{ kcal} \cdot \text{mol}^{-1}$ ). For the 3 mM  $\text{Mg}^{2+}$  condition, the reaction was again exothermic ( $\Delta H = -6.19 \pm 0.06 \text{ kcal} \cdot \text{mol}^{-1}$ ) with a positive change in  $T \cdot \Delta S$  ( $3.50 \pm 0.06 \text{ kcal} \cdot \text{mol}^{-1}$ ).

#### Discussion

This study provides novel information regarding the thermodynamics that underlie the interaction between cTnC and two physiologically prevalent divalent cations. Our results significantly advance the understanding of the mechanisms



**Figure 5. Binding of Ca<sup>2+</sup> and Mg<sup>2+</sup> to WT and D67A/D73A N-cTnC.** The effect of the D67A/D73A on Ca<sup>2+</sup> and Mg<sup>2+</sup> binding is assessed. ANOVA indicated that the binding affinity for both cations to human N-cTnC was lower when comparing the mutant and the WT ( $p < 0.0001$ ). Tukey's post hoc test indicated that all four test conditions were significantly different. The effect on Ca<sup>2+</sup> binding was more pronounced (11.9-fold reduction) compared with Mg<sup>2+</sup> (1.8-fold reduction), but this is reconcilable with the number of coordinating residues needed to bind Ca<sup>2+</sup> (6) versus Mg<sup>2+</sup> (5); having four available coordinating residues was expected to affect Ca<sup>2+</sup> binding to a greater extent. N-cTnC, cardiac troponin C with N domain.

and role of modifications in cellular Mg<sup>2+</sup> in control of the cTnC–Ca<sup>2+</sup> switch. Cellular-free Mg<sup>2+</sup> is known to change in pathological conditions in the heart (66), but the mechanisms of its effects on activation of the myofilaments remain incompletely understood.

As seen in previous reports, we found the binding of Ca<sup>2+</sup> to N-cTnC to be driven by entropy and unfavorable enthalpy (Table S2) (64, 65). The favorable  $\Delta S$  may be due in part to the hydration enthalpy of Ca<sup>2+</sup>, which is thought to be on the order of  $\sim 375$  kcal  $\cdot$  mol<sup>-1</sup> and slightly lower than that of Mg<sup>2+</sup> ( $\sim 460$  kcal  $\cdot$  mol<sup>-1</sup>) (67). It is also possible that the endothermic nature of these interactions results from other factors such as the exchange of protons that are transferred from the ligand to the buffer upon Ca<sup>2+</sup> binding (61).

Measurement of Ca<sup>2+</sup> binding to cTnC is often achieved indirectly by monitoring the fluorescence change and correlating this to the conformational change that results from the interaction. Fluorescent molecules such as 2-[4'-(iodoacetamido)anilino]naphthalene-6-sulfonic acid (59, 60, 68, 69) or reporters such as F27W (51, 70) can be used to quantify this binding interaction. At 21 °C, bovine F27W cTnC had a K<sub>d</sub> of  $\sim 5$   $\mu$ M, and 2-[4'-(iodoacetamido)anilino]naphthalene-6-sulfonic acid-labeled C35S cTnC had a K<sub>d</sub> of  $\sim 7$   $\mu$ M (20, 71). Through fluorescence-based measurement, the K<sub>d</sub> of N-cTnC for Ca<sup>2+</sup> was previously reported to be between 11.3 and 12.3  $\mu$ M (70, 72). These parameters agree with our measured Ca<sup>2+</sup> binding to apo-state N-cTnC and apo-state cTnC (Table S2), deviating only slightly, likely because of, different buffer and temperature conditions.

Normally, cytosolic [Mg<sup>2+</sup>]<sub>free</sub> is maintained around  $\sim 0.5$  to 1 mM (73). At these concentrations, Mg<sup>2+</sup> is known to

compete with Ca<sup>2+</sup> for sites III and IV. Circular dichroism has been used to show that Ca<sup>2+</sup> binding to sites III/IV increases the  $\alpha$ -helical content of cTnC, from 19 to 41% (27, 74) and causes conformational changes that remove nonpolar amino acids from the solvent-exposed environment (19). This contrasts with NMR-based visualization of N-cTnC, in which the apo-state and Ca<sup>2+</sup>-bound forms showed minimal structural deviation (30).

Ca<sup>2+</sup> has many times greater polarizability than Mg<sup>2+</sup> and lower hydration energy (75). The bare ion radius of Mg<sup>2+</sup> is smaller than Ca<sup>2+</sup> (0.65 versus 0.99 Å) (76); conversely, in its hydrated form, Mg<sup>2+</sup> is larger than Ca<sup>2+</sup> (4.3 versus 4.1 Å) (77). In other Ca<sup>2+</sup>-binding proteins such as calmodulin (CaM), metals with similar ionic radii are able to substitute for this cation (78, 79). Mg<sup>2+</sup> is able to bind to CaM but does not induce the conformational change associated with Ca<sup>2+</sup> binding; a phenomenon that is commonly observed in cell biology and is expected in cTnC (80, 81).

Normally, six oxygen atoms arranged in an octahedral geometry are thought to coordinate Mg<sup>2+</sup> (82). This is one less oxygen than needed to coordinate Ca<sup>2+</sup> through a pentagonal bipyramid (83). However, Ca<sup>2+</sup> can be coordinated by six to eight coordinating residues (but also by as many as 12) at a distance that can vary greatly (2.3–2.7 Å) compared with a much smaller variance for Mg<sup>2+</sup> coordination (2.0–2.2 Å) (84).

Ca<sup>2+</sup> and Mg<sup>2+</sup> are most often coordinated by oxygen atoms, and this is usually accomplished by a hydroxyl group for Mg<sup>2+</sup> and a carboxyl group for Ca<sup>2+</sup> (85). Ca<sup>2+</sup> is most frequently coordinated by side chains of aspartic acid, glutamic acid, asparagine, followed by serine/threonine, whereas Mg<sup>2+</sup> is most frequently coordinated by aspartic acid, glutamic acid, histidine, threonine, serine, or asparagine (86). EF-hand-containing proteins have also been shown to bind Mg<sup>2+</sup> when there are appropriately placed negatively charged amino residues (especially in the +z and -z positions) (54, 87, 88). In site II of mammalian cTnC, there is a polar serine at the +z position (residue 69) and a negatively charged glutamic acid at the -z position (residue 76) (Fig. S1).

Data from earlier studies suggested that Mg<sup>2+</sup> binds exclusively at sites III and IV of TnC (4). Shortly thereafter, a limited

**Table 1**  
Average calculated binding affinities for Ca<sup>2+</sup>/Mg<sup>2+</sup> interaction with site II of N-cTnC thermodynamic integration system

System	$\Delta G_{TI}$ (kcal $\cdot$ mol <sup>-1</sup> )
Ca <sup>2+</sup> to WT	-6.9 $\pm$ 1.3
Ca <sup>2+</sup> to D67A/D73A	-4.5 $\pm$ 2.4
Mg <sup>2+</sup> to WT	-0.6 $\pm$ 2.8
Mg <sup>2+</sup> to D67A/D73A	+0.4 $\pm$ 2.0

Averages were calculated over five independent runs.

## Ca and Mg binding to cTnC

series of equilibrium dialysis experiments did not show competition between  $Mg^{2+}$  and  $Ca^{2+}$  for the N-terminal sites of cTnC; instead, other binding sites were suggested (89). Later still, enthalpic titrations were unable to visualize a discernable change in  $Mg^{2+}$  binding to the low-affinity sites of sTnC (50, 90). However, assuming competitive binding, fluorescence assays at room temperature determined the  $K_d$  associated with  $Mg^{2+}$  binding to be about 4 mM (91). Moreover,  $Ca^{2+}$  sensitivity of the actomyosin ATPase and force production of skinned rat cardiac cells were unaltered when  $Mg^{2+}$  was increased from 1 to 8 mM (92). However, these findings were brought into question by studies that utilized metallochromic indicators to deduce sufficiently high  $Mg^{2+}$  affinity at the regulatory sites of sTnC (42).

The observation of  $Mg^{2+}$  binding to the low-affinity site of N-cTnC has led to the suggestion that differences in affinity may be due, at least in part, to  $Ca^{2+}$  buffering, and thus, the free concentration of the ion in these experiments. Given the kinetic rates associated with these interactions, it is difficult to have confidence in EGTA-determined rates of binding (93). Moreover, the temperature sensitivity of cTnC alone can alter experimental outcomes by orders of magnitude (53, 94). Change in sensitivity in the face of altered temperature has been suggested to result mostly from binding to the low-affinity sites and possibly through interactions with other members of the cTn complex (95–97).

Experiments testing the effects of alterations in free  $Mg^{2+}$  on  $Ca^{2+}$  activation of isolated myofibrils and skinned fiber bundles from different laboratories provide corroborative findings supporting the credibility of our postulate of a role for cytosolic  $Mg^{2+}$  as a controller of cTnC function at the N lobe. Fabiato and Fabiato (98) showed that increasing the concentration of free  $Mg^{2+}$  decreases myofilament  $Ca^{2+}$  sensitivity of skinned cardiomyocytes.  $[Mg^{2+}]$  affects the  $Ca^{2+}$  sensitivity of the myofibrillar ATPase as well as actomyosin tension development in both skeletal and cardiac muscle preparations (10, 39, 44, 99–103).

$Mg^{2+}$  affinity of sites III/IV alone is not sufficient to fully explain the change in the force–negative log of  $Ca^{2+}$  concentration relationship caused by  $Mg^{2+}$  in skinned skeletal muscle fibers (104). In rabbit fast skeletal muscle,  $Mg^{2+}$  competes with  $Ca^{2+}$  for low-affinity binding sites of TnC, where it binds with an affinity of  $1.9 \times 10^2 M^{-1}$  (much lower than the  $6.2 \times 10^6 M^{-1}$  seen for  $Ca^{2+}$ ). The  $K_A$  associated with sites III and IV was measured to be  $1.2 \times 10^6 M^{-1}$  for  $Ca^{2+}$  and  $1.1 \times 10^2 M^{-1}$  for  $Mg^{2+}$  in canine ventricular skinned myocytes (105).

In isolated cTnC,  $Mg^{2+}$  was found to interact with site II with an apparent binding constant of  $5.2 \times 10^2 M^{-1}$ . This was only slightly lower than the constant associated with  $Mg^{2+}$  binding to sites III/IV ( $\sim 10^3 M^{-1}$ ),  $Ca^{2+}$  binding to sites III/IV ( $\sim 10^6 M^{-1}$ ), and  $Ca^{2+}$  binding to site II ( $\sim 10^4 M^{-1}$ ) (42).

Fluorescent probes were used to measure the  $Mg^{2+}$  affinity of site II at 15 °C ( $\sim 1.2$ – $1.9$  mM) (20). In the presence of 3 mM  $Mg^{2+}$ , the  $K_d$  associated with binding of  $Ca^{2+}$  to site II of full-length cTnC was increased from 7  $\mu M$  in the apo-state to 24  $\mu M$  (20). Moreover, a system containing cTnC–cTnI had 2.5-fold lower  $Ca^{2+}$  affinity in the presence of 3 mM  $Mg^{2+}$  (106). Given these affinities, Tikunova and Davis (20)

hypothesized that site II would be 33 to 44% saturated by 1 mM cytosolic  $Mg^{2+}$  at diastolic  $Ca^{2+}$  concentrations.


In a recent ITC study, the  $Mg^{2+}$ -binding affinity of site II in lobster TnC isoforms, which are similar in sequence to human variants, was explored.  $Mg^{2+}$  affinity of site II was a single order of magnitude lower than that of  $Ca^{2+}$ , such that the cations would compete for binding under physiological conditions (62).

In our experiments on N-cTnC and full-length cTnC, site II-binding affinity of  $Mg^{2+}$  was an order of magnitude lower than seen for  $Ca^{2+}$  (Fig. 2 and 7 and Tables S2 and S3). At these affinities and given the relatively high cytosolic  $[Mg^{2+}]_{free}$  (79, 82), this cation would compete for binding to site II of cTnC (107). Competition experiments were also in agreement (Figs. 1 and 2) as were experiments that utilized a double mutant removing coordinating residues in site II (Figs. 4 and 5). Competition experiments were also analyzed using the model available on Origin, and the thermodynamic parameters obtained from this alternative method were comparable and within the range of errors obtained Fig. 3 and Table S2).

In order to further validate the ITC data, we also performed TI to calculate absolute binding affinities computationally. We performed these calculations for both  $Ca^{2+}$  and  $Mg^{2+}$  binding separately for both WT N-cTnC and D67A/D73A N-cTnC. For both sets of simulations, the structure of  $Ca^{2+}$ -bound N-cTnC (Protein Data Bank [PDB]: 1AP4) was used as the starting parameter and restrained throughout the simulation. ITC measures the thermodynamically quantifiable closed-to-open transition of the N-cTnC molecule. TI does not allow for such a transition, rather, it quantifies only the binding interaction. In the future, the closed structure of N-cTnC (PDB: 1SPY) can be simulated to quantify the presumably lower affinity it has for each of  $Ca^{2+}$  and  $Mg^{2+}$ . The difference between these sets of simulations could then be used to better corroborate the ITC data.

For  $Ca^{2+}$  binding, our TI results agreed very well with the binding affinities from ITC. For  $Mg^{2+}$  binding, the calculated absolute binding affinities were consistently underestimated by about 4 kcal  $\cdot$  mol $^{-1}$  but showed the same relative trends.  $Mg^{2+}$  was calculated to bind more weakly than  $Ca^{2+}$  in the WT and D67A/D73A mutant, in agreement with the ITC results. The  $Mg^{2+}$  absolute binding affinities were likely underestimated for multiple reasons. First, the crystal structure of WT N-cTnC (1AP4) was bound by  $Ca^{2+}$ , and no structure of  $Mg^{2+}$ -bound WT N-cTnC is available. We attempted to correct for this issue by minimizing the structure with  $Mg^{2+}$  bound WT N-cTnC. Because of the lack of an exact starting structure and restraints chosen, there is still likely some error. In addition, while we did try to choose the most accurate  $Mg^{2+}$  parameters for binding affinity calculations, there are well-documented difficulties in free energy calculations for  $Mg^{2+}$ , most notably that the free energy of solvation ( $\Delta G_{solvation}$ ) is consistently underestimated (108, 109). Even when using the same  $Mg^{2+}$  force field, solvation  $\Delta G$  values are also known to have large variations for  $Mg^{2+}$  depending on the exact simulation parameters used. For example, both Panteva *et al.* (109) and Li *et al.* tried to reproduce  $Mg^{2+}$  solvation-free energy using the





**Figure 6. Representative isotherms for binding of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  to full-length cTnC.** All titrations were carried out into 100  $\mu\text{M}$  full-length human WT cTnC. From left to right, the panels show the titration of 6 mM  $\text{Ca}^{2+}$  into apo-state full-length cTnC, 1 mM  $\text{Mg}^{2+}$  preincubated cTnC, and 3 mM  $\text{Mg}^{2+}$  preincubated cTnC. In the right most panel of this figure, 40 mM  $\text{Mg}^{2+}$  was titrated into full-length cTnC. The binding of  $\text{Mg}^{2+}$  occurs at two sets of different sites as seen in the isotherm, which contains both exothermic and endothermic components. All thermograms were fit to a “two sets-of-binding sites” model using Origin 7 MicroCal2000 ITC software package. cTnC, cardiac troponin C.


same parameters as Åqvist but saw variations on the order of 20 kcal  $\cdot$  mol $^{-1}$  (110). While this may be an extreme example, it illustrates the difficulty in the calculation of free-energy changes with  $\text{Mg}^{2+}$  ions involved. Given these potential errors in TI for  $\Delta G_{\text{solvation}}$  of  $\text{Mg}^{2+}$ , the fact that we still see relatively good agreement with the ITC data for absolute binding affinity of  $\text{Mg}^{2+}$  helps further validate the *in vitro* results.

The experiments outlined previously were designed with the intent to test the hypothesis that both  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  interact with all the functional EF-hand motifs in cTnC. The interaction with sites III/IV has been established for some time (91), but site II may also bind  $\text{Mg}^{2+}$ . Interestingly, a hypothesis that is reconcilable with our own was initially put forth; that of six binding sites (4). In this scenario, there were two  $\text{Ca}^{2+}$ -specific sites, 2  $\text{Mg}^{2+}$ -specific sites, and two sites that can bind both cations. During these experiments, only the absence of  $\text{Mg}^{2+}$  allowed for the binding sites in cTnC to be separated into low-affinity sites ( $\sim 10^5 \text{ M}^{-1}$ ) and high-affinity sites ( $\sim 10^7 \text{ M}^{-1}$ ) (4). That the presence of  $\text{Mg}^{2+}$  affects the affinity of  $\text{Ca}^{2+}$  binding

to TnC is also evident in a more recent study in which a fluorescence protein-based  $\text{Ca}^{2+}$  sensor was utilized to show the reorientation of both N and C domains of TnC upon  $\text{Mg}^{2+}$  binding at sites III and IV (111). In addition, studies using the ATPase activity on the strong-binding myosin heads also demonstrates the opening of more TF active sites upon  $\text{Mg}^{2+}$  binding to C-terminal domain, supporting the notion that  $\text{Mg}^{2+}$  binding causes structural changes in TnC (112).

Binding of  $\text{Mg}^{2+}$  to site II is not expected to induce significant structural changes in N-cTnC based on previous molecular dynamics simulation data (30, 61, 64). Therefore, it is likely that the favorable  $\Delta S$  associated with the interaction is due to increased degrees of freedom for water molecules that would result when stabilizing hydrogen bonds are transferred from the positively charged metal cation and the negatively charged amino acid side chains in the binding site II to the buffered environment (61).

Given that the binding of  $\text{Ca}^{2+}$  to site II of cTnC at systolic  $\text{Ca}^{2+}$  levels (0.5–1.2  $\mu\text{M}$ ) strengthens the interaction with cTnI and the rest of the cTn complex and the orders of magnitude



**Figure 7. Binding of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  to apo-state and preincubated full-length cTnC.** Left panel, the affinity of site II for  $\text{Ca}^{2+}$  is compared in the apo-state and with  $\text{Mg}^{2+}$  preincubation in full-length human cTnC. Right panel, the affinity of sites III/IV for  $\text{Ca}^{2+}$  is compared in the apo-state and with  $\text{Mg}^{2+}$  preincubation in full-length cTnC. At site II, preincubation with 1 and 3 mM  $\text{Mg}^{2+}$  caused a statistically insignificant reduction in the affinities for  $\text{Ca}^{2+}$  binding ( $p = 0.52$  and  $p = 0.14$ ). At sites III/IV, preincubation with 1 mM or 3 mM  $\text{Mg}^{2+}$  did not significantly change affinity for  $\text{Ca}^{2+}$  binding. Statistical differences were assessed through two-way ANOVA followed by Tukey's post hoc test. cTnC, cardiac troponin C.

## Ca and Mg binding to cTnC

differ between binding affinity at varying levels of filament complexity (4, 60, 72, 113), care must be taken when translating observations at the level of cTnC to more complex systems. We suggest that other proteins in the TF, particularly cTnI, may play a central role in the mechanism discussed here. Moreover, a further limitation may be highlighted in our approach, concerning the double mutant D67A/D73A. This mutation was able to reduce the binding of both  $\text{Ca}^{2+}$  (11.9-fold) and  $\text{Mg}^{2+}$  (1.8-fold) to site II of N-cTnC; however, the impact on binding might be expected to be greater. It is possible that the effect of this double mutant is to reduce the binding of these cations, especially  $\text{Mg}^{2+}$  through allosteric interactions. In CaM, mutation of  $\text{Ca}^{2+}$  coordinating residues within the EF-hand can have structural consequences leading to altered binding kinetics (114); this is conceivable in our double mutant. Similarly, it is possible that the competition observed between  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  for binding to site II of cTnC occurs through structural perturbations, which follow binding of  $\text{Mg}^{2+}$  to an allosteric site. Exploration of these limitations in future studies may shed light on the true nature of these interactions.

Our ITC results strongly suggest that  $\text{Mg}^{2+}$  binds to site III/IV and competes with  $\text{Ca}^{2+}$  for binding to site II. The amount of  $\text{Mg}^{2+}$  that binds the regulatory site II is likely to be highly dependent on the technique, biological system, and buffer conditions. In N-cTnC, occupation of site II by  $\text{Mg}^{2+}$  was again seen to reduce the amount of  $\text{Ca}^{2+}$ , which was able to bind this protein, at concentrations that may have physiologically relevant consequences under normal conditions and even more so in the face of diseases that alter the  $\text{Ca}^{2+}$  sensitivity of contraction.

Moreover, increases in cAMP in the cell through  $\alpha$ - and  $\beta$ -adrenergic stimulation elicit extrusion of  $\text{Mg}^{2+}$  from the cell in mammalian tissues (115–117) including cardiomyocytes (118, 119). If shown in the heart, both  $\text{Na}^+$ -dependent and  $\text{Na}^+$ -independent removal of  $\text{Mg}^{2+}$  from the cytosol under stressful conditions would lower cytosolic presence of this cation. Despite this, free  $\text{Mg}^{2+}$  does not fluctuate greatly under such stimulation, suggesting that buffered  $\text{Mg}^{2+}$  is removed from the cell (120). Nevertheless, this altered  $\text{Mg}^{2+}$  pool may affect the subset of ions available to compete with  $\text{Ca}^{2+}$  for binding to troponin.

Based on our binding experiments and given the previous studies cited herein,  $\text{Mg}^{2+}$  may also compete with  $\text{Ca}^{2+}$  in binding to the regulatory site II. Free  $\text{Ca}^{2+}$  is tightly regulated at rest ( $\sim 0.1 \mu\text{M}$ ) despite relatively high total cytosolic concentrations (2.1–2.6 mM) (84).  $\text{Mg}^{2+}$  is also abundant in the cell but is less tightly controlled. Binding of both  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  to site II is endothermic and thus driven by entropy. Relative to  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  binds site II with lower affinity; however, at physiological concentrations or with elevation of free  $\text{Mg}^{2+}$ , which accompanies states of energy depletion, it may reduce  $\text{Ca}^{2+}$  binding, leading to structural perturbations that modify the contractile function of the myofilament. Conversely,  $\text{Mg}^{2+}$  can be altered by diseased states such as secondary hyperparathyroidism, which results in hypomagnesemia and could potentially impact cardiac contractility (121).

## Conclusions

Our study provides insights regarding the thermodynamics of metal cation binding to cTnC. The interaction of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  with cTnC is characterized by differences consistent with dissimilar ionic radius, number of required coordinating residues, as well as the energetic cost of exposing hydrophobic amino acids to an aqueous environment. In the cell, these differences are functionally necessitated by dissimilar free cytosolic concentrations of each cation. Cellular  $\text{Mg}^{2+}$  is not necessarily prevalent enough to directly regulate contraction and is not thought to cause a conformational change upon binding to cTnC. However, given the affinities we have observed, its occupation of the binding site may restrict  $\text{Ca}^{2+}$  binding, disable key interactions with components of the cTn complex, such as cTnI, and prevent the subsequent conformational changes necessary for rigor-state formation. This competition for binding likely favors  $\text{Ca}^{2+}$  and is well tolerated; however, elevation of free  $\text{Mg}^{2+}$ , which may accompany states of ATP depletion, for example, during ischemic stress, could have relatively significant functional consequence for cardiac force production.

## Experimental procedures

### Construct preparation and protein expression

The human *TNNC1* gene (Uniprot ID: P63316) had previously been cloned into pET21a(+) vector and had a stop codon inserted at residue 90 to create the human N-cTnC construct using the Phusion site-directed mutagenesis protocol (Thermo Scientific). This construct was transformed into the BL21(DE3) *Escherichia coli* expression strain. The double-mutant D76A/D73A construct was made using site-directed mutagenesis carried out by GenScript. Expression and purification of all constructs were carried out as described previously (63, 64). In brief, 100 ml of lysogeny broth was supplemented with 50  $\mu\text{g}/\text{ml}$  ampicillin and a glycerol stock stab and grown overnight at a shaking speed of 250 rpm and 37 °C. The next day, the overnight grown culture was used to inoculate each liter of lysogeny broth with 1:100 back dilution supplemented with 50  $\mu\text{g}/\text{ml}$  ampicillin. Cell cultures were grown for  $\sim 3$  h until an absorbance reached to 0.8 to 1.0 at 600 nm followed by induction with 1 mM IPTG. After 3 h, the cells were harvested by centrifugation at 6000g for 6 min, and the collected cell pellets were stored at  $-80$  °C.

### Protein purification

The cell pellet was thawed and suspended in buffer A (50 mM Tris-HCl at pH 8.0 and 5 mM EDTA) and sonicated on ice at 50% amplitude with 30 s on and 30 s off for 5 min, or until there was no visible viscosity of the lysate solution. After sonication, the lysate was centrifuged at 30,000g for 15 min at 4 °C, and the supernatant was obtained. This centrifugation process was performed twice to ensure the removal of all cell debris before loading onto a fast-flow (FF) Q-Sepharose column (GE Healthcare). The FF Q-Sepharose column was connected with an AKTA FPLC system (GE Healthcare) and

pre-equilibrated with buffer A with the addition of 1 mM DTT. After applying the clear supernatant onto the column, the solution was run at 5 ml/min with a gradual gradient mixing with buffer A and buffer B (buffer A with 0.5 M NaCl), starting from 0% buffer B up to 100% buffer B by the end of the run. Following analysis by 12% SDS-PAGE, the fractions containing purified N-cTnC were pooled and concentrated using Amicon ultracentrifugal filter device (Millipore) with a 3-kDa molecular weight cutoff.

The full-length TnC was purified using the same protocol described previously, with the addition of 30% ammonium sulphate precipitation following the sonication step. After the addition of 30% ammonium sulphate, the solution was stirred on ice for 30 min and subsequently centrifuged at 28,900g for 30 min at 4 °C. The supernatant was obtained and dialyzed overnight against 4 l of buffer C (50 mM Tris-HCl at pH 8.0 and 100 mM NaCl).

After purification using the FF Q-Sepharose column, the fractions containing partially pure cTnC were concentrated to 3 ml using an Amicon ultracentrifugal filter device (Millipore) with a 10-kDa molecular weight cutoff. The concentrated protein sample was further purified by a HiPrep 26/60 Sephacryl S-100 column size-exclusion chromatography (GE Healthcare), which was equilibrated with buffer C. After confirming the purity of the protein on a 12% SDS-PAGE gel, all fractions containing the purified cTnC were combined, aliquoted, and stored at -80 °C prior to pre-ITC dialysis.

### Protein dialysis

To generate the apo-state protein, TnC was first dialyzed against 2 l of 50 mM Hepes at pH 7.2, 150 mM KCl, 2 mM EDTA, and 15 mM  $\beta$ -mercaptoethanol, followed by another dialysis against the same buffer with no EDTA added. Each of these dialysis steps was completed at 4 °C for a minimum of 4 h. A third dialysis was performed for a minimum of 16 h overnight against 2 l of 50 mM Hepes at pH 7.2, 150 mM KCl, and 2 mM  $\beta$ -mercaptoethanol. An extinction coefficient of 1490 and 4595  $M^{-1} cm^{-1}$  and a molecular weight of 10.1 and 18.4 kDa was used to determine protein concentration for the N-cTnC and full-length cTnC constructs, respectively, by 280 nm UV-visible spectroscopy using a NanoDrop 2000 spectrophotometer (Thermo Scientific). The final dialysis buffer was used to dilute the protein samples to a final concentration of 200  $\mu M$  for the N-terminal construct and 100  $\mu M$  for full-length cTnC as described previously (64).

Standard 1.0 M  $CaCl_2$  and  $MgCl_2$  stock solutions (Sigma) were serially diluted in the final dialysis buffer to produce 6 mM  $Ca^{2+}$  and 40 mM  $Mg^{2+}$ , respectively. The same standards were used to produce 4 mM  $Ca^{2+}$  and 20 mM  $Mg^{2+}$  titrants for the N-cTnC experiments. Given the key role of protein concentration in determination of affinity, we aimed to ensure consistency and did so through dilution of the protein from stock solutions with care taken to minimize human and pipetting error to fall at the minimum possible using recently calibrated instrumentation.

### Isothermal titration calorimetry

The ITC experiments were carried out in a MicroCal ITC200 instrument (Malvern). Repeat titrations were used to ensure reproducibility. The sample cell was set at 25 °C, 200  $\mu l$  of the protein was loaded, and the experiment was carried out at the same temperature. For the N-cTnC, 19 injections of the titrant were used with the first being a dummy injection of 0.8  $\mu l$  and the subsequent 18 injections, 2  $\mu l$  each. For these experiments, 4 mM  $Ca^{2+}$  was titrated into 200  $\mu l$  of 200  $\mu M$  apo-state N-cTnC as the baseline condition. For the full-length cTnC, 6 mM  $Ca^{2+}$  was titrated into 200  $\mu l$  of 100  $\mu M$  apo-state full-length cTnC with a dummy injection of 0.5  $\mu l$  and 38 injections of 1  $\mu l$ . The time interval between injections was 120 s, and stirring speed was set at 1000 rpm throughout each experiment.

### Analysis of results

Titration data were analyzed using MicroCal2000 for ITC through Origin 8.0 (OriginLab). Raw heats were integrated and fit by a least-squares algorithm using a “single-binding site” model for the N-cTnC titrations and a “two sets-of-binding sites” model for the full-length cTnC titrations to calculate the thermodynamic parameters. As a point of comparison, the “competition” model on Origin was also used to study the binding of  $Ca^{2+}$  and  $Mg^{2+}$  to the apo-state N-cTnC as well as in the presence of 1 mM of the counter ion (Fig. 3) (122). For the N-cTnC constructs (apart from the  $Ca^{2+}$  into apo-state N-cTnC condition), the N (number of binding sites) associated with each interaction was necessarily constrained to equal 1.00 to facilitate curve fitting without altering protein concentration. The baseline condition was repeated daily, and the consistency of the thermodynamic parameters in these sets of titrations indicates protein quality and function throughout the set of experiments. If multiple ligands were simultaneously present in the reaction mixture, an “apparent affinity” was determined for the injected titrant.

JMP 14.0 software package was used for statistical analysis. ANOVA was used to identify differences in each studied thermodynamic parameter from the N-cTnC and full-length cTnC titrations: in the apo-state and competition experiments, including both the WT and double-mutant proteins (N-cTnC only). Tukey's post hoc test was subsequently used to explore where the differences lie with  $p < 0.05$  considered the threshold for statistical significance.

### Thermodynamic integration

Starting from the representative model of PDB: 1AP4 (30), which contains N-cTnC with a single  $Ca^{2+}$  ion bound, the system was solvated with a 12 Å-padded transferable intermolecular potential 3P water box and neutralized with  $Na^+$  in Amber16 (123). The system was also prepared similarly for only the  $Ca^{2+}$  ion in a 12 Å-padded transferable intermolecular potential 3P water box. The alchemical thermodynamic cycle used for ligand binding was described in detail previously

## Ca and Mg binding to cTnC

(124). In short, TI was performed using the following three steps for Ca<sup>2+</sup> in protein: turn on restraints, turn off charge, and turn off van der Waals forces. The specific distance restraints used in all systems can be found in [Table S1](#). In addition, TI was performed for the following two steps for Ca<sup>2+</sup> in water: turn off charge and turn off van der Waals forces. Each step of the thermodynamic cycle was performed with the coupling parameter ( $\lambda$ ) ranging from 0.0 to 1.0 in increments of 0.1. For each simulation, the system was minimized (2000 cycles) and heated (0.5 ns) before the 5 ns production run at 300 K using the ff14SB force field (125). These calculations were also performed on the D67A/D73A mutated system. The mutations were imposed on the 1AP4 representative model using PyMOL (126).

For the calculation of Mg<sup>2+</sup>-binding affinity, Ca<sup>2+</sup> was replaced with Mg<sup>2+</sup> in the 1AP4 representative model since no Mg<sup>2+</sup>-bound N-cTnC structure was available in the PDB. In order to generate more accurate restraints and starting coordinates for the TI calculations, a minimization was performed on the structure in Amber (2000 cycles). Following the minimization, TI simulations were run similarly as for Ca<sup>2+</sup>. However, because of previously documented errors in the default Mg<sup>2+</sup> parameters, the  $\Delta G_{\text{solvation}}$ -optimized Mg<sup>2+</sup> parameters from Li *et al.* were used (109, 127). These calculations were also performed on the D67A/D73A mutated system.

To calculate absolute binding affinities for the ions, the change in free energy ( $\Delta G$ ) was calculated for each step in the thermodynamic cycle by integrating the potential energy with respect to the coupling parameter,  $\lambda$  (128). Two corrections were made to these calculated  $\Delta G$  values. The first correction was necessary because of the introduction of the distance restraints (as described in the study by Boresch *et al.* (129)), which quantified the free energy cost of restraining the ion to the binding site. The second correction was performed to correct the charged system (as described in the study by Rocklin *et al.* (130)) to revise the free energy for the fact that the system is charged during the disappearance of the charged ions. The overall  $\Delta G$  of binding was the change in free energy between the ions in complex with the protein (ion in protein steps 1, 2, and 3) and the ions in water (ion in water steps 1 and 2). For each system, five independent runs were performed, and results were averaged.

### Data availability

All data presented in this article are contained within the article.

*Supporting information*—This article contains [supporting information](#).

*Author contributions*—K. R. and A. Y. L. preliminary experiments; K. R., J. S., A. Y. L., J. P. D., A. M. S., F. V. P., S. L., and G. F. T. experimental design; K. R. and J. S. data collection; K. R., J. S., A. M. S., and S. L. data analysis; K. R., J. S., A. M. S., F. V. P., S. L., and G. F. T. manuscript preparation; K. R., A. Y. L., J. P. D., A. M. S., F. V. P., R. J. S., S. L., and G. F. T. manuscript review.

*Funding and additional information*—This work was primarily supported by a grant from the Canadian Institutes of Health Research (PJT-148964) to G. F. T. K. R. was supported by the National Sciences and Engineering Research Council Canada Graduate Scholarship—Doctoral fellowship.

*Conflict of interest*—The authors declare that they have no conflicts of interest with the contents of this article.

*Abbreviations*—The abbreviations used are: CaM, calmodulin; cTn, cardiac troponin; cTnC, cardiac troponin C; cTnI, cardiac troponin I; cTnT, cardiac troponin T; FF, fast-flow; ITC, isothermal titration calorimetry; PDB, Protein Data Bank; sTnC, skeletal muscle troponin TnC; TF, thin filament; TI, thermodynamic integration.

### References

1. Parmacek, M. S., and Solaro, R. J. (2004) Biology of the troponin complex in cardiac myocytes. *Prog. Cardiovasc. Dis.* **47**, 159–176
2. Kawasaki, Y., and van Eerd, J.-P. (1972) The effect of Mg<sup>2+</sup> on the conformation of the Ca<sup>2+</sup>-binding component of troponin. *Biochem. Biophys. Res. Commun.* **49**, 898–905
3. Murray, A. C., and Kay, C. M. (1972) Hydrodynamic and optical properties of troponin A. Demonstration of a conformational change upon binding calcium ion. *Biochemistry* **11**, 2622–2627
4. Potter, J. D., and Gergely, J. (1975) The calcium and magnesium binding sites on troponin and their role in the regulation of myofibrillar adenosine triphosphatase. *J. Biol. Chem.* **250**, 4628–4633
5. Filatov, V. L., Katrukha, A. G., Bulargina, T. V., and Gusev, N. B. (1999) Troponin: Structure, properties, and mechanism of functioning. *Biochemistry (Mosc)* **64**, 969–985
6. Strynadka, N. C., and James, M. N. (1989) Crystal structures of the helix-loop-helix calcium-binding proteins. *Annu. Rev. Biochem.* **58**, 951–998
7. Yap, K. L., Ames, J. B., Swindells, M. B., and Ikura, M. (1999) Diversity of conformational states and changes within the EF-hand protein superfamily. *Proteins* **37**, 499–507
8. Lewit-Bentley, A., and Rety, S. (2000) EF-hand calcium-binding proteins. *Curr. Opin. Struct. Biol.* **10**, 637–643
9. Seamon, K. B., Hartshorne, D. J., and Bothner-By, A. A. (1977) Ca<sup>2+</sup> and Mg<sup>2+</sup> dependent conformations of troponin C as determined by <sup>1</sup>H and <sup>19</sup>F nuclear magnetic resonance. *Biochemistry* **16**, 4039–4046
10. Ebashi, S., Nonomura, Y., Kohama, K., Kitazawa, T., and Mikawa, T. (1980) Regulation of muscle contraction by Ca ion. *Mol. Biol. Biochem. Biophys.* **32**, 183–194
11. van Eerd, J. P., and Takahashi, K. (1976) Determination of the complete amino acid sequence of bovine cardiac troponin C. *Biochemistry* **15**, 1171–1180
12. Farah, C. S., and Reinach, F. C. (1995) The troponin complex and regulation of muscle contraction. *FASEB J.* **9**, 755–767
13. Schober, T., Huke, S., Venkataraman, R., Gryshchenko, O., Kryshtal, D., Hwang, H. S., Baudenbacher, F. J., and Knollmann, B. C. (2012) Myofilament Ca sensitization increases cytosolic Ca binding affinity, alters intracellular Ca homeostasis, and causes pause-dependent Ca-triggered arrhythmia. *Circ. Res.* **111**, 170–179
14. Johnson, J. D., Charlton, S. C., and Potter, J. D. (1979) A fluorescence stopped flow analysis of Ca<sup>2+</sup> exchange with troponin C. *J. Biol. Chem.* **254**, 3497–3502
15. Johnson, J. D., Nakkula, R. J., Vasulka, C., and Smillie, L. B. (1994) Modulation of Ca<sup>2+</sup> exchange with the Ca<sup>2+</sup>-specific regulatory sites of troponin C. *J. Biol. Chem.* **269**, 8919–8923
16. Bers, D. M. (2000) Calcium fluxes involved in control of cardiac myocyte contraction. *Circ. Res.* **87**, 275–281
17. Leavis, P., and Kraft, E. L. (1978) Calcium binding to cardiac troponin C. *Arch. Biochem. Biophys.* **186**, 411–415
18. Robertson, S., Johnson, J. D., and Potter, J. (1981) The time-course of Ca<sup>2+</sup> exchange with calmodulin, troponin, parvalbumin, and myosin in response to transient increases in Ca<sup>2+</sup>. *Biophys. J.* **34**, 559

19. Sturtevant, J. M. (1977) Heat capacity and entropy changes in processes involving proteins. *Proc. Natl. Acad. Sci. U. S. A.* **74**, 2236–2240
20. Tikunova, S. B., and Davis, J. P. (2004) Designing calcium-sensitizing mutations in the regulatory domain of cardiac troponin C. *J. Biol. Chem.* **279**, 35341–35352
21. Sundaralingam, M., Bergstrom, R., Strasburg, G., Rao, S. T., Roychowdhury, P., Greaser, M., and Wang, B. C. (1985) Molecular structure of troponin C from chicken skeletal muscle at 3-angstrom resolution. *Science* **227**, 945–948
22. Sia, S. K., Li, M. X., Spyrapoulos, L., Gagné, S. M., Liu, W., Putkey, J. A., and Sykes, B. D. (1997) Structure of cardiac muscle troponin C unexpectedly reveals a closed regulatory domain. *J. Biol. Chem.* **272**, 18216–18221
23. Cheung, J. Y., Tillotson, D. L., Yelamarty, R., and Scaduto, R. (1989) Cytosolic free calcium concentration in individual cardiac myocytes in primary culture. *Am. J. Physiol. Cell Physiol.* **256**, C1120–C1130
24. Kirschenlohr, H. L., Grace, A. A., Vandenberg, J. I., Metcalfe, J. C., and Smith, G. A. (2000) Estimation of systolic and diastolic free intracellular  $\text{Ca}^{2+}$  by titration of  $\text{Ca}^{2+}$  buffering in the ferret heart. *Biochem. J.* **346 Pt 2**, 385–391
25. Gifford, J. L., Walsh, M. P., and Vogel, H. J. (2007) Structures and metal-ion-binding properties of the  $\text{Ca}^{2+}$ -binding helix–loop–helix EF-hand motifs. *Biochem. J.* **405**, 199–221
26. Bowman, J. D., and Lindert, S. (2018) Molecular dynamics and umbrella sampling simulations elucidate differences in troponin C isoform and mutant hydrophobic patch exposure. *J. Phys. Chem. B* **122**, 7874–7883
27. Herzberg, O., and James, M. N. (1985) Structure of the Calcium Regulatory Muscle Protein Troponin-C at 2.8 Å Resolution. *Nature* **313**, 653–659
28. Slupsky, C. M., and Sykes, B. D. (1995) NMR solution structure of calcium-saturated skeletal muscle troponin C. *Biochemistry* **34**, 15953–15964
29. Houdusse, A., Love, M. L., Dominguez, R., Grabarek, Z., and Cohen, C. (1997) Structures of four  $\text{Ca}^{2+}$ -bound troponin C at 2.0 Å resolution: Further insights into the  $\text{Ca}^{2+}$ -switch in the calmodulin superfamily. *Structure* **5**, 1695–1711
30. Spyrapoulos, L., Li, M. X., Sia, S. K., Gagné, S. M., Chandra, M., Solaro, R. J., and Sykes, B. D. (1997) Calcium-induced structural transition in the regulatory domain of human cardiac troponin C. *Biochemistry* **36**, 12138–12146
31. Li, M. X., Spyrapoulos, L., and Sykes, B. D. (1999) Binding of cardiac troponin-I147-163 induces a structural opening in human cardiac troponin-C. *Biochemistry* **38**, 8289–8298
32. Dai, L. J., Friedman, P. A., and Quamme, G. A. (1997) Cellular mechanisms of chlorothiazide and cellular potassium depletion on  $\text{Mg}^{2+}$  uptake in mouse distal convoluted tubule cells. *Kidney Int.* **51**, 1008–1017
33. Romani, A., and Scarpa, A. (1992) Regulation of cell magnesium. *Arch. Biochem. Biophys.* **298**, 1–12
34. Tessman, P. A., and Romani, A. (1998) Acute effect of EtOH on  $\text{Mg}^{2+}$  homeostasis in liver cells: Evidence for the activation of an  $\text{Na}^+/\text{Mg}^{2+}$  exchanger. *Am. J. Physiol.* **275**, G1106–G1116
35. Hongo, K., Konishi, M., and Kurihara, S. (1994) Cytoplasmic free  $\text{Mg}^{2+}$  in rat ventricular myocytes studied with the fluorescent indicator fura-2. *Jpn. J. Physiol.* **44**, 357–378
36. Laires, M. J., Monteiro, C. P., and Bicho, M. (2004) Role of cellular magnesium in health and human disease. *Front. Biosci.* **9**, 262–276
37. Godt, R. E. (1974) Calcium-activated tension of skinned muscle fibers of the frog. Dependence on magnesium adenosine triphosphate concentration. *J. Gen. Physiol.* **63**, 722–739
38. Godt, R. E., and Morgan, J. L. (1984) Contractile responses to  $\text{MgATP}$  and pH in a thick filament regulated muscle: Studies with skinned scallop fibers. *Adv. Exp. Med. Biol.* **170**, 569–572
39. Best, P. M., Donaldson, S. K., and Kerrick, W. G. (1977) Tension in mechanically disrupted mammalian cardiac cells: Effects of magnesium adenosine triphosphate. *J. Physiol.* **265**, 1–17
40. Li, M. X., Gagne, S. M., Tsuda, S., Kay, C. M., Smillie, L. B., and Sykes, B. D. (1995) Calcium binding to the regulatory N-domain of skeletal muscle troponin C occurs in a stepwise manner. *Biochemistry* **34**, 8330–8340
41. Potter, J. D., Robertson, S. P., and Johnson, J. D. (1981) Magnesium and the regulation of muscle contraction. *Fed. Proc.* **40**, 2653–2656
42. Ogawa, Y. (1985) Calcium binding to troponin C and troponin: Effects of  $\text{Mg}^{2+}$ , ionic strength and pH. *J. Biochem.* **97**, 1011–1023
43. Zot, A. S., and Potter, J. D. (1987) Structural aspects of troponin-tropomyosin regulation of skeletal muscle contraction. *Annu. Rev. Biochem. Biophys. Chem.* **16**, 535–559
44. Morimoto, S. (1991) Effect of myosin cross-bridge interaction with actin on the  $\text{Ca}^{2+}$ -binding properties of troponin C in fast skeletal myofibrils. *J. Biochem.* **109**, 120–126
45. Francois, J. M., Gerday, C., Prendergast, F. G., and Potter, J. D. (1993) Determination of the  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  affinity constants of troponin C from eel skeletal muscle and positioning of the single tryptophan in the primary structure. *J. Muscle Res. Cell Motil.* **14**, 585–593
46. She, M., Dong, W. J., Umeda, P. K., and Cheung, H. C. (1998) Tryptophan mutants of troponin C from skeletal muscle: An optical probe of the regulatory domain. *Eur. J. Biochem.* **252**, 600–607
47. Allen, T. S., Yates, L. D., and Gordon, A. M. (1992)  $\text{Ca}^{2+}$ -dependence of structural changes in troponin-C in demembrated fibers of rabbit psoas muscle. *Biophys. J.* **61**, 399–409
48. Yamada, K. (1978) The enthalpy titration of troponin C with calcium. *Biochim. Biophys. Acta* **535**, 342–347
49. Kometani, K., and Yamada, K. (1983) Enthalpy, entropy and heat capacity changes induced by binding of calcium ions to cardiac troponin C. *Biochem. Biophys. Res. Commun.* **114**, 162–167
50. Yamada, K., and Kometani, K. (1982) The changes in heat capacity and entropy of troponin C induced by calcium binding. *J. Biochem.* **92**, 1505–1517
51. Gillis, T. E., Blumenschein, T. M., Sykes, B. D., and Tibbits, G. F. (2003) Effect of temperature and the F27W mutation on the  $\text{Ca}^{2+}$  activated structural transition of trout cardiac troponin C. *Biochemistry* **42**, 6418–6426
52. Gillis, T. E., Liang, B., Chung, F., and Tibbits, G. F. (2005) Increasing mammalian cardiomyocyte contractility with residues identified in trout troponin C. *Physiol. Genomics* **22**, 1–7
53. Gillis, T. E., Moyes, C. D., and Tibbits, G. F. (2003) Sequence mutations in teleost cardiac troponin C that are permissive of high  $\text{Ca}^{2+}$  affinity of site II. *Am. J. Physiol. Cell Physiol.* **284**, C1176–C1184
54. Davis, J. P., Rall, J. A., Reiser, P. J., Smillie, L. B., and Tikunova, S. B. (2002) Engineering competitive magnesium binding into the first EF-hand of skeletal troponin C. *J. Biol. Chem.* **277**, 49716–49726
55. Yamada, K. (2003) Calcium binding to troponin C as a primary step of the regulation of contraction. A microcalorimetric approach. *Adv. Exp. Med. Biol.* **538**, 203–212. discussion 213
56. Wilcox, D. E. (2008) Isothermal titration calorimetry of metal ions binding to proteins: An overview of recent studies. *Inorg. Chim. Acta* **361**, 857–867
57. Grosseohme, N. E., Spuches, A. M., and Wilcox, D. E. (2010) Application of isothermal titration calorimetry in bioinorganic chemistry. *J. Biol. Inorg. Chem.* **15**, 1183–1191
58. Sacco, C., Skowronsky, R. A., Gade, S., Kenney, J. M., and Spuches, A. M. (2012) Calorimetric investigation of copper(II) binding to Abeta peptides: Thermodynamics of coordination plasticity. *J. Biol. Inorg. Chem.* **17**, 531–541
59. Davis, J. P., Norman, C., Kobayashi, T., Solaro, R. J., Swartz, D. R., and Tikunova, S. B. (2007) Effects of thin and thick filament proteins on calcium binding and exchange with cardiac troponin C. *Biophys. J.* **92**, 3195–3206
60. Li, A. Y., Stevens, C. M., Liang, B., Rayani, K., Little, S., Davis, J., and Tibbits, G. F. (2013) Familial Hypertrophic Cardiomyopathy Related Cardiac Troponin C L29Q Mutation Alters Length-dependent Activation and Functional Effects of Phosphomimetic Troponin I\*. *PLoS One* **8**, e79363
61. Skowronsky, R. A., Schroeter, M., Baxley, T., Li, Y., Chalovich, J. M., and Spuches, A. M. (2013) Thermodynamics and molecular dynamics simulations of calcium binding to the regulatory site of human cardiac

- troponin C: Evidence for communication with the structural calcium binding sites. *J. Biol. Inorg. Chem.* **18**, 49–58
62. Tanaka, H., Takahashi, H., and Ojima, T. (2013) Ca<sup>2+</sup>-binding properties and regulatory roles of lobster troponin C sites II and IV. *FEBS Lett.* **587**, 2612–2616
  63. Stevens, C. M., Rayani, K., Genge, C. E., Singh, G., Liang, B., Roller, J. M., Li, C., Li, A. Y., Tieleman, D. P., and van Petegem, F. (2016) Characterization of zebrafish cardiac and slow skeletal troponin C paralogs by MD simulation and ITC. *Biophys. J.* **111**, 38–49
  64. Stevens, C. M., Rayani, K., Singh, G., Lotfalimali, B., Tieleman, D. P., and Tibbits, G. F. (2017) Changes in the dynamics of the cardiac troponin C molecule explain the effects of Ca<sup>2+</sup>-sensitizing mutations. *J. Biol. Chem.* **292**, 11915–11926
  65. Johnson, R. A., Fulcher, L. M., Vang, K., Palmer, C. D., Grosseohme, N. E., and Spuches, A. M. (2019) In depth, thermodynamic analysis of Ca<sup>2+</sup> binding to human cardiac troponin C: Extracting buffer-independent binding parameters. *Biochim. Biophys. Acta* **1867**, 359–366
  66. Kolte, D., Vijayaraghavan, K., Khera, S., Sica, D. A., and Frishman, W. H. (2014) Role of magnesium in cardiovascular diseases. *Cardiol. Rev.* **22**, 182–192
  67. Smith, D. W. (1977) Ionic hydration enthalpies. *J. Chem. Edu.* **54**, 540
  68. Wang, S.-Q., Huang, Y.-H., Liu, K.-S., and Zhou, Z.-Q. (1997) Dependence of myocardial hypothermia tolerance on sources of activator calcium. *Cryobiology* **35**, 193–200
  69. Hazard, A. L., Kohout, S. C., Stricker, N. L., Putkey, J. A., and Falke, J. J. (1998) The kinetic cycle of cardiac troponin C: Calcium binding and dissociation at site II trigger slow conformational rearrangements. *Protein Sci.* **7**, 2451–2459
  70. Liang, B., Chung, F., Qu, Y., Pavlov, D., Gillis, T. E., Tikunova, S. B., Davis, J. P., and Tibbits, G. F. (2008) Familial hypertrophic cardiomyopathy-related cardiac troponin C mutation L29Q affects Ca<sup>2+</sup> binding and myofilament contractility. *Physiol. Genomics* **33**, 257–266
  71. Gillis, T. E., Marshall, C. R., Xue, X.-H., Borgford, T. J., and Tibbits, G. F. (2000) Ca<sup>2+</sup> binding to cardiac troponin C: Effects of temperature and pH on mammalian and salmonid isoforms. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **279**, R1707–R1715
  72. Pinto, J. R., Parvatiyar, M. S., Jones, M. A., Liang, J., Ackerman, M. J., and Potter, J. D. (2009) A functional and structural study of troponin C mutations related to hypertrophic cardiomyopathy. *J. Biol. Chem.* **284**, 19090–19100
  73. Romani, A. M. P. (2011) Intracellular magnesium homeostasis. In: Vink, R., Nechifor, M., eds. *Magnesium in the Central Nervous System*, University of Adelaide Press (c) 2011 The Authors., Adelaide, AU: 13–58
  74. Yumoto, F., Nara, M., Kagi, H., Iwasaki, W., Ojima, T., Nishita, K., Nagata, K., and Tanokura, M. (2001) Coordination structures of Ca<sup>2+</sup> and Mg<sup>2+</sup> in Akazara scallop troponin C in solution. FTIR spectroscopy of side-chain COO- groups. *Eur. J. Biochem.* **268**, 6284–6290
  75. Carafoli, E., and Krebs, J. (2016) Why calcium? How calcium became the best communicator. *J. Biol. Chem.* **291**, 20849–20857
  76. Lockless, S. W., Zhou, M., and MacKinnon, R. (2007) Structural and thermodynamic properties of selective ion binding in a K<sup>+</sup> channel. *PLoS Biol.* **5**, e121
  77. Maguire, M. E. (2006) Magnesium transporters: Properties, regulation and structure. *Front. Biosci.* **11**, 3149–3163
  78. Chao, S. H., Suzuki, Y., Zysk, J. R., and Cheung, W. Y. (1984) Activation of calmodulin by various metal cations as a function of ionic radius. *Mol. Pharmacol.* **26**, 75–82
  79. Malmendal, A., Linse, S., Evenas, J., Forsen, S., and Drakenberg, T. (1999) Battle for the EF-hands: Magnesium-calcium interference in calmodulin. *Biochemistry* **38**, 11844–11850
  80. Follenius, A., and Gerard, D. (1984) Fluorescence investigations of calmodulin hydrophobic sites. *Biochem. Biophys. Res. Commun.* **119**, 1154–1160
  81. Gilli, R., Lafitte, D., Lopez, C., Kilhoffer, M., Makarov, A., Briand, C., and Haiech, J. (1998) Thermodynamic analysis of calcium and magnesium binding to calmodulin. *Biochemistry* **37**, 5450–5456
  82. Linse, S., and Forsen, S. (1995) Determinants that govern high-affinity calcium binding. *Adv. Second Messenger Phosphoprotein Res.* **30**, 89–151
  83. Cates, M. S., Berry, M. B., Ho, E. L., Li, Q., Potter, J. D., and Phillips, G. N., Jr. (1999) Metal-ion affinity and specificity in EF-hand proteins: Coordination geometry and domain plasticity in parvalbumin. *Structure* **7**, 1269–1278
  84. Brini, M., Cali, T., Ottolini, D., and Carafoli, E. (2012) Calcium pumps: Why so many? *Compr. Physiol.* **2**, 1045–1060
  85. Harding, M. M. (2002) Metal-ligand geometry relevant to proteins and in proteins: Sodium and potassium. *Acta Crystallogr. D Biol. Crystallogr.* **58**, 872–874
  86. Dokmanic, I., Sikic, M., and Tomic, S. (2008) Metals in proteins: Correlation between the metal-ion type, coordination number and the amino-acid residues involved in the coordination. *Acta Crystallogr. D Biol. Crystallogr.* **64**, 257–263
  87. Reid, R. E., and Procyshyn, R. M. (1995) Engineering magnesium selectivity in the helix-loop-helix calcium-binding motif. *Arch. Biochem. Biophys.* **323**, 115–119
  88. Tikunova, S. B., Black, D. J., Johnson, J. D., and Davis, J. P. (2001) Modifying Mg<sup>2+</sup> binding and exchange with the N-terminal of calmodulin. *Biochemistry* **40**, 3348–3353
  89. Holroyde, M., Robertson, S., Johnson, J., Solaro, R., and Potter, J. (1980) The calcium and magnesium binding sites on cardiac troponin and their role in the regulation of myofibrillar adenosine triphosphatase. *J. Biol. Chem.* **255**, 11688–11693
  90. Li, M. X., Chandra, M., Pearlstone, J. R., Racher, K. I., Trigo-Gonzalez, G., Borgford, T., Kay, C. M., and Smillie, L. B. (1994) Properties of isolated recombinant N and C domains of chicken troponin C. *Biochemistry* **33**, 917–925
  91. Johnson, J. D., Collins, J. H., Robertson, S. P., and Potter, J. D. (1980) A fluorescent probe study of Ca<sup>2+</sup> binding to the Ca<sup>2+</sup>-specific sites of cardiac troponin and troponin C. *J. Biol. Chem.* **255**, 9635–9640
  92. Allen, K., Xu, Y. Y., and Kerrick, W. G. (2000) Ca<sup>2+</sup> measurements in skinned cardiac fibers: Effects of Mg<sup>2+</sup> on Ca<sup>2+</sup> activation of force and fiber ATPase. *J. Appl. Physiol.* (1985) **88**, 180–185
  93. Ebashi, S., and Ogawa, Y. (1988) Ca<sup>2+</sup> in contractile processes. *Biophys. Chem.* **29**, 137–143
  94. Kohama, K. (1980) Role of the high affinity Ca binding sites of cardiac and fast skeletal troponins. *J. Biochem.* **88**, 591–599
  95. Stephenson, D. G., and Williams, D. A. (1982) Effects of sarcomere length on the force-pCa relation in fast- and slow-twitch skinned muscle fibres from the rat. *J. Physiol.* **333**, 637–653
  96. Godt, R. E., and Lindley, B. D. (1982) Influence of temperature upon contractile activation and isometric force production in mechanically skinned muscle fibers of the frog. *J. Gen. Physiol.* **80**, 279–297
  97. Wnuk, W., Schoeclin, M., and Stein, E. A. (1984) Regulation of actomyosin ATPase by a single calcium-binding site on troponin C from crayfish. *J. Biol. Chem.* **259**, 9017–9023
  98. Fabiato, A., and Fabiato, F. (1975) Effects of magnesium on contractile activation of skinned cardiac cells. *J. Physiol.* **249**, 497–517
  99. Donaldson, S. K., and Kerrick, W. G. (1975) Characterization of the effects of Mg<sup>2+</sup> on Ca<sup>2+</sup>- and Sr<sup>2+</sup>-activated tension generation of skinned skeletal muscle fibers. *J. Gen. Physiol.* **66**, 427–444
  100. Kerrick, W. G. L., and Donaldson, S. K. B. (1975) The comparative effects of [Ca<sup>2+</sup>] and [Mg<sup>2+</sup>] on tension generation in the fibers of skinned frog skeletal muscle and mechanically disrupted rat ventricular cardiac muscle. *Pflügers Arch.* **358**, 195–201
  101. Solaro, R. J., and Shiner, J. S. (1976) Modulation of Ca<sup>2+</sup> control of dog and rabbit cardiac myofibrils by Mg<sup>2+</sup>. Comparison with rabbit skeletal myofibrils. *Circ. Res.* **39**, 8–14
  102. Ashley, C. C., and Moisescu, D. G. (1977) Effect of changing the composition of the bathing solutions upon the isometric tension-pCa relationship in bundles of crustacean myofibrils. *J. Physiol.* **270**, 627–652
  103. Donaldson, S. K., Best, P. M., and Kerrick, G. L. (1978) Characterization of the effects of Mg<sup>2+</sup> on Ca<sup>2+</sup>- and Sr<sup>2+</sup>-activated tension generation of skinned rat cardiac fibers. *J. Gen. Physiol.* **71**, 645–655
  104. Ebashi, S., and Endo, M. (1968) Calcium ion and muscle contraction. *Prog. Biophys. Mol. Biol.* **18**, 123–183
  105. Pan, B. S., and Solaro, R. J. (1987) Calcium-binding properties of troponin C in detergent-skinned heart muscle fibers. *J. Biol. Chem.* **262**, 7839–7849

106. Siddiqui, J. K., Tikunova, S. B., Walton, S. D., Liu, B., Meyer, M., de Tombe, P. P., Neilson, N., Kekenes-Huskey, P. M., Salhi, H. E., Janssen, P. M., Biesiadecki, B. J., and Davis, J. P. (2016) Myofilament calcium sensitivity: Consequences of the effective concentration of troponin I. *Front. Physiol.* **7**, 632
107. Nara, M., Morii, H., and Tanokura, M. (2013) Infrared study of synthetic peptide analogues of the calcium-binding site III of troponin C: The role of helix F of an EF-hand motif. *Biopolymers* **99**, 342–347
108. Steinbrecher, T., Joung, I., and Case, D. A. (2011) Soft-core potentials in thermodynamic integration: Comparing one- and two-step transformations. *J. Comput. Chem.* **32**, 3253–3263
109. Panteva, M. T., Giambasu, G. M., and York, D. M. (2015) Comparison of structural, thermodynamic, kinetic and mass transport properties of Mg<sup>2+</sup> ion models commonly used in biomolecular simulations. *J. Comput. Chem.* **36**, 970–982
110. Åqvist, J. (1990) Ion-water interaction potentials derived from free energy perturbation simulations. *J. Phys. Chem.* **94**, 8021–8024
111. Badr, M. A., Pinto, J. R., Davidson, M. W., and Chase, P. B. (2016) Fluorescent protein-based Ca<sup>2+</sup> sensor reveals global, divalent cation-dependent conformational changes in cardiac troponin C. *PLoS One* **11**, e0164222
112. Fuchs, F., and Grabarek, Z. (2011) The Ca<sup>2+</sup>/Mg<sup>2+</sup> sites of troponin C modulate crossbridge-mediated thin filament activation in cardiac myofibrils. *Biochem. Biophys. Res. Commun.* **408**, 697–700
113. Ramos, C. H. (1999) Mapping subdomains in the C-terminal region of troponin I involved in its binding to troponin C and to thin filament. *J. Biol. Chem.* **274**, 18189–18195
114. Wang, K., Brohus, M., Holt, C., Overgaard, M. T., Wimmer, R., and Van Petegem, F. (2020) Arrhythmia mutations in calmodulin can disrupt cooperativity of Ca<sup>2+</sup> binding and cause misfolding. *J. Physiol.* **598**, 1169–1186
115. Cefaratti, C., and Romani, A. M. (2007) Functional characterization of two distinct Mg<sup>2+</sup> extrusion mechanisms in cardiac sarcolemmal vesicles. *Mol. Cell. Biochem.* **303**, 63–72
116. Fagan, T. E., and Romani, A. (2001)  $\alpha$ 1-Adrenoceptor-induced Mg<sup>2+</sup> extrusion from rat hepatocytes occurs via Na<sub>+</sub>-dependent transport mechanism. *Am. J. Physiol. Gastrointest. Liver Physiol.* **280**, G1145–G1156
117. Wolf, F. I., Di Francesco, A., Covacci, V., Corda, D., and Cittadini, A. (1996) Regulation of intracellular magnesium in ascites cells: Involvement of different regulatory pathways. *Arch. Biochem. Biophys.* **331**, 194–200
118. Howarth, F., Waring, J., Hustler, B., and Singh, J. (1994) Effects of extracellular magnesium and beta adrenergic stimulation on contractile force and magnesium mobilization in the isolated rat heart. *Magnes. Res.* **7**, 187–197
119. Vormann, J., and Günther, T. (1987) Amiloride-sensitive net Mg<sup>2+</sup> efflux from isolated perfused rat hearts. *Magnesium* **6**, 220–224
120. Amano, T., Matsubara, T., Watanabe, J., Nakayama, S., and Hotta, N. (2000) Insulin modulation of intracellular free magnesium in heart: Involvement of protein kinase C. *Br. J. Pharmacol.* **130**, 731–738
121. Morsy, M. S., Dishmon, D. A., Garg, N., and Weber, K. T. (2017) Secondary hyperparathyroidism in heart failure. *Am. J. Med. Sci.* **354**, 335–338
122. Sigurskjold, B. W. (2000) Exact analysis of competition ligand binding by displacement isothermal titration calorimetry. *Anal. Biochem.* **277**, 260–266
123. Case, D. A., Betz, R., Cerutti, D., Cheatham, T., Darden, T., Duke, R., Giese, T., Gohlke, H., Goetz, A., and Homeyer, N. (2016) *AMBER 2016 Reference Manual*. University of California, San Francisco, CA: 1–923
124. Leelananda, S. P., and Lindert, S. (2016) Computational methods in drug discovery. *Beilstein J. Org. Chem.* **12**, 2694–2718
125. Maier, J. A., Martinez, C., Kasavajhala, K., Wickstrom, L., Hauser, K. E., and Simmerling, C. (2015) ff14SB: Improving the accuracy of protein side chain and backbone parameters from ff99SB. *J. Chem. Theor. Comput.* **11**, 3696–3713
126. L DeLano, W. (2002) *The PyMOL Molecular Graphics System (2002)*, DeLano Scientific, Palo Alto, CA
127. Li, P., Roberts, B. P., Chakravorty, D. K., and Merz, K. M., Jr. (2013) Rational design of particle mesh ewald compatible Lennard-Jones parameters for +2 metal cations in explicit solvent. *J. Chem. Theor. Comput.* **9**, 2733–2748
128. Shirts, M. R., Mobley, D. L., and Brown, S. P. (2010) Free-energy calculations in structure-based drug design. In: Merz, K. M., Ringe, D., Reynolds, C. H., eds. *Drug Design: Structure-and Ligand-Based Approaches*, Cambridge University Press, Cambridge, England: 61–86
129. Borech, S., Tettinger, F., Leitgeb, M., and Karplus, M. (2003) Absolute binding free energies: A quantitative approach for their calculation. *J. Phys. Chem. B* **107**, 9535–9551
130. Rocklin, G. J., Mobley, D. L., Dill, K. A., and Hunenberger, P. H. (2013) Calculating the binding free energies of charged species based on explicit-solvent simulations employing lattice-sum methods: An accurate correction scheme for electrostatic finite-size effects. *J. Chem. Phys.* **139**, 184103