



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Shuhan He, Debbie Lai, *Jarone Lee
jarone.lee@mgh.harvard.edu

Lab of Computer Science (SH), Massachusetts General Hospital, Harvard Medical School, Harvard University, Boston, MA 02114, USA (JL); Department of Psychology and Language Sciences, University College London, London, UK (DL)

- 1 Ranney ML, Griffith V, Jha AK. Critical supply shortages—the need for ventilators and personal protective equipment during the COVID-19 pandemic. *N Engl J Med* 2020; **382**: e41.
- 2 Mansoor S. California Gov Gavin Newsom says federal government sent '170 broken ventilators'. March 29, 2020. <https://time.com/5812147/california-coronavirus-broken-ventilators> (accessed Jan 5, 2021).
- 3 Proctor N, O'Reilly K. Hospital repair restrictions, manufacturer-imposed barriers to fixing medical equipment cause inefficiencies and delays. July 8, 2020. https://uspirg.org/sites/pirg/files/reports/Hospital_Repair_Restrictions_USPEF_7.8.20b.pdf (accessed Jan 5, 2021).
- 4 European Parliament. Towards a more sustainable single market for business and consumers. March 11, 2020. https://www.europarl.europa.eu/doceo/document/A-9-2020-0209_EN.html# (accessed Jan 5, 2021).
- 5 Robertson A. Massachusetts passes 'right to repair' law to open up car data. Nov 4, 2020. <https://www.theverge.com/2020/11/4/21549129/massachusetts-right-to-repair-question-1-wireless-car-data-passes> (accessed Jan 5, 2021).
- 6 Wyden R. Wyden and Clarke introduce bill to eliminate barriers to fixing critical medical equipment during the pandemic. Aug 6, 2020. <https://www.wyden.senate.gov/news/press-releases/wyden-and-clarke-introduce-bill-to-eliminate-barriers-to-fixing-critical-medical-equipment-during-the-pandemic-> (accessed Jan 2, 2021).

Malnutrition risk in hospitalised COVID-19 patients receiving CPAP

Continuous positive airway pressure (CPAP) that is delivered by face mask or hood is increasingly used in patients with COVID-19 who have been admitted to hospital, often on general or respiratory wards. Oral intake of food and drink in patients with COVID-19 might have been and will often continue to be poor due to disease-associated anorexia, nausea, and impairment of taste. Full-face or hood CPAP also makes it impossible to eat and drink without mask removal,¹ which can be associated with decreased arterial oxygen saturation.

Staff might also fear that the use of nasogastric feeding can cause mask air leaks or promote gastric distension and aspiration due to aerophagia.

Such issues are in fact readily managed, and the British Association for Parenteral and Enteral Nutrition has produced practical guidelines.² However, NHS England and NHS Improvement³ advocate opioid administration when CPAP is used to reduce the sensation of breathlessness and high tidal volumes, an intervention that can impair gut motility.

We received reports of three patients with COVID-19 who were treated with CPAP and developed starvation ketosis and other patients who have gone for extended periods (up to 25 days) without substantial oral intake or initiation of nutritional support. The negative effects of malnutrition can be worsened by the process of muscle wasting, which is common in patients with COVID-19 who are admitted to hospital, and by subsequent admission to an intensive care unit for mechanical ventilation, where gut function might be impaired (eg, by use of opioid analgesia), as dietitians have reported to the Critical Care Specialist Group of the British Dietetic Association.

We were also made aware of the use of 0.9% saline in some health-care centres as the routine (and sole) intravenous crystalloid. Such practice does not comply with the National Institute for Health and Care Excellence clinical guideline 174: although 0.9% saline can be used for replacement of gastrointestinal losses or as a bolus for acute resuscitation, its use is not recommended in terms of routine maintenance.⁴ Use as routine maintenance can increase sodium and chloride load in the body, potentially leading to bowel oedema and further impairment of gastrointestinal function.⁵

We recommend that health-care professionals with expertise in nutrition, especially dietitians, nutrition nurses, physicians, and pharmacists, should be engaged in the assessment

and care of all patients with COVID-19 who receive CPAP and patients who are subsequently admitted to intensive care units for mechanical ventilation. Appropriate nutritional support—including the introduction of parenteral nutrition, if necessary—improves outcomes in analogous cases.⁶

PT reports lecturing fees from Fresenius Kabi and Baxter Healthcare. All other authors declare no competing interests.

*Pete Turner, Hugh Montgomery, Mike Stroud, George Grimble, Trevor Smith

pete.turner@setrust.hscni.net

Department of Nutrition and Diet Therapy, South Eastern Health and Social Care Trust, Belfast BT16 1RH, UK (PT); Division of Medicine, University College London, London, UK (HM, GG); Department of Gastroenterology, University Hospital Southampton NHS Foundation Trust, Southampton, UK (MS, TS)

- 1 Reeves A, White H, Sosnowski K, Tran K, Jones M, Palmer M. Energy and protein intakes of hospitalised patients with acute respiratory failure receiving non-invasive ventilation. *Clin Nutr* 2014; **33**: 1068–73.
- 2 British Association for Parenteral and Enteral Nutrition, British Dietetic Association. Route of nutrition support in patients requiring NIV & CPAP during the COVID-19 response. April 17, 2020. <https://www.bapen.org.uk/pdfs/covid-19/nutrition-in-niv-21-04-20.pdf> (accessed Dec 4, 2020).
- 3 NHS England and NHS Improvement. Guidance for the role and use of non-invasive respiratory support in adult patients with coronavirus (confirmed or suspected). Redditch: NHS England, 2020.
- 4 Intravenous fluid therapy in adults in hospital. Clinical guideline [CG174]. May 5, 2017. <https://www.nice.org.uk/guidance/cg174> (accessed March 16, 2021).
- 5 Li H, Sun S, Yap JQ, Chen J, Qian Q. 0.9% saline is neither normal nor physiological. *J Zhejiang Univ Sci B* 2016; **17**: 181–87.
- 6 Faisy C, Lerolle N, Dachraoui F, et al. Impact of energy deficit calculated by a predictive method on outcome in medical patients requiring prolonged acute mechanical ventilation. *Br J Nutr* 2009; **101**: 1079–87.



Ima Fassbender/Getty Images



Published Online
March 17, 2021
[https://doi.org/10.1016/S0140-6736\(21\)00447-5](https://doi.org/10.1016/S0140-6736(21)00447-5)



Published Online
March 18, 2021
[https://doi.org/10.1016/S0140-6736\(21\)00467-0](https://doi.org/10.1016/S0140-6736(21)00467-0)

Free licensing of vaccines to end the COVID-19 crisis

The pace of COVID-19 vaccine development, authorisation, and production is unprecedented. Yet all three approved vaccines by Pfizer–BioNTech, Moderna, and AstraZeneca