



Published in final edited form as:

Atmosphere (Basel). 2011 December ; 2(4): 633–654. doi:10.3390/atmos2040633.

Emission Ratios for Ammonia and Formic Acid and Observations of Peroxy Acetyl Nitrate (PAN) and Ethylene in Biomass Burning Smoke as Seen by the Tropospheric Emission Spectrometer (TES)

Matthew J. Alvarado^{1,*}, Karen E. Cady-Pereira¹, Yaping Xiao¹, Dylan B. Millet², Vivienne H. Payne¹

¹Atmospheric and Environmental Research (AER), Lexington, MA 02421, USA

²Department of Soil, Water, and Climate, University of Minnesota, St. Paul, MN 55455-0213, USA

Abstract

We use the Tropospheric Emission Spectrometer (TES) aboard the NASA Aura satellite to determine the concentrations of the trace gases ammonia (NH₃) and formic acid (HCOOH) within boreal biomass burning plumes, and present the first detection of peroxy acetyl nitrate (PAN) and ethylene (C₂H₄) by TES. We focus on two fresh Canadian plumes observed by TES in the summer of 2008 as part of the Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS-B) campaign. We use TES retrievals of NH₃ and HCOOH within the smoke plumes to calculate their emission ratios (1.0% ± 0.5% and 0.31% ± 0.21%, respectively) relative to CO for these Canadian fires. The TES derived emission ratios for these gases agree well with previous aircraft and satellite estimates, and can complement ground-based studies that have greater surface sensitivity. We find that TES observes PAN mixing ratios of ~2 ppb within these mid-tropospheric boreal biomass burning plumes when the average cloud optical depth is low (<0.1) and that TES can detect C₂H₄ mixing ratios of ~2 ppb in fresh biomass burning smoke plumes.

Keywords

biomass burning; satellite; TES; formic acid; ammonia; PAN; ethylene

1. Introduction

Biomass burning is the second largest source of trace gases to the global atmosphere and is an important part of the interannual variability of atmospheric composition [1–3]. Trace gases from biomass burning can contribute to the secondary chemical formation of aerosol particles and global tropospheric ozone, both of which impact upon climate and human health. However, the emissions of trace gases from biomass burning are highly uncertain.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).0020

* Author to whom correspondence should be addressed; matthew.alvarado@aer.com; Tel.: +1-781-761-2330; Fax: +1-781-761-2299.

Emission factors for biomass burning are primarily based on airborne and ground field measurements and measurements from small fires conducted in laboratories. Laboratory studies have shown that emissions of trace gases and particles from biomass burning can vary widely based on the type of fuel burned as well as the phase of combustion (*i.e.*, whether the combustion is in the early “flaming” stages or the later “smoldering” stages) [4]. However, the size, fuel moisture, and combustion characteristics of laboratory fires may not be representative of large-scale wildfires. Aircraft and ground studies of biomass burning emissions can only sample a small number of fires infrequently, making it difficult to understand the impact that the regional variability of fuel type and combustion phase can have on biomass burning emissions. Satellite observations, with their extensive spatial and temporal coverage, provide the opportunity to sample a large number of fires in several different ecosystems, which will help to characterize the spatial and temporal variability of emissions within a region for use in models of atmospheric chemistry, air quality, and climate.

Recent investigations have focused on using nadir-viewing satellite observations to estimate biomass burning emissions of trace gases and particles and study their subsequent chemistry. Examples of these studies include: estimating the emission rate of fine particles with fire radiative power (FRP) and aerosol optical depth retrievals from the Moderate-Resolution Imaging Spectroradiometer (MODIS) [5,6]; estimating emissions of NO_x with MODIS FRP and tropospheric NO_2 columns from the Ozone Monitoring Instrument (OMI) [7]; constraining emissions of CO using retrievals from the Measurements Of Pollution In The Troposphere (MOPITT) instrument, the Atmospheric Infrared Sounder (AIRS), the Scanning Imaging Absorption Spectrometer for Atmospheric Cartography (SCIAMACHY), and the Tropospheric Emission Spectrometer (TES) [8]; detecting several trace gases within smoke plumes using the Infrared Atmospheric Sounding Interferometer (IASI) [9,10]; and estimating the correlation between CO and O_3 in smoke plumes with TES [11–13].

TES made multiple special observations during the summer of 2008 over eastern Siberia, the North Pacific, and North America as part of the Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS-B) campaign [14]. This data set includes several observations of smoke plumes from boreal fires in Siberia and Canada. Alvarado *et al.* previously analyzed the correlation between TES retrievals of CO and O_3 within these smoke plumes [12]. Here we use nadir observations from TES to determine the concentrations of the trace gases ammonia (NH_3) and formic acid (HCOOH) within two boreal biomass burning plumes over Canada and determine the emission ratio of these gases relative to CO. The use of emission ratios relative to CO allows us to build on previous studies of emissions of CO from biomass burning [8] to estimate emissions of these less well-studied trace gases. We also present the first TES detections of peroxy acetyl nitrate (PAN), an important reservoir species of nitrogen oxides (NO_x) that is formed chemically within biomass burning smoke plumes, and the first TES detection of ethylene (C_2H_4), a reactive hydrocarbon emitted by biomass burning.

Biomass burning is a significant source of NH_3 [15]; other anthropogenic sources are livestock and chemical fertilizers, while natural sources include oceans, wild animal respiration, and soil microbial processes [16]. NH_3 is an integral component of the nitrogen

cycle. It can combine with acidic gases like H_2SO_4 and HNO_3 to form secondary aerosol, which then can impact climate and human health. This reactivity leads to a very short lifetime (less than two weeks) and large temporal and spatial variability. Background summer ammonia mixing ratios in the United States can range from 0.05 to 47 ppbv [17]. *In situ* observations of atmospheric ammonia are sparse and infrequent, making satellite observations of tropospheric NH_3 highly desirable. Worden *et al.* derived molar ratios of NH_3 to CO in smoke plumes from forest fires near San Luis Obispo, California, on 15 August 1994 using the Airborne Emission Spectrometer (AES), the airborne prototype for TES [18]. Beer *et al.* reported the first satellite observations of boundary layer NH_3 using the TES instrument aboard Aura [19]. Shephard *et al.* extended this work to a detailed strategy for retrieving NH_3 using TES [20]. Pinder *et al.* showed that the TES NH_3 retrievals were able to capture the spatial and seasonal variability of NH_3 over eastern North Carolina and that the retrievals compared well with *in situ* surface observations of NH_3 [21]. In addition, Clarisse *et al.* have used the nadir viewing IASI instrument to retrieve mixing ratios and global distributions of tropospheric NH_3 [10,22].

Formic acid (HCOOH) is a significant contributor to the acidity of precipitation and is an important oxygenated volatile organic compound [23–25]. Formic acid is ubiquitous in the troposphere, with typical surface concentrations ranging from 0.1 ppbv for “clean” environments to over 10 ppbv in urban polluted environments (e.g., Table 1 of [26]). The HCOOH lifetime ranges from several hours in the boundary layer to a few weeks in the free troposphere with wet (precipitation) and dry deposition the primary sinks, and reaction with OH of lesser importance [27]. There is considerable uncertainty concerning the origin of formic acid in the atmosphere. Some identified HCOOH sources include biogenic emissions from vegetation and soils, emissions from motor vehicles, and secondary production from organic precursors [26–28].

Biomass burning is another major primary source of formic acid, with several airborne studies showing that secondary production of formic acid also takes place within the aging smoke plume as the initial organic gases in the smoke are oxidized [29]. Enhanced mixing ratios of formic acid were measured in TES prototype airborne measurements of western wildfires [18]. The limb-viewing Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS) observed formic acid in young and aged biomass burning plumes in the upper troposphere and derived emission ratios for formic acid to CO [30,31]. Similarly, Grutter *et al.* used the limb-viewing Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) to retrieve global distributions of formic acid in the upper troposphere and stratosphere [32]. Razavi *et al.* presented global distributions of formic acid retrieved using the nadir-viewing IASI instrument and showed that the retrieved formic acid is correlated with CO during the burning season in Brazil, the Congo, and Southeast Asia [33].

PAN is a thermally unstable reservoir for NO_x that can be transported over large distances before converting back into NO_x , thereby altering ozone formation far downwind from the original source [34–36]. The primary NO_x emissions from biomass burning are rapidly converted to PAN within biomass burning plumes [12,37]. Satellite retrievals of PAN could provide substantial information on the fate of NO_x emitted by biomass burning in the

atmosphere and the impact of these NO_x emissions on global tropospheric ozone. The limb-viewing sounders MIPAS [38] and ACE-FTS [39] and the nadir-viewing IASI instrument [9,10] have all previously identified PAN in biomass burning smoke, but this species had not been previously detected in TES spectra.

Ethylene (C_2H_4) is a reactive hydrocarbon that is emitted directly by biomass burning [3]. It has a short lifetime in the summer Arctic troposphere (14–35 h, [12]) due to rapid reaction with OH. As this rapid oxidation of ethylene impacts the ozone formation rate within young smoke plumes [40], better estimates of the emissions of ethylene from biomass burning could help to reduce the uncertainty in the impact of biomass burning on tropospheric ozone. Enhanced mixing ratios of ethylene were measured in TES prototype airborne measurements of western wildfires [18], and C_2H_4 has also been previously identified by ACE-FTS [39] and IASI [9,10], but it had not been previously detected in TES spectra.

Section 2 describes the methods we used to identify biomass burning plumes from TES spectra, to retrieve NH_3 and HCOOH within the smoke plumes, and to detect PAN and C_2H_4 in TES spectra. Section 3 presents the results of this study and Section 4 summarizes our conclusions.

2. Methods

2.1. TES Special Observations During ARCTAS-B

TES is a nadir-viewing Fourier-transform infrared (FTIR) spectrometer aboard the NASA Aura spacecraft with a high spectral resolution of 0.06 cm^{-1} and a nadir footprint of $5.3\text{ km} \times 8.3\text{ km}$. Here we use Level 1B spectra (V003) for the 1B2 ($950\text{--}1150\text{ cm}^{-1}$) and 2A1 ($1100\text{--}1325\text{ cm}^{-1}$) bands of the TES instrument [41].

TES retrievals of trace gas profiles are based on an optimal estimation approach (with *a priori* constraints) that minimizes the differences between the TES Level 1B spectra and a radiative transfer calculation that uses absorption coefficients calculated with the line-by-line radiative transfer model LBLRTM [20,42–45]. Current Level 2 products from TES (V004) include retrieved profiles of CO, O_3 , H_2O , HDO, and CH_4 . The NH_3 retrieval discussed below will be included in the upcoming V005 of TES products, while the HCOOH retrieval discussed below is a prototype retrieval being developed at AER. The averaging kernel matrix of the retrieval gives the vertical sensitivity of the retrieved profile to the true profile, while the trace of the averaging kernel gives the degrees of freedom for signal (DOFS), which represents the number of independent pieces of information contained in the retrieval [44]. Cloud properties are retrieved by assuming single layer clouds with an effective optical depth that accounts for both cloud absorption and scattering [46]. Due to their small size (count median diameters of $\sim 0.13\text{ }\mu\text{m}$ [47]), biomass burning aerosols are unlikely to significantly impact radiances in the thermal infrared regions detected by TES, and any impact from larger particles is accounted for by the retrieved effective cloud optical depth.

The TES special observations during ARCTAS-B included nadir observations over eastern Siberia, the North Pacific, and North America for every 0.4° latitude. Biomass burning plumes were identified following the procedure in Alvarado *et al.* [12], which we briefly

outline here. We used maps of Level 3 daily AIRS retrievals of CO at $1^\circ \times 1^\circ$ resolution to identify the transport of CO from major regions of boreal biomass burning. We then used TES Level 2 retrievals of CO (V003) to identify the corresponding plumes that were observed by TES. The CO retrievals for TES special observations between 15 June and 15 July 2008 were filtered for data quality as recommended in the TES Level 2 Data Users Guide [48]. In general, the retrievals had 1 DOFS below 250 hPa with the region of maximum sensitivity in the troposphere near 500 hPa. We defined a plume in the TES special observations as an area where the retrieved CO mixing ratio at 510 hPa exceeded 150 ppb. This threshold ensured that the CO retrievals were significantly different from the *a priori* values (~ 110 ppb). While this procedure will detect thick plumes that are transported between continents [49], it does not detect plumes near the surface (where the sensitivity is low) or very thin or dilute plumes. We then used HYSPLIT back-trajectories [50] to determine if the observed air masses came from boreal biomass burning regions in Siberia (17 plumes) and Canada (5 plumes). The CO and O₃ retrievals for these plumes were analyzed by Alvarado *et al.* [12]. In this paper, we restrict our analysis to the fresh plumes from Canadian biomass burning.

2.2. NH₃ Retrieval

NH₃ retrievals were performed using TES Level 1B spectra (V003) [41] following the method of Shephard *et al.* [20], which has been implemented in V005 of the TES Level 2 products. The *a priori* profiles and covariance matrices for TES NH₃ retrievals are derived from GEOS-Chem model simulations of the 2005 global distribution of NH₃. Figure 1a shows an observed TES brightness temperature spectrum in the region of strong NH₃ absorption for a scan of a fresh Canadian smoke plume. Figures 1b,c show the brightness temperature residuals (observed spectrum minus modeled spectrum) for the unpolluted background NH₃ profile and the retrieved NH₃ profile, respectively, while Figure 1d shows the modeled spectrum of NH₃, calculated as the difference between the modeled spectrum including the retrieved NH₃ profile and the modeled spectrum without any NH₃. We can see a strong residual (-0.8 K) in the background profile spectrum that is substantially reduced following retrieval of NH₃. (The second strong residual feature near 949 cm^{-1} in panels b and c appears to be due to the ν_7 Q-branch of ethylene (C₂H₄), as is discussed in Section 3.4 below.)

Figure 2a shows the averaging kernel for the same NH₃ retrieval. The retrievals are performed using 14 vertical levels. However, the number of DOFS is generally not greater than one, meaning that it is not possible to obtain information about the shape of the profile from these retrievals. In order to minimize the influence of the *a priori* constraints on the end result, it is desirable to report one retrieved quantity per DOFS. However, performing a “profile” retrieval offers a major advantage in that it allows the calculation of diagnostics, such as the averaging kernels, that enable characterization of the spatial and temporal variability of the vertical sensitivity of the measurements. Therefore, for this work, the retrievals are performed on 14 levels in order to take advantage of the sensitivity characterization that this enables, then post-processed to calculate a single quantity that better represents the information that is really available in the measurement and that is relatively insensitive to the *a priori* constraints. Shephard *et al.* developed a Representative

Volume Mixing Ratio (RVMR) metric for NH_3 [20] based on similar techniques used previously for CH_4 [51] and CH_3OH [19]. This RVMR represents a TES sensitivity weighted average value where the influence of the *a priori* profile is reduced as much as possible. (Note that for NH_3 , one of three *a priori* profiles is selected based on the signal-to-noise ratio in the TES NH_3 band [20]. For the scan shown in Figures 1 and 2, the polluted *a priori*, shown as a dashed blue line in Figure 2b, was selected.) Figure 2b shows a retrieved profile of NH_3 for a fresh Canadian smoke plume as a solid black line, while the red circle is the RVMR calculated from the profile. The horizontal error bars show the uncertainty in the RVMR due to noise in the TES spectrum, while the vertical bars encompass the region of TES sensitivity to NH_3 , defined as the full width at half maximum (FWHM) of the averaging kernel at the pressure level of the RVMR [20]. With the *a priori* assumption about profile shape that was chosen here the NH_3 RVMR is roughly 20% to 60% of the retrieved surface value for NH_3 . The estimated minimum detection level is an RVMR of approximately 0.3 ppb, corresponding to a profile with a surface mixing ratio of about 1–2 ppb [20]. However, it must be noted that this is merely a general estimate of the minimum detection level, which depends strongly on the location of peak NH_3 mixing ratio and the thermal structure of the atmosphere for a given scan [20], and thus retrieved NH_3 RVMRs of less than 0.3 ppb are considered valid and are included in our analysis.

The excess mixing ratio of a trace gas like NH_3 (EMR, NH_3) is defined as the mixing ratio of the gas in the smoke plume minus its mixing ratio in the background. This excess mixing ratio can be normalized using the excess mixing ratio of CO to give the normalized excess mixing ratio (NEMR, NH_3/CO). The emission ratio (ER) is a special case of the NEMR where the measurements are made in fresh smoke near the fire source [3]. The NEMR of a trace gas can be highly variable for reactive gases downwind of fires due to the different rates of deposition and secondary photochemical production and loss for the trace gas and CO. In this paper, we will refer to our derived NEMRs for fresh Canadian smoke observed by TES as emission ratios; however, these emission ratios are inherently convolutions of the initial NEMR and any secondary production and loss processes that have taken place within the smoke plume during plume lofting and transport from the fire source [28]. Furthermore, the lower sensitivity of the TES retrievals near the surface (see Figure 2a) means that TES will preferentially sample smoke from the flaming stages of combustion, as these emissions are more likely to be lofted well above the surface. This lower sensitivity near the surface is due to the physics of nadir thermal infrared sounding: when the surface and the layers of the atmosphere near the surface have similar temperatures (low thermal contrast), we cannot distinguish between radiation emitted by the surface and radiation emitted by the lowest layers of the atmosphere. Similar caution must be used in interpreting emission ratios measured from other platforms: for example, NEMRs of NH_3 measured at the ground are generally much higher than those measured by aircraft, as the aircraft does not sample the emissions from residual smoldering combustion very close to the ground (see Section 3.1 below).

In order to use the TES retrievals of NH_3 and CO to calculate the emission ratio of NH_3 , we first calculated the RVMR for NH_3 following the procedure of Shephard *et al.* [20]. Since the DOFS for the CO retrieval are generally higher than for NH_3 , in order to obtain a comparable metric we transform the TES CO retrieval using the same grid and weightings as

were used to generate the NH_3 RVMR to obtain a pseudo-RVMR for CO. Figure 2c shows the retrieved CO profile for a fresh Canadian smoke plume as a solid black line while the pseudo-RVMR for CO is shown in green. The emission ratio of NH_3 was then calculated as the slope of a least squares linear regression of the NH_3 RVMR and the CO pseudo-RVMR.

2.3. HCOOH Retrieval

We have developed a prototype retrieval for formic acid (HCOOH) from TES Level 1B spectra. The retrieval approach is similar to that used for NH_3 . The spectroscopic parameters for HCOOH were taken from the HITRAN 2008 database, which substantially improved the estimates of the strengths of HCOOH lines by removing interference from the formic acid dimer [52]. All other spectroscopic parameters were taken from v1.4 of the TES spectroscopic line parameters [53]. The *a priori* constraint and covariance matrix were compiled from GEOS-Chem model simulations of the 2004 global background mixing ratios of HCOOH; however, these background profiles may be biased low, as GEOS-Chem tends to underestimate HCOOH in the northern mid-latitudes [28]. For HCOOH, the initial guess profile is the same as the *a priori* constraint vector. LBLRTM was run using TES Level 2 (V004) retrievals of temperature, average cloud optical depth, emissivity, reflectivity, H_2O , CO, O_3 , and CH_4 . Profiles of CO_2 and N_2O were taken from the TES Level 2 supplemental data files. The top panel of Figure 3 shows the observed TES brightness temperature spectrum between $1090\text{--}1130\text{ cm}^{-1}$ for a scan of fresh smoke from a Canadian fire with the HCOOH retrieval microwindows shown in red. Note that only some sections of the HCOOH band have been retained in order to remove interfering lines from other species, principally water vapor. The second panel shows the residuals (observed spectrum minus modeled spectrum) for the background HCOOH profile. We see a strong (-2 K) residual in the HCOOH Q-branch ($\sim 1105\text{ cm}^{-1}$), which is removed after retrieval of HCOOH as seen in the third panel of Figure 3.

An RVMR for HCOOH and a corresponding CO pseudo-RVMR were calculated following the procedure of Shephard *et al.* for NH_3 [20], as illustrated in Figure 4. The RVMR for HCOOH is lower than the retrieved value at the level of maximum sensitivity to HCOOH, possibly because the TES sensitivity to HCOOH covers a wide pressure range. The emission ratio of HCOOH was calculated as the slope of the least squares linear regression of the HCOOH RVMR and the CO pseudo-RVMR.

2.4. Detection of PAN and C_2H_4

In order to use TES to detect PAN in boreal smoke plumes, we calculated the differences (residuals) between the TES Level 1B spectra (V003) and a forward run of LBLRTM using v1.4 of the TES spectroscopic line parameters for the region of strong absorption by PAN ($1140\text{--}1180\text{ cm}^{-1}$). As for the retrievals of HCOOH, the model was run using TES Level 2 (V004) retrievals of temperature, emissivity, reflectivity, average cloud optical depth, H_2O , CO, O_3 , and CH_4 and profiles of CO_2 and N_2O were taken from the TES Level 2 supplemental data files. Preliminary model runs using the spectrally resolved effective cloud optical depth retrieved by TES led to unphysical slopes in the residuals versus wavenumber between 1100 and 1200 cm^{-1} . We removed these slopes by setting cloud optical depth in this spectral region to the average effective cloud optical depth included in the TES Level 2

products; however, the spectral signature of PAN was detectable regardless of which TES cloud product was used. The absorption cross section of PAN was taken from HITRAN 2008 [52].

A similar procedure was used to detect C₂H₄ in TES spectra, with the addition that the surface emissivity, surface temperature, and NH₃ profile were taken from the final results of the NH₃ retrieval described in Section 2.2 above. The line parameters for C₂H₄ were taken from v1.4 of the TES spectroscopic line parameters.

3. Results and Discussion

3.1. Emission Ratio of NH₃

Figure 5 shows a map of the retrieved NH₃ RVMR values within smoke plumes from Canadian biomass burning in the summer of 2008. We have filtered the retrievals for quality following Shephard *et al.* [20]. First, NH₃ retrievals are rejected if the DOFS are less than 0.1 or if the average cloud optical depth is greater than 1. Second, retrievals with DOFS less than 0.5 are rejected if the thermal contrast between the surface and the atmospheric layer nearest the surface is between -7 K and 10 K.

Figure 6 shows the NH₃ RVMR versus the CO pseudo-RVMR for all of the retrievals shown on the map in Figure 5. The most elevated amounts of CO and NH₃ come from a fresh smoke plume from a single Canadian fire observed by TES between 52.5–54.1°N and 90.5–91.8°W at 19:03 UTC on 1 July 2008 (TES Run #7656). Due to the high retrieved values for CO (>500 ppb), this plume likely contains concentrated fresh smoke. Also included are retrievals of NH₃ within a more dilute smoke plume observed between 55.5–59.8°N and 88.8–91.4°W at 18:51 UTC on 17 June 2008 (TES Run #7472).

We calculate the NH₃ emission ratio (NH_3/CO) for these fires as $1.0\% \pm 0.5\%$. However, there is a large amount of scatter in this plot, which leads to a low correlation between CO and NH₃ ($r^2 = 0.30$). The scatter is likely due to the combination of: (1) variations in plume heights, which change the relative sensitivity of TES to CO and NH₃ in the smoke plume; (2) variations in the age of the smoke plumes, as older plumes could have lost more NH₃ to chemical reaction or deposition, and (3) variations in the initial emissions of ammonia from the fires, which are in turn related to differences in the relative fraction of flaming versus smoldering combustion as well as variations in fuel nitrogen content.

Table 1 compares our derived emission ratio for NH₃ to previous satellite, aircraft, and ground measurements of ammonia emission ratios for boreal biomass burning. Our value is close to the value of 1.3% derived from IASI retrievals in Siberian smoke over East Mongolia [9]. The results from the aircraft and ground studies are substantially different with ground studies showing values of NH_3/CO that are a factor of 5 higher than the aircraft or satellite studies. The higher ground values are likely due to emissions of NH₃ from residual smoldering combustion very close to the ground [3] where aircraft cannot sample and satellites have little sensitivity to NH₃ (see Figure 2). Thus, both aircraft and satellites are likely to underestimate the true emissions of NH₃ from biomass burning fires unless the emission factors from aircraft and satellite studies are carefully combined with

ground observations, such as in the review of Akagi *et al.* [3]. In addition, given the discrepancy between aircraft and ground observations, nadir-viewing satellite observations of NH_3 are most appropriately compared with aircraft observations, as satellites will not be very sensitive to the ground-level residual smoldering emissions detected in ground studies. Our NH_3 emission factor of $1.0\% \pm 0.5\%$ is similar to the values of $1.3\% \pm 0.7\%$ and $1.7\% \pm 0.9\%$ reported for Alaskan forest fires in the aircraft studies of Nance *et al.* [54] and Goode *et al.* [55], respectively, but is substantially higher than the value of $0.16\% \pm 0.14\%$ reported by Radke *et al.* for boreal forests [56]. Furthermore, our value lies within the uncertainty range of the average value of $1.0\% \pm 0.7\%$ (boreal forests, aircraft studies only) reported in the review of Akagi *et al.* [3], as well as the value of $2.6\% \pm 2.2\%$ (all extratropical forests, all study types) in the review of Andreae and Merlet [57].

3.2. Emission Ratio of HCOOH

Figure 7 shows a map of the retrieved HCOOH RVMR values within smoke plumes from boreal biomass burning over Canada. We have removed retrievals with less than 0.5 DOFS for HCOOH, which also eliminated retrievals with an HCOOH RVMR of less than 0.5 ppbv. The smoke plumes included in the analysis are the same as described above for NH_3 . Figure 8 shows the HCOOH RVMR versus the CO pseudo-RVMR for all of the retrievals shown on the map in Figure 7. We calculate the emission ratio HCOOH/CO for these fires as $0.31\% \pm 0.21\%$. As we saw for NH_3 , there is a large amount of scatter in this plot that leads to a low correlation between CO and HCOOH ($r^2 = 0.41$). As discussed above, this scatter is likely due to the combination of variations in (1) plume heights, (2) plume ages, and (3) initial smoke emissions of HCOOH. The large pressure range of TES sensitivity to HCOOH may also contribute to the low correlation of HCOOH with CO. It is worth noting that the ACE-FTS observations of HCOOH within boreal biomass burning plumes showed a much higher r^2 value (0.86), likely due to the fact that the higher vertical resolution of limb retrievals means that ACE is better able to separate the different vertical layers of the plume which are averaged together by nadir-viewing sounders like TES and IASI [31].

Table 2 compares our derived emission ratio for HCOOH to previous satellite, aircraft, and ground measurements of formic acid emission ratios for boreal biomass burning. Our HCOOH emission ratio is slightly lower than the ratio of $0.38\% \pm 0.06\%$ derived from ACE-FTS observations [31], possibly due to the fact that a limb-sounder like ACE is more sensitive to plumes that reach high altitudes; this type of plume is more common over more flaming fires since fire power output peaks during flaming combustion [4]. Table 2 shows that our value is similar to the emission ratios for HCOOH derived from aircraft and ground studies, with the exception of the value of $3.7\% \pm 2.0\%$ reported for Canadian forests by Lefer *et al.* [60], which is about a factor of 10 higher than the average value for boreal biomass burning in the review of Akagi *et al.* [3]. However, Lefer *et al.* note that the elevated enhancement ratios of HCOOH they observed (0.82% to 6.2%) are likely due to HCOOH production within the plumes in the first couple hours after emission, and do not necessarily reflect the initial emissions. When the Lefer *et al.* study is removed, the Andreae and Merlet average becomes $0.85\% \pm 0.53\%$, with the relatively higher value of HCOOH likely due to the inclusion of midlatitude fires as well as boreal fires in the average [57].

3.3. Detection of PAN

The solid black line of Figure 9a shows the brightness temperature residuals (data minus model) for the scan of fresh smoke from a Canadian fire at 53.32°N and 90.84°W.

The two lobes of PAN absorption are clearly visible on either side of the water line at $\sim 1165 \text{ cm}^{-1}$. The dotted red line shows the residuals when a hypothetical PAN profile with a peak concentration of 1.9 ppb at 560 hPa is added to the forward model, and Figure 9b shows the modeled PAN spectrum as a solid black line. Adding the PAN profile substantially reduces the mean residuals, but this peak concentration of PAN in the assumed profile is about a factor of 2 higher than the concentration of PAN observed 30–60 min downwind of the Lake McKay fire in Saskatchewan, Canada during ARCTAS-B [12]. We have also detected PAN in a scan of aged smoke from Siberian biomass burning observed near Kamchatka (not shown). These two scans shared several common features, including relatively high mixing ratios of CO (mixing ratio $> 250 \text{ ppbv}$ at 510 hPa) and low average cloud optical depth (< 0.1). This low cloud optical depth may be a necessary but insufficient criterion for detecting PAN in biomass burning smoke plumes with TES.

3.4. Detection of C₂H₄

As mentioned in Section 2.2 above, there is a second strong residual feature in Figure 1c near 949 cm^{-1} that appears to be due to absorption by C₂H₄.

Figure 10b shows this residual feature, which partially overlaps with the strong CO₂ line [39] that is visible in the TES spectrum shown in Figure 10a. Figure 10c shows that the addition of a hypothetical C₂H₄ profile to the model (with a peak concentration of 1.9 ppb at the surface) removes this feature. This peak C₂H₄ mixing ratio is approximately the 85th percentile of C₂H₄ observations observed downwind of the Lake McKay fire in Saskatchewan, Canada during ARCTAS-B [12]. While the scan in Figure 10 did have a low cloud optical depth, this criterion is likely to be less important for detecting C₂H₄ within fresh biomass burning plumes using TES, as the residual feature of C₂H₄ is both stronger and sharper than the PAN feature. However, the short lifetime of C₂H₄ may make it difficult to detect in smoke that is several hours old.

4. Conclusions

We have retrieved mixing ratios of ammonia (NH₃) and formic acid (HCOOH) within biomass burning smoke plumes over Canada from TES radiance measurements. We have combined these retrievals with the TES retrievals of CO to calculate molar ratios of NH₃ and HCOOH to CO within biomass burning plumes by calculating representative volume mixing ratios for NH₃ and HCOOH and then mapping the CO retrieval to the same vertical grid. Our estimated emission ratios for NH₃ ($1.0\% \pm 0.5\%$) and HCOOH ($0.31\% \pm 0.21\%$) for forest fires in Canada are within the range of values reported in the literature for airborne and satellite studies of boreal biomass burning emissions. This work thus provides a method for the use of TES spectra to study the emissions of NH₃ and HCOOH from biomass burning. This method, if applied to the entire TES data set, would help to estimate the spatial and temporal variability of these emissions. This information could then be used, in concert

with the other satellite observations discussed in Section 1 of our manuscript, to provide improved estimates of the emissions of these trace gases for use in models of atmospheric chemistry, air quality, and climate.

We have shown that TES can observe peroxy acetyl nitrate (PAN) and ethylene (C₂H₄) within boreal biomass burning plumes. A low cloud optical depth (<0.1) appears to be required for successful detection of PAN by TES within biomass burning smoke plumes. Continuing this line of research could lead to maps of tropospheric PAN concentrations near source regions, which would help to constrain the fate of NO_x emission within atmospheric chemistry models. These could be combined with satellite estimates of the emissions of ethylene from biomass burning to help reduce the uncertainty in the impact of biomass burning on tropospheric ozone.

Acknowledgments

This research was supported by NASA grant NNX10AG65G to the University of Minnesota and NASA grants to Atmospheric and Environmental Research (AER) including Grant NNH08CD52C. This work was supported in part by the University of Minnesota Supercomputing Institute. We thank Eli Mlawer of AER, P. F. Bernath of the University of York, and the anonymous reviewers for their helpful comments.

References

1. Crutzen PJ; Andreae MO Biomass burning in the tropics: Impact on atmospheric chemistry and biogeochemical cycles. *Science* 1990, 250, 1669–1678. [PubMed: 17734705]
2. Bond TC; Streets DG; Yarber KF; Nelson SM; Woo J-H; Klimont Z. A technology-based global inventory of black and organic carbon emissions from combustion. *J. Geophys. Res.* 2004, 109, doi:10.1029/2003JD003697.
3. Akagi SK; Yokelson RJ; Wiedinmyer C; Alvarado MJ; Reid JS; Karl T; Crounse JD; Wennberg PO Emission factors for open and domestic biomass burning for use in atmospheric models. *Atmos. Chem. Phys.* 2011, 11, 4039–4072.
4. Burling IR; Yokelson RJ; Griffith DWT; Johnson TJ; Veres P; Roberts JM; Warneke C; Urbanski SP; Reardon J; Weise DR; et al. Laboratory measurements of trace gas emissions from biomass burning of fuel types from the southeastern and southwestern United States. *Atmos. Chem. Phys.* 2010, 10, 11115–11130.
5. Wooster MJ; Roberts G; Perry GLW; Kaufman YJ Retrieval of biomass combustion rates and totals from fire radiative power observations: FRP derivation and calibration relationships between biomass consumption and fire radiative energy release. *J. Geophys. Res.* 2005, 110, doi:10.1029/2005JD006318.
6. Ichoku C; Kaufman YJ A method to derive smoke emission rates from MODIS fire radiative energy measurements. *IEEE Trans. Geosci. Rem. Sensing* 2005, 43, 2636–2649.
7. Mebust AK; Russell AR; Hudman RC; Valin LC; Cohen RC Characterization of wildfire NO_x emissions using MODIS fire radiative power and OMI tropospheric NO₂ columns. *Atmos. Chem. Phys.* 2011, 11, 5839–5851.
8. Kopacz M; Jacob DJ; Fisher JA; Logan JA; Zhang L; Megretskaia IA; Yantosca RM; Singh K; Henze DK; Burrows JP; et al. Global estimates of CO sources with high resolution by adjoint inversion of multiple satellite datasets (MOPITT, AIRS, SCIAMACHY, TES). *Atmos. Chem. Phys.* 2010, 10, 855–876.
9. Coheur P-F; Clarisse L; Turquety S; Hurtmans D; Clerbaux C. IASI measurements of reactive trace species in biomass burning plumes. *Atmos. Chem. Phys.* 2009, 9, 5655–5667.
10. Clarisse L; R'Honi Y; Coheur P-F; Hurtmans D; Clerbaux C. Thermal infrared nadir observations of 24 atmospheric gases. *Geophys. Res. Lett.* 2011, 38, doi:10.1029/2011GL047271.

11. Verma S; Worden J; Pierce B; Jones DBA; Al-Saadi J; Boersma F; Bowman K; Eldering A; Fisher B; Jourdain L; et al. Ozone production in boreal fire smoke plumes using observations from the Tropospheric Emission Spectrometer and the Ozone Monitoring Instrument. *J. Geophys. Res.* 2009, 114, doi:10.1029/2008JD010108.
12. Alvarado MJ; Logan JA; Mao J; Apel E; Riemer D; Blake D; Cohen RC; Min K-E; Perring AE; Browne EC; et al. Nitrogen oxides and PAN in plumes from boreal fires during ARCTAS-B and their impact on ozone: An integrated analysis of aircraft and satellite observations. *Atmos. Chem. Phys.* 2010, 10, 9739–9760.
13. Dupont R; Pierce B; Worden J; Hair J; Fenn M; Hamer P; Natarajan M; Schaack T; Lenzen A; Apel E; et al. Reconstructing ozone chemistry from Asian wild fires using models, satellite and aircraft measurements during the ARCTAS campaign. *Atmos. Chem. Phys. Discuss.* 2010, 10, 26751–26812.
14. Jacob DJ; Crawford JH; Maring H; Clarke AD; Dibb JE; Emmons LK; Ferrare RA; Hostetler CA; Russell PB; Singh HB; et al. The Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) mission: Design, execution, and first results. *Atmos. Chem. Phys.* 2010, 10, 5191–5212.
15. Hegg DA; Radke LF; Hobbs PV; Riggan PJ Ammonia emissions from biomass burning. *Geophys. Res. Lett.* 1988, 15, 335–337.
16. Henze DK; Seinfeld JH; Shindell DT Inverse modeling and mapping US airquality influences of inorganic PM_{2.5} precursor emissions using the adjoint of GEOS-Chem. *Atmos. Chem. Phys.* 2009, 9, 5877–5903.
17. Langford AO; Fehsenfeld FS; Zachariassen J; Schimel DS Gaseous ammonia fluxes and background concentrations in terrestrial ecosystems of the United States. *Glob. Biogeochem. Cycles* 1992, 6, 459–483.
18. Worden H; Beer R; Rinsland C. Airborne infrared spectroscopy of 1994 western wildfires. *J. Geophys. Res.* 1997, 102, 1287–1299.
19. Beer R; Shephard MW; Kulawik SS; Clough SA; Eldering A; Bowman KW; Sander SP; Fisher BM; Payne VH; Luo M; et al. First satellite observations of lower tropospheric ammonia and methanol. *Geophys. Res. Lett.* 2008, 35, doi:10.1029/2008GL033642.
20. Shephard MW; Cady-Pereira KE; Luo M; Henze DK; Pinder RW; Walker JT; Rinsland CP; Bash JO; Zhu L; Payne VH; et al. TES ammonia retrieval strategy and global observations of the spatial and seasonal variability of ammonia. *Atmos. Chem. Phys. Discuss.* 2011, 11, 16023–16074.
21. Pinder RW; Walker JT; Bash JO; Cady-Pereira KE; Henze DK; Luo M; Osterman GB; Shephard MW Quantifying spatial and seasonal variability in atmospheric ammonia with in situ and space-based observations. *Geophys. Res. Lett.* 2011, 38, doi:10.1029/2010GL046146.
22. Clarisse L; Clerbaux C; Dentener F; Hurtsmans D; Coheur P-F Global ammonia distribution derived from infrared satellite observations. *Nat. Geosci.* 2009, 2, 479–483.
23. Kawamura K; Steinberg S; Kaplan IR Homologous series of C1-C10 monocarboxylic acids and C1-C6 carbonyls in Los Angeles air and motor vehicle exhausts. *Atmos. Environ.* 2000, 34, 4175–4191.
24. Keene WC; Galloway JN The biogeochemical cycling of formic and acetic acids through the troposphere: An overview of current understanding. *Tellus B* 1988, 40B, 322–334.
25. Avery GB Jr.; Tang Y; Kieber RJ; Willey JD Impact of recent urbanization on formic and acetic acid concentrations in coastal North Carolina rainwater. *Atmos. Environ.* 2001, 35, 3353–3359.
26. Shephard MW; Goldman A; Clough SA; Mlawer EJ Spectroscopic improvements providing evidence of formic acid in AERI-LBLRTM validation spectra. *J. Quant. Spectrosc. Radiat. Transf.* 2003, 82, 383–390.
27. Rinsland CP; Mahieu E; Zander R; Goldman A; Wood S; Chiou L. Free tropospheric measurements of formic acid (HCOOH) from infrared ground-based solar absorption spectra: Retrieval approach, evidence for a seasonal cycle, and comparison with model calculations. *J. Geophys. Res.* 2004, 109, doi:10.1029/2004JD004917.
28. Paulot F; Wunch D; Crounse JD; Toon GC; Millet DB; DeCarlo PF; Vigouroux C; Deutscher NM; González Abad G; Notholt J; et al. Importance of secondary sources in the atmospheric budgets of formic and acetic acids. *Atmos. Chem. Phys.* 2011, 11, 1989–2013.

29. Yokelson RJ; Crounse JD; DeCarlo PF; Karl T; Urbanski S; Atlas E; Campos T;Shinozuka Y; Kapustin V; Clarke AD; et al. Emissions from biomass burning in the Yucatan. *Atmos. Chem. Phys.* 2009, 9, 5785–5812.
30. Rinsland CP; Boone CD; Bernath PF; Mahieu E; Zander R; Dufour G; Clerbaux C;Turquety S; Chiou L; McConnell JC; et al. First space-based observations of formic acid (HCOOH): Atmospheric Chemistry Experiment austral spring 2004 and 2005 Southern Hemisphere tropical-mid-latitude upper tropospheric measurements. *Geophys. Res. Lett.* 2006, 33, doi:10.1029/2006GL027128.
31. Tereszchuk KA; González Abad G; Clerbaux C; Hurtmans D; Coheur P-F; Bernath PFACE-FTS measurements of trace species in the characterization of biomass burning plumes. *Atmos. Chem. Phys. Discuss.* 2011, 11, 16611–16637.
32. Grutter M; Glatthor N; Stiller GP; Fischer H; Grabowski U; Höpfner M; Kellmann S;Linden A; von Clarmann T. Global distribution and variability of formic acid as observed by MIPAS-ENVISAT. *J. Geophys. Res.* 2010, 115, doi:10.1029/2009JD012980.
33. Razavi A; Karagulian F; Clarisse L; Hurtmans D; Coheur PF; Clerbaux C; Müller JF;Stavrakou T. Global distributions of methanol and formic acid retrieved for the first time from the IASI/MetOp thermal infrared sounder. *Atmos. Chem. Phys.* 2011, 11, 857–872.
34. Val Martín M; Honrath RE; Owen RC; Pfister G; Fialho P; Barata F. Significantenhancements of nitrogen oxides, black carbon, and ozone in the North Atlantic lower free troposphere resulting from North American boreal wildfires. *J. Geophys. Res.* 2006, 111, doi:10.1029/2006JD007530.
35. Leung F-YT; Logan JA; Park R; Hyer E; Kasischke E; Streets D; Yurganov L. Impacts of enhanced biomass burning in the boreal forests in 1998 on tropospheric chemistry and the sensitivity of model results to the injection height of emissions. *J. Geophys. Res.* 2007, 112, doi:10.1029/2006JD008132.
36. Real E; Law KS; Weinzierl B; Fiebig M; Petzold A; Wild O; Methven J; Arnold S;Stohl A; Huntrieser H; et al. Processes influencing ozone levels in Alaskan forest fire plumes during long-range transport over the North Atlantic. *J. Geophys. Res.* 2007, 112, doi:10.1029/2006JD007576.
37. Jacob D; Wofsy SC; Bakwin PS; Fan S-M; Harriss RC; Talbot RW; Bradshaw JD;Sandholm ST; Singh HB; Browell EV; et al. Summertime Photochemistry of the Troposphere at High Northern Latitudes. *J. Geophys. Res.* 1992, 97, 16421–16431.
38. Moore DP; Remedios JJ Seasonality of Peroxyacetyl nitrate (PAN) in the upper troposphereand lower stratosphere using the MIPAS-E instrument. *Atmos. Chem. Phys.* 2010, 10, 6117–6128.
39. Coheur P-F; Herbin H; Clerbaux C; Hurtmans D; Wespes C; Carleer M; Turquety S; Rinsland CP; Remedios J; Hauglustaine D; et al. ACE-FTS observation of a young biomass burning plume: First reported measurements of C₂H₄, C₃H₆O, H₂CO and PAN by infrared occultation from space. *Atmos. Chem. Phys.* 2007, 7, 5437–5446.
40. Alvarado MJ; Prinn RG Formation of ozone and growth of aerosols in young smoke plumesfrom biomass burning: 1. Lagrangian parcel studies. *J. Geophys. Res.* 2009, 114, doi:10.1029/2008JD011144.
41. Shephard MW; Worden HM; Cady-Pereira KE; Lampel M; Luo M; Bowman KW;Sarkissian E; Beer R; Rider DM; Tobin DC; et al. Tropospheric Emission Spectrometer nadir spectral radiance comparisons. *J. Geophys. Res.* 2008, 113, doi:10.1029/2007JD008856.
42. Clough SA; Shephard MW; Worden J; Brown PD; Worden HM; Luo M; Rodgers CD;Rinsland CP; Goldman A; Brown L; et al. Forward model and jacobians for tropospheric emission spectrometer retrievals. *IEEE Trans. Geosci. Remote* 2006, 44, 1308–1323.
43. Shephard MW; Clough SA; Payne VH; Smith WL; Kireev S; Cady-Pereira KEPerformance of the line-by-line radiative transfer model (LBLRTM) for temperature and species retrievals: IASI case studies from JAIVEx. *Atmos. Chem. Phys.* 2009, 9, 7397–7417.
44. Rodgers CD *Inverse Methods for Atmospheric Sounding: Theory and Practice*; World Scientific: Hackensack, NJ, USA, 2000.
45. Bowman KW; Rodgers CD; Sund-Kulawik S; Worden J; Sarkissian E; Osterman G;Steck T; Luo M; Eldering A; Shephard MW; et al. Tropospheric emission spectrometer: Retrieval method and error analysis. *IEEE Trans. Geosci. Remote* 2006, 44, 1297–1307.

46. Kulawik SS; Worden J; Eldering A; Bowman K; Gunson M; Osterman GB; Zhang L; Clough SA; Shephard MW; Beer R. Implementation of cloud retrievals for Tropospheric Emission Spectrometer (TES) atmospheric retrievals: Part 1. Description and characterization of errors on trace gas retrievals. *J. Geophys. Res.* 2006, 111, doi:10.1029/2005JD006733.
47. Reid JS; Koppmann R; Eck TF; Eleuterio DP A review of biomass burning emissions part II: Intensive physical properties of biomass burning particles. *Atmos. Chem. Phys.* 2005, 5, 799–825.
48. Bowman K; Eldering A; Fisher B; Herman R; Jacob D; Jourdain L; Kulawik S; Luo M; Monarrez R; et al. Earth Observing System (EOS) Tropospheric Emission Spectrometer (TES) Level 2 (L2) Data User's Guide; Osterman G, Ed.; Jet Propulsion Laboratory (JPL): Pasadena, CA, USA, 2008.
49. Zhang L; Jacob DJ; Boersma KF; Jaffe DA; Olson JR; Bowman KW; Worden JR; Thompson AM; Avery MA; Cohen RC; et al. Transpacific transport of ozone pollution and the effect of recent Asian emission increases on air quality in North America: An integrated analysis using satellite, aircraft, ozonesonde, and surface observations. *Atmos. Chem. Phys.* 2008, 8, 6117–6136.
50. Air Resources Laboratory. HYSPLIT—Hybrid Single Particle Lagrangian Integrated Trajectory Model. Available online: <http://ready.arl.noaa.gov/HYSPLIT.php> (accessed on 25 September 2008).
51. Payne VH; Clough SA; Shephard MW; Nassar R; Logan JA Information-centered representation of retrievals with limited degrees of freedom for signal: Application to methane from the Tropospheric Emission Spectrometer. *J. Geophys. Res.* 2009, 17, 1095.
52. Rothman L; Gordon IE; Barbe A; Benner DC; Bernath PF; Birk M; Boudon V; Brown LR; Campargue A; Champion J-P; et al. The HITRAN 2008 molecular spectroscopic database. *J. Quant. Spectrosc. Radiat. Transf.* 2009, 110, 533–572.
53. AER Line Parameter Database. What's New. Available online: http://rtweb.aer.com/line_param_frame.html (accessed on 1 August 2010).
54. Nance J; Hobbs P; Radke L; Ward D. Airborne measurements of gases and particles from an Alaskan wildfire. *J. Geophys. Res.* 1993, 98, 14873–14882.
55. Goode JG; Yokelson RJ; Ward DE; Susott RA; Babbitt RE; Davies MA; Hao WM Measurements of excess O₃, CO₂, CO, CH₄, C₂H₄, C₂H₂, HCN, NO, NH₃, HCOOH, CH₃COOH, HCHO and CH₃OH in 1997 Alaskan biomass burning plumes by airborne fourier transform infrared spectroscopy (AFTIR). *J. Geophys. Res.* 2000, 105, 22147–22166.
56. Radke LF; Hegg DA; Hobbs PV; Nance JD; Lyons JH; Laursen KK; Weiss RE; Riggan PJ; Ward DE Particulate and Trace Gas Emissions from Large Biomass Fires in North America. In *Global Biomass Burning: Atmospheric, Climatic, and Biospheric Implications*; Levine JS, Ed.; MIT Press: Cambridge, MA, USA, 1991; pp. 209–224.
57. Andreae M; Merlet P. Emission of trace gases and aerosols from biomass burning. *Glob. Biogeochem. Cycles* 2001, 15, 955–966.
58. Yokelson R; Susott R; Ward D; Reardon J; Griffith D. Emissions from smoldering combustion of biomass measured by open-path Fourier transform infrared spectroscopy. *J. Geophys. Res.* 1997, 102, 18865–18877.
59. Bertschi I; Yokelson RJ; Ward DE; Babbitt RE; Susott RA; Goode JG; Hao WM Trace gas and particle emissions from fires in large diameter and belowground biomass fuels. *J. Geophys. Res.* 2003, 108, doi:10.1029/2002JD002100.
60. Lefer B; Talbot RW; Harriss RH; Bradshaw JD; Sandholm ST; Olson JO; Sachse GW; Collins J; Shipham MA; Blake DR; et al. Enhancement of acidic gases in biomass burning impacted air masses over Canada. *J. Geophys. Res.* 1994, 99, 1721–1737.




Figure 1.

(a) Tropospheric Emission Spectrometer (TES) observed brightness temperature spectrum, (b) the background profile residuals, calculated as observed minus modeled spectrum for an unpolluted profile of NH₃, (c) the residuals after retrieval of NH₃, and (d) the modeled spectrum of NH₃ for the TES scan at 53.32° N and 90.84° W on 1 July 2008 near a Canadian fire. The microwindows used in the NH₃ retrieval are shown in red. Note that surface emissivity and temperature are adjusted during the retrieval, which requires microwindows outside of the NH₃ absorption band (~960–970 cm⁻¹).




Figure 2.

(a) NH_3 averaging kernel for the same TES scan as shown in Figure 1. (b) Background unpolluted NH_3 profile (open black squares), polluted *a priori* NH_3 profile (open blue diamonds), retrieved NH_3 profile (filled black squares) and NH_3 Representative Volume Mixing Ratio (RVMR) (red circle) for the same scan. The horizontal error bars show the uncertainty in the RVMR due to noise in the TES spectrum, while the vertical bars encompass the region of TES sensitivity to NH_3 . (c) Background (and *a priori*) CO profile (open black triangles), retrieved CO profile (filled black triangles) and CO pseudo-RVMR (green inverted triangle) for the same scan.




Figure 3. (a) TES observed brightness temperature spectrum, (b) background formic acid (HCOOH) profile residuals, calculated as observed minus modeled spectrum, (c) residuals after retrieval of HCOOH, and (d) the modeled spectrum of HCOOH for the same TES scan as shown in Figure 1. The microwindows used in the HCOOH retrieval are shown in red.




Figure 4.

(a) HCOOH averaging kernel for the same TES scan as shown in Figure 1. (b) Background (and *a priori*) HCOOH profile (open black squares), retrieved HCOOH profile (filled black squares), and HCOOH RVMR (intersection of red lines) for the same scan. The red dot was omitted so that the horizontal error bars would be visible. (c) Background (and *a priori*) CO profile (open black triangles), retrieved CO profile (filled black triangles) and CO pseudo-RVMR (green inverted triangle) for the same scan.




Figure 5. RVMR of NH_3 for the TES special observations for Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS-B) between 15 June 2008 and 15 July 2008. Only retrievals within biomass burning plumes over Canada are shown. Note that the red point is 7.3 ppb.




Figure 6. RVMR of NH₃ versus the pseudo-RVMR of CO for all of the retrievals shown in Figure 5.




Figure 7. RVMR of HCOOH for the TES special observations for ARCTAS-B between 15 June 2008 and 15 July 2008. Only retrievals within biomass burning plumes over Canada are shown.




Figure 8.
RVMR of HCOOH versus the pseudo-RVMR of CO for the retrievals shown in Figure 7.




Figure 9. (a) Brightness temperature residuals (data minus model) for the same TES scan as in Figure 1. The solid black line does not have PAN in the forward model, while the dotted red line does include PAN. (b) Difference between the model runs with and without PAN, which shows the modeled spectrum of PAN.




Figure 10.

(a) TES observed brightness temperature spectrum, (b) residuals without a C₂H₄ profile in the model, calculated as observed minus modeled spectrum, (c) residuals after addition of a hypothetical profile of C₂H₄, and (d) the modeled spectrum of C₂H₄ for the same TES scan as shown in Figure 1. The area of strong absorption by C₂H₄, corresponding to the ν_7 Q-branch, is shown in red.

Table 1.Ammonia emission ratios (NH_3/CO , mol/mol) for boreal biomass burning.

NH_3/CO	Ecosystem/Fuel	Study Type	Source
$1.0\% \pm 0.5\%$	Canadian Forest	Satellite (TES)	This Study
1.3%	Siberian Forest	Satellite (IASI)	[9]
$0.16\% \pm 0.14\%$	Boreal Forests	Aircraft	[56]
$1.3\% \pm 0.7\%$	Alaskan Forest	Aircraft	[54]
$1.7\% \pm 0.9\%$	Alaskan Forest	Aircraft	[55]
$6.9\% \pm 11.1\%$	Boreal Peat	Ground	[58]
$5.9\% \pm 3.6\%$	Boreal organic soil	Ground	[58]
$6.9\% \pm 3.2\%$	Boreal organic soil	Ground	[59]
$1.7\% \pm 2.2\%$	Boreal dead, woody material	Ground	[59]
3.5%	Alaskan Duff	Laboratory	[4]
$1.0\% \pm 0.7\%$	Boreal Forests	Review: Aircraft Only	[3]
$5.1\% \pm 5.6\%$	Boreal Forests	Review: Ground Only	[3]
$3.5\% \pm 3.2\%$	Boreal Forests	Review: Aircraft and Ground	[3]
$2.6\% \pm 2.2\%$	Extratropical Forests	Review: Aircraft, Satellite, and Ground	[57]

Table 2.Formic acid emission ratios (HCOOH/CO , mol/mol) for boreal biomass burning.

HCOOH/CO	Ecosystem/Fuel	Study Type	Source
$0.31\% \pm 0.21\%$	Canadian Forest	Satellite (TES)	This Study
$0.38\% \pm 0.06\%$	Canadian Forest	Satellite (ACE-FTS)	[31]
$0.31\% \pm 0.09\%$	Alaskan Forest	Aircraft	[55]
$0.26\% \pm 0.44\%$	Boreal Peat	Ground	[58]
$0.18\% \pm 0.12\%$	Boreal organic soil	Ground	[58]
$0.33\% \pm 0.29\%$	Boreal organic soil	Ground	[59]
$0.17\% \pm 0.18\%$	Boreal dead, woody material	Ground	[59]
0.35%	Alaskan Duff	Laboratory	[4]
$0.28\% \pm 0.24\%$	Boreal Forests	Review, Aircraft and Ground	[3]
$3.7\% \pm 2.0\%$	Canadian Forest	Aircraft	[60]
$1.4\% \pm 1.4\%$	Extratropical Forests	Review, Aircraft, Satellite and Ground	[57]