



Published in final edited form as:

Curr Phys Med Rehabil Rep. 2020 December ; 8(4): 461–468. doi:10.1007/s40141-020-00283-3.

Post-stroke cognitive impairments and responsiveness to motor rehabilitation: A review

Jennapher Lingo VanGilder, M.S.¹, Andrew Hooyman, Ph.D.¹, Daniel S. Peterson, Ph.D.^{2,3,4}, Sydney Y. Schaefer, Ph.D.^{1,4}

¹School of Biological and Health Systems Engineering, Arizona State University

²College of Health Solutions, Arizona State University

³U.S. Department of Veterans Affairs, Phoenix, AZ, USA.

⁴Department of Physical Therapy and Athletic Training, University of Utah

Abstract

a) Purpose of review: This review discusses the prevalence of cognitive deficits following stroke and their impact on responsiveness to therapeutic intervention within a motor learning context.

b) Recent findings: Clinical and experimental studies have established that post-stroke cognitive and motor deficits may impede ambulation, augment fall risk, and influence the efficacy of interventions. Recent research suggests the presence of cognitive deficits may play a larger role in motor recovery than previously understood.

c) Summary: Considering that cognitive impairments affect motor relearning, post-stroke motor rehabilitation therapies may benefit from formal neuropsychological testing. For example, early work suggests that in neurotypical adults, cognitive function may be predictive of responsiveness to motor rehabilitation and cognitive training may improve mobility. This sets the stage for investigations probing these topics in people post-stroke. Moreover, the neural basis for and extent to which these cognitive impairments influence functional outcome remains largely unexplored and require additional investigation.

Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. <http://www.springer.com/gb/open-access/authors-rights/aam-terms-v1>

Corresponding author: Sydney Y. Schaefer, PhD, Arizona State University, 501 E. Tyler Mall, MC 9709 Tempe, AZ 85287, sydney.schaefer@asu.edu.

Compliance with Ethics Guidelines

Human and Animal Rights and Informed Consent

This article does not contain any studies with human or animal subjects performed by any of the authors.

Publisher's Disclaimer: This Author Accepted Manuscript is a PDF file of a an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

Conflict of Interest

Jennapher Lingo VanGilder, Andrew Hooyman, Daniel Peterson and Sydney Schaefer declare no conflicts of interest relevant to this manuscript.

Keywords

cognitive decline; rehabilitation; stroke; falls; gait; function

Introduction

Motor learning is important for motor rehabilitation as individuals must often relearn lost motor skills [1]. Evidence from clinical and experimental studies have long supported that specific cognitive abilities such as attention, working memory, and visuospatial ability are related to both performance and performance improvement on novel motor tasks (i.e., procedural learning) [2–10]. This may be problematic for stroke survivors, as these same domains are often impacted post-stroke. While the reporting of cognitive data has become more customary in stroke rehabilitation studies, these data are often limited to global cognitive screening tools that provide only a cursory glimpse into cognitive function and are typically used only to exclude individuals with low scores. Moreover, a comprehensive understanding of specific cognitive impairments in stroke-survivors, and the extent to which they interfere with gait rehabilitation remains a critical knowledge gap.

Here, we will briefly summarize key points regarding the effects of stroke on gait and posture, and discuss a novel, less-studied area of research regarding the extent to which cognitive impairments may interfere with motor learning and rehabilitation. We first review common neuropsychological assessments used to evaluate cognition, particularly those relevant to the stroke survivor population, and discuss the limitations and broader implications of using these assessments. We then discuss the role of stroke-specific cognitive deficits in gait and postural control, and how these deficits may influence the degree of improvement in motor behaviors. Finally, we discuss our understanding of how specific cognitive deficits may impact stroke rehabilitation and propose that thorough neuropsychological evaluation be integrated in stroke protocols (rather than serve as exclusion criteria) to clarify cognition's role in recovery and relearning of motor skills.

What are common cognitive impairments associated with stroke and how can they be measured?

While cognitive impairment following stroke is certainly linked to the lesion location and/or size [11–13], clinical studies indicate that impairments in attention, executive function, and processing speed are the most prevalent across stroke survivors [12, 14]. In fact, stroke survivors are tenfold more likely to show impaired memory, orientation, language, and attention, compared to age-matched non-stroke individuals, with prevalence rates of 35% vs 3%, respectively [15]. Moreover, these impairments may be differentially impacted throughout neurological recovery. For instance, attention and executive functions may be most susceptible to impairment at the time of stroke diagnosis [16], whereas impairments in memory, executive, and visuospatial functions are notable three months post-stroke [17]. Interestingly, these cognitive deficits persisted in patients with no apparent physical or cognitive disability as screened by the modified Rankin Scale and Mini-Mental State Exam, respectively [17], suggesting subtle cognitive impairments may remain undetected if evaluated with a brief global cognitive screen.

However, there is no gold standard for the diagnosis of post-stroke cognitive impairment (i.e., vascular cognitive impairment) [12], which obfuscates the selection of clinical cognitive screening and assessments. For instance, the Mini-Mental State Exam (MMSE) has been widely used as a clinical diagnostic tool of cognitive impairment since its advent in 1975, despite its authors' warnings it cannot be used exclusively to diagnose impairment [18]. In fact, it excludes an evaluation of executive function and has poor sensitivity in Mild Cognitive Impairment detection [19]. The Montreal Cognitive Assessment (MoCA) is also commonly used, and although it provides a measure of executive function and may be more reliable in Mild Cognitive Impairment detection [19], it has poor sensitivity in quantifying cognitive function of the domains it asserts to evaluate [20]. Moreover, these brief clinical assessments do not provide standardized age-adjusted scoring (although recent work has attempted to address this limitation [21]), which may be particularly important considering the preponderance of older adults among the stroke population [22].

The National Institutes of Health (NIH) developed the National Institutes of Health Stroke Scale [23], a measure of activities of consciousness, movement, sensation, response and advanced neurological function in stroke patients, and is a reliable indicator of stroke severity [24]. Similar to other global cognitive screens, however, it yields a global measure of cognition that may be insensitive to cognitive deficits [25]; it is also susceptible to floor effects and biased towards hemisphere-specific lesions [26]. The NIH has proposed a validated, standardized, robust measure of cognitive function, namely the NIH Toolbox Cognitive Battery [27], however, it is only appropriate for research applications and does not serve as a substitution for formal neuropsychological or other clinical testing. At present, the battery has only been validated in healthy populations while the work to validate it in traumatic brain injury, spinal cord injury, and stroke cases remains ongoing [28–30]. In the interim, implementing formal neuropsychological testing, specifically tests that thoroughly evaluate cognitive domains particularly relevant to motor-related outcomes (e.g., executive function and functional outcome in stroke patients [31]; spatial working memory with procedural learning [32]), may provide critical prognostic insight into stroke rehabilitation outcomes. This aligns with the first recommendation by the Cognition Working Group, which convened as part of the second international Stroke Recovery and Rehabilitation Roundtable [33].

How is cognition typically measured?

Global cognition broadly encompasses six domains of cognitive function, namely attention, executive function, learning and memory, language, perceptual-motor function, and social cognition, with each domain being further stratified into subdomains [34]. As previously mentioned, the MoCA and MMSE are ubiquitous clinical tools used to quickly screen global cognitive status by cursorily evaluating attention, language, memory, and visuospatial/executive functions. Although these global screening tools are relatively quick (approximately 5–10 minutes) and easy to administer, they may have differential sensitivity to premorbid abilities [34], and are subject to age, educational, and cultural background confounds [35]. More extensive neuropsychological assessments that rigorously test each cognitive domain may provide a more robust estimation of cognitive status (e.g., Repeatable Battery for Neuropsychological Status [36], Wechsler Adult Intelligence Scale [37]), yet

often require costly instrumentation, appropriate licensure, and longer administration periods (approximately 30 and 75 minutes, respectively). Unlike the MoCA or MMSE (or similar), these assessments can evaluate individual domains (e.g., complex attention, language) and subdomains (e.g., long-term memory, working memory). For instance, the Repeatable Battery for Neuropsychological Status and Weschler Adult Intelligence Scale yield an age-adjusted composite score comprising multiple index scores, each validated to represent a specific cognitive domain.

If the function of a single cognitive domain is of interest to a clinician, they can utilize individual neuropsychological tests. For example, the Rey-Osterrieth Complex Figure test is a widely used paper-and-pencil examination that measures visuospatial construction, immediate visuospatial memory, and delayed visuospatial memory [38], and has also been shown to evaluate latent constructs such as graphomotor function, object use and planning, visuo-motor transformation, and visuospatial perception [39]. One major advantage of using standardized cognitive assessments is that normative data, user qualifications, administration instructions, and results reporting are generally well-documented, and reputable online databases that thoroughly review important considerations of individual neuropsychological assessments (such as cost, test/retest reliability, cutoff scores, normative data, etc.) are publicly accessible, much like many physical/motor assessments that physical and occupational therapists use (e.g., <https://www.sralab.org/rehabilitation-measures> Shirley Ryan AbilityLab). We take the time to summarize this point here to encourage future studies to utilize standardized, validated assessments alongside (or in place of) novel experimental methods for characterizing cognition post-stroke.

When formal neuropsychological testing is unavailable or infeasible, the American Psychiatric Association's Diagnostic and Statistical Manual of Mental Disorders provides a list of brief assessments that can provide insight into each subdomain. For example, to evaluate planning ability (a subdomain of executive function), the examinee should demonstrate the ability to find the exit to a maze and/or interpret a sequential picture or object arrangement [34]. Similarly, while researchers may develop experimental approaches that provide insight into a specific cognitive domain (e.g., [32]), one caveat to experimenter-derived assessments is that they are not necessarily standardized (i.e., generalizable), potentially complicating comparison and replication of research findings among laboratories.

We also want to briefly acknowledge a recent study demonstrated that years of education explained differences in cognitive factors such as executive function, working memory, global cognition, and alertness, as well as motor function (as measured by the modified Rankin Scale) among stroke survivors with right-hemispheric lesion [40]. Education served as a proxy for cognitive reserve (i.e., the brain's resilience to neuropathological damage [41]), which is a very feasible variable to collect, and may be an important factor to consider or control for in motor rehabilitation trials. Notably, results of this study also highlight the complex interplay between cognitive function, cognitive reserve, and motor behavior.

What is the effect of stroke and cognitive impairment on gait and balance?

Gait and balance deficits are common post-stroke, and directly contribute to poor mobility, increased falls, and reduced quality of life [42, 43]. Given that a stroke can result in heterogeneous sensory and motor deficits such as muscle weakness, altered movement selection, spasticity, and altered sensation and integration of proprioceptive signals, the severity of balance and/or gait changes observed post-stroke is largely variable among individuals. In general though, slower ambulation, increased variability [44] and asymmetry [45], shorter, wider steps [46], and large anterior-posterior and lateral deviations of the trunk and pelvis [47], are common gait impairments following stroke. Like gait, impairments in balance are similarly broad in scope and may include asymmetric and increased sway [48, 49], poor weight transfer [50], a reduced limit of stability [51], and poor reactive postural control (i.e., the ability to quickly react to imbalance) [52]. As proprioceptive capacity is typically diminished following stroke, many stroke survivors may also become more reliant on visual information to maintain appropriate postural control [48].

Nearly 60% of older adults with cognitive impairment experience at least one fall annually [53], more than twofold that of their cognitively intact peers [54]. Indeed, the link between cognition and motor behavior has been well-established, but to what extent do stroke-related cognitive impairments affect gait and balance? Recent research has shown that the Stroop Color Word Test (a measure of inhibition of cognitive interference) and errors made on part B of the Trail Making Test, (a measure of attention switching) predict fall risk in stroke survivors [55]. Further, when compared to age-matched controls, stroke survivors tend to have the highest levels of cognitive-motor interference (i.e., the relative cost of dual-tasking) when performing concurrent working memory and balance tasks [56]; results indicated that working memory, but not semantic memory, had a disproportionately negative impact on cognitive-motor interference for the stroke group compared to controls. Interestingly, the stroke survivors had similar cognitive scores (i.e., score >10 on the Short Orientation–Memory–Concentration Test of Cognitive Impairment [57]) as age-matched controls, suggesting that individuals with stroke may require greater attentional resources due to their motor impairment(s), to perform as well as their age-matched counterparts. Collectively, these findings suggest that working memory, attention switching, and inhibition may be the most pertinent cognitive domains for proper balance control, and encourages future work to discern if the presence (or absence) of selective cognitive impairments following stroke may, in part, explain inter-individual differences in motor behavior (e.g., balance and gait).

Does cognitive impairment interfere with motor rehabilitation?

Given that an estimated 40–50% of all physical rehabilitation patients currently receiving care in the United States are over age 65 [58], it is imperative that today's therapies are effective for older adults [59]. And while it is difficult to precisely quantify the amount of elderly stroke survivors seeking some form of motor rehabilitation, the Centralized Open-Access Rehabilitation Database for Stroke (SCOAR) [22] approximates that ~33% of motor rehabilitation trials for stroke have an average participant age of 65 or older. Carefully chosen rehabilitation interventions can improve mobility, even in older stroke survivors [60, 61]. However, improvements through rehabilitation can be variable, with some survivors improving more than others or at different rates. Thus, an important factor to consider is the

likelihood that many older patients may present with cognitive impairments. Rigorous assessment of cognitive capacity may improve rehabilitative care for at least two reasons. First, it can instruct the patient interaction, including the types and modalities of instructions given to the patient. Second, cognitive impairments may interfere with their ability to learn or regain motor skills after stroke or neurological injury. For example, as a proof-of-concept, we recently demonstrated that regardless of primary diagnosis, overall cognitive status affected the extent to which transitional care patients improved their gait speed over their length of stay [62] (see also [63]). This trend persisted in a stroke-specific sample as well over one-year post-stroke [64]. Importantly, this does not suggest that cognitively impaired individuals should not participate in motor therapy, given they may still experience significant gains [65–67]. Instead, it advocates for 1) developing more personalized or targeted physical therapeutic interventions that are effective in cases of specific cognitive impairments post-stroke (e.g., [68]), and 2) conducting additional research that investigates which post-stroke cognitive impairments interfere most with motor skill learning. However, the field of stroke rehabilitation has only recently begun to investigate the impact of cognitive deficits on therapeutic responsiveness. In a recent meta-analysis of 215 stroke rehabilitation randomized controlled clinical trials (RCTs) from SCOAR, only 31% of the studies reported collecting cognitive information from participants (as measured by the MMSE), and nearly half of those studies used this information to *exclude* participants with cognitive deficits [22]. Overall, the use of cognitive assessments is encouraging, as it indicates that cognitive data are being collected in stroke motor rehabilitation, and could theoretically be used in retrospective, secondary analyses of clinical trial data. For example, Dobkin et al. [69] conducted secondary analyses on the Locomotor Experience Applied Post Stroke (LEAPS) RCT [70] and reported that attentional switching (measured as the difference in performance on Trails A and B) at baseline predicted participants' change in gait speed in response to a partial bodyweight-supported intervention involving both treadmill and over-ground walking. These analyses occurred retrospectively, after the initial LEAPS trial reported equivalent walking outcomes for both the treadmill and over-ground walking intervention and home-based exercise (that did not emphasize walking). However, given that the most common cognitive assessments reported among stroke rehabilitation RCTs are global cognitive screens, there remains opportunity for scientific inquiry regarding which, when, and to what extent specific cognitive deficits affect the motor rehabilitative process.

Are there specific cognitive impairments that can affect motor skill learning after stroke?

As summarized above, considerable work has demonstrated the relationship between cognition and gait/posture performance. However, an equally pertinent question for clinicians is whether cognitive factors affect *responsiveness* to gait training (which is driven by mechanisms of motor learning) are less understood. Although there remains a relative dearth of information on this topic, several recent studies have begun to provide insight into this knowledge gap. For example, evidence from a small (6-study) meta-analysis by Mullick et al. [71] suggests that both executive function and attention deficits after stroke can affect the amount of improvement in upper-extremity motor function following training, although there may be stronger cognitive-motor learning associations when kinematic outcomes (i.e., peak velocity or endpoint accuracy [72]) are used rather than clinical scales (e.g., the Action

Research Arm Test or the Wolf Motor Function Test [73]). Visuospatial impairments also influence how much motor task performance improves following upper extremity task-specific training [6–8, 74], as well as the amount and rate of functional improvement (measured with the Functional Independence Measure motor subscale) [31]. Provocative findings from Schweighofer and colleagues [10, 75] also implicate visuospatial working memory in upper extremity motor learning after stroke, particularly the role of contextual interference. Moreover, the post-stroke integrity of functional networks that are critical for visuospatial function (namely, frontoparietal) can predict how responsive individuals will be to upper-extremity motor training [76].

However, it is likely that the cognitive processes underlying the learning of upper vs. lower extremity movements are distinct. For example, while evidence supports the impact of visuospatial deficits on upper-extremity motor learning, such deficits may be less detrimental to gait and/or postural training. At present, there are limited studies that investigate cognitive factors related to lower- vs. upper-extremity motor learning, and no studies that have systematically compared across cognitive domains and body effectors. Moreover, it is plausible that different types of motor learning (implicit vs. explicit, [77]) are more reliant on different cognitive domains, which suggests the cognitive impairments that interfere with learning discrete upper extremity skills (like reaching, grasping, or object manipulation) would likely differ from those that interfere with adaptations of gait and posture.

McDowd et al. [78] suggests that attention (divided and switching, specifically) may be the most critical for determining the amount of improvement made during gait training in stroke. This has also led to an important area of research regarding whether engaging in concurrent cognitive tasks during gait training is more efficacious than simply walking (see [79]). For example, gait training while solving a problem using visual feedback has been shown to improve both gait and some aspects of cognition (namely, backwards visual digit span) but not others (auditory digit span) [80]. Such findings do not, however, directly address the question of whether attentional deficits post-stroke result in poorer gait relearning or slower recovery of balance, per se, although there is evidence of this in upper-extremity recovery, (see [81] for review). If so, therapists could use different strategies, such as internal or external loci of attention [82], to enhance gait and balance rehabilitation via an attention mechanism/intervention.

Can cognitive rehabilitation improve motor rehabilitation after stroke?

The correlative relationship observed between cognition and balance suggests it is plausible that cognitive training (or, motor rehabilitation paired with a cognitive task) could enhance motor performance improvement (i.e., relearning). However, while early evidence in healthy adults suggests cognitive training may improve some aspects of mobility [83], evidence in stroke survivors is limited and mixed. For example, Helm and colleagues [84] had two groups of stroke survivors perform locomotor training with either constant or variable practice structure, where variable practice requires greater attentional demands compared to constant practice; results indicated there was no difference in performance or retention of the locomotor task between either group. Whereas, Liu and colleagues [85] evaluated if various

dual-task training (i.e., motor-motor: tray-carrying and walking, or cognitive-motor: serial subtraction and walking) would affect different gait parameters in individuals with stroke; results indicated that each dual-task had differential effects on stride length and dual-task cost, suggesting that specific dual-task training may address selective cognitive deficits. At present, no studies have reported the effect of specialized training to target select cognitive deficits and its subsequent effect on gait in a neuropathological population (i.e., stroke). Future work is necessary to better understand if cognitive training impacts motor performance and learning.

Conclusions

Cognition is important for motor rehabilitation, particularly domains shown to underlie procedural learning such as attention, working memory, visuospatial abilities. Global cognitive testing is insensitive to subtle deficits in cognition and may only narrowly establish if an individual is cognitively intact. Neuropsychological tests that thoroughly evaluate the function of select cognitive domains may provide critical prognostic insight into rehabilitation outcomes. For example, recent work suggests that visuospatial deficits are associated with poorer upper-extremity motor recovery. Further, other studies have linked stroke-specific cognitive impairments not only with altered gait and balance performance, but with the degree to which patients improve gait with training as well. Stroke-related research studies should consider incorporating comprehensive neuropsychological testing to further our understanding of cognition's role in recovery and relearning of motor skills.

References

Papers of particular interest, published recently, have been highlighted as:

- Of importance
 - Of major importance
1. Bastian AJ (2008) Understanding sensorimotor adaptation and learning for rehabilitation. *Curr Opin Neurol* 21:628–633 [PubMed: 18989103]
 2. Song J-H (2019) The role of attention in motor control and learning. *Curr Opin Psychol* 29:261–265 [PubMed: 31491612]
 3. Buszard T, Farrow D, Verswijveren SJJM, Reid M, Williams J, Polman R, Ling FCM, Masters RSW (2017) Working Memory Capacity Limits Motor Learning When Implementing Multiple Instructions. *Front Psychol* 8:1350 [PubMed: 28878701]
 4. Bo J, Seidler RD (2009) Visuospatial working memory capacity predicts the organization of acquired explicit motor sequences. *J Neurophysiol* 101:3116 [PubMed: 19357338]
 5. Langan J, Seidler RD (2011) Age differences in spatial working memory contributions to visuomotor adaptation and transfer. *Behav Brain Res* 225:160–168 [PubMed: 21784106]
 6. Schaefer SY, Duff K (2017) Within-session and one-week practice effects on a motor task in amnesic mild cognitive impairment. *J Clin Exp Neuropsychol* 39:473–484 [PubMed: 27690745]
 7. Lingo VanGilder J, Hengge CR, Duff K, Schaefer SY (2018) Visuospatial function predicts one-week motor skill retention in cognitively intact older adults. *Neurosci Lett* 664:139–143 [PubMed: 29154858]
 8. Lingo VanGilder J, Walter CS, Hengge CR, Schaefer SY (2019) Exploring the relationship between visuospatial function and age-related deficits in motor skill transfer. *Aging Clin Exp Res*. 10.1007/s40520-019-01345-w

9. Raw RK, Wilkie RM, Allen RJ, Warburton M, Leonetti M, Williams JHG, Mon-Williams M (2019) Skill acquisition as a function of age, hand and task difficulty: Interactions between cognition and action. *PLoS One* 14:e0211706
10. Schweighofer N, Lee J-Y, Goh H-T, Choi Y, Kim SS, Stewart JC, Lewthwaite R, Winstein CJ (2011) Mechanisms of the contextual interference effect in individuals poststroke. *J Neurophysiol* 106:2632 [PubMed: 21832031]
11. Corbetta M, Ramsey L, Callejas A, et al. (2015) Common behavioral clusters and subcortical anatomy in stroke. *Neuron* 85:927–941 [PubMed: 25741721]
12. Teasell R, Salter K, Faltynek P, Cotoi A, Eskes G (2018) Post-Stroke Cognitive Disorders. In: Evidence-Based Rev. Stroke Rehabil., 18th ed. London, Ontario, pp 1–86.* An evidenced-based review of stroke rehabilitation that emphasizes recent reports of post-stroke cognitive impairments and their (white matter) neural correlates.
13. Geschwind N (1965) Disconnexion syndromes in animals and man. I. *Brain* 88:237–294 [PubMed: 5318481]
14. Nys GMS, van Zandvoort MJE, de Kort PLM, Jansen BPW, de Haan EHF, Kappelle LJ (2007) Cognitive disorders in acute stroke: prevalence and clinical determinants. *Cerebrovasc Dis* 23:408–416 [PubMed: 17406110]
15. Tatemichi TK, Desmond DW, Stern Y, Paik M, Sano M, Bagiella E (1994) Cognitive impairment after stroke: frequency, patterns, and relationship to functional abilities. *J Neurol Neurosurg Psychiatry* 57:202–7 [PubMed: 8126506]
16. Al-Qazzaz NK, Ali SH, Ahmad SA, Islam S, Mohamad K (2014) Cognitive impairment and memory dysfunction after a stroke diagnosis: a post-stroke memory assessment. *Neuropsychiatr Dis Treat* 10:1677–1691 [PubMed: 25228808]
17. Jokinen H, Melkas S, Ylikoski R, Pohjasvaara T, Kaste M, Erkinjuntti T, Hietanen M (2015) Post-stroke cognitive impairment is common even after successful clinical recovery. *Eur J Neurol* 22:1288–1294 [PubMed: 26040251]
18. Folstein MF, Folstein SE, McHugh PR (1975) “Mini-Mental State”. A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 12:189–198 19. [PubMed: 1202204]
19. Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, Cummings JL, Chertkow H (2005) The Montreal Cognitive Assessment, MoCA: A Brief Screening Tool For Mild Cognitive Impairment. *J Am Geriatr Soc* 53:695–699 [PubMed: 15817019]
20. Moafmashhadi P, Koski L (2013) Limitations for interpreting failure on individual subtests of the Montreal Cognitive Assessment. *J Geriatr Psychiatry Neurol* 26:19–28 [PubMed: 23385364]
21. Malek-Ahmadi M, O’Connor K, Schofield S, Coon DW, Zamrini E (2018) Trajectory and variability characterization of the Montreal Cognitive Assessment in older adults. *Aging Clin Exp Res* 30:993–998 [PubMed: 29188578]
22. Lohse KR, Schaefer SY, Raikes AC, Boyd LA, Lang CE (2016) Asking New Questions with Old Data: The Centralized Open-Access Rehabilitation Database for Stroke. *Front Neurol* 7:153 [PubMed: 27703445]
23. Ortiz GA L. Sacco (2014) National Institutes of Health Stroke Scale (NIHSS). Wiley StatsRef Stat Ref Online. 10.1002/9781118445112.stat06823
24. Zhao X-J, Li Q-X, Liu T-J, et al. (2018) Predictive values of CSS and NIHSS in the prognosis of patients with acute cerebral infarction: A comparative analysis. *Medicine (Baltimore)* 97:e12419
25. Abzhandadze T, Reinholdsson M, Stibrant Sunnerhagen K (2020) NIHSS is not enough for cognitive screening in acute stroke: A cross-sectional, retrospective study. *Sci Rep* 10:534 [PubMed: 31953508]
26. Gottesman RF, Kleinman JT, Davis C, Heidler-Gary J, Newhart M, Hillis AE (2010) The NIHSS-plus: improving cognitive assessment with the NIHSS. *Behav Neurol* 22:11–15 [PubMed: 20543454]
27. Weintraub S, Dikmen SS, Heaton RK, et al. (2013) Cognition assessment using the NIH Toolbox. *Neurology* 80:S54–S64 [PubMed: 23479546]
28. . Tulsky DS, Holdnack JA, Cohen ML, Heaton RK, Carlozzi NE, Wong AWK, Boulton AJ, Heinemann AW (2017) Factor structure of the NIH Toolbox Cognition Battery in individuals with

acquired brain injury. *Rehabil Psychol* 62:435–442. [PubMed: 29265864] * Results of this study suggest the NIH Toolbox Cognition Battery may be validated for use in populations with acquired brain injury (i.e., stroke).

29. Carlozzi NE, Goodnight S, Casaletto KB, Goldsmith A, Heaton RK, Wong AWK, Baum CM, Gershon R, Heinemann AW, Tulskey DS (2017) Validation of the NIH Toolbox in Individuals with Neurologic Disorders. *Arch Clin Neuropsychol* 32:555–573 [PubMed: 28334392]
30. Babakhanyan I, Carlozzi NE, McKenna BS, Casaletto KB, Heinemann AW, Heaton RK (2019) National Institutes of Health Toolbox Emotion Battery: Application of Summary Scores to Adults With Spinal Cord Injury, Traumatic Brain Injury, and Stroke. *Arch Phys Med Rehabil* 100:1863–1871 [PubMed: 31153854]
31. Toglia J, Fitzgerald KA, O'Dell MW, Mastrogiovanni AR, Lin CD (2011) The Mini-Mental State Examination and Montreal Cognitive Assessment in Persons With Mild Subacute Stroke: Relationship to Functional Outcome. *Arch Phys Med Rehabil* 92:792–798 [PubMed: 21530727]
32. Bo J, Borza V, Seidler RD (2009) Age-related declines in visuospatial working memory correlate with deficits in explicit motor sequence learning. *J Neurophysiol* 102:2744–2754 33.
33. McDonald MW, Black SE, Copland DA, et al. (2019) Cognition in Stroke Rehabilitation and Recovery Research: Consensus-Based Core Recommendations From the Second Stroke Recovery and Rehabilitation Roundtable. *Neurorehabil Neural Repair* 33:943–950. [PubMed: 31660787] ** This paper highlights key recommendations for guiding future research and clinical care regarding cognition and stroke rehabilitation, and advocates for assessing specific cognitive domains rather than just global measures.
34. Diagnostic and Statistical Manual of Mental Disorders: DSM-5. (2013) 5th ed. American Psychiatric Association, Arlington, VA
35. Gluhm S, Goldstein J, Loc K, Colt A, Van Liew C, Corey-Bloom J (2013) Cognitive performance on the Mini-Mental State Examination and the Montreal Cognitive Assessment across the healthy adult lifespan. *Cogn Behav Neurol* 26:1–5 [PubMed: 23538566]
36. Randolph C, Tierney MC, Mohr E, Chase TN (1998) The Repeatable Battery for the Assessment of Neuropsychological Status (RBANS): Preliminary Clinical Validity. *J Clin Exp Neuropsychol* 20:310–319 [PubMed: 9845158]
37. Wechsler D (1955) Manual for the Wechsler Adult Intelligence Scale. Psychological Corp., Oxford, England
38. Osterrieth PA (1944) Le test de copie d'une figure complexe. *Arch Psychol* 30:206–356
39. Chen H, Pan X, Lau JKL, Bickerton W-L, Pradeep B, Taheri M, Humphreys G, Rotshtein P (2016) Lesion-symptom mapping of a complex figure copy task: A large-scale PCA study of the BCoS trial. *NeuroImage Clin* 11:622–634 [PubMed: 27182489]
40. Umarova RM, Sperber C, Kaller CP, Schmidt CSM, Urbach H, Kloppel S, Weiller C, Karnath H-O (2019) Cognitive reserve impacts on disability and cognitive deficits in acute stroke. *J Neurol* 266:2495–2504 [PubMed: 31254064]
41. Stern Y (2009) Cognitive reserve. *Neuropsychologia* 47:2015–2028 [PubMed: 19467352]
42. Mackintosh S, Goldie P, Hill K (2005) Falls incidence and factors associated with falling in older, community-dwelling, chronic stroke survivors (>1 year after stroke) and matched controls. *Aging Clin Exp Res* 17:74–81 [PubMed: 15977453]
43. Schmid AA, Van Puymbroeck M, Altenburger PA, Miller KK, Combs SA, Page SJ (2013) Balance is associated with quality of life in chronic stroke. *Top Stroke Rehabil* 20:340–346 [PubMed: 23893833]
44. Balasubramanian CK, Neptune RR, Kautz SA (2009) Variability in spatiotemporal step characteristics and its relationship to walking performance post-stroke. *Gait Posture* 29:408–414 [PubMed: 19056272]
45. Lewek MD, Braun CH, Wutzke C, Giuliani C (2018) The role of movement errors in modifying spatiotemporal gait asymmetry post stroke: A randomized controlled trial. *Clin Rehabil* 32:161–172 [PubMed: 28750549]
46. Stimpson KH, Heitkamp LN, Embry AE, Dean JC (2019) Post-stroke deficits in the step-by-step control of paretic step width. *Gait Posture* 70:136–140 [PubMed: 30856525]

47. Van Criekinge T, Saeys W, Halleman A, Velghe S, Viskens P-J, Vereeck L, De Hertogh W, Truijten S (2017) Trunk biomechanics during hemiplegic gait after stroke: A systematic review. *Gait Posture* 54:133–143. [PubMed: 28288334] * This systematic review highlights the impact of stroke-related changes in trunk biomechanics.
48. de Haart M, Geurts AC, Huidekoper SC, Fasotti L, van Limbeek J (2004) Recovery of standing balance in postacute stroke patients: A rehabilitation cohort study. *Arch Phys Med Rehabil* 85:886–895 [PubMed: 15179641]
49. Sackley CM (1991) Falls, sway, and symmetry of weight-bearing after stroke. *Int Disabil Stud* 13:1–4 [PubMed: 1917796]
50. Rogers MW, Hedman LD, Pai YC (1993) Kinetic analysis of dynamic transitions in stance support accompanying voluntary leg flexion movements in hemiparetic adults. *Arch Phys Med Rehabil* 74:19–25 [PubMed: 8420514]
51. Goldie PA, Matyas TA, Evans OM, Galea M, Bach TM (1996) Maximum voluntary weight-bearing by the affected and unaffected legs in standing following stroke. *Clin Biomech (Bristol, Avon)* 11:333–342
52. de Kam D, Roelofs JMB, Bruijnes AKBD, Geurts ACH, Weerdesteyn V (2017) The Next Step in Understanding Impaired Reactive Balance Control in People With Stroke: The Role of Defective Early Automatic Postural Responses. *Neurorehabil Neural Repair* 31:708–716. [PubMed: 28691582] * This paper reports the deficits in and importance of reactive postural control for stroke survivors.
53. van Dijk PT, Meulenberg OG, van de Sande HJ, Habbema JD (1993) Falls in dementia patients. *Gerontologist* 33:200–204 [PubMed: 8468012]
54. Burns E, Kakara R (2018) Deaths from falls among persons aged 65 years — United States, 2007–2016. *Morb Mortal Wkly Rep* 67:509–514
55. Saverino A, Waller D, Rantell K, Parry R, Moriarty A, Playford ED (2016) The Role of Cognitive Factors in Predicting Balance and Fall Risk in a Neuro-Rehabilitation Setting. *PLoS One* 11:e0153469
56. Bhatt T, Subramaniam S, Varghese R (2016) Examining interference of different cognitive tasks on voluntary balance control in aging and stroke. *Exp Brain Res* 234:2575–2584 [PubMed: 27302401]
57. Katzman R, Brown T, Fuld P, Peck A, Schechter R, Schimmel H (1983) Validation of a short Orientation-Memory-Concentration Test of cognitive impairment. *Am J Psychiatry* 140:734–739 [PubMed: 6846631]
58. Bell A (2015) 9 Physical Therapist Tips to Help You Age Well. In: *Am. Phys. Ther. Assoc* <https://www.moveforwardpt.com/Resources/Detail/9-physical-therapist-tips-to-help-you-agewell>. Accessed 22 Mar 2018
59. Gatchel RJ, Schultz IZ, Ray CT, Hanna M, Choi JY (2018) Functional Rehabilitation in Older Adults: Where Are We Now and Where Should We Be Going? In: *Handb. Rehabil. Older Adults*. Springer, Cham, pp 561–567
60. Arienti C, Lazzarini SG, Pollock A, Negrini S (2019) Rehabilitation interventions for improving balance following stroke: An overview of systematic reviews. *PLoS One* 14:e0219781. * The authors present a systematic review that outlines the impact of physical rehabilitation to improve balance following stroke.
61. Schroder J, Truijten S, Van Criekinge T, Saeys W (2019) Feasibility and effectiveness of repetitive gait training early after stroke: A systematic review and meta-analysis. *J Rehabil Med* 51:78–88 [PubMed: 30516821]
62. Schaefer SY, Sullivan JM, Peterson DS, Fauth EB (2019) Cognitive function at admission predicts amount of gait speed change in geriatric physical rehabilitation. *Ann Phys Rehabil Med*. 10.1016/j.rehab.2019.08.004
63. Friedman PJ, Baskett JJ, Richmond DE (1989) Cognitive impairment and its relationship to gait rehabilitation in the elderly. *N Z Med J* 102:603–606 [PubMed: 2594278]
64. Sagnier S, Renou P, Olindo S, Debruxelles S, Poli M, Rouanet F, Munsch F, Tourdias T, Sibon I (2017) Gait Change Is Associated with Cognitive Outcome after an Acute Ischemic Stroke. *Front Aging Neurosci* 9:153. [PubMed: 28572768] * By following over 200 stroke survivors for one

year after discharge, this study found that changes in global cognition were independently associated with 10 meter walk performance regardless of stroke severity.

65. Rabadi MH, Rabadi FM, Edelstein L, Peterson M (2008) Cognitively impaired stroke patients do benefit from admission to an acute rehabilitation unit. *Arch Phys Med Rehabil* 89:441–448 [PubMed: 18295621]
66. Poynter L, Kwan J, Sayer AA, Vassallo M (2011) Does cognitive impairment affect rehabilitation outcome? *J Am Geriatr Soc* 59:2108–2111 [PubMed: 22092047]
67. Vassallo M, Poynter L, Kwan J, Sharma JC, Allen SC (2016) A prospective observational study of outcomes from rehabilitation of elderly patients with moderate to severe cognitive impairment. *Clin Rehabil* 30:901–908. [PubMed: 27496699] ** This paper demonstrates that patients with moderate or severe cognitive impairment still benefit from physical rehabilitation, and that physical interventions may themselves have cognitive benefits. This paper can be used to refute the argument that if cognitive impairment interferes with motor rehabilitation, then should it even be administered? Further, it challenges the finding noted in [22] that patients with low MMSE scores should be excluded from physical intervention studies.
68. Fasoli SE, Adans-Dester CP (2019) A Paradigm Shift: Rehabilitation Robotics, Cognitive Skills Training, and Function After Stroke. *Front Neurol* 10:1088 [PubMed: 31681154]
69. Dobkin BHK, Nadeau SE, Behrman AL, Wu SS, Rose DK, Bowden M, Studenski S, Lu X, Duncan PW (2014) Prediction of responders for outcome measures of locomotor Experience Applied Post Stroke trial. *J Rehabil Res Dev* 51:39–50 [PubMed: 24805892]
70. Duncan PW, Sullivan KJ, Behrman AL, Azen SP, Wu SS, Nadeau SE, Dobkin BH, Rose DK, Tilson JK, Team TLI (2007) Protocol for the Locomotor Experience Applied Post-stroke (LEAPS) trial: a randomized controlled trial. *BMC Neurol* 7:39 [PubMed: 17996052]
71. Mullick AA, Subramanian SK, Levin MF (2015) Emerging evidence of the association between cognitive deficits and arm motor recovery after stroke: A meta-analysis. *Restor Neurol Neurosci* 33:389–403 [PubMed: 26410581]
72. Cirstea CM, Pfitter A, Levin MF (2006) Feedback and cognition in arm motor skill reacquisition after stroke. *Stroke* 37:1237–1242 [PubMed: 16601218]
73. Boe EW, Pedersen AD, Pedersen AR, Nielsen JF, Blicher JU (2014) Cognitive status does not predict motor gain from post stroke constraint-induced movement therapy. *NeuroRehabilitation* 34:201–207 [PubMed: 24284455]
74. Wang P, Infurna FJ, Schaefer SY (2019) Predicting Motor Skill Learning in Older Adults Using Visuospatial Performance. *J Mot Learn Dev* 1–14
75. Kim S, Oh Y, Schweighofer N (2015) Between-Trial Forgetting Due to Interference and Time in Motor Adaptation. *PLoS One* 10:e0142963
76. Zhou RJ, Hondori HM, Khademi M, et al. (2018) Predicting Gains With Visuospatial Training After Stroke Using an EEG Measure of Frontoparietal Circuit Function. *Front Neurol* 9:597 [PubMed: 30087653]
77. Schmidt R, Lee T, Winstein C, Wulf G, Zelaznik H (2018) Motor Control and Learning: A Behavioral Emphasis, 6th Edition (online access included). *ProtoView* 2018:1–552
78. McDowd JM, Fillion DL, Pohl PS, Richards LG, Stiers W (2003) Attentional Abilities and Functional Outcomes Following Stroke. *Journals Gerontol Ser B* 58:P45–P53
79. Silsupadol P, Shumway-Cook A, Lugade V, van Donkelaar P, Chou L-S, Mayr U, Woollacott MH (2009) Effects of single-task versus dual-task training on balance performance in older adults: A double-blind, randomized controlled trial. *Arch Phys Med Rehabil* 90:381–387 [PubMed: 19254600]
80. Chung S, Wang X, Fieremans E, Rath JF, Amorapanth P, Foo F-YA, Morton CJ, Novikov DS, Flanagan SR, Lui YW (2019) Altered Relationship between Working Memory and Brain Microstructure after Mild Traumatic Brain Injury. *Am J Neuroradiol*. 10.3174/ajnr.A6146
81. Doron N, Rand D (2019) Is Unilateral Spatial Neglect Associated With Motor Recovery of the Affected Upper Extremity Poststroke? A Systematic Review. *Neurorehabil Neural Repair* 33:179–187 [PubMed: 30784364]

82. Kal E, Prosee R, Winters M, van der Kamp J (2018) Does implicit motor learning lead to greater automatization of motor skills compared to explicit motor learning? A systematic review. *PLoS One* 13:e0203591
83. Marusic U, Verghese J, Mahoney JR (2018) Cognitive-Based Interventions to Improve Mobility: A Systematic Review and Meta-analysis. *J Am Med Dir Assoc* 19:484–491.e3 [PubMed: 29680203]
84. Helm EE, Pohlig RT, Kumar DS, Reisman DS (2019) Practice Structure and Locomotor Learning after Stroke. *J Neurol Phys Ther* 43:85–93 [PubMed: 30883495]
85. Liu YC, Yang YR, Tsai YA, Wang RY (2017) Cognitive and motor dual task gait training improve dual task gait performance after stroke - A randomized controlled pilot trial. *Sci Rep* 7:4070. [PubMed: 28642466] * This paper shows that cognitive-motor and motor-motor dual-task training can have differential effects on dual-task capability in chronic stroke survivors when compared to conventional physical therapy.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript