

Review

# Synthetic Applications of Aziridinium Ions

Jala Ranjith and Hyun-Joon Ha \*

Department of Chemistry, Hankuk University of Foreign Studies, Yongin 17035, Korea; ranjithreddyict@gmail.com

\* Correspondence: hjha@hufs.ac.kr; Tel.: +82-31-330-4369

**Abstract:** Nonactivated aziridine with an electron-donating group at the ring nitrogen should be activated to an aziridinium ion prior to being converted to cyclic and acyclic nitrogen-containing molecules. This review describes ways to generate aziridinium ions and their utilization for synthetic purposes. Specifically, the intra- and intermolecular formation of aziridinium ions with proper electrophiles are classified, and their regio- and stereoselective transformations with nucleophiles are described on the basis of recent developments.

**Keywords:** aziridine; nonactivated aziridine; aziridinium ion; ring-opening; synthesis; nitrogen-containing molecules; regio- and stereoselective

## 1. Introduction

Aziridines are three-membered cyclic organic heterocyclic compounds with one nitrogen atom in the ring. They are valuable and versatile due to the reactive three-membered ring [1–9]. Highly ring-strained aziridines, e.g., other three-membered ring compounds such as cyclopropane and oxirane, render various nitrogen-containing compounds through ring-opening reactions with nucleophiles [10–13]. However, their stability and reactivity depend on substituents at the ring nitrogen (whether they are electron-withdrawing or -donating). Aziridines are bifurcated into “activated” ones bearing electron-withdrawing substituents at the ring nitrogen and “nonactivated” ones with electron-donating substituents [14] (Figure 1).



**Citation:** Ranjith, J.; Ha, H.-J. Synthetic Applications of Aziridinium Ions. *Molecules* **2021**, *26*, 1774. <https://doi.org/10.3390/molecules26061774>

Academic Editor: Alexander F. Khlebnikov

Received: 2 March 2021


Accepted: 17 March 2021

Published: 22 March 2021

**Publisher’s Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.




**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).



**Figure 1.** Chemical structures of “activated” and “nonactivated” aziridines and electrostatic potential maps of their representative examples *N*-methylaziridine and *N*-acetylaziridine. EDG = electron-donating group; EWG = electron-withdrawing group.


Activated aziridines are quite reactive toward most nucleophiles, while nonactivated aziridines are inert unless they are activated as aziridinium ions or their equivalents by proper electrophiles including alkyl, acyl, trimethylsilyl, and Lewis acids [15,16]. When the simple aziridine without any substituent on the ring is protonated as an aziridinium ion, the ring strain is increased by 47 kJ/mol [17]. Most ring-opening reactions of nonactivated aziridines proceed with breakage of the bond between C and N, such as **A** or **B**, with the assistance of electrophiles after the formation of aziridinium ions. Experimental and theoretical studies have shown that nonactivated aziridine is possibly not serving as a 1,3-dipole, as shown in **C** and **D**, while activated aziridines can be utilized for a cyclization

reaction with cleavage of the C–C bond for dipolarophile [17–19]. This difference is due to bond energy differences between C–N and C–C of nonactivated aziridines (Scheme 1).



**Scheme 1.** Comparison of two different ring openings via C–N and C–C bond cleavages of nonactivated aziridines.

Thus, most reactions with nonactivated aziridines are carried out with the formation of aziridinium ion as a quaternary ammonium at first, which is created via the chemical bond between nucleophilic and basic ring nitrogen and an applied electrophile [15,16]. The aziridinium ion as an ionic intermediate is a well-known chemical species called “nitrogen mustard”, including the notorious chemical warfare agent VX [O-ethyl S-[2-(diisopropylamino) ethyl] methylphosphonothioate [20]. In most cases, this aziridinium ion intermediate has a classical ammonium ion character, albeit strained [21]. However, there are many ways to generate aziridinium ions with various electrophiles. Extensive calculation showed that reactivity differences were ordered as shown in Figure 2. Acyl aziridinium is the most active one, followed by alkoxyacetyl, trimethylsilyl, alkyl, and protonyl aziridinium ions. The Lewis acid-coordinated complex is the least active one [14].




**Figure 2.** Order of reactivities of aziridinium ions by different electrophiles.

There are several protocols to generate aziridinium ions from nonactivated aziridines, as shown in Figure 2. Direct addition of external electrophiles is the most convenient and easiest way for the preparation of an aziridinium ion as a quaternary ammonium salt. Electrophiles needed for correspondents shown in Figure 2 include acid halides, haloformates, trimethylsilyl halide, halomethane, proton, and Lewis acid (Equation (1), in Scheme 2). Aziridinium ions can also be generated upon expulsion of the leaving group at the  $\beta$ -position of amine of the acyclic starting substrate (Equation (2i), in Scheme 2) [18,19]. The same strategy to expulse the leaving group from the pendant of aziridine or 2-leaving group substituted methyl aza-cycle yielded a bicyclic aziridinium ion (Equation (2ii), in Scheme 2) [22,23]. The same bicyclic aziridinium ion can also be derived from removal of the leaving group at the  $\beta$ -site of cyclic starting substrates through aza-ring contraction (Equation (2iii), in Scheme 2) [24,25]. Lastly, aziridinium ylides schematized below (Equations (3i) and (3ii), in Scheme 2) can be created by addition of a ring-nitrogen to alkyne (Equation (3i), in Scheme 2) or diazo compound (Equation (3ii), in Scheme 2).



ring opening via “a” is kinetic while the ring opening via “b” is thermodynamic, yielding reaction products 3 and 4 as regioisomers, respectively (Scheme 3).



**Scheme 3.** Two different ring-opening reaction pathways of nonactivated aziridine: “a” (kinetic) and “b” (thermodynamic) with the formation of an aziridinium ion with electrophile.

The two different regiochemical pathways are bifurcated into kinetic and thermodynamic pathways with substituents attached to the aziridine ring [27,28]. When 2-substituted nonactivated aziridine is activated and then treated with various nucleophiles in the same medium, some nucleophiles such as bromine and iodine can lead the reaction into the thermodynamic pathway, while hydride and fluorine yield the kinetic product. In particular, when the hydride nucleophile is applied, only a kinetic product is afforded without formation of the regioisomer. The nucleophile “chlorine” behaves in between, showing that the kinetic product is formed first. It is then diminished with elapsed time to give the thermodynamic product from the equilibrated aziridinium ion intermediate [29]. This big regiochemical difference can also be observed between acetate and water, although both are oxygen nucleophiles that can attack aziridinium ions [22,23]. The reactivity is also influenced by the solvent as in the case of most nucleophilic substitution reactions. The medium effects are also quite big for ring-opening reactions of nonactivated aziridines due to different activation energies [15,16].


## 2. Synthetic Application of Aziridinium Ions

Many studies have reported synthetic applications of aziridines to provide a versatile entry to various nitrogen-containing molecules. In addition, recent advances in preparative enantiopure nonactivated aziridines warrant a streamlined synthesis of biologically important molecules, including natural products such as aza-sugars, alkaloids, and others [30–32]. However, a few problems need to be solved. Firstly, an efficient and general method is needed to prepare starting aziridines bearing diverse substitutes at three different sites, including N1 nitrogen and two carbons, C2 and/or C3 [33–36]. For a more efficient and diverse use of nonactivated aziridines, a better understanding of the regiochemical pathway is needed. It is known that the regiochemical pathway of nonactivated aziridine is more diverse than that of activated aziridine [1–9]. In this short review, we describe synthetic developments to build important nitrogen-containing cyclic and acyclic molecules according to a few reports published recently.

### 2.1. Aziridinium Ions by Addition of External Electrophiles


The most popular and general synthetic method to generate an aziridinium ion is by adding electrophiles to nonactivated aziridines, as shown in Eq 1 of Scheme 2. Representative examples are described below. When chiral (2*R*,1'*R*)-2-acyl-(1'-phenylethyl)aziridines (5) and chiral nonactivated aziridines were treated with various acid chlorides such as acetyl chloride, methoxymethyl formate, and oxalyl chloride, the corresponding *N*-acylaziridinium ion (6) intermediates were generated by acylation of the nucleophilic aziridine ring nitrogen. These *N*-acylaziridinium ion intermediates then reacted with

chloride released from the acid chloride, an electrophile, to give rise to ring-opened  $\beta$ -amino- $\beta$ -chlorocarbonyl compounds (7). When we carried out the reaction in  $\text{CHCl}_3$ , chlorinated compounds were observed [37]. However, under most reaction media including  $\text{CH}_3\text{CN}$ , subsequent displacement of chloride with an internal oxygen nucleophile from methylchloroformate, acetyl chloride, and methyl chlorooxoacetate proceeded, yielding oxazolidin-2-ones (8),  $\beta$ -amino- $\alpha$ -acetyloxypropionates (9), and morpholin-2,3-diones (10), respectively (Scheme 4) [38].



**Scheme 4.** Reactions of 2-acylaziridines with acid chlorides, specifically including methyl chloroformate, acetyl chloride, and methyl chlorosuccinate.

Another useful external electrophile applicable to nonactivated aziridines is trimethylsilyl iodide with the formation of an aziridinium ion attached to the ring nitrogen as shown in Scheme 5. When enantiopure (1'-phenylethyl)aziridines (11) were treated with TMSI, the aziridinium ion (12) was formed. Subsequent ring opening by the released iodide gave rise to an iodinated product (13), whose iodine was replaced by amine to afford enantiopure diamines (14) (Scheme 5) [39,40].




**Scheme 5.** Formation of the trimethylsilyl (TMS) aziridinium ion with nonactivated aziridine and its subsequent ring opening by iodine and amine for the synthesis of 1,2-diamines.

This method is quite good to introduce a mild nucleophile that is not reactive enough to break the aziridine ring. In addition, TMS used for the activation of aziridine through an aziridinium ion is removed after addition of a nucleophile to replace iodine when the reaction proceeds. This method has been used for the synthesis of many biologically important molecules in our lab [40].

As shown in Scheme 1, an aziridinium ion for the activation of nonactivated aziridine is generated by adding an alkyl group to aziridine nitrogen with the formation of quaternary amines. However, there is not an efficient method to introduce an alkyl group without breakage of the highly ring-strained aziridine ring. In addition, this aziridinium ion as a quaternary amine formed by an external alkyl group should be inert so that it does not react with the counter anion of electrophiles after alkylation of the amine, i.e., the counter anion of the ammonium ion should not have the nucleophilicity to break down three-membered ammonium ions. We successfully achieved formation of a methylated aziridinium ion (16) by treating starting aziridine (15) with methyl triflate, taking advantage



bromine was released and reacted to give a brominated product. Its regiochemical outcome was derived from the thermodynamic pathway. When the same substrate was treated with HBr, a protonated aziridinium ion was generated, and subsequent ring opening occurred at C3 without any substituent, assuming that the regiochemical pathway was dominated by the kinetic coordinate. These two contrasting regiochemical pathways stemmed from different aziridinium ions by the addition of a benzyl group and a proton as electrophiles to the same aziridine (Scheme 7) [46]. This distinctive regiochemistry taught us that it would be possible to control reaction pathways according to the formation mechanism of the aziridinium ion [47,48].



**Scheme 7.** Ring-opening reactions of *N*-benzylaziridine as a nonactivated aziridine by BnBr and HBr with formation of the aziridinium ions (20 and 22) and their products (21 and 23).


Many studies have reported the regioselectivity of ring-opening reactions of nonactivated aziridinium ions via pathway “a” (nonsubstituted carbon C3, kinetically favored) or “b” (substituted carbon C2, thermodynamically favored) depending on chemical aspects. When the aziridine ring has an allylic or benzylic C2–N1 bond as in 2-vinyl or 2-phenylaziridines, ring-opening reactions proceed at the C2 position (pathway “b”) regardless of electrophiles or nucleophiles. Ring opening with 2-acylaziridines takes pathway “b” with a few exceptions of electrophiles and nucleophiles applied [47,48]. With 2-alkylaziridines, their reaction pathways are diversified depending on the nature of electrophiles and nucleophiles. However, the reaction would warrant a kinetically controlled ring-opened product at C3 (pathway “a”). According to experimental and theoretical data, including ours, a general overview is provided in Table 1 as a practical guide to predict the regiochemical pathway of nucleophilic nonactivated aziridines using different substrates, electrophiles, and nucleophiles. Bearing this in mind, it might also be possible to predict a regioselective preference for unexamined ring-opening reactions of nonactivated 2-substituted aziridines [49].

In the literature, there are ample examples for the activation for ring-opening reactions using a Lewis acid, one of which is shown in Scheme 8. This scheme showed that an azide nucleophile could drive the aziridine ring (24) with assistance of Lewis acid AlCl<sub>3</sub> in aqueous medium to yield  $\alpha$ -azido- $\beta$ -aminopropionate (25) [50], which was utilized for the synthesis of natural product biemamide (B) (26) [51].

**Table 1.** Regioselectivity in ring opening of nonactivated 2-substituted aziridines either at C2 (pathway “b”) with a substituent or at C3 (pathway “a”) without any substituent depending on the substrate, the electrophile, and the nucleophile.

$R^1$	$E^+$	Lewis Acid	$RCO^+$	$H^+$	$R^+$	$TMS^+$
$-CH=CH_2$ , $-CH=CH-COOEt$		C2 <sup>c</sup>	C2	C2 <sup>c</sup>	C2	C2
Aryl		C2	C2	C2	C2 <sup>c</sup>	C2 <sup>c</sup>
COR, COOR, CONH <sub>2</sub>		C3	C2	C2	C2	C2
Alkyl		C3	C3 and/or C2 <sup>a</sup>	C3	C3 <sup>b</sup>	C3

<sup>a</sup> Depending on the nucleophile (e.g., azide attacks at C2; alcohol attacks at C3). <sup>b</sup> Among halides, only the fluoride ion attacks the C3 position to a major extent. <sup>c</sup> Proposed regioselectivity (no experimental data available).



**Scheme 8.** Azide-induced ring opening of enantiopure *N*-phenylethylaziridine-2-carboxylate and its use for the synthesis of biemamide (B).


## 2.2. Aziridinium Ions upon Expulsion of the Leaving Group

For the last few years, Cossy's group has extensively studied the synthesis of various compounds from the formation of aziridinium ions, some of which were reported in a review article [52,53]. A typical method for aziridinium ions comes from expulsion of the leaving group of hydroxides at  $\beta$ -hydroxy amines [54]. One big advantage of  $\beta$ -hydroxyamine is that its enantiopure form is readily available from a rich chiral pool of amino acids. Its stereoselective transformation can realize diverse enantiopure products starting from diverse chiral amino acids [55]. Transforming the alcohol moiety into a good leaving group has allowed the rearrangement of these  $\beta$ -amino alcohols (27) to yield the aziridinium ion (28), which can then readily react with a large number of nucleophiles to afford ring-opened products (29). An overview of recent progress realized for the rearrangement of these  $\beta$ -amino alcohols in the presence of catalytic amount  $(CF_3CO)_2O$  and  $H_2SO_4$  or *N,N*-diethylaminosulfur trifluoride (DAST) has been reported [56]. This method has been applied for the synthesis of drug candidate LY-503430 (30) for Parkinson's disease (Scheme 9) [57].


A similar reaction with the fluoride nucleophile yielded a fluorinated product after the formation of an aziridinium ion driven by DAST [58]. The synthesis of various optically active  $\alpha$ -trifluoromethyl amines (33) was realized from  $\beta$ -amino- $\alpha$ -trifluoromethylalcohols (31) via an aziridinium ion intermediate (32) under a kinetic condition (Scheme 10) [59].

Recently, a report revealed that the same strategy is applicable to realize stereoselective and regioselective synthesis of  $\alpha$ -amino- $\beta$ -fluorophosphonates [60].






**Scheme 9.** Formation of aziridinium ion from  $\beta$ -hydroxyamine followed by nucleophilic ring opening in the presence of cat trifluoroacetate.



**Scheme 10.** Preparation of  $\alpha$ -trifluoromethyl amines or  $\alpha$ -amino- $\beta$ -fluorophosphonates bearing extra fluorine at the  $\alpha$ -position from the kinetically favored fluoride attack of an aziridinium ion intermediate formed from  $\beta$ -amino- $\alpha$ -trifluoromethyl alcohols.


Most aforementioned cases for the formation of an aziridinium ion took ionic pathways. Recently, photo-induced single-electron transfer for olefin-diamination with alkylamines was successful (Scheme 11) [61]. A useful protocol to introduce two different amines at each site of olefin with an aziridinium ion as an intermediate was introduced [61].




**Scheme 11.** Photo-induced single-electron transfer for olefin-diamination with alkylamines.

The synthetic protocol consists of the formation of an aziridinium ion (**36**) from an amine (**34**) and olefin (**35**) via in situ activation with *N*-chlorosuccinimide (NCS), protonation, and photo-induced SET (single-electron transfer) reduction in the presence of a Brønsted acid and  $\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$  (1 mol%) as the photocatalyst in  $\text{CH}_2\text{Cl}_2$  solvent under






**Scheme 13.** Formation of bicyclic aziridinium ion from aza-ring by removing the hydroxy group of hydroxymethyl at the  $\alpha$ -position and the subsequent ring opening by a nucleophile either at the suitably disposed bridgehead (*a*) or at the bridge position (*b*) to lead products.



**Figure 5.** Representative examples of Scheme 13 following pathway “*a*” or “*b*” to prepare various aza-cyclic valuables.


This reaction, generation, and ring opening of a bisaziridinium ion (52) from 3-hydroxy-3-trifluoromethyl piperidine (51) with removal of the hydroxy group could also be applicable for the synthesis of biologically important  $\alpha$ -trifluoromethyl pyrrolidines (53) (Scheme 14) [73,74].




**Scheme 14.** Preparation of 2-substituted methyl 2-(trifluoromethyl)pyrrolidines from ring contraction via bisaziridinium ion with removal of the hydroxy group of 3-hydroxy-3-trifluoromethylpiperidine.

### 2.5. Aziridinium Ylides

Recently, aziridinium ylides were developed by the addition of electrons at the aziridine nitrogen in the starting substrates to carbene or alkyne, followed by subsequent addition of an anion located at the  $\alpha$ -position of the carboxylate to the olefin to afford new rings via intramolecular ring-opening pathways. The schemes below (Schemes 15 and 16) show the formation of aziridinium ylides (55 and 59) via intramolecular or intermolecular addition of diazoacetate (54 or 58) to aziridine (54 or 57) with metal catalysts. Subsequent ring openings initiated by the anion at the  $\alpha$ -position of carboxylate give ring-expanded products (56) and (60) [75–77].




**Scheme 15.** Formation and ring expansion of aziridinium ion generated from intramolecular Cu(II)-catalyzed aziridinium ylides.



**Scheme 16.** Formation and ring expansion of aziridinium ion generated from intermolecular Rh-catalyzed aziridinium ylides.

The addition of carbene generated by Rh in an intermolecular manner was reported by Rowland with limited examples [78]. This has been developed to a great extent by Schomaker's group, whose representative example is shown in Scheme 17 [79]. The formation of aziridinium ylide (63) was created by adding carbene from diazoacetate (62) to aziridine (61), whose aziridine ring was opened to give an expanded aza-ring (64).



**Scheme 17.** Ring expansion of aziridines via aziridinium ylides to dehydropiperidines.

Another interesting formation of aziridinium ylide (67) was reported by Yudin's lab using the reaction of aryne from 1-iodo-2-sulfonyloxybenzene (66) with 2-alkenylaziridine (65). This was followed by anionic rearrangement to produce benzazepine (68) with a good yield (Scheme 18) [80].

It has been disclosed that the addition of electron-rich nitrogen at the nonactivated aziridine (69) to alkyne (70) can make an aziridinium ylide (71) whose anionic olefin can attack the vinyl group, thus giving rise to various substituted benzazepines (72) (Scheme 19) [81].



### 3. Conclusions

Nonactivated aziridines are valuable starting materials for the synthesis of densely functionalized nitrogen-containing compounds in a highly regio- and stereocontrolled manner. Their chemical diversity stems from the intrinsic properties of nonactivated aziridines to be activated with electrophiles including acyl halide, TMS halide, alkylhalide, sulfonyloxyalkanes, protic acid, and a Lewis acid. In addition to the backbone of starting substrates and applicable nucleophiles, the chemical reactivity and regiochemical pathways depend highly on the electronic characteristics of aziridinium ions. In this review, we summarized the ways of generating and classifying various types of aziridinium ions for synthetic purposes. The hope is that readers find the value of nonactivated aziridines and their synthetic utilities.

**Author Contributions:** Investigation and collection of references, J.R. and H.-J.H.; Writing—Original draft preparation, J.R. and H.-J.H.; writing and editing, J.R. and H.-J.H.; supervision, H.-J.H.; funding acquisition, H.-J.H. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was supported by the National Research Foundation of Korea (NRF-2012M3A7B4049645 and 2021R1A5A6002803) with the Center for New Directions in Organic Synthesis. It was also supported by an NRF grant (2020R1A2C1007102) and an HUFs Grant 2021 (for the preparation of this manuscript).

**Conflicts of Interest:** The authors declare no conflict of interest.

### References

1. Lindstrom, U.M.; Somfai, P. Aminolysis of vinyl epoxides as an efficient entry to N-H vinylaziridines. *Synthesis* **1998**, 109–117. [[CrossRef](#)]
2. Zwanenburg, B.; ten Holte, P. The synthetic potential of three-membered ring aza-heterocycles. *Top. Curr. Chem.* **2001**, *216*, 93–124.
3. Sweeney, J.B. Aziridines: Epoxides' ugly cousins? *Chem. Soc. Rev.* **2002**, *31*, 247–258. [[CrossRef](#)] [[PubMed](#)]
4. Hu, X.E. Nucleophilic ring opening of aziridines. *Tetrahedron* **2004**, *60*, 2701–2743. [[CrossRef](#)]
5. Tanner, D. Chiral aziridines—Their synthesis and use in stereoselective transformations. *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 599–619. [[CrossRef](#)]
6. McCoull, W.; Davis, F.A. Recent synthetic applications of chiral aziridines. *Synthesis* **2000**, 1347–1365. [[CrossRef](#)]
7. Watson, I.D.G.; Yu, L.; Yudin, A.K. Advances in nitrogen transfer reactions involving aziridines. *Acc. Chem. Res.* **2006**, *39*, 194–206. [[CrossRef](#)]
8. Padwa, A.; Murphree, S.S. Epoxides and aziridines—A mini review. *Arkivoc* **2006**, *3*, 6–33.
9. Singh, G.S.; D'hooghe, M.; De Kimpe, N. Synthesis and reactivity of C-heteroatom-substituted aziridines. *Chem. Rev.* **2007**, *107*, 2080–2135. [[CrossRef](#)]
10. Pitzer, K.S. Strain energies of cyclic hydrocarbons. *Science* **1945**, *101*, 672. [[CrossRef](#)]
11. Lewis, L.L.; Turner, L.L.; Salter, E.A.; Magers, D.H. Computation of the conventional strain energy in oxaziridine. *J. Mol. Struct. Theochem.* **2002**, *592*, 161–171. [[CrossRef](#)]
12. Dudev, T.; Lim, C. Ring strain energies from ab initio calculations. *J. Am. Chem. Soc.* **1998**, *120*, 4450–4458. [[CrossRef](#)]
13. Morgan, K.M.; Ellis, J.A.; Lee, J.; Fulton, A.; Wilson, S.L.; Dupart, P.S.; Dastoori, R. Thermochemical studies of epoxides and related compounds. *J. Org. Chem.* **2013**, *78*, 4303–4311. [[CrossRef](#)] [[PubMed](#)]
14. D'hooghe, M.; Ha, H.-J. *Synthesis of 4- to 7-Membered Heterocycles by Ring Expansion*; Springer: Berlin/Heidelberg, Germany, 2016.
15. Cherni, E.; Essalah, K.; Besbes, N.; Abderrabba, M.; Ayadi, S. Theoretical investigation of the regioselective ring opening of 2-methylaziridine. *Lewis acid effect. J. Mol. Modeling* **2018**, *24*, 309. [[CrossRef](#)] [[PubMed](#)]
16. Baruah, B.; Deuri, S.; Phukan, P. Reactivity and regioselectivity in the ring opening of 2-substituted non-activated aziridines: A density functional theory based analysis. *Comput. Theor. Chem.* **2014**, *1027*, 197–202. [[CrossRef](#)]
17. Paasche, A.; Arnone, M.; Fink, R.F.; Schirmeiste, T.; Engels, B. Origin of the reactivity differences of substituted aziridines: CN vs. CC bond breakages. *J. Org. Chem.* **2009**, *74*, 5244–5249. [[CrossRef](#)] [[PubMed](#)]
18. Cossy, J.; Pardo, D.G.; Dumas, C.; Mirguet, O.; Dechamps, I.; Metro, T.-X.; Burger, B.; Roudeau, R.; Appenzeller, J.; Cochi, A. Rearrangement of  $\beta$ -amino alcohols and application to the synthesis of biologically active compounds. *Chirality* **2009**, *21*, 850–856. [[CrossRef](#)]
19. Chen, Y.; Sun, X.; Wu, N.; Li, J.; Jin, S.; Zhong, Y.; Liu, Z.; Rogachev, A.; Chong, H.-S. Synthetic and theoretical investigation on the one-pot halogenation of  $\beta$ -amino alcohols and nucleophilic ring opening of aziridinium ions. *Org. Biomol. Chem.* **2016**, *14*, 920–939. [[CrossRef](#)]
20. Black, R. Development, Historical Use and Properties of Chemical Warfare Toxicology. In *Chemical Warfare Toxicology: Volume 1: Fundamental Aspects*; Worek, F., Jenner, J., Thiermann, H., Eds.; Royal Society of Chemistry: Cambridge, UK, 2016; Volume 1, pp. 1–28.

21. Lodewyk, M.W.; Tantillo, D.J. Nonclassical ammonium ions as intermediates in cinchona alkaloid rearrangement? *Chirality* **2020**, *32*, 484–488. [[CrossRef](#)]
22. Ji, M.-K.; Hertsen, D.; Yoon, D.-H.; Eum, H.; Goossens, H.; Waroquier, M.; Van Speybroeck, V.; D'hooghe, M.; De Kimpe, N.; Ha, H.-J. Nucleophile-dependent regio- and stereoselective ring opening of 1-azoniabicyclo [3.1.0]hexane tosylate. *Chem. Asian J.* **2014**, *9*, 1060–1067. [[CrossRef](#)]
23. Choi, J.; Yadav, N.N.; Ha, H.-J. Preparation of a stable bicyclic aziridinium ion and its ring expansion toward piperidines and azepanes. *Asian J. Org. Chem.* **2017**, *6*, 1292–1307. [[CrossRef](#)]
24. Dechamps, I.; Gomez Pardo, D.; Cossy, J. Synthesis of optically active substituted 3-fluoropiperidines from prolinols by using DAST. *Synlett* **2007**, 263–267. [[CrossRef](#)]
25. Dechamps, I.; Pardo, D.G.; Cossy, J. Ring expansion induced by DAST: Synthesis of substituted 3-fluoropiperidines from prolinols and 3-fluoroazepanes from 2-hydroxymethylpiperidines. *Eur. J. Org. Chem.* **2007**, 4224–4234. [[CrossRef](#)]
26. Silva, M.A.; Goodman, J.M. Aziridinium ring opening: A simple ionic reaction pathway with sequential transition states. *Tetrahedron Lett.* **2005**, *46*, 2067–2069. [[CrossRef](#)]
27. Cochi, A.; Pardo, D.G.; Cossy, J. Access to optically active 3-aminopiperidines by ring expansion of prolinols: Thermodynamic versus kinetic control. *Eur. J. Org. Chem.* **2012**, 2023–2040. [[CrossRef](#)]
28. Boydas, E.B.; Tanriver, G.; D'hooghe, M.; Ha, H.-J.; Van Speybroeck, V.; Catak, S. Theoretical insight into the regioselective ring-expansions of bicyclic aziridinium ions. *Org. Biomol. Chem.* **2018**, *16*, 796–806. [[CrossRef](#)] [[PubMed](#)]
29. D'hooghe, M.; Catak, S.; Stankovic, S.; Waroquier, M.; Kim, Y.; Ha, H.-J.; Van Speybroeck, V.; De Kimpe, N. Systematic study of halide-induced ring opening of 2-substituted aziridinium salts and theoretical rationalization of the reaction pathways. *Eur. J. Org. Chem.* **2010**, *25*, 4920–4931. [[CrossRef](#)]
30. Akhtar, R.; Naqvi, S.A.R.; Zahoor, A.F.; Saleem, S. Nucleophilic ring opening reactions of aziridines. *Mol. Divers.* **2018**, *22*, 447–501. [[CrossRef](#)]
31. Glowacka, I.E.; Trocha, A.; Wroblewski, A.E.; Piotrowska, D.G. *N*-(1-Phenylethyl)aziridine-2-carboxylate esters in the synthesis of biologically relevant compounds. *Beilstein J. Org. Chem.* **2019**, *15*, 1722–1757. [[CrossRef](#)] [[PubMed](#)]
32. Macha, L.; D'hooghe, M.; Ha, H.-J. Deployment of aziridines for the synthesis of alkaloids and their derivatives. *Synthesis* **2019**, *51*, 1491–1515.
33. Lee, W.-K.; Ha, H.-J. High light of the chemistry of enantiomerically pure aziridine-2-carboxylates. *Aldrichimica Acta* **2003**, *36*, 57–63.
34. Pellissier, H. Recent Developments in Asymmetric Aziridination. *Adv. Synth. Catal.* **2014**, *356*, 1899–1935. [[CrossRef](#)]
35. Pellissier, H.; Lattanzi, A.; Dalpozzo, R. Asymmetric Synthesis of Three-Membered Rings. Wiley-VCH: Weinheim, Germany, 2017; pp. 205–378.
36. Dogan, O.; Polat-Cakir, S. Recent progress in asymmetric synthesis of aziridine derivatives (microreview). *Chem. Heterocycl. Compd.* **2018**, *54*, 397–399. [[CrossRef](#)]
37. Park, C.S.; Kim, M.S.; Sim, T.B.; Pyun, D.K.; Lee, C.H.; Chang, J.-W.; Ha, H.-J. Novel stereoselective synthesis of functionalized oxazolidinone from chiral aziridine. *J. Org. Chem.* **2003**, *68*, 43–49. [[CrossRef](#)] [[PubMed](#)]
38. Kim, Y.; Ha, H.-J.; Yun, H.; Lee, B.K.; Lee, W.K. Ring opening of 2-acylaziridines by acid chlorides. *Tetrahedron* **2006**, *62*, 8844–8849. [[CrossRef](#)]
39. Lee, B.K.; Kim, M.S.; Hahm, H.S.; Kim, D.S.; Lee, W.K.; Ha, H.-J. An efficient synthesis of chiral terminal 1,2-diamines using an enantiomerically pure [1-(1'*R*)-methylbenzyl]aziridine-2-yl]methanol. *Tetrahedron* **2006**, *62*, 8393–8397. [[CrossRef](#)]
40. Shin, S.-H.; Han, E.Y.; Park, C.S.; Lee, W.K.; Ha, H.-J. Stereoselective synthesis of enantiomerically pure D-threo-PDMP; Manipulation of a core unit of 1,2-diaminoalcohol. *Tetrahedron* **2000**, *11*, 3293–3301. [[CrossRef](#)]
41. Kim, Y.; Ha, H.-J.; Yun, S.Y.; Lee, W.K. Preparation of stable aziridinium ions and their ring openings. *Chem. Commun.* **2008**, 36, 4363–4365. [[CrossRef](#)]
42. Kim, Y.; Yoon, D.-H.; Ha, H.-J.; Kang, K.Y.; Lee, W.K. *N*-Methylative aziridine ring opening and asymmetric synthesis of MeBmt. *Tetrahedron Lett.* **2011**, *52*, 5918–5920. [[CrossRef](#)]
43. Yoon, D.-H.; Kang, P.; Lee, W.K.; Kim, Y.; Ha, H.-J. Alkylative ring opening of *N*-methylaziridinium ions and a formal synthesis of tyroscherin. *Org. Lett.* **2012**, *14*, 429–431. [[CrossRef](#)]
44. Lee, J.; Lee, J.-E.; Ha, H.-J.; Son, S.I.; Lee, W.K. *N*-methylative aziridine ring opening: Asymmetric synthesis of hygroline, pseudo-hygroline and hygrine. *Tetrahedron Lett.* **2015**, *56*, 856–858. [[CrossRef](#)]
45. Jung, J.-H.; Kim, S.; Eum, H.; Lee, W.K.; Ha, H.-J. *N*-methylative aziridine ring opening and the synthesis of (S)-3-methylamino-3-[(R)-pyrrolidin-3-yl]propanenitrile. *Tetrahedron* **2017**, *73*, 5993–5999. [[CrossRef](#)]
46. Yun, S.Y.; Catak, S.; Lee, W.K.; D'hooghe, M.; De Kimpe, N.; Van Speybroeck, V.; Waroquier, M.; Kim, Y.; Ha, H.-J. Nucleophile-dependent regioselective ring opening of 2-substituted *N,N*-dibenzylaziridinium ions: Bromide versus hydride. *Chem. Commun.* **2009**, *18*, 2508–2510. [[CrossRef](#)] [[PubMed](#)]
47. Berger, G. Using conceptual density functional theory to rationalize regioselectivity: A case study on the nucleophilic ring-opening of activated aziridines. *Comp. Theor. Chem.* **2013**, *1010*, 11–18. [[CrossRef](#)]
48. Catak, S.; D'hooghe, M.; Verstraelen, T.; Hemelsoet, K.; Van Nieuwenhove, A.; Ha, H.-J.; Waroquier, M.; Van Speybroeck, V.; De Kimpe, N. Opposite regiospecific ring opening of 2-(Cyanomethyl)aziridines by hydrogen bromide and benzyl bromide: Experimental study and theoretical rationalization. *J. Org. Chem.* **2010**, *75*, 4530–4541. [[CrossRef](#)] [[PubMed](#)]

49. Stankovic, S.; D'hooghe, M.; Catak, S.; Eum, H.; Waroquier, M.; Van Speybroeck, V.; De Kimpe, N.; Ha, H.-J. Regioselectivity in the ring opening of non-activated aziridines. *Chem. Soc. Rev.* **2012**, *41*, 643–665. [[CrossRef](#)] [[PubMed](#)]
50. Kim, Y.; Ha, H.-J.; Han, K.; Ko, S.W.; Yun, H.; Yoon, H.J.; Kim, M.S.; Lee, W.K. Preparation of 2,3-diaminopropionate from ring opening of aziridine-2-carboxylate. *Tetrahedron Lett.* **2005**, *46*, 4407–4409. [[CrossRef](#)]
51. Srivastava, N.; Macha, L.; Ha, H.-J. Total synthesis and stereochemical revision of biemamides B and D. *Org. Lett.* **2019**, *21*, 8992–8996. [[CrossRef](#)]
52. Metro, T.-X.; Duthion, B.; Pardo, D.G.; Cossy, J. Rearrangement of  $\beta$ -amino alcohols via aziridiniums: A review. *J. Chem. Soc. Rev.* **2010**, *39*, 89–102. [[CrossRef](#)]
53. Pardo, D.G.; Cossy, J. Access to optically active 3-substituted piperidines by ring expansion of prolinols and derivatives. *Chem. Eur. J.* **2014**, *20*, 4516–4525. [[CrossRef](#)]
54. Metro, T.-X.; Pardo, D.G.; Cossy, J. Highly enantioselective synthesis of  $\alpha$ -amino alcohols: A catalytic version. *J. Org. Chem.* **2007**, *72*, 6556–6561. [[CrossRef](#)]
55. Metro, T.-X.; Pardo, D.G.; Cossy, J. Highly Enantioselective Synthesis of Linear  $\beta$ -Amino Alcohols. *Chem. Eur. J.* **2009**, *15*, 1064–1070. [[CrossRef](#)]
56. Metro, T.-X.; Pardo, D.G.; Cossy, J. Stereospecific rearrangement of  $\beta$ -amino alcohols catalyzed by  $H_2SO_4$ . *Synlett* **2007**, *4*, 2888–2890.
57. Duthion, B.; Pardo, D.G.; Cossy, J. Enantioselective synthesis of  $\beta$ -fluoroamines from  $\beta$ -amino alcohols: Application to the synthesis of LY503430. *Org. Lett.* **2010**, *12*, 4620–4623. [[CrossRef](#)]
58. Duthion, B.; Pardo, D.G.; Cossy, J. Synthesis of LY503430 by using a selective rearrangement of  $\beta$ -amino alcohols induced by DAST. *ARKIVOC* **2014**, *3*, 239–255. [[CrossRef](#)]
59. Neouchy, Z.; Pardo, D.G.; Cossy, J. Synthesis of optically active  $\alpha$ -trifluoromethylamines by rearrangement of  $\beta$ -amino- $\alpha$ -trifluoromethyl alcohols. *Org. Lett.* **2018**, *20*, 6017–6021. [[CrossRef](#)]
60. Kaźmierczak, M.; Dutkiewicz, G.; Cytłaka, T. Application of  $\alpha$ -amino- $\beta$ -fluorophosphonates in construction of their dipeptide analogues. *Synthesis* **2020**, *52*, 2364–2372. [[CrossRef](#)]
61. Govaerts, S.; Angelini, L.; Hampton, C.; Malet-Sanz, L.; Ruffoni, A.; Leonori, D. Photoinduced olefin diamination with alkylamines. *Angew. Chem. Int. Ed.* **2020**, *59*, 15021–15028. [[CrossRef](#)] [[PubMed](#)]
62. Dolfen, J.; Yadav, N.N.; De Kimpe, N.; D'hooghe, M.; Ha, H.-J. Bicyclic aziridinium ions in azaheterocyclic chemistry—Preparation and synthetic application of 1-azoniabicyclo[n.1.0]alkanes. *Adv. Synth. Catal.* **2016**, *358*, 3483–3511. [[CrossRef](#)]
63. Goossens, H.; Hertsen, D.; Mollet, K.; Catak, S.; D'hooghe, M.; De Proft, F.; Geerlings, P.; De Kimpe, N.; Waroquier, M.; Van Speybroeck, V. *Structure, Bonding and Reactivity of Heterocyclic Compounds, Topics in Heterocyclic Chemistry 38*; De Proft, F., Geerlings, P., Eds.; Springer: Berlin/Heidelberg, Germany, 2014.
64. Eum, H.; Choi, J.; Cho, C.-G.; Ha, H.-J. Regiochemistry-Directed Syntheses of Polyhydroxylated Alkaloids from Chiral Aziridines. *Asian J. Org. Chem.* **2015**, *4*, 1399–1409. [[CrossRef](#)]
65. Yadav, N.N.; Ha, H.-J. Preparation of stable bicyclic aziridinium ions and their ring-opening for the synthesis of azaheterocycles. *J. Vis. Exp.* **2018**, *138*, e57572. [[CrossRef](#)]
66. Yadav, N.N.; Lee, Y.G.; Srivastava, N.; Ha, H.-J. Alkylative ring-opening of bicyclic aziridinium ion and its application for alkaloid synthesis. *Front. Chem.* **2019**, *7*, 1–9. [[CrossRef](#)] [[PubMed](#)]
67. Riouton, S.; Pardo, D.G.; Cossy, J. Synthesis of substituted  $\alpha$ -trifluoromethyl piperidinic derivatives. *Molecules* **2017**, *22*, 483. [[CrossRef](#)] [[PubMed](#)]
68. Arico, F.; Aldoshin, A.S.; Musolino, M.; Crisma, M.; Tundo, P.  $\beta$ -Aminocarbonates in Regioselective and Ring Expansion Reactions. *J. Org. Chem.* **2018**, *83*, 236–243. [[CrossRef](#)] [[PubMed](#)]
69. Cochi, A.; Pardo, D.G.; Cossy, J. Access to optically active 3-azido- and 3-aminopiperidine derivatives by enantioselective ring expansion of prolinols. *Org. Lett.* **2011**, *13*, 4442–4445. [[CrossRef](#)]
70. Anxionnat, B.; Robert, B.; George, P.; Ricci, G.; Perrin, A.; Pardo, D.G.; Cossy, J. Ring expansion of cyclic  $\beta$ -amino alcohols induced by diethylaminosulfurtrifluoride: Synthesis of cyclic amines with a tertiary fluorine at C3. *J. Org. Chem.* **2012**, *77*, 6087–6099. [[CrossRef](#)]
71. Lam, Y.-h.; Houk, K.H.; Cossy, J.; Pardo, D.G.; Cochi, A. Fluorine as a Regiocontrol Element in the Ring Opening of Bicyclic Aziridiniums. *Helv. Chim. Acta* **2012**, *95*, 2265–2277. [[CrossRef](#)]
72. Orliac, A.; Routier, J.; Charvillon, F.B.; Sauer, W.H.B.; Bombrun, A.; Kulkarni, S.S.; Pardo, D.G.; Cossy, J. Enantioselective synthesis and physicochemical properties of libraries of 3-amino- and 3-amidofluoropiperidines. *Chem. Eur. J.* **2014**, *20*, 3813–3824. [[CrossRef](#)]
73. Xypolia, A.F.; Pardo, D.G.; Cossy, J. Ring contraction of 3-hydroxy-3-(trifluoromethyl)piperidines: Synthesis of 2-substituted 2-(Trifluoromethyl) pyrrolidines. *Chem. Eur. J.* **2015**, *21*, 12876–12880. [[CrossRef](#)]
74. Riouton, S.; Orliac, A.; Antoun, Z.; Bidault, R.; Pardo, D.G.; Cossy, J. Stereoselective rearrangement of (Trifluoromethyl)prolinols to enantioenriched 3-substituted 2-(Trifluoromethyl) piperidines. *Org. Lett.* **2015**, *17*, 2916–2919. [[CrossRef](#)]
75. Dequina, H.J.; Schomaker, J.M. Aziridinium ylides: Underused intermediates for complex amine synthesis. *Trends Chem.* **2020**, *2*, 874–887. [[CrossRef](#)] [[PubMed](#)]
76. Schmid, S.C.; Guzei, I.A.; Schomaker, J.M. A stereoselective [3 + 1] ring expansion for the synthesis of highly substituted methylene azetidines. *Angew. Chem. Int. Ed.* **2017**, *56*, 12229–12233. [[CrossRef](#)] [[PubMed](#)]



77. Schmid, S.C.; Guzei, I.A.; Fernandez, I.; Schomaker, J.M. A stereoselective [3 + 1] ring expansion of bicyclic methyleneaziridines via concerted, near barrierless [2,3]-stevens rearrangements of aziridinium ylides. *ACS Catal.* **2018**, *8*, 7907–7914. [[CrossRef](#)] [[PubMed](#)]
78. Barnes, W.K.; Rowlands, G.J. Studies on the [2,3]-Stevens rearrangement of aziridinium ions. *Tetrahedron Lett.* **2004**, *45*, 5347–5350.
79. Esho, J.; Nicastri, K.A.; Schmid, S.C.; Raskopf, W.T.; Guzei, I.A.; Fernández, I.; Schomaker, J.M. Intermolecular [3 + 3] ring expansion of aziridines to dehydropiperidines through the intermediacy of aziridinium ylides. *Nat. Comm.* **2020**, *11*, 1273. [[CrossRef](#)] [[PubMed](#)]
80. Kaldas, S.J.; Kran, E.; Meck-Lichtenfeld, C.; Yudin, A.K.; Studer, A. Reaction of vinyl aziridines with arynes: Synthesis of benzazepines and branched allyl fluorides. *Chem. Eur. J.* **2020**, *26*, 1501–1505. [[CrossRef](#)]
81. Deepak, S.; Ha, H.-J. Metal-free aza-Claisen type ring expansion of vinyl aziridine: An expeditious synthesis of seven Membered N-heterocycles. *Org. Biomol. Chem.* **2019**, *17*, 3093–3097.
82. Wang, T.; Daniliuc, C.G.; Muck-Lichtenfeld, C.; Kehr, G.; Erker, G. Formation of reactive  $\pi$ -conjugated frustrated N/B pairs by borane-induced propargyl amine rearrangement. *J. Am. Chem. Soc.* **2018**, *140*, 3635–3643. [[CrossRef](#)] [[PubMed](#)]