

HHS Public Access

Author manuscript JAMA. Author manuscript; available in PMC 2021 October 27.

Published in final edited form as:

JAMA. 2020 October 27; 324(16): 1609–1610. doi:10.1001/jama.2020.18941.

Nutrient warnings on unhealthy foods

Marissa G. Hall, PhD^{1,2}, Anna H. Grummon, PhD^{3,4}

¹.Department of Health Behavior, University of North Carolina Gillings School of Global Public Health, Chapel Hill, NC, USA

² Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

³ Harvard Center for Population and Development Studies, Harvard TH Chan School of Public Health, Cambridge, MA, USA

⁴ Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, MA, USA

Poor diet is one of the leading contributing factors for death in the US and worldwide.¹ Unhealthy diets, characterized by overconsumption of ultra-processed foods and sugary drinks, increase the risk of obesity, type 2 diabetes, and heart disease.¹ A nutrient warning policy is a common-sense response to rising rates of diet-related disease. The US should require prominent warning labels on the front of product packaging to alert consumers when products contain high levels of unhealthy nutrients.

Since 2016, five countries (Chile, Israel, Mexico, Peru, and Uruguay) have passed legislation requiring nutrient warnings, with the goal of addressing obesity and other diet-related chronic diseases. These policies focus on products with excessive levels of unhealthy nutrients, and require that these products display front-of-package warning labels such as "WARNING: High in added sugar" or "WARNING: High in sodium" to inform consumers of the high content of potentially unhealthy nutrients. These simple, prominent warnings contrast with the status quo in the US, in which most nutrition information is communicated via the Nutrition Facts Panel, an elaborate, numeric label with small print required on the back of product packaging.

Mounting experimental and observational data suggests that nutrient warnings inform consumers and could help them make healthier choices. Meta-analyses of experimental studies show that warnings on unhealthy foods² and sugary drinks³ help people understand which products are unhealthy and choose healthier options. Warnings also could be applied widely and permanently, in contrast to educational campaigns and weight loss programs, which are typically individually-focused and temporary. Evidence from countries that have already implemented nutrient warning policies supports the effectiveness of warnings. For example, in Chile, sugary drink purchases decreased from 122 mL per capita per day in 2015–2016 to 86 mL per capita per day in 2017 (one year after implementation of nutrient

Corresponding author: Anna H. Grummon, PhD, Harvard Center for Population and Development Studies, 9 Bow St, Room 306, Cambridge, MA 02138, (617) 867-4982, agrummon@hsph.harvard.edu.

Hall and Grummon

warnings), a 24% reduction relative to expected purchases.⁴ Given the recent implementation of the law, it is not yet known if these behavior changes will affect prevalence of obesity or type 2 diabetes. However, research on the physiology of weight loss indicates that even modest reductions in calorie intake (about 55 calories/day) can lead to meaningful weight loss (about 1 pound after 1 year).⁵ These changes could yield substantial benefits at the population level. For example, a modeling study estimated that implementing a national warning policy for sugary drinks could reduce US adults' calorie intake by about 30 calories per day, yielding a projected reduction in obesity prevalence of 3.1 percentage points (from 40.7% to 37.6%) over 5 years.⁶ It is possible that benefits could be more substantial if warnings were applied not only to sugary drinks, but also across many unhealthy foods and beverages.

The benefits of warnings could also extend to the food supply, as warning policies could prompt manufacturers to reformulate products to make them healthy enough to avoid mandated warnings. A recent analysis of Chile's front-of-package nutrient warnings found that the proportion of products high in unhealthy nutrients (i.e., above the thresholds that trigger a mandatory warning under the law) decreased by 7 percentage points in the year after the warnings were implemented, from 51% in 2015–2016 to 44% in 2017.⁷ These changes to the food supply are important because they can improve public health without consumers needing to notice or be influenced by food labels.

Nutrient warnings may be an important step toward ensuring equitable access to nutrition information. Many consumers have difficulty understanding the quantitative information presented on the Nutrition Facts Panel, currently the primary source of nutrition information about foods. In contrast, studies have demonstrated that warnings work well across diverse populations, and simple, prominent nutrient warnings could therefore enhance equitable access to nutrition information. Warnings accompanied by icons or other images could be particularly useful for people with lower literacy or limited English proficiency. Moreover, warnings may also help to counteract industry marketing of unhealthy foods, which is disproportionately targeted toward Black and Latino communities (for example, in a study from 2017, Black children (ages 2–11) were exposed to 86% more food ads on television than their White counterparts, 16.4 vs. 8.8 ads per day as estimated using Nielsen gross rating points).⁸

When crafting a new nutrient labeling regulation, US regulatory agencies can benefit from existing international examples to develop the specifics of the law. The US Food and Drug Administration (FDA) would likely oversee this rulemaking process, similar to their role regulating cigarette pack warnings and the Nutrition Facts Panel for foods and beverages. First, FDA would decide which nutrients to target. The existing laws in Chile, Israel, Mexico, Peru, and Uruguay all require warnings for products with high levels of added sugar, sodium, and saturated fat, given the strong evidence that consuming foods with high levels of these nutrients contributes to chronic disease risk.¹ Some of the existing laws also include warnings about calories or trans fats. Second, FDA would need to set the nutrient thresholds above which a warning would be required. To set these thresholds, FDA could apply one of several available international nutrient profile models that specify unhealthy amounts of specific nutrients. FDA could also take into account their existing guidance that

JAMA. Author manuscript; available in PMC 2021 October 27.

Hall and Grummon

Health warning policies in the US have encountered legal challenges from industry on First Amendment grounds. Warnings are considered "compelled commercial speech," meaning that, to be legally viable, they must meet certain requirements. One requirement is that warnings must be reasonably related to government interests.¹⁰ A Ninth Circuit ruling in 2017 on San Francisco's sugary drink warning ordinance provides case law to suggest nutrient warnings are likely to be deemed as advancing the government's interests of informing consumers and improving health.¹⁰ Warnings must also be factual and uncontroversial.¹⁰ There is reason to be optimistic that nutrient warnings would be judged by the courts as factual and uncontroversial because the warnings would be applied using objective, quantitative nutrient thresholds, and because warnings' thresholds can be linked with existing regulations and guidelines. Moreover, nutrient warnings do not make claims about disease risk that industry can contest in legal battles. In addition, to be allowable, warnings must be unduly burdensome and avoid "chilling" protected commercial speech;¹⁰ FDA could examine recent case law from sugary drink and tobacco warnings to design nutrient warnings that are effective without being unduly burdensome.

Congress should authorize FDA to require nutrient warnings, just as they did in 2009 for cigarette graphic warnings and in 2010 for calorie labels on restaurant menus. Action on warnings is beginning at the local and state levels, promising efforts that should be complemented by action at the federal level. While challenging, the federal pathway would offer several important benefits. First, a federal policy would reach all people living in the US, instead of only those who live in cities and states with local requirements in place. Second, the federal pathway would be the simplest option for industry because companies would not have to comply with a patchwork system of local and state policies. This could reduce administrative costs, ultimately making the policy more cost effective. Third, in contrast with state and local laws, which are subject to preemption, a federal policy would not have to confront preemption challenges. Any federal policy could also include language allowing complementary warning approaches at the state and local level.

Requiring simple, prominent nutrient warnings is a common-sense policy. Whether this approach will be effective in the US is not yet known, but the unabated increase in obesity in the US is one of the countries' greatest health challenges. Given the magnitude of diet-related diseases, no single policy is likely to be the sole answer to solving poor diet and obesity. Instead, multiple interventions across many sectors are needed. Nutrient warnings are an important strategy that should be leveraged as soon as possible. Nutrient warnings could help inform consumers, encourage the food industry to make healthier products, benefit public health, counteract certain industry marketing practices, and potentially improve health equity.

Acknowledgement

The authors have no conflicts of interest to disclose. Marissa Hall was supported by K01HL147713 from the National Heart, Lung, and Blood Institute of the National Institutes of Health. The content is solely the responsibility of the authors and does not necessarily represent the official views of the funders. We thank Sabrina Adler, JD, Jason P. Block, MD, Mark A. Hall, JD, and Lindsey Smith Taillie, PhD, for their thoughtful and constructive feedback on this article.

References

- 1. Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. Lancet. 2019;393(10184):1958–1972. [PubMed: 30954305]
- Clarke N, Pechey E, Kos te D, et al. Impact of Health Warning Labels on Selection and Consumption of Food and Alcohol Products: Systematic Review with Meta-analysis. Health Psychol Rev. 2020:1–39.
- Grummon AH, Hall MG. Sugary drink warnings: A meta-analysis of experimental studies. PLoS Med. 2020;17(5):e1003120. [PubMed: 32433660]
- Taillie LS, Reyes M, Colchero MA, Popkin B, Corvalan C. An evaluation of Chile's Law of Food Labeling and Advertising on sugar-sweetened beverage purchases from 2015 to 2017: A beforeand-after study. PLoS Med. 2020;17(2):e1003015. [PubMed: 32045424]
- Hall KD, Schoeller DA, Brown AW. Reducing Calories to Lose Weight. JAMA. 2018;319(22):2336–2337.
- Grummon AH, Smith NR, Golden SD, Frerichs L, Taillie LS, Brewer NT. Health Warnings on Sugar-Sweetened Beverages: Simulation of Impacts on Diet and Obesity Among U.S. Adults. Am J Prev Med. 2019;57(6):765–774. [PubMed: 31630966]
- Reyes M, Taillie LS, Popkin B, Kanter R, Vandevijvere S, Corvalan C. Changes in the amount of nutrient of packaged foods and beverages after the initial implementation of the Chilean Law of Food Labelling and Advertising: A nonexperimental prospective study. PLoS Med. 2020;17(7):e1003220. [PubMed: 32722710]
- Harris J, Frazier W, Kumanyika S, Ramirez A. Increasing disparities in unhealthy food advertising targeted to Hispanic and Black youth. Hartford, CT: UConn Rudd Center for Food Policy and Obesity;2019.
- 9. A food labeling guide: Guidance for industry. US Department of Health and Human Services, Food and Drug Administration, Center for Food Safety and Applied Nutrition;2013.
- 10. Pomeranz JL, Mozaffarian D, Micha R. Can the government require health warnings on sugarsweetened beverage advertisements? JAMA. 2018;319(3):227–228. [PubMed: 29340685]