




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Evaluation of stress, burnout and hair cortisol levels in health workers at a University Hospital during COVID-19 pandemic

Carolina Ibar^{a,b,1}, Federico Fortuna^{a,1}, Diego Gonzalez^{b,c,d}, Juan Jamaro^c, Dario Jacobsen^{b,c}, Lucas Pugliese^b, Laura Giraud^a, Veronica Ceres^a, Cynthia Mendoza^c, Esteban M. Repetto^{b,e}, Graciela Reboredo^f, Silvia Iglesias^g, Sergio Azzara^h, Gabriela Berg^{b,d,e}, Damian Zopatti^{i,1}, Bibiana Fabre^{b,c,d,*,1}

^a Universidad de Buenos Aires, Facultad de Farmacia y Bioquímica, Residencia en Bioquímica Clínica, Argentina

^b Universidad de Buenos Aires, Facultad de Farmacia y Bioquímica, Departamento de Bioquímica Clínica, Cátedra de Bioquímica Clínica I, Buenos Aires, Argentina

^c Universidad de Buenos Aires, Departamento de Bioquímica Clínica, Hospital de Clínicas José de San Martín, Argentina

^d Universidad de Buenos Aires, Facultad de Farmacia y Bioquímica, Instituto de Fisiopatología y Bioquímica Clínica (INFIBIOC), Buenos Aires, Argentina

^e Universidad de Buenos Aires, Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Facultad de Farmacia y Bioquímica, Buenos Aires, Argentina

^f Universidad de Buenos Aires, Hospital de Clínicas José de San Martín, Área Ambulatoria del tratamiento y seguimiento del HIV-SIDA, Argentina

^g Cátedra de Química Analítica Instrumental, Facultad de Farmacia y Bioquímica, Universidad de Buenos Aires, Buenos Aires, Argentina

^h Cátedra de Metodología para la Investigación Psicológica I, Facultad de Psicología, Universidad de Buenos Aires, Buenos Aires, Argentina

ⁱ Universidad de Buenos Aires, Facultad de Medicina, Dirección de Estadística y Archivo Médico, Hospital de Clínicas José de San Martín, Argentina

ARTICLE INFO

Keywords:

Hair cortisol
Burnout
Stress
COVID-19
Healthcare workers

ABSTRACT

In the critical context of COVID-19 pandemic, healthcare workers are on the front line, participating directly in the care, diagnosis, and treatment of patients with COVID-19. This exposes them to a higher risk of developing chronic stress, psychological distress, and any other mental health symptoms.

Objective: to evaluate stress and burnout in a health workers population and, in addition, to measure hair cortisol concentration as a current biomarker of stress.

Materials and methods: 234 health workers from Hospital de Clínicas “José de San Martín”, Buenos Aires University, were included in this study. In this population hair samples were obtained from the posterior vertex as close to the scalp as possible and the individuals completed the following surveys: perceived stress, social support, burnout scale, life event scale, and sociodemographic data. Hair cortisol was measured by an automated chemiluminescent method. The studied population was divided into three groups considering those individuals below the healthy reference sample range (< 40 pg/mg hair), within the healthy reference range (40–128 pg/mg hair) and above the reference range (> 128 pg/mg hair). This study used a transversal and observational design.

Results: Our results show that 40% of the studied population presented hair cortisol values outside of the healthy reference range. In the whole studied population, a direct correlation was found between hair cortisol concentration and perceived stress as well as between hair cortisol concentration and the emotional exhaustion component of burnout ($r = 0.142$, $p = 0.030$; $r = 0.143$, $p = 0.029$, respectively). 12% of the studied population showed Burnout (52% doctors and residents, 19% nurses, 19% administrative personnel). Higher values in hair cortisol levels were found in the group with burnout versus individuals without burnout ($p = 0.034$). Finally, a mediation analysis was performed, finding that depersonalization is a mediating variable in the relationship between self-perceived stress and hair cortisol level ($F = 4.86$, $p = 0.0086$; indirect effect IC: 0.0987–1.8840).

Conclusion: This is the first study in which a stress biomarker such as hair cortisol is evaluated in this population and in this context. Healthcare workers are subjected to increased levels of stress and burnout. High depersonalization, emotional exhaustion, and decreased personal sense of accomplishment characterize this population. It is the responsibility of the health authorities to implement strategies to manage this psychological emergency.

* Correspondence to: Universidad de Buenos Aires, Facultad de Farmacia y Bioquímica, Departamento de Bioquímica Clínica, Junín 956, C1113AAD Buenos Aires, Argentina.

E-mail address: bfabre@ffyba.uba.ar (B. Fabre).

¹ These authors contributed equally to this paper.

<https://doi.org/10.1016/j.psyneuen.2021.105213>

Received 27 January 2021; Received in revised form 5 March 2021; Accepted 26 March 2021

Available online 1 April 2021

0306-4530/© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

During December 2019, in Wuhan China, cases of severe atypical viral pneumonia emerged, spreading rapidly all over the country, causing a significant number of deaths (WHO, 2020; Zhu et al., 2020). Subsequently, the virus genome was sequenced, determining that it was a new coronavirus called SARS-CoV-2. This virus caused a new disease called COVID-19 that quickly spread worldwide. The World Health Organization (WHO) declared COVID-19 a pandemic disease on March 11th.

Today, it has caused more than 1.900.000 deaths worldwide and has affected up to 191 countries (Hopkins, 2020). Thus, this pandemic is today considered an international public health emergency (Jee, 2020).

In this critical context, healthcare workers are on the front line, participating directly in the care, diagnosis, and treatment of patients with COVID-19. This exposes them to a higher risk of developing chronic stress, psychological distress, and any other mental health symptom. The increasing number of confirmed and suspected cases, overwhelming workload, lack of personal protective equipment, widespread media coverage, lack of specific treatment and feelings of inadequate support can all contribute to the mental burden of these health workers (Mauder et al., 2008).

Previous studies conducted during the 2003 SARS outbreak showed adverse psychological reactions among healthcare workers. In this population, high levels of stress, anxiety, and symptoms of depression (Lee et al., 2007) associated with uncertainty, stigmatization, reluctance to work and resignation (Bai et al., 2004; Mauder et al., 2003) were observed. Several studies found that the emergency nursery department was more likely to develop distress and behavioral disconnection (Lai et al., 2020; Shih et al., 2007; Wong et al., 2005).

At present, facing the pandemic caused by COVID-19, health workers who take care of these patients experience similar effects on their mental health. The public health crisis creates an environment of uncertainty and stress that demands both assessment and containment of health personnel. So far, most of the studies about COVID-19 have focused on epidemiological investigations, prevention, diagnosis, and treatment. Few studies have investigated the mental health problems that affect health workers during the pandemic (Spoorthy et al., 2020; Zhang et al., 2020).

In hospital settings, the burnout syndrome has been extensively studied, even though the most widely accepted definition of burnout is that proposed by Maslach & Jackson (1996) who identified it as a three-dimensional syndrome involving: a) high emotional exhaustion (tiredness and diminished personal resources to face the demands of the task), b) high depersonalization (development of a negative and insensitive attitude towards the recipients of the service), and c) diminished sense of personal accomplishment (perception that professional achievements fall below personal expectations and low personal self-worth). The study of burnout is important because its negative effects can impact both on the professional who suffers it, causing different signs and symptoms, and also on the health institution itself, by increasing staff absenteeism, and on the quality of care provided by increasing medical errors and diminishing patient safety.

Some studies have shown that younger physicians, those performing high-risk procedures and/or experiencing work-life conflicts were at greatest risk (Lacy and Chan, 2018). The phenomena of physician burnout were adequately studied and have a direct negative impact on stress, anxiety, depression, poor patient quality care, among others (Patel et al., 2018; Shah et al., 2020). Besides, chronic stress related to work environment, known as burnout, leads to a hypothalamic-pituitary-adrenal (HPA) axis deregulation which leads to an altered cortisol release (Wingenfeld et al. 2009). Several studies report that chronic stress alters HPA axis activity, which contributes to the development of psychiatric disorders such as depression, anxiety, and burnout (Chrousos, 2009; Miller et al., 2007; Tsigos and Chrousos, 2002). However, most of studies evaluating HPA axis alterations in burnout are controversial. Some studies report increased cortisol values

(Grossi et al., 2005), others do not show any modification in cortisol levels (Bellingrath et al., 2008) and others even show reduced activity of the axis (Pruessner et al., 1999).

In the last years, hair cortisol evaluation arose as a suitable biomarker for adrenal axis evaluation, reflecting the individual's exposure to chronic stressful events. Given that on average, hair grows 1 cm per month, 3 cm of hair would reflect the cortisol levels to which the individual was exposed in the last 3 months (Sauvé et al., 2007).

Currently, worldwide hair cortisol measurement is performed mostly by mass spectrometry or by manual methods with low analytical precision and high cost (e.g., ELISA). However, in the last years, Gonzalez et al. (2019) have developed and evaluated the analytical performance of a procedure that enables hair cortisol measurement by an automated system that allows a fast response time with a greater number of samples with high analytical precision.

In the present study, we propose to explore the deregulation of the HPA axis through a mediation model including hair cortisol concentrations and psychological variables to evaluate the effect of third variables.

Taking this into account, the aim of our study was to evaluate chronic stress and burnout in health workers during COVID-19 pandemic in a university public hospital, as well as its association with a biological biomarker such as hair cortisol.

2. Materials and methods

Health workers ($n = 234$; 68 men (39.6 ± 10 years) and 166 women (41 ± 12 years)) from Hospital de Clínicas "José de San Martín", Buenos Aires University (UBA) were included in this study. They were recruited via email. Individuals under treatment with glucocorticoids, psychotropic drugs, with HPA axis alterations or a previous diagnosis of mental health disorders were excluded. Additionally, individuals with less than 3 cm hair length on the posterior vertex were also excluded. This study used a transversal observational design. The normal reference interval was obtained in a previous study using percentile (P) 2.5 and 97.5 from hair cortisol levels measured in 213 healthy individuals (Gonzalez et al. 2019). The participants did not receive any kind of compensation for participating in the study and all of them gave written prior informed consent. The study was approved in advance by the Ethics Committee of the Hospital and was performed following the Helsinki Declaration for medical studies in humans.

2.1. Hair sample collection and hair cortisol measurement

Hair samples were obtained with scissors from the posterior vertex as close to the scalp as possible. Considering that hair grows approximately 1 cm per month, 3 cm were obtained in order to evaluate hair cortisol levels of the last 3 months. Each sample was stored in a paper envelope at room temperature until it was processed. Once the samples were obtained, three centimeters were measured from the root segment adjacent to the cutting. Then, each sample was weighed, and cortisol was extracted and processed by an automated chemiluminescent method (Immulite 2000 autoanalyzer, Siemens, LA, USA) according to the patented procedure developed in our laboratory. The results were expressed in pg/mg. Hair cortisol concentration reference interval in healthy individuals with low levels of stress is 40–128 pg/mg hair (P2.5–P97.5) (Gonzalez et al., 2019). The studied population was divided into three groups considering this healthy reference sample range: below the lower value (< 40 pg/mg hair), within the healthy reference sample range (40–128 pg/mg hair) and above the reference range (> 128 pg/mg hair).

2.2. Epidemiological data and psychological tests

All participants were asked to complete an epidemiological sheet including data about: age, sex, weight, height, family nucleus,

medication, dye hair, used shampoo, any cosmetic hair treatment, smoking, profession, workplace, and pre-existing pathologies. In addition, they completed the following surveys: perceived stress (Cohen et al., 1988), social support (Timmerman et al., 2000), burnout scale (Maslach et al., 1997), and life event scale Homes-Rahe (Holmes and Rahe, 1967). Perceived stress survey consists in 4 items, answered on a 5-point Likert scale ranging from “never” to “very often”. Social support survey consists in 5 items, answered using a Likert scale ranging from 1 to 4 each item. Holmes-Rahe life events scale consist in an inventory of 45 life events with different scores from 11 to 100 each event. For the analysis, median values were used as follows, for social support (median score 14.5), perceived stress (median score 6) and life events (median score 180).

The reliability analysis of the psychological tools used yielded a Cronbach’s alpha of 0.708 for perceived stress survey, 0.805 for burnout survey and 0.815 for social support scale.

2.2.1. Burnout

The multidimensional concept of burnout was measured by Maslach Burnout Inventory - Human Services Survey (MBI - HSS) This scale consists of 22 items distributed into three subscales: emotional exhaustion (EE) scoring from 0 to 54, depersonalization (DP) scoring from 0 to 30 and personal accomplishment (PA) scoring from 0 to 48. Item responses are from 0 (never) to (6) every day (Maslach and Leiter, 2016), the definition of burnout requires the presence of the three subscales: high EE (score > 26), high DP (score > 9) and low PA (score < 34).

Since its development in 1981, the MBI - HSS (Maslach and Jackson, 1981) has a long history of reliability and validity testing in human services workers (Schaufeli et al., 1996) and in nurses from Europe and the United States (Poghosyan et al., 2009).

2.3. Statistical analyses

We first tested the distribution of variables using normality tests (kurtosis and skewness). Results were expressed as mean \pm standard deviation (SD) or median (range), according to data distribution. Correlations between variables were calculated using Pearson (parametric distribution data) or Spearman test (non-parametric distribution data). Median differences were performed by *t*-test or Mann-Whitney test according to data distribution. We performed the Kruskal Wallis analysis to evaluate differences between medians when we had more than two groups. We then used a binary logistic regression analysis to test whether hair cortisol, perceived stress, social support, and life events were predictors of burnout, controlling for necessary confounders, age, BMI. Finally, the mediation analysis was performed using Hayes PRO-CESS macro in SPSS statistical software (Bolín, 2014). All analyses were conducted with SPSS 19.0 statistical software.

3. Results

3.1. Sociodemographic and psychological characteristics of the study population

Samples from 234 health workers from Hospital de Clínicas José de San Martín were analyzed. Table 1 shows the sociodemographic data of the analyzed population and median (range) to score perceived stress; social support, life events scale, depersonalization (DP), Emotional exhaustion (EE) and Personal accomplishment (AP).

3.2. Stress and hair cortisol concentration

Our results show that 40% (n = 94) of the studied population presented hair cortisol values outside of the healthy reference range. Out of this 40%, 63% showed values higher than the upper limit of the reference value for the methodology (128 pg/mg of hair) and 37% presented

Table 1

Sociodemographic and psychological characteristics of the study population.

Variable	Results
Age (years, mean \pm SD)	41 \pm 11
Gender % (n)	F:71 (166); M: 29 (68)
BMI (Kg/m ²)	25.7 (18–39)
Smoker (%)	Yes (76) No (24)
Dyed hair (%)	Yes (55) No (45)
Cosmetic hair treatment (%)	Yes (17) No (83)
Anti-dandruff shampoo (%)	Yes (88) No (12)
Profession/Occupation (n, %)	Physician (n = 34, %=14) Nurses (n = 25, %=11) Residents (n = 35, %=15) Other health professionals (n = 39, %=17) Administrative staff (n = 63, %=27) Maintenance assistants (n = 27, %=11) Auxiliary health technicians (n = 11, %=5)
Psychological scale (score)	Median (range)
Perceived stress	6 (0–15)
Social support	14.5 (2–20)
Life events scale	180 (0–1002)
Personal accomplishment (AP)	37 (0–48)
Personalisation (DP)	6 (0–27)
Emotional exhaustion (EE)	23 (0–49)

F: female; M: male. Results are expressed as mean \pm standard deviation (SD) or median (range), according to the data distribution.

values below the lower limit of reference 40 pg/mg of hair.

Considering the whole population, hair cortisol concentration directly correlated with perceived stress ($r = 0.142$, $p = 0.030$) as well as with the EE ($r = 0.143$, $p = 0.029$). In addition, an inverse correlation was found between hair cortisol concentration and age ($r = -0.182$, $p = 0.005$).

When dividing the population into three groups according to their hair cortisol concentration (< 40 pg/mg of hair, between 40 and 128 pg/mg, and > 128 pg mg), an association was observed between hair cortisol concentration and DP only in those individuals with hair cortisol levels over 128 pg/mg ($r = 0.297$, $p = 0.021$). In this group, perceived stress was also associated to life events ($r = 0.462$, $p < 0.0001$), as well as with DP and EE ($r = 0.367$, $p = 0.004$; $r = 0.680$, $p < 0.0001$, respectively). Besides, in these subject’s social support showed an inverse association with life events and a direct association with AP ($r = -0.263$, $p = 0.042$; $r = 0.299$, $p = 0.020$, respectively).

When considering the group of individuals with altered hair cortisol levels (< 40 and > 128 pg/mg of hair, n = 94) an association was observed between hair cortisol and perceived stress ($r = 0.230$, $p = 0.026$) as well as with DP ($r = 0.221$, $p = 0.032$).

Table 2 shows hair cortisol concentration according to the profession/occupation of the population. Physicians were those with higher hair cortisol levels and they were significantly different from health workers who performed administrative and maintenance activities ($p = 0.019$, $p = 0.015$, respectively).

When the studied population was divided according to interaction with patients, hair cortisol levels were significantly higher in those who were in direct contact with patients (physicians, nurses, residents) than in those who were not in direct contact with patients (administrative, maintenance personnel), (Table 2, $p = 0.04$).

3.3. Burnout and hair cortisol concentration

12% of the studied population showed burnout with the following distribution: 52% physicians and residents, 19% nurses, 19% administrative personnel and the rest were technical and maintenance personnel. High DP and EE and low PA were more frequent among healthcare workers who were in direct contact with patients compared to those workers who performed tasks that do not involve direct contact with patients. When comparing hair cortisol levels, individuals with burnout presented significantly higher levels than individuals without

Table 2
Hair cortisol concentration and Burnout in the studied population.

	Profession/ occupation	Hair cortisol concentration (pg/mg Hair)	Hair cortisol concentration (pg/mg Hair)	Burnout subscales
Workers in direct contact with patients (n = 133)	Physician (n = 34)	98 (40–348)*	89 (40–423) ¹	AP: 30 (4–48) ⁺
	Nurses (n = 25)	83 (40–280)		
	Residents (n = 35)	85 (40–423)		DP: 9 (0–24) ^x
	Other professional health workers (n = 39)	87 (40–304)		EE: 29 (0–47) ^{&}
Workers not in direct contact with patients (n = 101)	Administrative staff (n = 63)	73 (40–290)#	70 (40–290) ²	AP: 37 (0–48) ⁺⁺
	Maintenance assistants (n = 27)	65 (40–152)**		DP: 5 (0–27) ^{xx}
	Auxiliary health technicians (n = 11)	70 (40–190)		EE: 18 (0–49) ^{&&}

Kruskall Wallis test $p = 0.139$, *vs# $p = 0.019$ *vs** $p = 0.015$
Mann Whitney test 1vs2 $p = 0.014$; +vs++ $p = 0.04$
Mann Whitney test x vs xx $p < 0.0001$; & vs && $p < 0.0001$

burnout, (91 (40–280) vs 77 (40–423) pg /mg hair; $p = 0.037$).
A binary logistic regression, considering the presence of burnout as dependent variable, and hair cortisol concentration, perceived stress, social support, and life events as independent variables was performed. Hair cortisol concentration ($b = 0.006$, OR = 1.006 (CI 95%: 1.000–1.011) $p = 0.047$), and perceived stress ($b = 0.235$, OR = 1.265 (95% CI: 1.075–1.488) $p = 0.002$) explained 15% of the variance of the


dependent variable, even after adjusting for BMI and age.
Even though only 12% of the studied population ($n = 28$) presented burnout (2 of 3 burnout altered components were present in 22% of the subjects ($n = 51$)). Furthermore, it was observed that 36% ($n = 85$) of the studied population presented decreased PA (score: 0–33), 33% ($n = 79$) increased DP (score: 10–30) and 38% ($n = 90$) increased EE (score: 27–54). Moreover, significant differences were observed in hair cortisol levels when comparing individuals with low PA vs those with high PA (91 vs 60 pg/mg of hair, $p = 0.05$). The same is observed when comparing individuals with low DP vs individuals with high DP (70 vs 90 pg/mg of hair, $p = 0.05$), while in the case of individuals with low EE vs individuals with high EE only a tendency was observed (75 vs 90 pg /mg of hair, $p = 0.07$).

3.4. Analysis of mediation

The statistical mediation analysis supported that DP mediated the relationship between self-perceived stress and hair cortisol concentration (Indirect effect: $b = 0.69$, SE = 0.42, IC: 0.0987–1.8840) as shown in Fig. 1.

4. Discussion

In this study, stress and burnout were evaluated in health workers at “Hospital de Clínicas José de San Martín” in the context of COVID-19 pandemic. To our knowledge, this is the first study in which a stress biomarker such as hair cortisol is evaluated in this population and in this context. Our results show that 40% of the studied population presented altered hair cortisol values. In addition, an association between perceived stress and hair cortisol levels was observed, particularly in



Indirect effects: $B=0.69, SE=0.428, IC [0.055-1.6952]$

Fig. 1. Proposed mediation model. a: $B = 0.41, SE = 0.14, t = 2.62, p = 0.005, IC [0.129-0.6903]$; b: $B = 1.71, S = 0.67, t = 2.53, p = 0.012, IC [0.3834-3.0379]$; c: $B = 2.67, SE = 1.49, t = 1.78, p = 0.070, IC [-0.2117 to 5.6060]$; c': $B = 1.97, SE = 1.499, t = 1.3148, p = 0.189, IC [-0.9830 to 4.9262]$. IV: Independent variable; DV: Dependent variable; M: Mediator.

individuals with altered cortisol levels. These findings highlight the usefulness of the measurement of hair cortisol as a suitable biomarker in the evaluation of stress (Iglesias et al., 2015).

Recent studies have similarly shown that COVID-19 affects mental health outcomes such as anxiety, depression, and post-traumatic stress symptoms (Salari et al., 2020). Furthermore, healthcare workers constitute a population particularly vulnerable to develop adverse mental health events due to long working hours, risk of infection, lack of personal protection elements, physical fatigue and separation from their families (Kang et al., 2020).

In this study, a direct correlation was also observed between hair cortisol levels and the EE. Sagherian et al. (2020) found higher levels of EE and some degree of DP in nursing staff, additionally, an increased sense of DP was observed among those who cared for patients with COVID-19. Different studies show that certain work-related characteristics (work schedule, type of work environment, and years of experience) affect levels of poor sleep, fatigue, recovery between shifts, and burnout in nurses (Dall'Ora et al., 2015; Hirsch Allen et al., 2014; Sagherian et al., 2017). We found an association between hair cortisol and DP, only in individuals with altered cortisol levels. So far, no associations have been reported between this biomarker and any of the burnout components in healthcare workers.

When analysing the studied population according to their profession, it was observed that physicians presented higher and significantly different hair cortisol levels compared to other healthcare workers who perform administrative or maintenance tasks. There is evidence of adverse psychological effects on medical personnel during infectious diseases outbreaks such as SARS, MERS, and H1N1 influenza (Brooks et al., 2020; Khalid et al., 2016; Koh et al., 2005; Maunder et al., 2003). Cai et al. (2020) showed that the main stressors for medical personnel were the perceived risk of infection for themselves and their families, patient mortality, and the lack of a specific treatment. In 2005, Wong et al. found that during SARS outbreak psychological stress was higher in nurses, followed by emergency physicians and then auxiliary health personnel, and the most important variables associated with stress were also vulnerability to infection and fear of spreading the new virus.

To our knowledge, this is the first study to analyse health personnel, including not only physicians and nurses but also non-medical health personnel (biochemists, kinesiologists, psychologists, radiologists), maintenance personnel, medical and non-medical residents, technical staff and workers who perform administrative tasks.

The prevalence of burnout in our population was lower than the reported by Takahiro et al. (2020) who found that more than 40% of nurses and more than 30% of radiological technologists and pharmacists met the criteria for burnout. Nevertheless, we found that burnout was more frequent in health personnel who were in direct contact with patients (physicians, nurses and residents), as well as higher values of EE and DP and lower values of AP than those obtained by workers who were not in direct contact with patients (Maslach and Leiter, 2016). Decreased AP values have been reported to be associated with a negative evaluation of the abilities to carry out work tasks. The increase in burnout parameters in workers who are in direct contact with patients may be due to the fact that physicians and nurses are facing a high workload in providing health services. Burnout among physicians has been found to be associated with a wide range of occupational stressors, which are likely to increase during COVID-19 pandemic (Pappa et al., 2020; Rajkumar, 2020).

The binary logistic regression analysis showed that hair cortisol and perceived stress explain 15% of burnout. This result is consistent with the scientific literature that reports that burnout and depressive symptoms are associated with self-perceived stress and exposure to stressful events (Sawatzky et al., 2012). Furthermore, according to binary logistic regression analysis hair cortisol and self-perceived stress are associated with burnout.

Besides, the mediation analysis carried out in our study revealed that depersonalization is a mediating variable between perceived stress and

hair cortisol concentration. This finding becomes relevant since depersonalization has been associated in different studies with anxiety, and significant levels of stress and emotional hyperactivity (Michal et al., 2009). Several studies have shown that when subjects do not have personal resources to face conflict situations, depersonalization acts as a dysfunctional coping strategy (Magdalena Escamilla-Quintal, 2008; Sierra-Siegert, 2008; Gil Monte et al., 1998). Thompson and Jaque (2017) state that to counteract the negative effects of greater emotionality, people with high depersonalization need to acquire and use even more task-oriented strategies to compensate their emotional tendencies. In this sense, our interdisciplinary work team has implemented an Online Training Program to Improve Stress Management aimed to health workers belonging to the Hospital de Clínicas "José de San Martín", Buenos Aires University (UBA).

Considering the lack of data on burnout prevalence in health workers during COVID-19 pandemic, it would be interesting to evaluate this especially susceptible population in the future (post-pandemic) to know more precisely the effects of the pandemic on them. On that account, to establish health strategies with the aim of improving their mental health and life quality.

There are some limitations in our study. One limitation is that only one single measure of hair cortisol concentration has been performed during the pandemic context. Secondly, it would be important to evaluate hair cortisol levels in a non-hospital population, as well as stress and burnout, in pandemic to observe possible associations. Finally, even though we have implemented programs to reduce stress levels, we have not yet been able to evaluate the impact of these programs on stress and burnout.

In summary, our study demonstrates the importance of evaluating stress and burnout in health workers, a vulnerable population in the context of COVID-19 pandemic. As occurred during the MERS and SARS pandemics, an impact on mental health was observed in this population, being more important in those who work in direct contact with COVID patients. Additionally hair cortisol concentrations has been a valuable screening instrument for stress and burnout.

5. Conclusion

In this historically unexpected period, healthcare workers are under higher levels of stress and burnout. Given the current pandemic context, personnel in direct contact with patients face an increased risk of exposure to the disease, a greater workload, moral dilemmas during care and fears about their own personal health. From our research, hair cortisol evaluated by an automated method fulfills the requirement for evaluating stress in this vulnerable population.

CRedit authorship contribution statement

All authors should have made substantial contributions to all of the following: **Ibar C**: acquisition of data, analysis and interpretation of data, drafting the article, final approval of the version to be submitted. **Fortuna F**: acquisition of data, analysis and interpretation of data, drafting the article, final approval of the version to be submitted. **Gonzalez D**: acquisition of data, analysis and interpretation of data, drafting the article or revising it critically for important intellectual content, final approval of the version to be submitted. **Jamardo J**: acquisition of data, analysis and interpretation of data, final approval of the version to be submitted. **Jacobsen D**: acquisition of data, analysis and interpretation of data, drafting the article or revising it critically for important intellectual content, final approval of the version to be submitted. **Pugliese L**: acquisition of data, final approval of the version to be submitted. **Giraud L**: acquisition of data, final approval of the version to be submitted. **Ceres V**: acquisition of data, final approval of the version to be submitted. **Mendoza C**: acquisition of data, final approval of the version to be submitted. **Repetto EM**: analysis and interpretation of data, revising it critically for important intellectual

content, final approval of the version to be submitted. **Reboredo G:** revising it critically for important intellectual content, final approval of the version to be submitted. **Iglesias S:** revising it critically for important intellectual content, final approval of the version to be submitted. **Azzara S:** analysis and interpretation of data, drafting the article or revising it critically for important intellectual content, final approval of the version to be submitted. **Berg G:** analysis and interpretation of data, drafting the article or revising it critically for important intellectual content, final approval of the version to be submitted. **Zopatti D:** the conception and design of the study, acquisition of data, analysis and interpretation of data, drafting the article or revising it critically for important intellectual content, final approval of the version to be submitted. **Fabre B:** the conception and design of the study, acquisition of data, analysis and interpretation of data, drafting the article or revising it critically for important intellectual content, final approval of the version to be submitted.

Conflict of interests

None declared.

Acknowledgements

We thank Dr. Angel Zaninovich for providing comments on an earlier draft.

References

- Bai, Y., Lin, C.-C., Lin, C.-Y., Chen, J.-Y., Chue, C.-M., Chou, P., 2004. Survey of Stress Reactions Among Health Care Workers Involved With the SARS Outbreak. *Psychiatr. Serv.* 55. <https://doi.org/10.1176/appi.ps.55.9.1055>.
- Bellingrath, S., Weigl, T., Kudielka, B.M., 2008. Cortisol dysregulation in school teachers in relation to burnout, vital exhaustion, and effort-reward-imbalance. *Biol. Psychol.* 78. <https://doi.org/10.1016/j.biopsycho.2008.01.006>.
- Bolin, J.H., 2014. Review of Introduction to mediation, moderation, and conditional process analysis: A regression-based approach [Review of the book Introduction to mediation, moderation, and conditional process analysis: A regression-based approach, by A. F. Hayes]. *J. Educ. Meas.* 51 (3), 335–337. <https://doi.org/10.1111/jedm.12050>.
- Brooks, S.K., Webster, R.K., Smith, L.E., Woodland, L., Wessely, S., Greenberg, N., Rubin, G.J., 2020. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 395. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8).
- Cai, H., et al., 2020. Psychological impact and coping strategies of frontline medical staff in Huanan between January and March 2020 during the outbreak of coronavirus disease 2019 (COVID-19) in Hubei, China. *Med. Sci.* <https://doi.org/10.12659/MSM.924171>.
- Chrousos, G.P., 2009. Stress and disorders of the stress system. *Nat. Rev. Endocrinol.* 5. <https://doi.org/10.1038/nrendo.2009.106>.
- Cohen, L.H., Towbes, L.C., Flocco, R., 1988. Effects of induced mood on self-reported life events and perceived and received social support. *J. Personal. Soc. Psychol.* 55. <https://doi.org/10.1037/0022-3514.55.4.669>.
- Dall’Ora, C., Griffiths, P., Ball, J., Simon, M., Aiken, L.H., 2015. Association of 12h shifts and nurses’ job satisfaction, burnout and intention to leave: findings from a cross-sectional study of 12 European countries. *BMJ Open* 5. <https://doi.org/10.1136/bmjopen-2015-008331>.
- Gil Monte, P.R., Peiro, J.M., Valcárcel, P., 1998. A model of burnout process development: an alternative from appraisal models of stress. *COEG* 4, 165–179.
- Gonzalez, D., Jacobsen, D., Ibar, C., Pavan, C., Monti, J., Fernandez Machulsky, N., Balbi, A., Fritzier, A., Jamaro, J., Repetto, E.M., Berg, G., Fabre, B., 2019. Hair cortisol measurement by an automated method. *Sci. Rep.* 9. <https://doi.org/10.1038/s41598-019-44693-3>.
- Grossi, G., Perski, A., Ekstedt, M., Johansson, T., Lindström, M., Holm, K., 2005. The morning salivary cortisol response in burnout. *J. Psychosom. Res.* 59. <https://doi.org/10.1016/j.jpsychores.2005.02.009>.
- Hirsch Allen, A.J., Park, J.E., Adhami, N., Sirounis, D., Tholin, H., Dodek, P., Rogers, A.E., Ayas, N., 2014. Impact of work schedules on sleep duration of critical care nurses. *Am. J. Crit. Care* 23. <https://doi.org/10.4037/ajcc2014876>.
- Holmes, T.H., Rahe, R.H., 1967. The social readjustment rating scale. *J. Psychosom. Res.* 11. [https://doi.org/10.1016/0022-3999\(67\)90010-4](https://doi.org/10.1016/0022-3999(67)90010-4).
- Iglesias, S., Jacobsen, D., Gonzalez, D., Azzara, S., Repetto, E.M., Jamaro, J., Gómez, S.G., Mesch, V., Berg, G., Fabre, B., 2015. Hair cortisol: a new tool for evaluating stress in programs of stress management. *Life Sci.* 141. <https://doi.org/10.1016/j.lfs.2015.10.006>.
- Jee, Y., 2020. WHO international health regulations emergency committee for the COVID-19 outbreak. *Epidemiol. Health* 42. <https://doi.org/10.4178/epih.e2020013>.
- Johns Hopkins Coronavirus Resource Center, 2020. (<https://coronavirus.jhu.edu/map.html>) [WWW Document].
- Kang, L., Li, Y., Hu, S., Chen, M., Yang, C., Yang, B.X., Wang, Y., Hu, J., Lai, J., Ma, X., Chen, J., Guan, L., Wang, G., Ma, H., Liu, Z., 2020. The mental health of medical workers in Wuhan, China dealing with the 2019 novel coronavirus. *Lancet Psychiatry* 7. [https://doi.org/10.1016/S2215-0366\(20\)30047-X](https://doi.org/10.1016/S2215-0366(20)30047-X).
- Khalid, I., Khalid, T.J., Qabajah, M.R., Barnard, A.G., Qushmaq, I.A., 2016. Healthcare workers emotions, perceived stressors and coping strategies during a MERS-CoV outbreak. *Clin. Med. Res.* 14, 7–14. <https://doi.org/10.3121/cmr.2016.1303>.
- Koh, D., Lim, M.K., Chia, S.E., Ko, S.M., Qian, F., Ng, V., Tan, B.H., Wong, K.S., Chew, W.M., Tang, H.K., Ng, W., Muttakin, Z., Emmanuel, S., Fong, N.P., Koh, G., Kwa, C.T., Tan, K.B.-C., Fones, C., 2005. Risk perception and impact of Severe Acute Respiratory Syndrome (SARS) on work and personal lives of healthcare workers in Singapore. *Med. Care* 43. <https://doi.org/10.1097/01.mlr.0000167181.36730.cc>.
- Lacy, B.E., Chan, J.L., 2018. Physician burnout: the hidden health care crisis. *Clin. Gastroenterol. Hepatol.* 16. <https://doi.org/10.1016/j.cgh.2017.06.043>.
- Lai, J., Ma, S., Wang, Y., Cai, Z., Hu, J., Wei, N., Wu, J., Du, H., Chen, T., Li, R., Tan, H., Kang, L., Yao, L., Huang, M., Wang, H., Wang, G., Liu, Z., Hu, S., 2020. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Netw. Open* 3. <https://doi.org/10.1001/jamanetworkopen.2020.3976>.
- Lee, A.M., Wong, J.G., McAlonan, G.M., Cheung, V., Cheung, C., Sham, P.C., Chu, C.-M., Wong, P.-C., Tsang, K.W., Chua, S.E., 2007. Stress and psychological distress among SARS survivors 1 year after the outbreak. *Can. J. Psychiatry* 52. <https://doi.org/10.1177/070674370705200405>.
- Escamilla-Quintal, M., Rodríguez-Molina, I., Peiró, J.M., Tomás Marco, I., 2008. Cynicism: a differential coping strategy as a function of gender. *Psicothema* 20, 592–602.
- Maslach, C., Jackson, S.E., 1981. The measurement of experienced burnout. *J. Organ. Behav.* 2. <https://doi.org/10.1002/job.4030020205>.
- Maslach, C., Jackson, S.E., 1996. *Maslach Burnout Inventory Manual*, 3rd. The Scaerocrow. Palo Alto, CA: Consulting Psychologists Press.
- Maslach, C., Leiter, M.P., 2016. Understanding the burnout experience: recent research and its implications for psychiatry. *World Psychiatry* 15. <https://doi.org/10.1002/wps.20311>.
- Maslach, C., Jackson, S., Leiter, M., 1997. *The Maslach burnout inventory manual, in: Evaluating Stress: A Book of Resources*. The Scaerocrow press, Lanham, Maryland, United States, pp. 191–218.
- Mauder, R., Hunter, J., Vincent, L., Bennett, J., Peladeau, N., Leszcz, M., Sadavoy, J., Verhaeghe, L.M., Steinberg, R., Mazzulli, T., 2003. The immediate psychological and occupational impact of the 2003 SARS outbreak in a teaching hospital. *CMAJ: Can. Med. Assoc. J. J. Assoc. Med. Can.* 168, 1245–1251.
- Mauder, R.G., Leszcz, M., Savage, D., Adam, M.A., Peladeau, N., Romano, D., Rose, M., Schulman, R.B., 2008. Applying the lessons of SARS to pandemic influenza. *Can. J. Public Health* 99. <https://doi.org/10.1007/BF03403782>.
- Michal, M., Wiltink, J., Subic-Wrana, C., Zwerenz, R., Tuin, I., Lichy, M., Brähler, E., Beutel, M.E., 2009. Prevalence, correlates, and predictors of depersonalization experiences in the German general population. *J. Nerv. Ment. Dis.* 197. <https://doi.org/10.1097/NMD.0b013e3181aacd94>.
- Miller, G.E., Chen, E., Zhou, E.S., 2007. If it goes up, must it come down? Chronic stress and the hypothalamic-pituitary-adrenocortical axis in humans. *Psychol. Bull.* 133. <https://doi.org/10.1037/0033-2909.133.1.25>.
- Pappa, S., Ntella, V., Giannakakos, T., Giannakoulis, V.G., Papoutsi, E., Katsaounou, P., 2020. Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: a systematic review and meta-analysis. *Brain Behav. Immun.* 88. <https://doi.org/10.1016/j.bbi.2020.05.028>.
- Patel, R., Bachu, R., Adikey, A., Malik, M., Shah, M., 2018. Factors related to physician burnout and its consequences: a review. *Behav. Sci.* 8. <https://doi.org/10.3390/bs8110098>.
- Poghosyan, L., Aiken, L.H., Sloane, D.M., 2009. Factor structure of the Maslach burnout inventory: an analysis of data from large scale cross-sectional surveys of nurses from eight countries. *Int. J. Nurs. Stud.* 46. <https://doi.org/10.1016/j.ijnurstu.2009.03.004>.
- Pruessner, J.C., Hellhammer, D.H., Kirschbaum, C., 1999. Burnout, perceived stress, and cortisol responses to awakening. *Psychosom. Med.* 61. <https://doi.org/10.1097/00006842-199903000-00012>.
- Rajkumar, R.P., 2020. COVID-19 and mental health: a review of the existing literature. *Asian J. Psychiatry* 52. <https://doi.org/10.1016/j.ajp.2020.102066>.
- Sagherian, K., Clinton, M.E., Abu-Saad Huijjer, H., Geiger-Brown, J., 2017. Fatigue, work schedules, and perceived performance in bedside care nurses. *Workplace Health Saf.* 65, 304–312. <https://doi.org/10.1177/2165079916665398>.
- Sagherian, K., Steege, L.M., Cobb, S.J., Cho, H., 2020. Insomnia, fatigue and psychosocial well-being during COVID-19 pandemic: a cross-sectional survey of hospital nursing staff in the United States. *J. Clin. Nurs.* <https://doi.org/10.1111/jocn.15566>.
- Salari, N., Hosseini-Far, A., Jalali, R., Vaisi-Raygani, A., Rasoulpoor, Shna, Mohammadi, M., Rasoulpoor, Shabnam, Khaledi-Paveh, B., 2020. Prevalence of stress, anxiety, depression among the general population during the COVID-19 pandemic: a systematic review and meta-analysis. *Glob. Health* 16. <https://doi.org/10.1186/s12992-020-00589-w>.
- Sauvé, B., Koren, G., Walsh, G., Tokmakejian, S., van Uum, S.H., 2007. Measurement of cortisol in human hair as a biomarker of systemic exposure. *Clin. Invest. Med.* 30, 183. <https://doi.org/10.25011/cim.v30i5.2894>.
- Sawatzky, R.G., Ratner, P.A., Richardson, C.G., Washburn, C., Sudmant, W., Mirwaldt, P., 2012. Stress and depression in students. *Nurs. Res.* 61. <https://doi.org/10.1097/NNR.0b013e31823b1440>.

- Schaufeli, W., Leiter, M., Maslach, C., Jackson, S., 1996. Maslach Burnout Inventory – General Survey (GS). Maslach Burnout Inventory Manual.
- Shah, K., Chaudhari, G., Kamrai, D., Lail, A., Patel, R.S., 2020. How essential is to focus on physician's health and burnout in coronavirus (COVID-19) pandemic? *Cureus*. <https://doi.org/10.7759/cureus.7538>.
- Shih, F.-J., Gau, M.-L., Kao, C.-C., Yang, C.-Y., Lin, Y.-S., Liao, Y.-C., Sheu, S.-J., 2007. Dying and caring on the edge: Taiwan's surviving nurses' reflections on taking care of patients with severe acute respiratory syndrome. *Appl. Nurs. Res.* 20. <https://doi.org/10.1016/j.apnr.2006.08.007>.
- Sierra-Siegert, M., 2008. La despersonalización: aspectos clínicos y neurobiológicos. *Rev. Colomb. Psiquiatr.* 37, 40–55.
- Spoorthy, M.S., Pratapa, S.K., Mahant, S., 2020. Mental health problems faced by healthcare workers due to the COVID-19 pandemic—a review. *Asian J. Psychiatry* 51. <https://doi.org/10.1016/j.ajp.2020.102119>.
- Takahiro, M., et al., 2020. Prevalence of Health Care Worker Burnout During the Coronavirus Disease 2019 (COVID-19) Pandemic in Japan. *JAMA Netw. Open*. <https://doi.org/10.1001/jamanetworkopen.2020.17271>.
- Thompson, P., Jaque, S.V., 2017. Depersonalization, adversity, emotionality, and coping with stressful situations. *J. Trauma Dissociation* 19 (2), 143–161.
- Timmerman, I.G.H., Emanuels-Zuurveen, E.S., Emmelkamp, P.M.G., 2000. The social support inventory (SSI): a brief scale to assess perceived adequacy of social support. *Clin. Psychol. Psychother.* 7. [https://doi.org/10.1002/1099-0879\(200011\)7:5<401.AID-CPP253>3.0.CO;2-I](https://doi.org/10.1002/1099-0879(200011)7:5<401.AID-CPP253>3.0.CO;2-I).
- Tsigos, C., Chrousos, G.P., 2002. Hypothalamic–pituitary–adrenal axis, neuroendocrine factors and stress. *J. Psychosom. Res.* 53. [https://doi.org/10.1016/S0022-3999\(02\)00429-4](https://doi.org/10.1016/S0022-3999(02)00429-4).
- WHO, Novel coronavirus-China, 2020. (<https://www.who.int/csr/don/12-january-2020-novel-coronavirus-china/en>) (Jan 12, 2020) (Accessed Dec 19, 2020) [WWW Document]. Novel coronavirus-China.
- Wingenfeld, K., Schulz, M., Damkroeger, A., Rose, M., Driessen, M., 2009. Elevated diurnal salivary cortisol in nurses is associated with burnout but not with vital exhaustion. *Psychoneuroendocrinology* 34. <https://doi.org/10.1016/j.psyneuen.2009.02.015>.
- Wong, T.W., Yau, J.K.Y., Chan, C.L.W., Kwong, R.S.Y., Ho, S.M.Y., Lau, C.C., Lau, F.L., Lit, C.H., 2005. The psychological impact of severe acute respiratory syndrome outbreak on healthcare workers in emergency departments and how they cope. *Eur. J. Emerg. Med.* 12. <https://doi.org/10.1097/00063110-200502000-00005>.
- Zhang, W., Wang, K., Yin, L., Zhao, W., Xue, Q., Peng, M., Min, B., Tian, Q., Leng, H., Du, J., Chang, H., Yang, Y., Li, W., Shangguan, F., Yan, T., Dong, H., Han, Y., Wang, Y., Cosci, F., Wang, H., 2020. Mental health and psychosocial problems of medical health workers during the COVID-19 epidemic in China. *Psychother. Psychosom.* 89. <https://doi.org/10.1159/000507639>.
- Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R., Niu, P., Zhan, F., Ma, X., Wang, D., Xu, W., Wu, G., Gao, G.F., Tan, W., 2020. A novel coronavirus from patients with pneumonia in China, 2019. *N. Engl. J. Med.* 382. <https://doi.org/10.1056/NEJMoa2001017>.