



Published in final edited form as:

Curr Hepatol Rep. 2020 December ; 19(4): 462–469. doi:10.1007/s11901-020-00554-6.

Racial and Sex Disparities in Hepatocellular Carcinoma in the USA

Faith Ajayi¹, Jenny Jan¹, Amit G. Singal^{1,2,3}, Nicole E. Rich^{1,2}

¹Division of Digestive and Liver Diseases, Department of Internal Medicine

²Harold C. Simmons Comprehensive Cancer Center

³Department of Population and Data Sciences, UT Southwestern Medical Center

Abstract

Purpose of Review: In this review, we aim to provide a summary of the current literature on race and gender disparities in hepatocellular carcinoma (HCC) incidence, stage at diagnosis, treatment and prognosis in the United States.

Recent Findings: HCC incidence rates are rising in the U.S. in all racial/ethnic groups except for Asian/Pacific Islanders, with disproportionate rises and the highest rates among Hispanics compared to Blacks and non-Hispanic whites. There are striking sex disparities in HCC incidence and mortality; however, with the shifting epidemiology of HCC risk factors in the U.S, there is recent evidence that HCC is trending towards less male predominance, particularly among younger birth cohorts. Despite significant advances in HCC treatment over the past decade, disparities in HCC surveillance and treatment receipt persist among racial and ethnic minorities and the socioeconomically disadvantaged. Black patients continue to experience worse survival outcomes than non-Black patients with HCC.

Summary: There are significant racial and gender disparities in HCC incidence, treatment, and mortality in the U.S. Though these disparities are well-documented, data are still limited on the specific determinants driving disparities in HCC. To achieve health equity for all patients with HCC, we must advance beyond simply reporting on disparities and begin implementing targeted interventions to eliminate disparities.

Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. <http://www.springer.com/gb/open-access/authors-rights/aam-terms-v1>

Correspondence: Nicole E. Rich, MD MS, Division of Digestive and Liver Diseases, Department of Internal Medicine, UT Southwestern Medical Center, 5959 Harry Hines Blvd, POB 1, Suite 420, Dallas TX 75390-8887, Tel: 214-645-6111, Fax: 214-645-6114, Nicole.rich@utsouthwestern.edu.

Author Contributions: Study concept and design (all authors); Drafting of the manuscript (all authors); Critical revision of the manuscript for important intellectual content (all authors)

Publisher's Disclaimer: This Author Accepted Manuscript is a PDF file of a an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

Conflicts of Interest: Amit Singal has been on advisory boards and served as a consultant for Wako Diagnostics, Roche, Exact Sciences, Glycotest, Bayer, Eisai, Exelixis, BMS, Merck Genentech, and TARGET-Pharmasolutions. The other authors have no relevant conflicts of interest.

Keywords

Ethnicity; racial; sex; disparities; liver cancer

Introduction

Hepatocellular carcinoma (HCC) is one of the fastest rising causes of cancer-related death in the U.S. and a leading cause of death in patients with cirrhosis¹; however, the disease burden of HCC is unequally distributed, with men and racial/ethnic minorities disproportionately affected compared to women and non-minority populations. As has been observed in other cancers, disparities can occur at any point along the HCC care continuum, including under-recognition of cirrhosis, screening underuse, and delayed diagnosis or treatment, all of which may lead to poorer outcomes and increased mortality. HCC prognosis is significantly impacted by stage at diagnosis, with curative treatments available for patients diagnosed at an early stage. Though racial and gender disparities in HCC incidence and prognosis have been well-described over the past decade, data are limited on the specific determinants driving these disparities. The epidemiology of HCC in the U.S. is shifting, with fewer cases attributed to viral hepatitis, given the impact of effective direct-acting antivirals for hepatitis C and vaccination against hepatitis B, whereas the epidemic of non-alcoholic fatty liver disease (NAFLD) is rising exponentially.² Understanding the current landscape of racial and gender disparities in HCC is the first step to identify targets for intervention to reduce these disparities and promote equity for all patients with this deadly cancer. In this review, we will summarize the literature on the current landscape of racial and gender disparities in HCC.

Disparities in HCC Incidence

Racial and Ethnic Disparities

HCC is one of the few cancers in the U.S. with rising incidence and mortality rates³, and there is notable variation in disease burden by race/ethnicity. Historically, the highest HCC incidence rates were observed in Asian/Pacific Islanders⁴, due in large part to high prevalence of hepatitis B virus (HBV) infection among immigrants from HBV-endemic areas; however, these rates plateaued around 2007 and began to decline in recent years (annual percent change [APC] -1.59% and -2.2% among men and women, respectively).⁵ These declines are attributed to lower rates of HBV infection among subsequent generations born in the U.S., HBV vaccination programs and improvements in antiviral therapy for HBV.⁶ Conversely, HCC incidence rates have risen in all other racial/ethnic groups, with disproportionate rises and the highest rates among Hispanic men (4.7% per year since 2000) compared to Blacks and non-Hispanic whites.^{7, 8} A study by Petrick et al forecasted HCC incidence rates will continue to increase through the year 2030 in all racial/ethnic groups except Asian/Pacific Islanders.⁵ This rise is largely attributed to aging of the baby-boomer birth cohort. Among younger adults (aged <50), HCC incidence rates have begun to decline since the mid-2000s, with largest decreases noted among middle-aged Blacks (-17.2% per year since 2012) compared to adults of similar age in other racial and groups.⁹ Alaskan Natives have also historically had high prevalence of HCC due in large part to high rates of HBV infection in the 1970s¹⁰, though incidence rates, particularly among young adults, have

sharply declined in recent years due to a universal newborn and catch-up HBV vaccination program established in the 1980s.^{11, 12}

There is substantial geographic variation in HCC incidence in the U.S., with Texas having the highest rates¹³, followed by Hawaii, New Mexico, and California.¹⁴ This is due in part to racial/ethnic diversity of citizens living in these states, with age-adjusted incidence rates highest among Hispanics in Texas (21.2 per 100,000) compared to 9.3 per 100,000 for Whites, 16.3 per 10,000 for Blacks and 15.1 per 100,000 for Asian/Pacific Islanders.¹⁴ Nativity, or country of birth, also appears to impact HCC risk.¹⁵ Foreign-born Asian/Pacific Islanders have higher HCC risk whereas U.S.-born Hispanics have higher HCC incidence rates than their foreign-born counterparts.¹⁵ The reasons for this have not been elucidated but are thought to be due to higher rates of known HCC risk factors (e.g. HCV infection, diabetes, metabolic syndrome and non-alcoholic fatty liver disease [NAFLD]) in U.S.-born Hispanics compared to those born outside the U.S. Disparities in HCC incidence have also been observed based on neighborhood-level socioeconomic status^{16, 17} and within subgroups of Asian/Pacific Islanders and Hispanics in California living in neighborhoods based on ethnic enclave status.¹⁸

Sex Disparities

Sex disparities have been observed for most cancers; in fact, there are only a few examples of malignancies with higher incidence rates in women compared to men (i.e. thyroid, bladder, brain cancers).¹⁹ Sex disparities in HCC incidence are well-documented, with men having 2-8 times higher incidence rates than women across geographic regions, time period, and all racial/ethnic groups.^{20, 21} There is some recent evidence that HCC is trending towards less male predominance, particularly among younger birth cohorts.²² Since the year 2000, the male-to-female incidence rate ratio (IRR) has decreased in younger adults of all races/ethnicities but has remained stable in adults over 50 years.²²

However, despite this consistent observation, the specific factors driving this sex disparity are poorly understood. While there are gender differences in the distribution of certain HCC risk factors (e.g. alcohol use, viral hepatitis, diabetes, obesity)²³, these risk factors have varied in prevalence over time in the general population while the sex disparity has remained striking and consistent, suggesting that biological factors (e.g. sex hormones) also play a role in HCC risk. Though the role of sex hormones in HCC pathogenesis has primarily been investigated in animal models, there are data suggesting estrogen may be protective (via suppression of pro-inflammatory cytokines such as IL-6²⁴) and androgens deleterious (via signaling pathways leading to upregulation of vascular endothelial growth factor [VEGF]²⁵). A retrospective cohort study of 928 men with chronic HBV and diabetes found higher serum testosterone levels were associated with increased HCC risk²⁶, whereas a small case-control study of 234 women found exogenous estrogen use was associated with reduced risk of HCC.²⁷ A large population-based study of circulating sex hormones in stored serum found no significant association between estrogen levels and HCC risk, however sex hormone binding globulin (SHBG) and 4-androstendione levels were associated with higher and lower risk of HCC, respectively.²⁸ Further studies in large prospective cohorts of men and

women with cirrhosis are needed to untangle the relative contribution of sex- and gender-related factors in HCC risk in order to address this sex disparity.

Disparities in HCC Risk Factors, Surveillance, and Stage at Diagnosis

Risk Factors

Between 80-90% of HCC in the U.S. occur in a background of cirrhosis.²⁹ The prevalence of underlying risk factors for HCC, including underlying liver disease etiology, smoking, obesity, and diabetes, differ by race/ethnicity and gender.^{23, 30} For example, an estimated 75-100 million Americans have non-alcoholic fatty liver disease (NAFLD), which has surpassed viral hepatitis and become the most common cause of chronic liver disease in the U.S.³¹ There are significant disparities in NAFLD prevalence and severity, with Hispanics having the highest prevalence of NAFLD and highest risk of progression to non-alcoholic steatohepatitis (NASH) compared to other racial/ethnic groups.³² Makarova-Rusher et al evaluated the population attributable fraction (PAF) for various HCC risk factors using SEER-Medicare data, and found the PAF for metabolic disorders was highest among Hispanics (39.3%, 95% CI 31.9 – 40.4%) and whites (34.8%, 95% CI 33.1 – 33.6%); whereas the PAF for HCV was highest among Black patients with HCC (36.1%, 95% CI 31.8 – 40.4%).³⁰ There are also differences in the risk factors and presentation for HCC between men and women. Women tend to be older than men at time of HCC diagnosis with higher rates of NAFLD and metabolic syndrome, whereas viral hepatitis, alcohol use, and smoking are more commonly seen in men.^{33–35} Targeted intervention at the underlying etiology of liver disease can guide HCC prevention strategies. It cannot be neglected that social determinants of health (e.g. poverty and environmental stressors) may lead to higher rates of HCC risk factors such as obesity and alcohol use in racial and ethnic minorities and the socioeconomically disadvantaged.

Surveillance

HCC surveillance is associated with early tumor detection and improved survival in patients with cirrhosis.³⁶ Unfortunately, despite professional society recommendations, HCC surveillance continues to be underutilized in the U.S.³⁷, particularly among racial/ethnic minorities and the socioeconomically disadvantaged.³⁸ There do not appear to be significant sex-related differences in HCC surveillance utilization.³⁹ Lower SES has been associated with decreased rates of cancer screening⁴⁰, even among patients receiving care in an integrated health system. Singal et al found that only 13% of patients with cirrhosis received annual HCC surveillance at an urban safety-net hospital, with lower rates of consistent surveillance among Black patients and the underinsured.⁴¹ Patient-reported barriers to surveillance include issues with scheduling, transportation, and financial burden.⁴² Receipt of consistent and timely HCC surveillance is a key target for interventions to reduce disparities in HCC outcomes, as patients diagnosed at later tumor stages have limited treatment options and poor prognosis.⁴³

Tumor Stage at Diagnosis

Given disparities in surveillance, it is not unexpected that Black patients are more often diagnosed with HCC at an advanced stage compared to non-Black patients.⁴⁴ In an analysis

of the SEER database, Ha et al found Black patients had significantly higher odds (OR 1.20, 95% CI 1.10 – 1.30) and Asians had significantly lower odds (OR 0.87, 95% CI 0.80 – 0.94) of having advanced stage HCC at time of diagnosis compared to whites.⁴⁵ Similarly, in a study of 1117 patients with HCC at two large health systems in Texas, we found Hispanic (OR 0.75, 95% CI 0.55–1.00) and Black patients (OR 0.74, 95% CI 0.56–0.98) were less likely to be diagnosed with early stage HCC than white patients.⁴⁶ Lastly, among 999 patients with HCC in Florida, Jones et al found Blacks presented with more advanced stages with larger tumors, with 65.9% of Blacks beyond Milan criteria at diagnosis, compared to 54.6% of Asians, 49.7% of Hispanics, and 39.9% of Whites ($p < 0.01$).⁴⁷ One of the major contributing factors to late stage HCC detection among Black patients is that they are less likely to be adequately screened for HCC compared to other racial and ethnic groups, and patients connected with a hepatologist or gastroenterologist are more likely to receive routine surveillance.⁴⁸ Several studies have demonstrated sex disparities in tumor stage at diagnosis.^{34, 35} In a retrospective study of 1100 patients with HCC, we found women with HCC were more likely than men to be detected at an early tumor stage (50.8% vs 42.1%; OR 1.55, 95% CI 1.16 – 2.08), even after adjusting for age, race/ethnicity, liver disease etiology and liver function.³⁶

Disparities in HCC Treatment

HCC treatment is underutilized in the U.S., including among patients with early stage disease eligible for curative therapy (i.e. liver transplantation, surgical resection, and ablation)⁴⁹, with the some of the lowest rates of curative treatment receipt observed among patients with low socioeconomic status⁵⁰ and racial and ethnic minorities.⁵¹ Several population-based studies using data from SEER and the Nationwide Inpatient Sample (NIS) demonstrated Blacks and Hispanics have significantly lower rates of curative treatment receipt compared to whites, even after adjusting for tumor stage at diagnosis.^{45, 52, 53} Similarly, in a study of patients with HCC at two large health systems in Texas, we found Hispanics (OR 0.51, 95% CI 0.37 – 0.70) and Blacks (OR 0.60, 95% CI 0.4 - 0.81) were less likely to receive curative treatment than whites, even after adjusting for insurance status and tumor stage at diagnosis.⁵⁴ Significant racial and ethnic disparities in liver transplantation for HCC have also been reported, with Black^{51, 55} and Hispanic⁵⁶ patients having less access to transplant compared to whites. In a study of data from the Texas Cancer Registry, Alawadi et al also found Hispanics and Blacks were less likely to receive surgical treatment for HCC compared whites, but unexpectedly, rural residency was not associated with disparities in surgical treatment for HCC.⁵⁷ Finally, Stewart et al found substantial inter-ethnic disparities in receipt of curative HCC treatment and survival among various Asian ethnic subgroups, even after accounting for tumor stage and socioeconomic status.⁵⁸

Women with HCC appear to receive curative treatment at similar or higher rates compared to men, which may be due in part to women being more likely to receive regular HCC surveillance⁴⁸ and therefore having a higher proportion of early stage tumors, as well as lower rates of hepatic decompensation at time of HCC diagnosis.⁵⁹ In a study analyzing SEER data, Yang et al found a higher proportion of women received surgical treatment than men (44% vs 36%, $p < 0.001$).⁶⁰ Sobotka et al analyzed data from the Nationwide Inpatient Sample and found women had higher odds of treatment with surgical resection (OR 1.31)

and ablation (OR 1.22) compared to men; they found no significant difference in the rates of transarterial chemoembolization (TACE) between men and women.⁶¹ In another study, Cauble et al found women with HCC presented less often with decompensated cirrhosis (OR 0.79, $p < 0.001$), however, even among the subgroup of patients with compensated cirrhosis, women were offered curative therapy more often than men.⁵⁹ Women are less likely than men to undergo liver transplantation, both generally as well as for HCC.^{62, 63} There are limited data on sex disparities in locoregional and systemic therapy in HCC.

Potential causes of treatment disparities are complex and likely related to a combination of patient-, provider-, and/or system-level factors. Patients may have misconceptions about cancer treatment that may impact their willingness to proceed with treatment. Financial toxicity, difficulty with transportation, language barriers, mistrust of providers and health systems, and other patient beliefs/preferences may all impact an individual patient's decision to proceed with treatment.^{64–66} Provider biases and system-level factors (e.g. hospital volume and facilities) also play a role. For instance, Mokdad et al found that 42% of patients in Texas with HCC received treatment at a safety-net health systems, facilities which care for a disproportionate number of racial and ethnic minorities and the socioeconomically disadvantaged.⁶⁷ Patients were less likely to receive curative treatment at the safety-net hospitals, even when diagnosed at an early stage (OR 0.51, 95% CI 0.40-0.66) and experienced worse overall survival (HR 1.30, 95% CI 1.22 – 1.39) than patients at non-safety-net hospitals.⁶⁷ Similarly, Hoehn et al found liver transplantation and resection were performed less often for HCC at safety-net hospitals (50.7% vs 66.7%).⁶⁸ While patient insurance status is associated with more advanced tumor stage at diagnosis and lower odds of HCC treatment receipt⁶⁹, “equal” health care coverage or insurance may still not translate into equitable access to care or outcomes. For example, in a study of Medicaid/Medicare enrollees with colorectal cancer, disparities in surgical treatment still persisted with fewer Black patients undergoing resection compared to whites.⁷⁰

Disparities in HCC Prognosis and Survival

Black patients experience poorer survival than whites for many cancers⁷¹, including HCC. In the U.S., studies demonstrating that Black patients and the socioeconomically disadvantaged experience worse survival than non-Black patients are ubiquitous.⁵⁷ One explanation for this finding is disparate use of HCC surveillance resulting in delayed detection, later stage diagnosis and disparities in curative treatment as outlined above. Other factors driving disparities in outcomes may be liver disease etiology, liver disease severity, and other comorbid conditions. Black patients may also experience more fragmented care, which has also been associated with worse outcomes in HCC.⁷² However, even among the subset of patients undergoing potentially curative therapy (e.g. surgical resection, liver transplantation, ablation), Black patients have significantly worse survival compared to other races and ethnicities.⁷³ In a study of the SEER database, Mathur et al found Blacks had a 12% higher mortality rate (HR, 1.11; 95% CI, 1.03-1.20), Hispanics had a similar mortality rate (0.97; 0.91-1.04), and Asians had a 16% lower mortality rate (0.84; 0.79-0.89) compared to whites.⁷⁴ In a study among 1117 patients with HCC from two U.S. health systems, we found Black patients had worse mortality (HR 1.12, 95% CI 1.10 – 1.14) and Hispanics had lower mortality (HR 0.83, 95% CI 0.74 – 0.94) compared to whites, even

after adjusting for tumor stage, Child Pugh score and HCC treatment receipt.⁵⁴ Njei et al found persistent disparities in survival even after liver transplantation, with Blacks having worse survival and Asian/Pacific Islanders having better survival compared to whites.⁷⁵ Geographic disparities in HCC outcomes have also been reported, with worse outcomes in Southern states.⁷²

Age-adjusted mortality rates are higher in men than in women for most GI malignancies, including colorectal and esophageal cancers⁷⁶, and women also appear to have a survival advantage in HCC. In an analysis of SEER data, Yang et al found women had better overall survival compared to men, independent of age, race, disease stage, or treatment.⁷⁷ Notably, the largest difference in overall survival was among patients aged 18 to 44 in which women had a 4-month survival benefit relative to age-matched men (HR 0.75, P<001).⁷⁷ Similarly, in a study of 1110 patients with HCC from two large health systems in the United States, we found women had better overall survival than men (17.1 vs 12.0 months; HR 0.81, 95% CI 0.68 – 0.97), after adjusting for tumor stage, liver disease etiology, and liver function.³⁴ This sex disparity was consistent across subgroups including liver disease etiology, BCLC stage, and type of HCC treatment, however the survival benefit seen in women differed by age, with younger women having a survival benefit versus younger men (18.3 vs 11.2 months, p=0.02), whereas no difference in survival was observed between older women and men (15.7 vs 15.5 months, p=0.45).³⁴ These findings suggest that sex hormones may play a role in HCC prognosis, however further studies are needed.

While some believe differences in HCC prognosis may extend beyond disparities in healthcare access, studies evaluating the doubling time of HCC tumors have not demonstrated significant sex or racial/ethnic differences in tumor biology or “aggressiveness”, except more rapid growth in Asian populations with predominately HBV-related HCC.^{78, 79}

Future Directions Toward Eliminating Disparities

Though racial and gender differences in incidence and mortality have been described for many cancers, including HCC, few studies have evaluated the root causes and mechanisms underlying these disparities. Most studies on disparities in HCC are limited in scope and rely on administrative datasets lacking granular information on liver disease etiology, liver function, socioeconomic status, and psychosocial or behavioral factors. Further, few studies in HCC have investigated the intersectionality of race and gender or race and socioeconomic status.⁸⁰ There are several challenges inherent to the study of cancer disparities including accurate ascertainment of race and ethnicity and classification of multi-ethnic individuals. Social determinants of health (e.g. poverty, environmental stress, health literacy, social support) are likely to heavily influence disparities in HCC incidence and mortality.⁸¹ Systemic racial inequity in the U.S. represents a public health crisis and large-scale interventions are needed to mitigate and reverse its effects on the health of vulnerable populations. Racism and discrimination are indeed social determinants of health that have negative downstream effects on health and access to healthcare.⁸² Implicit biases and stereotypes may impact decision making regarding screening and treatment, compounding disparities in marginalized populations with decreased healthcare access already facing other

barriers.⁸³ Furthermore, it is increasingly recognized that seemingly “neutral” diagnostic and prognostic algorithms in medicine may perpetuate, rather than mitigate, racial and gender biases and disparities.^{84, 85} For example, estimated glomerular filtration rate (eGFR) may overestimate kidney function in Black patients, potentially delaying referral for treatment or kidney transplantation⁸⁶, and the use of the Model for End Stage Liver Disease (MELD) score for organ allocation has disadvantaged women resulting in lower odds of liver transplantation and higher rates of waitlist death.^{87, 88} While there is growing body of research on healthcare disparities, it is not sufficient to merely continue to identify and prove that disparities exist; rather, we must collectively take the critical next step to determine their root causes, implement interventions, and advocate for system-level change to eliminate disparities.

Conclusions

There are significant racial and gender disparities in the disease burden, treatment, and outcomes of HCC in the U.S, with men and racial/ethnic minorities disproportionately impacted. Though some progress has been made, data are limited on the specific determinants driving disparities in HCC. As a scientific community, we must advance beyond simply reporting on disparities and instead begin implementing interventions to eliminate disparities to achieve equitable care for all patients with HCC.

Acknowledgments

Grant Support: Drs. Singal and Rich are supported by National Cancer Institute R01 MD12565. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

- *1. Kulik L, El-Serag HB. Epidemiology and Management of Hepatocellular Carcinoma. *Gastroenterology* 2019; 156:477–491.e1. [PubMed: 30367835] Comprehensive, up-to-date review on trends in risk factors, diagnosis, and therapeutic advances in HCC.
2. Estes C, Razavi H, Loomba R, et al. Modeling the epidemic of nonalcoholic fatty liver disease demonstrates an exponential increase in burden of disease. *Hepatology* 2018;67:123–133. [PubMed: 28802062]
3. Torre LA, Siegel RL, Ward EM, et al. Global Cancer Incidence and Mortality Rates and Trends—An Update. *Cancer Epidemiology Biomarkers & Prevention* 2016;25:16.
4. Chang ET, Keegan THM, Gomez SL, et al. The burden of liver cancer in Asians and Pacific Islanders in the Greater San Francisco Bay Area, 1990 through 2004. *Cancer* 2007;109:2100–2108. [PubMed: 17385214]
- *5. Petrick JL, Kelly SP, Altekruse SF, et al. Future of Hepatocellular Carcinoma Incidence in the United States Forecast Through 2030. *J Clin Oncol* 2016;34:1787–94. [PubMed: 27044939] A summary of trends in HCC incidence among all racial/ethnic groups and forecasts expected trends through the next decade.
6. Wasley A, Kruszon-Moran D, Kuhnert W, et al. The Prevalence of Hepatitis B Virus Infection in the United States in the Era of Vaccination. *The Journal of Infectious Diseases* 2010;202:192–201. [PubMed: 20533878]
7. El-Serag HB, Lau M, Eschbach K, et al. Epidemiology of Hepatocellular Carcinoma in Hispanics in the United States. *Archives of Internal Medicine* 2007;167:1983–1989. [PubMed: 17923599]
8. El-Serag HB. Epidemiology of Viral Hepatitis and Hepatocellular Carcinoma. *Gastroenterology* 2012; 142:1264–1273.e1. [PubMed: 22537432]

9. Rich NE, Yopp AC, Singal AG, et al. Hepatocellular Carcinoma Incidence Is Decreasing Among Younger Adults in the United States. *Clin Gastroenterol Hepatol* 2020; 18:242–248.e5. [PubMed: 31042582]
10. Jim MA, Perdue DG, Richardson LC, et al. Primary liver cancer incidence among American Indians and Alaska Natives, US, 1999–2004. *Cancer* 2008;113:1244–1255. [PubMed: 18720380]
11. McMahon BJ, Bulkow L, Harpster A, et al. Screening for hepatocellular carcinoma in Alaska natives infected with chronic hepatitis B: a 16-year population-based study. *Hepatology* 2000;32:842–6. [PubMed: 11003632]
12. McMahon BJ, Bulkow LR, Singleton RJ, et al. Elimination of hepatocellular carcinoma and acute hepatitis B in children 25 years after a hepatitis B newborn and catch-up immunization program. *Hepatology* 2011;54:801–807. [PubMed: 21618565]
13. El-Serag HB, Sardell R, Thrift AP, et al. Texas Has the Highest Hepatocellular Carcinoma Incidence Rates in the USA. *Digestive Diseases and Sciences* 2020.
14. White DL, Thrift AP, Kanwal F, et al. Incidence of Hepatocellular Carcinoma in All 50 United States, From 2000 Through 2012. *Gastroenterology* 2017; 152:812–820.e5. [PubMed: 27889576]
15. Setiawan VW, Wei PC, Hernandez BY, et al. Disparity in liver cancer incidence and chronic liver disease mortality by nativity in Hispanics: The Multiethnic Cohort. *Cancer* 2016;122:1444–52. [PubMed: 26916271]
16. Shebl FM, Capo-Ramos DE, Graubard BI, et al. Socioeconomic status and hepatocellular carcinoma in the United States. *Cancer Epidemiology and Prevention Biomarkers* 2012;21:1330–1335.
17. Danos D, Leonardi C, Gilliland A, et al. Increased Risk of Hepatocellular Carcinoma Associated With Neighborhood Concentrated Disadvantage. *Front Oncol* 2018;8:375. [PubMed: 30254987]
18. Yang B, Liu JB, So SK, et al. Disparities in hepatocellular carcinoma incidence by race/ethnicity and geographic area in California: Implications for prevention. *Cancer* 2018;124:3551–3559. [PubMed: 30113700]
19. Cook MB, Dawsey SM, Freedman ND, et al. Sex disparities in cancer incidence by period and age. *Cancer Epidemiol Biomarkers Prev* 2009; 18:1174–82. [PubMed: 19293308]
20. McGlynn KA, Petrick JL, El-Serag HB. Epidemiology of Hepatocellular Carcinoma. *Hepatology*; n/a.
21. Mittal S, El-Serag HB. Epidemiology of HCC: consider the population. *Journal of clinical gastroenterology* 2013;47:S2. [PubMed: 23632345]
22. Zhang X, El-Serag HB, Thrift AP. Sex and Race Disparities in the Incidence of Hepatocellular Carcinoma in the United States Examined through Age-Period-Cohort Analysis. *Cancer Epidemiology Biomarkers & Prevention* 2019.
23. Welzel TM, Graubard BI, Quraishi S, et al. Population-attributable fractions of risk factors for hepatocellular carcinoma in the United States. *Am J Gastroenterol* 2013; 108:1314–21. [PubMed: 23752878]
24. Naugler WE, Sakurai T, Kim S, et al. Gender disparity in liver cancer due to sex differences in MyD88-dependent IL-6 production. *Science* 2007;317:121–124. [PubMed: 17615358]
25. Kanda T, Jiang X, Yokosuka O. Androgen receptor signaling in hepatocellular carcinoma and pancreatic cancers. *World Journal of Gastroenterology: WJG* 2014;20:9229. [PubMed: 25071315]
26. Yip TC-F, Wong GL-H, Chan HL-Y, et al. Elevated testosterone increases risk of hepatocellular carcinoma in men with chronic hepatitis B and diabetes mellitus. *Journal of Gastroenterology and Hepatology* 2020;n/a.
27. Hassan MM, Botrus G, Abdel-Wahab R, et al. Estrogen Replacement Reduces Risk and Increases Survival Times of Women With Hepatocellular Carcinoma. *Clinical Gastroenterology and Hepatology* 2017;15:1791–1799. [PubMed: 28579181]
28. Petrick JL, Florio AA, Zhang X, et al. Associations between Prediagnostic Concentrations of Circulating Sex Steroid Hormones and Liver Cancer Among Post-Menopausal Women. *Hepatology*; n/a.
29. Mittal S, El-Serag HB. Epidemiology of hepatocellular carcinoma: consider the population. *J Clin Gastroenterol* 2013;47 Suppl:S2–6. [PubMed: 23632345]

30. Makarova-Rusher OV, Altekrouse SF, McNeel TS, et al. Population attributable fractions of risk factors for hepatocellular carcinoma in the United States. *Cancer* 2016;122:1757–65. [PubMed: 26998818]
31. Younossi ZM. Non-alcoholic fatty liver disease – A global public health perspective. *Journal of Hepatology* 2019;70:531–544. [PubMed: 30414863]
- *32. Rich NE, Oji S, Mufti AR, et al. Racial and Ethnic Disparities in Nonalcoholic Fatty Liver Disease Prevalence, Severity, and Outcomes in the United States: A Systematic Review and Meta-analysis. *Clin Gastroenterol Hepatol* 2018;16:198–210.e2. [PubMed: 28970148] A large meta-analysis describing disparities in NAFLD/NASH., with the highest prevalence and severity in Hispanics compared to non-Hispanics.
33. Wu EM, Wong LL, Hernandez BY, et al. Gender differences in hepatocellular cancer: disparities in nonalcoholic fatty liver disease/steatohepatitis and liver transplantation. *Hepatoma Res* 2018;4. [PubMed: 29479565]
- **34. Rich NE, Murphy CC, Yopp AC, et al. Sex disparities in presentation and prognosis of 1110 patients with hepatocellular carcinoma. *Alimentary Pharmacology & Therapeutics* 2020;52:701–709. [PubMed: 32598091] Large cohort study describing sex disparities in HCC prognosis at two large US health systems; demonstrated women with HCC have better overall survival compared to men even after adjusting for tumor stage and liver disease severity.
35. Ladenheim MR, Kim NG, Nguyen P, et al. Sex differences in disease presentation, treatment and clinical outcomes of patients with hepatocellular carcinoma: a single-centre cohort study. *BMJ Open Gastroenterology* 2016;3:e000107.
36. Singal AG, Pillai A, Tiro J. Early detection, curative treatment, and survival rates for hepatocellular carcinoma surveillance in patients with cirrhosis: a meta-analysis. *PLoS Med* 2014;11:e1001624. [PubMed: 24691105]
37. Wolf E, Rich NE, Marrero JA, et al. Utilization of hepatocellular carcinoma surveillance in patients with cirrhosis: A systematic review and meta-analysis. *Hepatology*;n/a.
38. Singal AG, Yopp A, Skinner CS, et al. Utilization of hepatocellular carcinoma surveillance among American patients: a systematic review. *Journal of general internal medicine* 2012;27:861–867. [PubMed: 22215266]
39. Singal AG, Tiro J, Li X, et al. Hepatocellular Carcinoma Surveillance Among Patients With Cirrhosis in a Population-based Integrated Health Care Delivery System. *J Clin Gastroenterol* 2017;51:650–655. [PubMed: 27870642]
40. Warren Andersen S, Blot WJ, Lipworth L, et al. Association of Race and Socioeconomic Status With Colorectal Cancer Screening, Colorectal Cancer Risk, and Mortality in Southern US Adults. *JAMA Network Open* 2019;2:e1917995–e1917995. [PubMed: 31860105]
41. Singal AG, Li X, Tiro J, et al. Racial, Social, and Clinical Determinants of Hepatocellular Carcinoma Surveillance. *The American Journal of Medicine* 2015;128:90.e1–90.e7.
42. Farvardin S, Patel J, Khambaty M, et al. Patient-reported barriers are associated with lower hepatocellular carcinoma surveillance rates in patients with cirrhosis. *Hepatology (Baltimore, Md.)* 2017;65:875–884.
43. Llovet JM, Fuster J, Bruix J. The Barcelona approach: diagnosis, staging, and treatment of hepatocellular carcinoma. *Liver transplantation* 2004;10:S115–S120. [PubMed: 14762851]
44. Momin BR, Pinheiro PS, Carreira H, et al. Liver cancer survival in the United States by race and stage (2001–2009): Findings from the CONCORD-2 study. *Cancer* 2017;123 Suppl 24:5059–5078. [PubMed: 29205306]
45. Ha J, Yan M, Aguilar M, et al. Race/ethnicity-specific disparities in hepatocellular carcinoma stage at diagnosis and its impact on receipt of curative therapies. *Journal of clinical gastroenterology* 2016;50:423–430. [PubMed: 26583267]
- **46. Rich NE, Hester C, Odewole M, et al. Racial and Ethnic Differences in Presentation and Outcomes of Hepatocellular Carcinoma. *Clin Gastroenterol Hepatol* 2019;17:551–559 e1. [PubMed: 29859983] Large cohort study conducted at two urban health systems demonstrating Black patients with HCC suffer worse outcomes compared to Non-Hispanic Black and Non-Hispanic White counterparts.

47. Jones PD, Diaz C, Wang D, et al. The Impact of Race on Survival After Hepatocellular Carcinoma in a Diverse American Population. *Digestive Diseases & Sciences*;63:515–528. [PubMed: 29275448]
48. Davila JA, Morgan RO, Richardson PA, et al. Use of surveillance for hepatocellular carcinoma among patients with cirrhosis in the United States. *Hepatology* 2010;52:132–41. [PubMed: 20578139]
49. Tan D, Yopp A, Beg MS, et al. Meta-analysis: underutilisation and disparities of treatment among patients with hepatocellular carcinoma in the United States. *Alimentary pharmacology & therapeutics* 2013;38:703–712. [PubMed: 23957569]
50. Shah SA, Smith JK, Li Y, et al. Underutilization of therapy for hepatocellular carcinoma in the medicare population. *Cancer* 2011;117:1019–1026. [PubMed: 20945363]
51. Yu JC, Neugut AI, Wang S, et al. Racial and insurance disparities in the receipt of transplant among patients with hepatocellular carcinoma. *Cancer* 2010;116:1801–1809. [PubMed: 20143441]
52. Sonnenday CJ, Dimick JB, Schulick RD, et al. Racial and Geographic Disparities in the Utilization of Surgical Therapy for Hepatocellular Carcinoma. *Journal of Gastrointestinal Surgery* 2007;11:1636–1646. [PubMed: 17912593]
53. Sobotka LA, Hinton A, Conteh LF. African Americans are less likely to receive curative treatment for hepatocellular carcinoma. *World journal of hepatology* 2018;10:849–855. [PubMed: 30533185]
- *54. Rich NE, Hester C, Odewole M, et al. Racial and Ethnic Differences in Presentation and Outcomes of Hepatocellular Carcinoma. *Clinical Gastroenterology and Hepatology* 2019;17:551–559.e1. [PubMed: 29859983] Recent large, multicenter study demonstrating racial disparities in liver transplant for HCC.
55. Dakhoul L, Gawrieh S, Jones KR, et al. Racial Disparities in Liver Transplantation for Hepatocellular Carcinoma Are Not Explained by Differences in Comorbidities, Liver Disease Severity, or Tumor Burden. *Hepatology Communications* 2019;3:52–62. [PubMed: 30619994]
56. Mathur AK, Schaubel DE, Gong Q, et al. Racial and ethnic disparities in access to liver transplantation. *Liver Transpl* 2010;16:1033–40. [PubMed: 20818740]
57. Alawadi ZM, Phatak UR, Kao LS, et al. Race not rural residency is predictive of surgical treatment for hepatocellular carcinoma: Analysis of the Texas Cancer Registry. *Journal of Surgical Oncology* 2016;113:84–88. [PubMed: 26696033]
58. Stewart SL, Kwong SL, Bowlus CL, et al. Racial/ethnic disparities in hepatocellular carcinoma treatment and survival in California, 1988–2012. *World J Gastroenterol* 2016;22:8584–8595. [PubMed: 27784971]
59. Cauble S, Abbas A, Balart L, et al. United States women receive more curative treatment for hepatocellular carcinoma than men. *Digestive diseases and sciences* 2013;58:2817–2825. [PubMed: 23812858]
60. Yang D, Hanna DL, Usher J, et al. Impact of sex on the survival of patients with hepatocellular carcinoma: a Surveillance, Epidemiology, and End Results analysis. *Cancer* 2014;120:3707–16. [PubMed: 25081299]
61. Sobotka L, Hinton A, Conteh L. Women receive more inpatient resections and ablations for hepatocellular carcinoma than men. *World J Hepatol* 2017;9:1346–1351. [PubMed: 29359018]
62. Sarkar M, Watt KD, Terrault N, et al. Outcomes in liver transplantation: does sex matter? *Journal of Hepatology*;62:946–55. [PubMed: 25433162]
63. Mathur AK, Schaubel DE, Gong Q, et al. Sex-based disparities in liver transplant rates in the United States. *Am J Transplant* 2011;11:1435–43. [PubMed: 21718440]
64. Halbert CH, Weathers B, Delmoor E, et al. Racial differences in medical mistrust among men diagnosed with prostate cancer. *Cancer* 2009;115:2553–61. [PubMed: 19296516]
65. Karliner LS, Hwang ES, Nickleach D, et al. Language barriers and patient-centered breast cancer care. *Patient Educ Couns* 2011;84:223–8. [PubMed: 20685068]
66. Abbott DE, Voils CL, Fisher DA, et al. Socioeconomic disparities, financial toxicity, and opportunities for enhanced system efficiencies for patients with cancer. *J Surg Oncol* 2017;115:250–256. [PubMed: 28105638]

67. Mokdad AA, Murphy CC, Pruitt SL, et al. Effect of hospital safety net designation on treatment use and survival in hepatocellular carcinoma. *Cancer* 2018;124:743–751. [PubMed: 29072773]
68. Hoehn RS, Hanseman DJ, Dhar VK, et al. Opportunities to Improve Care of Hepatocellular Carcinoma in Vulnerable Patient Populations. *Journal of the American College of Surgeons*;224:697–704. [PubMed: 28069526]
69. Wang J, Ha J, Lopez A, et al. Medicaid and Uninsured Hepatocellular Carcinoma Patients Have More Advanced Tumor Stage and Are Less Likely to Receive Treatment. *J Clin Gastroenterol* 2018;52:437–443. [PubMed: 28723861]
70. Rogers SO, Ray WA, Smalley WE. A population-based study of survival among elderly persons diagnosed with colorectal cancer: does race matter if all are insured? (United States). *Cancer Causes & Control* 2004;15:193–199. [PubMed: 15017132]
71. Zeng C, Wen W, Morgans AK, et al. Disparities by Race, Age, and Sex in the Improvement of Survival for Major Cancers: Results From the National Cancer Institute Surveillance, Epidemiology, and End Results (SEER) Program in the United States, 1990 to 2010. *JAMA Oncology* 2015; 1:88–96. [PubMed: 26182310]
72. Hester CA, Karbhari N, Rich NE, et al. Effect of fragmentation of cancer care on treatment use and survival in hepatocellular carcinoma. *Cancer* 2019;125:3428–3436. [PubMed: 31299089]
73. Artinyan A, Mailey B, Sanchez-Luege N, et al. Race, ethnicity, and socioeconomic status influence the survival of patients with hepatocellular carcinoma in the United States. *Cancer*; 116:1367–77.
74. Mathur AK, Osborne NH, Lynch RJ, et al. Racial/ethnic disparities in access to care and survival for patients with early-stage hepatocellular carcinoma. *Archives of Surgery*; 145:1158–63.
75. Njei B, Ditah I, Lim JK. Persistent racial disparities in survival among u.s. Adults with hepatocellular carcinoma after liver transplantation: the paradox of all-cause and cause-specific mortality. *Gastrointest Cancer Res* 2013;6:73–4. [PubMed: 23936546]
76. Cook MB, McGlynn KA, Devesa SS, et al. Sex disparities in cancer mortality and survival. *Cancer Epidemiol Biomarkers Prev* 2011;20:1629–37. [PubMed: 21750167]
77. Yang D, Hanna DL, Usher J, et al. Impact of sex on the survival of patients with hepatocellular carcinoma: a Surveillance, Epidemiology, and End Results analysis. *Cancer*;120:3707–16. [PubMed: 25081299]
78. Rich NE, John BV, Parikh ND, et al. Hepatocellular carcinoma demonstrates heterogeneous growth patterns in a multi-center cohort of patients with cirrhosis. *Hepatology*;n/a.
79. Nathani P, Gopal P, Rich N, et al. Hepatocellular carcinoma tumour volume doubling time: a systematic review and meta-analysis. *Gut* 2020:gutjnl-2020-321040.
80. Williams DR, Kontos EZ, Viswanath K, et al. Integrating Multiple Social Statuses in Health Disparities Research: The Case of Lung Cancer. *Health Services Research* 2012;47:1255–1277. [PubMed: 22568674]
- **81. Alcaraz KI, Wiedt TL, Daniels EC, et al. Understanding and addressing social determinants to advance cancer health equity in the United States: A blueprint for practice, research, and policy. *CA: A Cancer Journal for Clinicians* 2020;70:31–46. [PubMed: 31661164] Provides a framework to improve our understanding of social determinants of health to improve health equity for all patients with cancer; presents actionable recommendations for research health policy.
82. Paradies Y, Ben J, Denson N, et al. Racism as a Determinant of Health: A Systematic Review and Meta-Analysis. *PLoS One* 2015;10:e0138511. [PubMed: 26398658]
83. Penner LA, Dovidio JF, Gonzalez R, et al. The Effects of Oncologist Implicit Racial Bias in Racially Discordant Oncology Interactions. *Journal of Clinical Oncology* 2016;34:2874–2880. [PubMed: 27325865]
84. Obermeyer Z, Powers B, Vogeli C, et al. Dissecting racial bias in an algorithm used to manage the health of populations. *Science* 2019;366:447–453. [PubMed: 31649194]
85. Vyas DA, Eisenstein LG, Jones DS. Hidden in Plain Sight — Reconsidering the Use of Race Correction in Clinical Algorithms. *New England Journal of Medicine* 2020.
86. Levey AS, Tighiouart H, Titan SM, et al. Estimation of Glomerular Filtration Rate With vs Without Including Patient Race. *JAMA Intern Med* 2020;180:793–5. [PubMed: 32176270]
- *87. Verna EC, Lai JC. Time for Action to Address the Persistent Sex-Based Disparity in Liver Transplant Access. *JAMA Surgery* 2020;155:545–547. [PubMed: 32432664] Timely editorial on

the need to reduce disparities in access to life-saving treatments for patients with advanced liver disease and HCC, such as liver transplantation.

88. Locke JE, Shelton BA, Olthoff KM, et al. Quantifying Sex-Based Disparities in Liver Allocation. *JAMA Surgery* 2020;155:e201129–e201129. [PubMed: 32432699]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript