

RESEARCH ARTICLE

Clusterin levels are increased in Alzheimer's disease and influence the regional distribution of A β

J. Scott Miners; Polly Clarke; Seth Love

Dementia Research Group, Institute of Clinical Neurosciences, School of Clinical Sciences, University of Bristol, United Kingdom.

Keywords

Alzheimer's disease, amyloid- β , amyloid- β clearance, apoJ, cerebral amyloid angiopathy, clusterin, plaque.

Corresponding author:

Seth Love, PhD, FRCP, FRCPath, School of Clinical Sciences, University of Bristol, Learning and Research level 2, Southmead Hospital, Bristol BS10 5NB, UK
(E-mail: seth.love@bris.ac.uk)

Received 01 April 2016

Accepted 19 May 2016

Published Online Article Accepted

01 June 2016

doi:10.1111/bpa.12392

Abstract

Clusterin, also known as apoJ, is a lipoprotein abundantly expressed within the CNS. It regulates A β fibril formation and toxicity and facilitates amyloid- β (A β) transport across the blood-brain barrier. Genome-wide association studies have shown variations in the clusterin gene (*CLU*) to influence the risk of developing sporadic Alzheimer's disease (AD). To explore whether clusterin modulates the regional deposition of A β , we measured levels of soluble (NP40-extracted) and insoluble (guanidine-HCl-extracted) clusterin, A β 40 and A β 42 by sandwich ELISA in brain regions with a predilection for amyloid pathology—mid-frontal cortex (MF), cingulate cortex (CC), parahippocampal cortex (PH), and regions with little or no pathology—thalamus (TH) and white matter (WM). Clusterin level was highest in regions with plaque pathology (MF, CC, PH and PC), approximately mirroring the regional distribution of A β . It was significantly higher in AD than controls, and correlated positively with A β 42 and insoluble A β 40. Soluble clusterin level rose significantly with severity of cerebral amyloid angiopathy, and in MF and PC regions was highest in *APOE* ϵ 4 homozygotes. In the TH and WM (areas with little amyloid pathology) clusterin was unaltered in AD and did not correlate with A β level. There was a significant positive correlation between the concentration of clusterin and the regional levels of insoluble A β 42; however, the molar ratio of clusterin : A β 42 declined with insoluble A β 42 level in a region-dependent manner, being lowest in regions with predilection for A β plaque pathology. Under physiological conditions, clusterin reduces aggregation and promotes clearance of A β . Our findings indicate that in AD, clusterin increases, particularly in regions with most abundant A β , but because the increase does not match the rising level of A β 42, the molar ratio of clusterin : A β 42 in those regions falls, probably contributing to A β deposition within the tissue.

INTRODUCTION

Alzheimer's disease (AD) is believed to be initiated by the accumulation and aggregation of amyloid- β (A β) peptides [the so-called amyloid cascade hypothesis (1)]. The steady-state level of A β reflects the balance between its production and removal from the brain (2). A β peptides are produced by sequential cleavage of A β protein precursor (APP) and mostly end at amino acid 40 or 42. A β 42 is the more amyloidogenic form—relatively insoluble in the interstitial fluid and prone to parenchymal deposition. A β 40 is more soluble, less prone to parenchymal deposition but more likely to accumulate in the walls of cortical and leptomeningeal blood vessels (3, 4). Most mutations in familial AD are associated with increased amyloidogenic processing of APP and elevated A β 42 or an increase in the A β 42 : A β 40 ratio (5–8). In sporadic AD, which accounts for most cases, the accumulation of A β is thought largely to reflect alterations in the pathways responsible for the removal of A β [reviewed in (2)] or altered expression of chaperone proteins, such as apoE and clusterin (also known as apoJ) that regulate the structure, toxicity and clearance of A β [reviewed in (9)].

A β peptides are produced throughout life (10, 11) but begin to accumulate and aggregate in the brain more than a decade before the onset of AD (12, 13). Risk factors for AD, such as ageing and *APOE* genotype (14), accelerate the parenchymal deposition of A β . The deposition of A β within the brain follows a hierarchical sequence first appearing in the neocortex and spreading to limbic areas, deep cerebral gray matter and brain stem regions and finally the cerebellum (15). The determinants of regional variability in the susceptibility of different brain regions to A β deposition remain unclear. No link was found between the distribution of plaque pathology and the regional distribution of enzymes involved in the amyloidogenic processing of APP (APP, APP-CRFB, BACE-1 and PS-1; (16–19)). However, Shinohara *et al* (19) found a strong inverse relationship between apolipoprotein E (apoE) level and A β deposition in brain tissue from cognitively normal elderly people and those with mild cognitive impairment (MCI). The authors suggested that apoE had a role in preventing A β accumulation and was reduced in brain regions that would later develop significant plaque pathology.

Clusterin, also known as apolipoprotein-J, is a 78–80 kDa heterodimeric glycoprotein that is abundantly expressed in the CNS (20). Genome-wide association studies have identified several single nucleotide polymorphisms within the clusterin gene (*CLU*) that are risk factors for AD (21–24). Clusterin is up-regulated in the brain in AD (25, 26) and is present in plaques (27, 28). *In vitro* studies suggest that clusterin influences A β fibril formation and neurotoxicity (reviewed in (29, 30)) and can facilitate the transport of A β across the blood-brain barrier (31). We have undertaken a comprehensive analysis of the regional distribution of clusterin, soluble and insoluble A β 40 and A β 42 in post-mortem brain tissue across a number of brain regions that vary in their predilection to amyloid pathology. Our findings indicate that clusterin level rises with the accumulation of insoluble A β 42 but the molar ratio of clusterin : A β 42 falls, which probably influences the regional distribution of A β deposition.

MATERIALS AND METHODS

Case selection

Brain tissue was obtained from the South West Dementia Brain Bank (SWDBB), University of Bristol, UK with local research ethics committee approval. All brains had been retrieved within 72 h of death. The right cerebral hemisphere had been fixed in 10% formalin for three weeks before the tissue was processed and paraffin blocks were taken for pathological assessment. The left cerebral hemisphere had been sliced and frozen at -80°C until used for biochemical assessment. According to the NIA-AA guidelines AD neuropathological change was considered an adequate explanation for the dementia in all cases in the AD group (32). Controls were defined by an absence of clinical history of cognitive decline or other neurological disease and a lack of neuropathological abnormalities apart from sparse neuritic or diffuse plaques in some of the older cases, all of which were of Braak tangle stage III or lower. *APOE* genotyping and assessment of severity of cerebral amyloid angiopathy (CAA) had been performed as previously reported (33, 34). Demographic information, neuropathological findings and MRC identifiers for each case are shown in Supporting Information Tables S1 and S2. β

Brain tissue

Brain tissue (200 mg) was dissected from the midfrontal, cingulate, parahippocampal and medial parietal cortex, thalamus (pulvinar) and white matter (WM) underlying the parietal cortex. Brain tissue samples were prepared using a Precellys 24 homogenizer (Stretton Scientific, Derbyshire, UK) with 2.3 mm ceramic beads (Biospec, Stratech, Suffolk, UK) as previously described for A β measurements in human postmortem tissue (10, 11, 35, 36). Soluble and insoluble extracts were prepared sequentially following initial homogenization in 1% NP-40 buffer containing 140 mM NaCl, 3 mM KCl, 25 mM TRIS, 5 mM ethylenediaminetetraacetic acid (EDTA) and 2 mM 1,10 phenanthroline. The homogenates were spun at $13\,000 \times g$ for 15 min at 4°C and the supernatant was removed and stored at -80°C . Insoluble extracts were prepared by homogenization of pelleted insoluble material in 6 M GuHCl and were left for 4 h at room temperature (RT) before storage at -80°C .

Measurement of clusterin levels by sandwich ELISA

Clusterin level was measured by sandwich ELISA (duoset kit # DY5874, R&D systems, Oxford, UK) according to the manufacturer's guidelines; 96-well Maxisorp plates (R&D systems, Oxford, UK) were coated at RT overnight with mouse anti-human clusterin. We washed the plates in phosphate-buffered saline (PBS)/tween-20 (0.01%), added 1% PBS/BSA at RT for 1 h to block nonspecific binding, then added recombinant human clusterin (62.5–4000 pg/mL) and tissue homogenates (2.5 μL supernatant diluted in 3.125 mL PBS, and 1.8 μL insoluble extract diluted in 10 mL PBS) for 2 h at RT. After a further wash, the plate was incubated for 2 h at RT with biotinylated mouse anti-human clusterin. The plate was again washed and incubated with streptavidin-horseradish peroxidase (HRP; 1:200 in 0.01% PBS : Tween-20) for 20 minute at RT in the dark, washed, and incubated for 10 minute with chromogenic substrate (TMBS, R&D systems, Oxford, UK). Absorbance was read at 450 nm in a FLUOstar Optima plate reader (BMG Labtech, Aylesbury, UK) after the addition of 50 μL of 2 N sulfuric acid. Measurements were repeated in duplicate and across two plates to ensure that there was minimal plate-to-plate variation.

Measurement of A β 40 and A β 42 by sandwich ELISA

We measured A β 40 and A β 42 level in both soluble (NP1-40) and insoluble (guanidine-HCl-extracted) brain tissue fractions by sandwich ELISA as previously described (10, 11, 34–39). For the A β 40 ELISA, mouse monoclonal anti-human A β (clone 6E10, raised against amino acids 1–16; Covance, Harrogate, UK), 2 $\mu\text{g}/\text{mL}$ in PBS, was incubated overnight at RT, washed and then blocked with 300 μL protein-free PBS blocking buffer (Thermo Fisher Scientific, Loughborough, UK) for 2 h at RT. Samples of brain homogenate (diluted 1:49 for guanidine extracts and 1:3 for soluble extracts) or recombinant human A β 1-40 (Sigma Aldrich, Dorset, UK) diluted in PBS containing 1% 1,10 phenanthroline (Sigma Aldrich, Dorset, UK) to prevent degradation of A β (40), were incubated for 2 h at RT on a rocking platform. After a further wash, the plates were incubated for 2 h at RT with mouse anti-human A β 1-40 (1 $\mu\text{g}/\text{mL}$; 11A50-B10; Covance, Harrogate, UK) that had been biotinylated using Lightning-Link Biotinylation Kit (Innova Biosciences, Cambridge, UK) according to the manufacturer's guidelines. Streptavidin-HRP (R&D Systems Europe, Abingdon, UK) diluted 1:200 was added to each well for 20 min at RT before they were washed and substrate solution (TMB; R&D Systems Europe, Abingdon, UK) was added for 30 min in the dark. The reaction was stopped with 2N sulfuric acid (R&D Systems Europe, Abingdon, UK) and the optical density of each well read at 450 nm in a FLUOstar plate reader (BMG Labtech, Aylesbury, UK).

For the A β 42 ELISA, anti-human A β 1-42 (12 F4, Covance) diluted 0.5 $\mu\text{g}/\text{mL}$ in PBS was used as the capture antibody. Tissue samples (insoluble extracts diluted 1:9, soluble extracts diluted 1:3) were incubated at RT for 4 h. Biotinylated anti-human A β (Thermo Fisher Scientific) diluted to 0.1 $\mu\text{g}/\text{mL}$ in PBS was used for detection and left overnight at 4°C . After washing, streptavidin-HRP was added for 1 h and chromogenic substrate for 20 min in the dark after a further wash. A β 1-42 concentration in brain tissue was




Figure 1. Bar charts showing regional levels of A β level in control (black bars) and AD brains (red bars), in MF, CC, PH, medial parietal cortex (PC), thalamus (TH) and parietal WM. Bars indicate the mean and SEM. * $P < 0.05$ ** $P < 0.01$ *** $P < 0.001$ **** $P < 0.0001$. Lines

indicate significant differences between regions, in the controls (black lines) and AD groups (red lines); the thickness of the line indicates the significance of the difference between the two regions, ranging from $P < 0.01$ to $P < 0.0001$.

interpolated from a standard curve generated by serial dilution (16 000 to 1.024 nM) of recombinant human A β 1–42 (Sigma Aldrich). Each sample was assayed in duplicate. The A β 1–42 ELISA did not detect A β 1–40, and the A β 1–40 ELISA did not detect A β 1–42.

Statistical analysis

Unpaired two-tailed t-test or ANOVA with Dunnett's *post hoc* analysis was used for comparisons between groups, and Pearson's or Spearman's test was used to assess linear or rank order correlation, as appropriate, with the help of SPSS version 16 (SPSS, Chicago) and GraphPad Prism version 6 (GraphPad Software, La Jolla, CA). P -values < 0.05 were considered statistically significant.

RESULTS

Regional distribution of soluble and insoluble A β 40 and A β 42 in AD

We examined the regional distribution of soluble (NP-40-soluble) and insoluble (after guanidine-HCl extraction) A β 40 and A β 42 in sequentially extracted brain homogenates from the following regions in AD and age-matched controls: midfrontal cortex (MF), cingulate cortex (CC), parahippocampal cortex (PH), medial parietal cortex (PC), thalamus (TH) and parietal WM (Figure 1). In CC,

MF, PH and PC, the concentrations of insoluble A β 40 and A β 42 were significantly higher in AD than age-matched controls. The differences between AD and control brains in TH and WM did not reach significance.

In both control and AD groups, the concentration of insoluble A β 40 and A β 42 tended to decrease in the following order: CC > MF > PH > PC > TH and WM (Figure 1A,B). In AD, the concentration of insoluble A β 40 and A β 42 was significantly higher in neocortex (MF and CC) than in other regions (Figure 1A,B).

The regional distribution of soluble A β 40 and A β 42 differed substantially from that of insoluble A β 40 and A β 42. The concentration of the soluble forms of A β was lowest in MF and CC and tended to be higher in PC, PH, TH and WM (Figure 1C,D). Soluble A β level did not differ to a statistically significant extent between AD and controls, with the exception of increased A β 42 in AD within WM. In general the concentration of A β in gray matter regions was much lower in the soluble than the insoluble tissue fractions. In contrast, A β 40 and A β 42 concentration was higher in the soluble than the insoluble tissue fractions of WM, and soluble A β level was several-fold higher in WM than cortex. Soluble A β 42 level was also relatively high in PH.

The relative contribution of soluble and insoluble A β 40 and A β 42 to "total" A β load in all regions studied is shown in Supporting Information Figure S1. "Total" A β was highest in MF and CC and tended to decrease in PH > PC > TH. Total A β in MF and CC




Figure 2. Bar chart showing regional levels of clusterin level in control (black bars) and AD brains (red bars), in the soluble and insoluble brain tissue fractions. Bars indicate the mean and SEM. * $P < 0.05$ ** $P < 0.01$ *** $P < 0.001$. Lines indicate significant differences between regions, in the controls (black lines) and AD groups (red lines); the thickness of the solid lines indicates the significance of the difference between the two regions, ranging from $P < 0.01$ to $P < 0.0001$. The interrupted horizontal lines indicate differences significant at the $P < 0.5$ level.

consisted almost entirely of insoluble A β 40 and A β 42. In contrast, in WM, most of the A β consisted of soluble A β 40 with a small amount of insoluble A β 40 and negligible A β 42.

Regional distribution of clusterin in AD

We examined the regional distribution of clusterin in the same soluble and insoluble brain fractions in AD and age-matched controls that we had used to measure A β levels (Figure 2). In AD, clusterin level within both the soluble and insoluble extracts was highest in CC, MF and PH and PC and lowest in the TH and WM (approximately mirroring the regional distribution of “total” A β). The level of soluble clusterin was less variable between regions in the controls but was significantly higher in CC than PC or TH (Figure 2A). Clusterin level in the insoluble extract was significantly higher in all gray matter regions than in the WM (Figure 2A).

The level of soluble clusterin was significantly higher in AD than controls in most regions (CC, MF, PH and PC; Figure 2A) and was increased in the insoluble extract in MF and PH (Figure 2B). Clusterin level within the soluble and insoluble extracts correlated significantly with soluble and insoluble A β 42 and A β 40 in MF (with the exception of insoluble A β 42; Table 1). A similar trend was observed between clusterin and A β levels within the

soluble extract in the PC but not the insoluble extract. A strong correlation was observed in the CC between clusterin and soluble and insoluble A β 40. There was less correlation between clusterin and A β in the TH and WM (Table 1).

Regional association of clusterin and a β

To assess whether variations in clusterin concentration might influence the regional distribution of soluble and insoluble A β 40 or A β 42, we looked at the correlation between A β level and clusterin concentration across all regions. There was a significant correlation between the concentration of soluble clusterin and the level of insoluble A β 42 within the AD cohort and a weaker, nonsignificant trend in the controls (Figure 3A,B). A trend approaching significance was observed between clusterin in the insoluble extract and insoluble A β 42 in both controls and AD (Figure 3C,D). We did not find significant correlations between clusterin concentration and the level of soluble A β 42, soluble A β 40 or insoluble A β 40.

To investigate further whether clusterin might promote the accumulation of insoluble A β 42, we calculated the molar ratio of insoluble clusterin to insoluble A β 42 in the different regions. In both controls and AD, the molar ratio of insoluble clusterin : insoluble A β 42 was lowest in regions with the highest concentration of insoluble A β 42 (Figure 4A,B) and *vice versa*.

Clusterin levels influenced by APOE genotype

Clusterin level was highest in *APOE* $\epsilon 4$ homozygotes in MF and PC (Figure 5) and rose significantly with severity of CAA. *Post-hoc* analysis revealed significantly higher clusterin level in $\epsilon 4/4$ than $\epsilon 3/3$ brains in PC ($P < 0.05$), and in $\epsilon 4/4$ than $\epsilon 3/4$ ($P < 0.01$) or $\epsilon 3/3$ ($P < 0.05$) in MF. *Post-hoc* analysis also showed clusterin level in MF and PC to be significantly higher in brains with severe than absent CAA ($P < 0.05$ for both regions). Insoluble clusterin level did not vary significantly in relation to *APOE* genotype or CAA severity.

DISCUSSION

We have examined the relationship between clusterin/apoJ level and the regional distribution of A β within the brain. Although the concentration of clusterin was elevated in AD and was highest in cortical regions with the most abundant A β deposition, the molar ratio of clusterin : A β was lowest in those regions and was highest in parts of the brain with little or no amyloid pathology, such as in the thalamus and WM. These findings in human brain tissue support experimental studies indicating that (i) clusterin level rises in association with increasing A β , (ii) within the physiological range of clusterin : A β , clusterin reduces aggregation and promotes clearance of A β , but (iii) when, despite a rise in clusterin, A β level increases to an extent that causes clusterin : A β to fall below the physiological range, A β -clusterin complexes tend to aggregate and deposit within the brain parenchyma. We have also shown that clusterin concentration is influenced by *APOE* genotype, being highest in brain tissue from $\epsilon 4$ homozygotes, and rises in relation to the severity of CAA.

Clusterin is highly expressed in the CNS, within which it is present at a similar concentration to that of apoE (20). Variations in the clusterin gene (*CLU*) are associated with sporadic AD (21), and previous studies showed that clusterin is increased in the CSF (41, 42)

Table 1. Correlation between concentrations of clusterin and A β in soluble and insoluble tissue fractions in different regions.

Region		Insoluble A β 42	Soluble A β 42	Insoluble A β 40	Insoluble A β 40
Mid-frontal	Soluble clusterin	$r = 0.359$ $P = 0.048$	$r = 0.507$ $P = 0.008$	$r = 0.496$ $P = 0.005$	$r = 0.354$ $P = 0.090$
	Insoluble clusterin	NS (not significant)	$r = 0.68$ $P = 0.0001$	$r = 0.462$ $P = 0.009$	$r = 0.374$ $P = 0.072$
Cingulate	Soluble clusterin	NS	NS	$r = 0.484$ $P = 0.080$	NS
	Insoluble clusterin	NS	NS	$r = 0.537$ $P = 0.04$	NS
PH	Soluble clusterin	$r = 0.428^*$ $P = 0.018$	NS	NS	NS
	Insoluble clusterin	NS	NS	NS	NS
Parietal cortex	Soluble clusterin	$r = 0.640$ $P < 0.000001$	$r = 0.236$ $P = 0.043$	$r = 0.521$ $P = 0.000002$	NS
	Insoluble clusterin	$r = 0.549$ $P = 0.002$	NS	NS	NS
TH	Soluble clusterin	NS	NS	NS	NS
	Insoluble clusterin	$r = 0.302$ $P = 0.028$	NS	NS	NS
Parietal WM	Soluble clusterin	NS	$r = 0.369$ $P = 0.002$	NS	NS
	Insoluble clusterin	NS	NS	NS	NS

* r = coefficient of correlation

and brain tissue (25, 26) in AD. Clusterin is detectable immunohistochemically within plaques (27, 28) and increases in association with neuritic plaque density (43–45). In a transgenic APP/PS1 mouse model, clusterin level was elevated in plasma and brain tissue and found to co-localize with amyloid plaques (42). Present findings show that the concentration of clusterin in human brain tissue in AD is highest in regions with the greatest concentration of A β . Within those regions, clusterin concentration correlates closely with A β level [as was also shown in transgenic mouse models of AD expressing mutant APP (46)]. These findings are in keeping with clinical evidence of a correlation between raised plasma clusterin level and accelerated clinical progression of disease (41, 42), and imaging studies showing that increased plasma clusterin levels were a strong predictor of brain amyloid load in AD patients (42). Within subcortical regions that have a much lower level of A β , clusterin concentration does not differ significantly between AD and controls.

In vitro studies indicate that clusterin binds to A β and influences both fibril formation and toxicity (47, 48). Yerbury *et al* (49) reported that clusterin coprecipitates with A β as insoluble aggregates when A β is present in large molar excess. We have found the molar ratio of clusterin : A β 42 to be lowest in regions that have the greatest accumulation of A β (almost entirely in the insoluble tissue fraction) even though clusterin levels are highest in those regions. In contrast, within the WM and thalamus, which had the highest ratio of clusterin : A β 42, A β was almost all in a soluble form. It seems that when the clusterin : A β ratio falls low enough, clusterin actually promotes rather than simply fails to prevent the precipitation of A β , as evidenced by *in vitro* data (49). These data are consistent with experimental evidence in PDAPP mice (homozygous for the APP^{V717F} transgene) showing that clusterin stimulates amyloid aggregation when A β is present in excess. PDAPP mice homozygous for knock-out of the clusterin gene have significantly fewer fibrillar A β deposits and dystrophic neurites than PDAPP

mice expressing clusterin (50). The rise in clusterin concentration that occurs with increasing A β seems likely to be a consequence of the latter. Thamsbiety *et al* (42) reported that plasma clusterin level was elevated almost 10 years in advance of fibrillary A β deposition, suggesting that clusterin production is raised at an early stage in the disease process, although we know from other studies that A β starts to accumulate even earlier (51, 52).

ApoE is also highly expressed within the CNS and has been implicated in the pathogenesis of AD. The *APOE* gene is a strong risk factor for sporadic AD and individuals possessing the $\epsilon 4$ allele have more abundant plaque and cerebrovascular deposition of A β and a higher level of this peptide (33, 34, 53). *In vitro* studies demonstrated that apoE interacts with and influences A β fibrillogenesis and clearance (54–58). However, apoE and clusterin play somewhat divergent roles in the progression of AD. In contrast to clusterin, apoE concentration shows a strong inverse correlation with regional A β load (19), and while fibril formation was reduced in clusterin-deficient PDAPP mice (50) it was significantly increased in PDAPP mice deficient in both clusterin and apoE (59). It is of interest that no regional association was found between A β level and molecules involved in APP processing (APP, APP-CTF β , BACE-1 or PS-1) or enzymes involved in A β clearance (neprilysin and insulin-degrading enzyme). Together, these data suggest that the regional distribution of A β is influenced to a greater extent by apoE and clusterin expression than by pathways involved in the production or enzymatic degradation of A β .

Clusterin has also been shown to facilitate the clearance of A β at the blood-brain barrier (31) and is localized not only to plaques but also arterioles and capillaries within the brain (43–45). A recent immunohistochemical study showed that clusterin was associated with vascular A β , particularly A β 40, in CAA (60). Co-localization of clusterin with perivascular A β deposits and our finding of increased clusterin level in relation to CAA severity is supportive




Figure 3. Regional association between clusterin and insoluble A β 42 levels. The concentration of clusterin in soluble (A, B) and insoluble (C, D) brain tissue fractions was plotted against insoluble A β 42 level in each region in controls and AD cases. The solid circles and thin

bars indicate the mean values and SEM for clusterin (horizontal bars) and A β 42 (vertical bars). The thick solid and dotted lines indicate the best-fit linear regression and 95% confidence intervals.

of a role in the perivascular drainage of A β , which is impaired in CAA (61, 62). Craggs *et al* (60) also reported increased clusterin in the frontal WM in cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL), suggest-

ing that clusterin may also accumulate as a consequence of failed perivascular drainage of interstitial fluid.

In conclusion, we have shown that clusterin level is elevated in AD in regions with a predilection for plaque deposition. Yet despite




Figure 4. The ratio of clusterin : insoluble A β 42 was lowest in regions with a predilection for A β 42 deposition. The solid circles and bars indicate the mean values and SEM.




Figure 5. Bar charts showing clusterin level in relation to APOE genotype (A–C) and severity of CAA. Bars indicate the mean and SEM. * $P < 0.05$ ** $P < 0.01$. Clusterin level was highest in APOE ϵ_4 homozygotes in MF and PC and increased with severity of CAA.

that elevation, the molar ratio of clusterin : A β is lowest in those same regions, which is likely to influence the regional distribution of A β by promoting its aggregation and precipitation.

ACKNOWLEDGEMENTS

This work was supported by Alzheimer's Research UK (ART-PG2011-1). The South West Dementia Brain Bank is part of the Brains for Dementia Research program, jointly funded by Alzheimer's Research UK and Alzheimer's Society, and is supported by BRACE (Bristol Research into Alzheimer's and Care of the Elderly) and the Medical Research Council.

REFERENCES

- Hardy J, Allsop D (1991) Amyloid deposition as the central event in the aetiology of Alzheimer's disease. *Trends Pharmacol Sci* **12**:383–388.
- Miners JS, Baig S, Palmer J, Palmer LE, Kehoe PG, Love S (2008) A β -degrading enzymes in Alzheimer's disease. *Brain Pathol* **18**:240–252.
- Gravina SA, Ho L, Eckman CB, Long KE, Otvos L, Jr., Younkin LH, *et al* (1995) Amyloid β protein (A β) in Alzheimer's disease brain. Biochemical and immunocytochemical analysis with antibodies specific for forms ending at A β 40 or A β 42(43). *J Biol Chem* **270**:7013–7016.
- Iwatsubo T, Odaka A, Suzuki N, Mizusawa H, Nukina N, Ihara Y (1994) Visualization of A β 42(43) and A β 40 in senile plaques with end-specific A β monoclonals: Evidence that an initially deposited species is A β 42(43). *Neuron* **13**:45–53.
- Price DL, Sisodia SS (1998) Mutant genes in familial Alzheimer's disease and transgenic models. *Annu Rev Neurosci* **21**:479–505.
- Scheuner D, Eckman C, Jensen M, Song X, Citron M, Suzuki N, *et al* (1996) Secreted amyloid β -protein similar to that in the senile plaques of Alzheimer's disease is increased in vivo by the presenilin 1 and 2 and APP mutations linked to familial Alzheimer's disease. *Nat Med* **2**:864–870.
- Selkoe DJ (1998) The cell biology of β -amyloid precursor protein and presenilin in Alzheimer's disease. *Trends Cell Biol* **8**:447–453.
- Wolfe MS (2007) When loss is gain: Reduced presenilin proteolytic function leads to increased A β 42/A β 0. Talking Point on the role of presenilin mutations in Alzheimer disease. *EMBO Rep* **8**:136–140.
- Holtzman DM (2004) In vivo effects of ApoE and clusterin on amyloid- β metabolism and neuropathology. *J Mol Neurosci* **23**:247–254.
- van Helmond Z, Miners JS, Kehoe PG, Love S (2010) Higher soluble amyloid β concentration in frontal cortex of young adults than in normal elderly or Alzheimer's disease. *Brain Pathol* **20**:787–793.

11. Miners JS, Jones R, Love S (2014) Differential changes in A β 42 and A β 40 with age. *J Alzheimers Dis* **40**:727–735.
12. Ballard C, Gauthier S, Corbett A, Brayne C, Aarsland D, Jones E (2011) Alzheimer's disease. *Lancet* **377**:1019–1031.
13. Jack CR, Jr., Knopman DS, Jagust WJ, Shaw LM, Aisen PS, Weiner MW, *et al* (2010) Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *Lancet Neurol* **9**:119–128.
14. Morris JC, Roe CM, Xiong C, Fagan AM, Goate AM, Holtzman DM, Mintun MA (2010) *APOE* predicts amyloid- β but not tau Alzheimer pathology in cognitively normal aging. *Ann Neurol* **67**:122–131.
15. Thal DR, Rub U, Orantes M, Braak H (2002) Phases of A β -deposition in the human brain and its relevance for the development of AD. *Neurology* **58**:1791–1800.
16. Benkovic SA, McGowan EM, Rothwell NJ, Hutton M, Morgan DG, Gordon MN (1997) Regional and cellular localization of presenilin-2 RNA in rat and human brain. *Exp Neurol* **145**:555–564.
17. Page K, Hollister R, Tanzi RE, Hyman BT (1996) In situ hybridization analysis of presenilin 1 mRNA in Alzheimer disease and in lesioned rat brain. *Proc Natl Acad Sci U S A* **93**:14020–14024.
18. Takami K, Terai K, Matsuo A, Walker DG, McGeer PL (1997) Expression of presenilin-1 and -2 mRNAs in rat and Alzheimer's disease brains. *Brain Res* **748**:122–130.
19. Shinohara M, Petersen RC, Dickson DW, Bu G (2013) Brain regional correlation of amyloid- β with synapses and apolipoprotein E in non-demented individuals: Potential mechanisms underlying regional vulnerability to amyloid- β accumulation. *Acta Neuropathol* **125**:535–547.
20. Roheim PS, Carey M, Forte T, Vega GL (1979) Apolipoproteins in human cerebrospinal fluid. *Proc Natl Acad Sci U S A* **76**:4646.
21. Bertram L, McQueen MB, Mullin K, Blacker D, Tanzi RE (2007) Systematic meta-analyses of Alzheimer disease genetic association studies: The AlzGene database. *Nat Genet* **39**:17–23.
22. Butler AW, Ng MY, Hamshere ML, Forabosco P, Wroe R, Al-Chalabi A, *et al* (2009) Meta-analysis of linkage studies for Alzheimer's disease—a web resource. *Neurobiol Aging* **30**:1037–1047.
23. Harold D, Abraham R, Hollingworth P, Sims R, Gerrish A, Hamshere ML, *et al* (2009) Genome-wide association study identifies variants at *CLU* and *PICALM* associated with Alzheimer's disease. *Nat Genet* **41**:1088–1093.
24. Lambert JC, Heath S, Even G, Campion D, Sleegers K, Hiltunen M, *et al* (2009) Genome-wide association study identifies variants at *CLU* and *CR1* associated with Alzheimer's disease. *Nat Genet* **41**:1094–1099.
25. Bertrand P, Poirier J, Oda T, Finch CE, Pasinetti GM (1995) Association of apolipoprotein E genotype with brain levels of apolipoprotein E and apolipoprotein J (clusterin) in Alzheimer disease. *Brain Res Mol Brain Res* **33**:174–178.
26. Lidstrom AM, Bogdanovic N, Hesse C, Volkman I, Davidsson P, Blennow K (1998) Clusterin (apolipoprotein J) protein levels are increased in hippocampus and in frontal cortex in Alzheimer's disease. *Exp Neurol* **154**:511–521.
27. Calero M, Rostagno A, Matsubara E, Zlokovic B, Frangione B, Ghiso J (2000) Apolipoprotein J (clusterin) and Alzheimer's disease. *Microsc Res Tech* **50**:305–315.
28. May PC, Lampert-Etchells M, Johnson SA, Poirier J, Masters JN, Finch CE (1990) Dynamics of gene expression for a hippocampal glycoprotein elevated in Alzheimer's disease and in response to experimental lesions in rat. *Neuron* **5**:831–839.
29. Kanekiyo T, Xu H, Bu G (2014) ApoE and A β in Alzheimer's disease: Accidental encounters or partners? *Neuron* **81**:740–754.
30. Yu JT, Tan L (2012) The role of clusterin in Alzheimer's disease: Pathways, pathogenesis, and therapy. *Mol Neurobiol* **45**:314–326.
31. Zlokovic BV, Martel CL, Matsubara E, McComb JG, Zheng G, McCluskey RT, *et al* (1996) Glycoprotein 330/megalin: Probable role in receptor-mediated transport of apolipoprotein J alone and in a complex with Alzheimer disease amyloid β at the blood-brain and blood-cerebrospinal fluid barriers. *Proc Natl Acad Sci U S A* **93**:4229–4234.
32. Montine TJ, Phelps CH, Beach TG, Bigio EH, Cairns NJ, Dickson DW, *et al* (2012) National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease: A practical approach. *Acta Neuropathol* **123**:1–11.
33. Chalmers K, Wilcock GK, Love S (2003) *APOE* ϵ 4 influences the pathological phenotype of Alzheimer's disease by favouring cerebrovascular over parenchymal accumulation of A β protein. *Neuropathol Appl Neurobiol* **29**:231–238.
34. van Helmond Z, Miners JS, Kehoe PG, Love S (2010) Oligomeric A β in Alzheimer's disease: Relationship to plaque and tangle pathology, *APOE* genotype and cerebral amyloid angiopathy. *Brain Pathol* **20**:468–480.
35. Swirski M, Miners JS, de Silva R, Lashley T, Ling H, Holton J, *et al* (2014) Evaluating the relationship between amyloid- β and α -synuclein phosphorylated at Ser129 in dementia with Lewy bodies and Parkinson's disease. *Alzheimers Res Ther* **6**:77.
36. Thomas T, Miners S, Love S (2015) Post-mortem assessment of hypoperfusion of cerebral cortex in Alzheimer's disease and vascular dementia. *Brain* **138**:1059–1069.
37. Ashby EL, Miners JS, Kumar S, Walter J, Love S, Kehoe PG (2015) Investigation of A β phosphorylated at serine 8 (pA β) in Alzheimer's disease, dementia with Lewy bodies and vascular dementia. *Neuropathol Appl Neurobiol* **41**:428–444.
38. Barua NU, Miners JS, Bienemann AS, Wyatt MJ, Welsch K, Tabor AB, *et al* (2012) Convection-enhanced delivery of neprilysin: A novel amyloid- β -degrading therapeutic strategy. *J Alzheimers Dis* **32**:43–56.
39. Miners JS, Palmer JC, Love S. (in press) Pathophysiology of hypoperfusion of the precuneus in early Alzheimer's disease. *Brain Pathol*. doi: 10.1111/bpa.12331
40. Qiu WQ, Ye Z, Kholodenko D, Seubert P, Selkoe DJ (1997) Degradation of amyloid β -protein by a metalloprotease secreted by microglia and other neural and non-neural cells. *J Biol Chem* **272**:6641–6646.
41. Schrijvers EM, Koudstaal PJ, Hofman A, Breteler MM (2011) Plasma clusterin and the risk of Alzheimer disease. *JAMA* **305**:1322–1326.
42. Thambisetty M, Simmons A, Velayudhan L, Hye A, Campbell J, Zhang Y, *et al* (2010) Association of plasma clusterin concentration with severity, pathology, and progression in Alzheimer disease. *Arch Gen Psychiatry* **67**:739–748.
43. Giannakopoulos P, Kovari E, French LE, Viard I, Hof PR, Bouras C (1998) Possible neuroprotective role of clusterin in Alzheimer's disease: A quantitative immunocytochemical study. *Acta Neuropathol* **95**:387–394.
44. Howlett DR, Hortobagyi T, Francis PT (2013) Clusterin associates specifically with A β 40 in Alzheimer's disease brain tissue. *Brain Pathol* **23**:623–632.
45. McGeer PL, Kawamata T, Walker DG (1992) Distribution of clusterin in Alzheimer brain tissue. *Brain Res* **579**:337–341.
46. Matarin M, Salih DA, Yasvoina M, Cummings DM, Guelfi S, Liu W, *et al* (2015) A genome-wide gene-expression analysis and database in transgenic mice during development of amyloid or tau pathology. *Cell Rep* **10**:633–644.
47. Matsubara E, Soto C, Governale S, Frangione B, Ghiso J (1996) Apolipoprotein J and Alzheimer's amyloid β solubility. *Biochem J* **316**:671–679.
48. Oda T, Wals P, Osterburg HH, Johnson SA, Pasinetti GM, Morgan TE, *et al* (1995) Clusterin (apoJ) alters the aggregation of amyloid β -peptide (A β 1–42) and forms slowly sedimenting A β complexes that cause oxidative stress. *Exp Neurol* **136**:22–31.
49. Yerbury JJ, Poon S, Meehan S, Thompson B, Kumita JR, Dobson CM, Wilson MR (2007) The extracellular chaperone clusterin influences

- amyloid formation and toxicity by interacting with prefibrillar structures. *Faseb J* **21**:2312–2322.
50. DeMattos RB, O'Dell MA, Parsadanian M, Taylor JW, Harmony JA, Bales KR, *et al* (2002) Clusterin promotes amyloid plaque formation and is critical for neuritic toxicity in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A* **99**:10843–10848.
 51. Benzinger TL, Blazey T, Jack CR, Jr., Koeppe RA, Su Y, Xiong C, *et al* (2013) Regional variability of imaging biomarkers in autosomal dominant Alzheimer's disease. *Proc Natl Acad Sci U S A* **110**:E4502–E4509.
 52. Binnewijzend MA, Kuijjer JP, Benedictus MR, van der Flier WM, Wink AM, Wattjes MP, *et al* (2013) Cerebral blood flow measured with 3D pseudocontinuous arterial spin-labeling MR imaging in Alzheimer disease and mild cognitive impairment: A marker for disease severity. *Radiology* **267**:221–230.
 53. Yamaguchi H, Sugihara S, Ogawa A, Oshima N, Ihara Y (2001) Alzheimer β amyloid deposition enhanced by apoE 4 gene precedes neurofibrillary pathology in the frontal association cortex of nondemented senior subjects. *J Neuropathol Exp Neurol* **60**:731–739.
 54. Castano EM, Prelli F, Wisniewski T, Golabek A, Kumar RA, Soto C, Frangione B (1995) Fibrillogenesis in Alzheimer's disease of amyloid β peptides and apolipoprotein E. *Biochem J* **306**:599–604.
 55. LaDu MJ, Falduto MT, Manelli AM, Reardon CA, Getz GS, Frail DE (1994) Isoform-specific binding of apolipoprotein E to β -amyloid. *J Biol Chem* **269**:23403–23406.
 56. Ma J, Yee A, Brewer HB, Jr., Das S, Potter H (1994) Amyloid-associated proteins alpha 1-antichymotrypsin and apolipoprotein E promote assembly of Alzheimer β -protein into filaments. *Nature* **372**:92–94.
 57. Sanan DA, Weisgraber KH, Russell SJ, Mahley RW, Huang D, Saunders A, *et al* (1994) Apolipoprotein E associates with β amyloid peptide of Alzheimer's disease to form novel monofibrils. Isoform apoE4 associates more efficiently than apoE3. *J Clin Invest* **94**:860–869.
 58. Strittmatter WJ, Weisgraber KH, Huang DY, Dong LM, Salvesen GS, Pericak-Vance M, *et al* (1993) Binding of human apolipoprotein E to synthetic amyloid β peptide: Isoform-specific effects and implications for late-onset Alzheimer disease. *Proc Natl Acad Sci U S A* **90**:8098–8102.
 59. DeMattos RB, Cirrito JR, Parsadanian M, May PC, O'Dell MA, Taylor JW, *et al* (2004) ApoE and clusterin cooperatively suppress A β levels and deposition: Evidence that ApoE regulates extracellular Abeta metabolism in vivo. *Neuron* **41**:193–202.
 60. Craggs L, Taylor J, Slade JY, Chen A, Hagel C, Kuhlenbaeumer G, *et al* (2016) Clusterin/Apolipoprotein J immunoreactivity is associated with white matter damage in cerebral small vessel diseases. *Neuropathol Appl Neurobiol* **42**:194–209.
 61. Hawkes CA, Jayakody N, Johnston DA, Bechmann I, Carare RO (2014) Failure of perivascular drainage of β -amyloid in cerebral amyloid angiopathy. *Brain Pathol* **24**:396–403.
 62. Weller RO, Subash M, Preston SD, Mazanti I, Carare RO (2008) Perivascular drainage of amyloid-beta peptides from the brain and its failure in cerebral amyloid angiopathy and Alzheimer's disease. *Brain Pathol* **18**:253–266.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Figure S1. Stacked bar chart illustrating regional differences in the relative contributions of soluble and insoluble A β 40 and A β 42 to 'total' A β load in AD and control brains in MF, CC, PH, PC, TH, and WM.