

Corrigendum

Corrigendum to article “A programmable omnipotent Argonaute nuclease from mesophilic bacteria *Kurthia massiliensis*”

Yang Liu, Wenqiang Li, Xiaoman Jiang, Yaping Wang, Zhiwei Zhang, Qi Liu, Ruyi He, Quan Chen, Jun Yang, Longyu Wang, Fei Wang* and Lixin Ma*

State Key Laboratory of Biocatalysis and Enzyme Engineering, Hubei Collaborative Innovation Center for Green Transformation of Bio-resources, Hubei Key Laboratory of Industrial Biotechnology, School of Life Sciences, Hubei University, Wuhan, Hubei 430062, China

Nucleic Acids Research, Volume 49, Issue 3, 22 February 2021, Pages 1597–1608, <https://doi.org/10.1093/nar/gkaa1278>

In Figure 3 the lower-left panel should be labelled as B, and the upper-right panel should be labelled as C.

This correction does not affect the results, discussion and conclusions presented in the article. The published article has been updated.




Figure 3. Effects of the guide length on KmAgo activity. (A) Effects of the 5'P-gDNA length on DNA cleavage activity. (B) Effects of the 5'P-gDNA length on RNA cleavage activity. (C) Effects of the 5'P-gRNA length on RNA cleavage activity. (D) Effects of the 5'P-gRNA length on DNA cleavage activity. Black triangle indicates the 30 nt 5'P-gRNA. All experiments were performed at the 4:2:1 KmAgo:guide:target molar ratio in reaction buffer containing Mn^{2+} ions for 15 min at 37°C.

*To whom correspondence should be addressed. Tel: +86 27 50865628; Fax: +86 27 88666349; Email: malixing@hubu.edu.cn
Correspondence may also be addressed to Fei Wang. Tel: +86 27 50865628; Fax: +86 27 88666349; Email: wangfei@hubu.edu.cn