



Published in final edited form as:

Tob Control. 2022 January ; 31(1): 107–111. doi:10.1136/tobaccocontrol-2020-055739.

Changes in responses to nicotine vaping product warnings and leaflets in England compared with Canada, the US and Australia: findings from the 2016–2018 ITC Four Country Smoking and Vaping Surveys

Eve Violet Taylor¹, Katherine A East^{1,2}, Ann McNeill^{1,3}, Michael Cummings^{4,5}, James Thrasher^{6,7}, Geoffrey T Fong^{2,8,9}, Anne C K Quah⁸, Máirtín McDermot¹, Grace Li⁸, Ron Borland^{10,11}, David Hammond², Sarah Aleyan^{1,2}, Sara C Hitchman¹

¹Addictions Department, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

²School of Public Health and Health Systems, University of Waterloo, Waterloo, Ontario, Canada

³Shaping Public Health Policies to Reduce Inequalities and Harm (SPECTRUM), London, UK

⁴Psychiatry and Behavioral Sciences, Medical University of South Carolina, Charleston, South Carolina, USA

⁵Hollings Cancer Centre, Medical University of South Carolina, Charleston, South Carolina, USA

⁶Health Promotion, Education and Behavior, School of Public Health, University of South Carolina, Columbia, South Carolina, USA

⁷Tobacco Research Department, Centre for Population Health Research, National Institute of Public Health, Cuernavaca, Mexico

⁸Department of Psychology, University of Waterloo, Waterloo, Ontario, Canada

⁹Ontario Institute for Cancer Research, Toronto, Ontario, Canada

¹⁰Melbourne Centre for Behaviour Change, School of Psychological Sciences, University of Melbourne, Melbourne, Victoria, Australia

¹¹Cancer Council Victoria, Melbourne, Victoria, Australia

Correspondence to: Eve Violet Taylor, Addictions Department, King's College London, London WC2R 2LS, UK; eve.v.taylor@kcl.ac.uk.

Contributors This paper was supervised by SCH and AM. SA and KAE assisted with statistical analysis, planning and revisions of the paper. MM reviewed and revised the paper. MC, JT, GTF, ACKQ, GL, RB and DH were involved in the design of the ITC study and reviewed and revised the paper.

Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/tobaccocontrol-2020-055739>).

Twitter Ann McNeill @kingsNRG

Ethics approval Study questionnaires and materials were reviewed and provided clearance by Research Ethics Committees at the following institutions: University of Waterloo (Canada, ORE#20803/30570, ORE#21609/30878); King's College London, UK (RESCM-17/18–2240); Cancer Council Victoria, Australia (HREC1603); University of Queensland, Australia (2016000330/HREC1603); and Medical University of South Carolina (waived due to minimal risk). All participants provided consent to participate.

Provenance and peer review Not commissioned; externally peer reviewed.

Abstract

Background and aims—In May 2017, black-and-white text nicotine addiction warning labels (‘warnings’) and health and safety leaflets (‘leaflets’) became mandatory for nicotine vaping products (NVPs) in England, in accordance with the European Union’s Tobacco Products Directive. We compared changes over time in noticing warnings and leaflets, recall of warnings about nicotine and concerns about using NVP due to noticing warnings in England, compared with Canada, the US and Australia, where no warnings and leaflets were mandated.

Design—19 005 adult (aged 18+) NVP users, smokers and quitters of cigarettes and NVP from the 2016 and 2018 International Tobacco Control Four Country Smoking and Vaping Surveys in England, Canada, the US and Australia, recruited via probability and non-probability sampling.

Findings—Noticing warnings increased in England from 4.9% (2016) to 9.4% (2018) (adjusted OR/AOR=1.64, 95% CI=1.15–2.36); this change was larger than changes in Canada (AOR=2.51, 95% CI=1.71–3.69) and the US (AOR=2.22, 95% CI=1.45–3.39). Recall of a nicotine warning increased in England from 86% (2016) to 94.9% (2018) (AOR=5.50, 95% CI=1.57–19.27) but not significantly elsewhere. Noticing leaflets increased in England from 14.6% (2016) to 19.1% (2018) (AOR=1.42, 95% CI=1.15–1.74); this change was larger than in Canada (AOR=1.42, 95% CI=1.12–1.79), the US (AOR=1.55, 95% CI=1.17–2.06) and Australia (AOR=1.51, 95% CI=1.02–2.22). Among those noticing warnings, concern about NVP use did not change significantly between 2016 and 2018 (all countries $p>0.081$).

Conclusions—Introduction of mandatory NVP warnings and leaflets in England was associated with small increases in noticing them but not with changes in concerns about NVP use.

INTRODUCTION

The European Union Tobacco Products Directive (EU TPD) requires the inclusion of health and safety leaflets (‘leaflets’) and health warning labels (‘warnings’) on packaging for nicotine vaping products (NVPs) and e-liquids containing nicotine.¹ England implemented the EU TPD in May 2016, mandating warnings by May 2017. The current 30% black-and-white text warning in England reads ‘this product contains nicotine which is a highly addictive substance’ (online supplemental figure 1).²³ There is no set design for the leaflets, however, they must state that the product is not recommended for young people and non-smokers.¹ In addition to warnings, NVP packaging must include hazard symbols in accordance with the EU classification labelling and packaging of substances regulation.²³ EU legislation was the first to require NVP warnings, with no warnings or leaflets required in Canada, the US or Australia during the current study period (2016–2018), although manufacturers sometimes voluntarily added warnings or toxin symbols⁴⁵ (online supplemental figure 2).

Little is known about the effectiveness and salience of NVP warnings and leaflets, or how they might influence concerns about NVP use. It has been suggested that they could increase concerns about use and perceptions of harm among non-smokers.^{6–8} By contrast, qualitative research about heated tobacco products conducted with UK smokers and ex-smokers suggests that text-only warnings might decrease perceptions of harm relative to

cigarettes because of the contrast between text-only warnings and the pictorial cigarette warnings.⁹

Comparisons of changes in noticing NVP warnings and leaflets and concerns about NVPs between countries with (England) and without labelling policies (Canada, US, Australia) could provide timely evidence regarding the impact of mandated warnings.

Objective

This study used data from the 2016 and 2018 International Tobacco Control (ITC) Policy Evaluation Project Four Country Smoking and Vaping (4CV) Surveys to investigate, in England compared with countries without labelling policies (Canada, the US and Australia), the impact of NVP warnings and leaflets on noticing warnings/leaflets, recall of a nicotine warning and concerns about using NVPs due to warnings.

METHODS

Participants and design

This study used data from wave 1 (2016) and wave 2 (2018) of the longitudinal ITC 4CV Survey. Detailed methods can be found elsewhere.^{10–13} Briefly, the sample comprised smokers, vapers and former smokers and vapers from England, Canada, the US and Australia, recruited through probability and non-probability sampling frames (via random-digit dialling or email invitations from web-based or address-based panels). At wave 2, respondents from wave 1 were invited to participate; respondents lost to attrition were replaced by newly recruited respondents using the same recruitment strategy. Data were collected July–November 2016 (wave 1) and February–July 2018 (wave 2).

At wave 1, n=13 099 participants were recruited. At wave 2, n=5985 were successfully followed up and n=7650 were added for replenishment, providing a sample of N=20 749. The following participants were then excluded: those recruited via the Australian Dedicated Vapers (not representative of Australian NVP users and smokers; n=896); had never heard of NVP (n=214); long term quitters (those who quit at least 5 years ago; n=231); did not respond to questions about education (n=236) or ethnicity (n=143) or responded don't know to ethnicity (n=167). The final sample comprised N=19 005 participants.

Measures

The survey development process is outlined in the ITC technical reports, with measures chosen previously used by McDermott *et al.*¹⁰¹¹¹³

Outcomes

Noticing warnings.: Participants were asked: 'Now thinking about e-cigarettes, in the last 30 days, have you noticed any health warnings on packaging for e-cigarettes, cartridges or e-liquid containers?' Responses were coded 'Yes' versus 'Other' ('No', 'Refused', 'Don't know').

Noticing leaflets.: Participants were asked ‘As far as you know, is there health and product safety information contained on leaflets inside the packaging for e-cigarettes or on boxes of components?’ Responses were coded ‘Yes’ versus ‘Other’ (‘No’, ‘Don’t know’, ‘Refused’).

Recall a nicotine warning.: Participants were asked ‘In the last 30 days, have you read any of the health warnings?’ Those who stated that ‘Yes’ were asked ‘What do you recall the health warning(s) saying?’ followed by a list of potential warnings (online supplemental table 1), including the current England EU TPD warning. Responses were coded ‘Yes’ versus ‘Other’ (‘No’, ‘Don’t know’, ‘Refused’).

Effects of warnings on concern about NVP use.: Participants who noticed warnings were asked ‘What effect have the health warnings had on your thoughts about using e-cigarettes?’ Responses were coded ‘Concerned’ (‘Made me concerned about using them’) or ‘Not concerned’ (‘Had no effect’, ‘Reassured me about using them’, ‘Don’t know’, ‘Refused’).

Smoking/vaping behavior

Smoking and vaping status: Participants were defined as either exclusive ‘Daily NVP users’, ‘Daily smokers’, ‘Non-daily NVP’ users, ‘Non-daily smokers’, or ‘Concurrent’ users (currently smoking and using NVP) or ‘Quitters of NVP and/or smoking’ (no longer using either product). Groups were mutually exclusive (online supplemental table 2).¹³

Additionally, a new variable was coded, ‘collapsed smoking/vaping status’, ‘Daily NVP’, ‘Non-daily NVP’ and ‘Concurrent’ users were coded ‘Vapers’. ‘Daily smokers’ and ‘Non-daily smokers’ were coded ‘Exclusive smokers’. Groups were mutually exclusive. Quitters of both products were excluded.

Analysis

Logistic regression models employing Generalised Estimating Equations (GEE) (model 1) were conducted to assess changes from wave 1 to 2 in: noticing warnings and leaflets, recall of a nicotine warning (among those who read warnings, n=1521) and concern about NVPs due to warnings (among those who noticed warnings, n=2320). Interactions between country and wave were used to examine changes in outcomes over time in England compared with Canada, the US and Australia.

A second set of GEE logistic regression models (model 2) were conducted to examine changes over time in the three outcome measures by smoking and vaping status. The same analyses were also conducted in the sample that excluded quitters and used the ‘collapsed smoking/vaping’ status variable (model 3). In model 3, interactions between exclusive smokers and vapers and wave were used to examine changes in outcomes over time between exclusive smokers and vapers.

Analyses controlled for age, gender, ethnicity, income, education, wave of recruitment, having a friend/relative who uses an NVP and smoking and vaping status. Having a friend/relative who uses an NVP and smoking and vaping status were treated as time-varying covariates. All other variables were treated as time-invariant. Analyses were weighted and conducted in SPSS V.25.¹⁰¹¹

RESULTS

See online supplemental table 3 for participant demographics.

Noticing warnings

Between waves 1 and 2, noticing warnings increased significantly in England (adjusted OR/AOR=1.64, 95% CI=1.15–2.36), increased non-significantly in Australia, decreased significantly in Canada and did not change in the US. The change in England was greater than in Canada (AOR=2.51, 95% CI=1.71–3.69) and the US (AOR=2.22, 95% CI=1.45–3.39), but no different from Australia (model 1, table 1). Noticing increased among daily NVP users, but decreased among daily smokers (model 2, table 1). There was a decrease in noticing warnings between waves 1 and 2 among exclusive smokers, and an increase in noticing warnings among vapers, and the change among vapers was greater than among exclusive smokers (model 3, table 1).

Recall of warnings

Among those who had read warnings (online supplemental table 1), a significant increase was found in recall of a nicotine warning in England between waves 1 and 2 (86.0%–94.9%, AOR=5.50; 95% CI=1.57–19.27, $p=0.008$). Increases were observed in Canada (79.6%–87.0%, AOR=2.00; 95% CI=0.75–5.34, $p=0.168$), the US (71.9%–88.1%, AOR=3.39; 95% CI=0.90–12.75, $p=0.071$) and Australia (71.1%–73.5%, AOR=2.16; 95% CI=0.04–2.46, $p=0.783$) although they were not significant.

Noticing leaflets

Between waves 1 and 2, noticing leaflets increased significantly in England, (AOR=1.42, 95% CI=1.15–1.74), with small and non-significant increases in Australia and small non-significant decreases in Canada and the US. The change between waves in England was greater than in Canada, (AOR=1.42, 95% CI=1.12–1.79), the US (AOR=1.55, 95% CI=1.17–2.06) and Australia (AOR=1.51, 95% CI=1.02–2.22) (model 1, table 1). Noticing leaflets increased among daily NVP users (model 2, table 1). There was an increase in noticing leaflets between waves 1 and 2 among vapers, and this change was greater than that among exclusive smokers (model 3, table 1).

Concern about using NVPs after noticing NVP warnings

Concern about NVP after noticing warnings decreased in England between waves 1 and 2, although this was non-significant when adjusted for covariates. Concern also decreased non-significantly in Canada and Australia and did not change in the US (model 1, table 1). Concern decreased significantly among NVP users and increased, but not significantly, among exclusive smokers (model 2, table 1). The change in concern among NVP users was not significantly different to changes in concern among exclusive smokers (model 3, table 1).

DISCUSSION

This study reports on changes in noticing NVP warnings and leaflets among adult smokers and NVP users before and after mandating of NVP packaging policies in England compared with other countries without such policy. Noticing warnings and leaflets increased between 2016 and 2018 in England, compared with the other countries. Moreover, noticing warnings or leaflets increased among vapers with the changes being significantly different to those among exclusive smokers. Overall, findings suggest that the enforcement of mandatory warnings and leaflets led to increases in noticing them.

Noticing warnings and leaflets was low across all countries (4.8%–9.4%); this was expected in Canada, the US and Australia where warnings and leaflets were not mandatory. However, noticing was relatively low in England in 2018 even after enforcement of mandatory warnings and leaflets. There may be several reasons. First, the warning is only present on NVP and e-liquid packaging, so consumers may only notice them at initial purchase and when refilling e-liquids, which may limit exposure. Also, the warning message on NVPs may not be especially salient since most consumers already know nicotine is addictive and is in the NVP that they are using.¹⁴ Moreover, compared with the warnings found on cigarette packs, the warnings on NVP packaging and e-liquids are smaller and text-only. It is well established that, for cigarettes, larger pictorial warnings are more likely to be noticed,^{15,16} although such warnings might not be warranted on NVP given their likely lower health risks compared with cigarettes.^{17–19}

While noticing leaflets was low across all countries, it was higher than expected in Canada and the US given they were not mandatory in these countries at the time of this study. However, this may be due to voluntary safety information placed on product information leaflets by manufacturers.²⁰ Concern over the use of NVP due to warnings decreased between 2016 and 2018 in England, however when adjusted for covariates this change was not significant. There was also a decrease in concern between 2016 and 2018 among vapers, suggesting that the nicotine warnings may not influence concern about NVP among current users. Increases in concern about NVP were seen among smokers, this increase was not significant when adjusting for covariates, possibly due to small cell counts. Future research should further investigate warning perceptions among smokers, who might use NVP to help them quit smoking, and among non-smokers, who the warnings are aiming to deter from using NVP.

Study limitations included that our analysis was not a clear pre-post comparison since NVP companies in England may have applied warnings and included leaflets during the 2016 implementation year. Even with this limitation we observed a small, but significant, increase in noticing warnings and leaflets in England. We acknowledge that small sample sizes, especially for Australia, and analysis of concerns about NVP, limit the robustness of our conclusions. Finally, measures were derived from well-established measures used to examine the impact of cigarette health warnings but have limited use in the context of NVP.¹³

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We are grateful to the editor and anonymous reviewers for their guidance and comments which helped in improving this article.

Funding This study was supported by grants from the US National Cancer Institute (P01 CA200512), the Canadian Institutes of Health Research (FDN-148477), and by the National Health and Medical Research Council of Australia (APP 1106451). GTF was supported by a Senior Investigator Grant from the Ontario Institute for Cancer Research.

Competing interests MC has received payment as a consultant to Pfizer for service on an external advisory panel to assess ways to improve smoking cessation delivery in healthcare settings. MC also has served as paid expert witness in litigation filed against the tobacco industry. GTF, DH and JT have served as expert witnesses on behalf of governments in litigation involving the tobacco industry. AM is a UK National Institute for Health Research (NIHR) Senior Investigator.

REFERENCES

1. European Parliament. Tobacco products directive, 2014. Available: https://ec.europa.eu/health/sites/health/files/tobacco/docs/dir_201440_en.pdf
2. Health and Safety Executive. The CLP regulation - chemical classification. Available: <http://www.hse.gov.uk/chemical-classification/legal/clp-regulation.htm>
3. Medicines and Healthcare products Regulatory Agency (MHRA). E-cigarettes: regulations for consumer products - GOV.UK, 2019. Available: <https://www.gov.uk/guidance/e-cigarettes-regulations-for-consumer-products>
4. Fagan P, Pokhrel P, Herzog TA, et al. Warning statements and safety practices among manufacturers and Distributors of electronic cigarette liquids in the United States. *Nicotine Tob Res* 2018;20:970–6. [PubMed: 28520985]
5. Kong AY, Derrick JC, Abrantes AS, et al. What is included with your online e-cigarette order? an analysis of e-cigarette shipping, product and packaging features. *Tob Control* 2018;27:699–702. [PubMed: 27357936]
6. Wackowski OA, Hammond D, O'Connor RJ, O'Connor R, et al. Considerations and future research directions for e-cigarette Warnings-Findings from expert interviews. *Int J Environ Res Public Health* 2017; 14:781.
7. Kimber C, Frings D, Cox S, et al. Communicating the relative health risks of e-cigarettes: an online experimental study exploring the effects of a comparative health message versus the EU nicotine addiction warnings on smokers' and non-smokers' risk perceptions and behavioural intentions. *Addict Behav* 2020;101:106177. [PubMed: 31753541]
8. Cox S, Frings D, Ahmed R, et al. Messages matter: the tobacco products Directive nicotine addiction health warning versus an alternative relative risk message on smokers' willingness to use and purchase an electronic cigarette. *Addict Behav Rep* 2018;8:136–9. [PubMed: 30263928]
9. Tompkins CNE, Burnley A, McNeill A, et al. Factors that influence smokers' and ex-smokers' use of IQOS: a qualitative study of IQOS users and ex-users in the UK. *Tob Control* 2020. doi:10.1136/tobaccocontrol-2019-055306. [Epub ahead of print: 15 Jan 2020].
10. ITC. ITC four county smoking and Vaping survey wave 2 (4CV2) technical report, 2020. Available: https://itcproject.s3.amazonaws.com/uploads/documents/4CV2_Technical_Report_15Jan202.pdf
11. ITC. ITC four county smoking and vaping survey, wave 1 (4CV1) technical report, 2018. Available: https://itcproject.s3.amazonaws.com/uploads/documents/4CV1_Technical_Report_Nov23201.pdf
12. Thompson ME, Fong GT, Boudreau C, et al. Methods of the ITC four country smoking and Vaping survey, wave 1 (2016). *Addiction* 2019; 114 Suppl 1:6–14. [PubMed: 30548345]

13. McDermott MS, Li G, McNeill A, et al. Exposure to and perceptions of health warning labels on nicotine vaping products: findings from the 2016 international tobacco control four country smoking and Vaping survey. *Addiction* 2019; 114 Suppl 1:134–43. [PubMed: 30618081]
14. Wackowski O, Hammond D, O'Connor R, et al. Smokers' and e-cigarette users' perceptions about e-cigarette warning statements. *Int J Environ Res Public Health* 2016;13:655.
15. Hammond D Health warning messages on tobacco products: a review. *Tob Control* 2011;20:327–37. [PubMed: 21606180]
16. Noar SM, Francis DB, Bridges C, et al. The impact of strengthening cigarette pack warnings: systematic review of longitudinal observational studies. *Soc Sci Med* 2016;164:118–29. [PubMed: 27423739]
17. McNeill A, Brose LS, Calder R, et al. Evidence review of e-cigarettes and heated tobacco products 2018 a report commissioned by public health England, 2018. Available: www.facebook.com/PublicHealthEngland
18. NASEM. Public health consequences of e-cigarettes conclusions by outcome constituents of e-cigarettes, 2018. Available: <https://www.nap.edu/resource/24952/012318ecigaretteConclusionsbyOutcome.pdf>
19. World Health Organisation. Electronic nicotine and non-nicotine delivery systems a brief, 2020. Available: <http://www.euro.who.int/pubrequest>
20. Shang C, Chaloupka FJ. The trend of voluntary warnings in electronic nicotine delivery system magazine advertisements. *Int J Environ Res Public Health* 2017;14. Doi:10.3390/ijerph14010062. [Epub ahead of print: 10 Jan 2017].

What this paper adds

- Following the 2014 European Union Tobacco Products Directive (EU TPD), nicotine addiction warning labels and health and safety leaflets on nicotine vaping products (NVPs) became mandatory in England in 2017.
- This is the first study to investigate and compare changes in noticing and recall of warnings and leaflets on NVPs in England, to countries where no warning or leaflets were mandatory (Canada, the US and Australia).
- The findings show that noticing warnings and leaflets on NVPs was higher in England after EU TPD implementation than before. Moreover, concerns about using NVPs did not change after warnings on labels and leaflets were introduced.

Change over time in noticing NVP health warning labels, health and safety leaflets, and concern about NVPs between 2016 (wave 1) and 2018 (wave 2) *

Table 1

Waves	Noticed a health warning label (n=19 005)			Noticed a health and safety leaflet (n=19005)			Concern about NVP among those who noticed a health warning label (n=2320) †			
	N (%)	AOR (95% CI)	P value	N (%)	AOR (95% CI)	P value	N (%)	AOR (95% CI)	P value	
Model 1: changes within each country between wave 1 and wave 2										
England	1	201 (4.9)	ref	599 (14.6)	ref	ref	52 (25.8)	ref	ref	
	2	430 (9.4)	1.64 (1.15–2.36)	0.007	877 (19.1)	1.42 (1.15–1.74)	63 (14.7)	0.52 (0.25–1.08)	0.081	
Canada	1	254 (7.0)	ref	680 (18.7)	ref	ref	95 (37.3)	ref	ref	
	2	188 (5.1)	0.65 (0.50–0.86)	0.003	643 (17.4)	1.03 (0.85–1.19)	62 (33.1)	0.68 (0.40–1.16)	0.157	
US	1	188 (7.0)	ref	398 (14.9)	ref	ref	55 (29.2)	ref	ref	
	2	180 (6.5)	0.74 (0.54–1.02)	0.068	385 (13.8)	0.91 (0.73–1.15)	51 (28.5)	0.93 (0.46–1.90)	0.843	
Australia	1	39 (2.7)	ref	156 (10.6)	ref	ref	5 (13.9)	ref	ref	
	2	71 (4.8)	0.94 (0.56–1.60)	0.832	186 (12.5)	0.94 (0.67–1.33)	8 (11.2)	1.58 (0.38–6.56)	0.526	
Changes between wave 1 and 2 by country interaction §										
		Change between waves		Change between waves		Change between waves		Change between waves		
England		4.50%		4.50%		-11.10%		-11.10%		
Canada		-1.90%	2.51 (1.71–3.69)	<0.001	-1.30%	1.42 (1.12–1.79)	0.004	-4.20%	0.76 (0.36–1.60)	0.470
US		-0.50%	2.22 (1.45–3.39)	<0.001	-1.10%	1.55 (1.17–2.06)	0.002	-0.70%	0.56 (0.23–1.36)	0.198
Australia		2.10%	1.74 (0.95–3.21)	0.075	1.90%	1.51 (1.02–2.22)	0.039	-2.70%	0.33 (0.07–1.58)	0.165
Model 2: changes within smoking and vaping status between wave 1 and wave 2										
Concurrent	1	299 (9.7)	ref	729 (23.8)	ref	ref	111 (37.3)	ref	ref	
	2	328 (11.5)	1.02 (0.82–1.28)	0.867	712 (24.9)	1.10 (0.94–1.28)	93 (28.3)	0.60 (0.27–0.98)	0.042	
Daily NVP	1	107 (17.4)	ref	191 (30.9)	ref	ref	4 (3.8)	ref	ref	

Waves	Noticed a health warning label (n=19 005)			Noticed a health and safety leaflet (n=19005)			Concern about NVP among those who noticed a health warning label (n=2320) †		
	N (%)	AOR (95% CI)	P value	N (%)	AOR (95% CI)	P value	N (%)	AOR (95% CI)	P value
2	315 (37)	2.40 (1.51–3.81)	<0.001	416 (49)	2.11 (1.46–3.05)	<0.001	21 (6.6)	1.67 (0.38–7.28)	0.493
Non-daily NVP									
1	49 (8.8)	ref		128 (23.1)	ref		15 (30.9)	ref	
2	55 (10.5)	0.91 (0.43–1.90)	0.795	140 (26.7)	1.35 (0.73–2.13)	0.416	15 (27.3)	0.70 (1.18–2.75)	0.604
Daily smokers									
1	110 (2.7)	ref		424 (10.3)	ref		31 (28)	ref	
2	106 (2.1)	0.69 (0.48–0.98)	0.028	472 (9.3)	0.89 (0.74–1.07)	0.209	36 (34.1)	1.09 (0.47–2.52)	0.846
Non-daily smokers									
1	55 (5.4)	ref		106 (11.3)	ref		13 (24.5)	ref	
2	36 (4.8)	0.73 (0.41–1.30)	0.287	83 (10.8)	1.11 (0.75–1.63)	0.609	9 (24.2)	0.88 (0.24–3.30)	0.855
Quitters									
1	63 (2.5)	ref		253 (10.1)	ref		32 (50.8)	ref	
2	31 (1.2)	0.43 (0.17–1.06)	0.066	268 (10.7)	1.06 (0.74–1.53)	0.747	11 (37.4)	0.43 (0.09–1.96)	0.267
Model 3: changes within exclusive smokers or vapers between wave 1 and wave 2									
(n=17923) †									
Exclusive smoker									
1	165 (3.2)	ref		530 (10.3)	ref		44 (26.9)	ref	
2	142 (2.4)	0.68 (0.50–0.93)	0.015	556 (9.6)	0.93 (0.78–1.10)	0.367	45 (31.6)	1.13 (0.53–2.39)	0.754
Vapers									
1	455 (10.7)	ref		1049 (24.8)	ref		130 (28.7)	ref	
2	698 (16.5)	1.26 (1.01–1.59)	0.049	1289 (30)	1.28 (1.10–1.50)	0.002	129 (18.40%)	0.58 (0.36–0.92)	0.022
Changes between wave 1 and 2 by exclusive smokers or vapers interaction †									
		Change between waves		Change between waves		Change between waves		Change between waves	
Exclusive smokers		–0.80%	ref	–0.70%	ref	4.70%	ref	ref	
Vapers		5.80%	1.85 (1.33–2.58)	<0.001	5.20%	1.38 (1.13–1.69)	0.002	–10.30%	0.51 (0.24–1.11)
									0.089

Bold represents p<.05

* Models used weighted data. Country, age, gender, education, income, wave of recruitment, smoking and vaping status and having a friend who uses NVP were all included as covariates within modelling procedures. AOR represent the odds at wave 2 compared with wave 1.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

⁷The interaction explored the difference in difference between current smoking or vaping status over time, comparing the change in vapers to the change observed among exclusive smokers. Quitters were excluded from this analysis. Vapers included daily NVP users, non-daily NVP users and concurrent users. Exclusive smokers include daily smokers and non-daily smokers.

⁷Sample only included those who reported noticing a health warning label.

⁸The interaction explored the difference in difference between countries over time, comparing changes in key outcome measures over time within England relative to the changes seen in Canada, the US and Australia.

⁹Quitters were excluded from this analysis. Vapers included daily NVP users, non-daily NVP users and concurrent users. Exclusive smokers include daily smokers and non-daily smokers. AOR, adjusted OR; NVP, nicotine vaping product.