

Effect of media distraction (audio-visual and music) for pain and anxiety control in patients undergoing shock-wave lithotripsy: A systematic review and meta-analysis

WEIHUA HU¹, KE YANG¹, LI ZHANG¹ and XU LU²

Departments of ¹Urology and ²Laboratory Hunan Provincial People's Hospital, The First Affiliated Hospital of Hunan Normal University, Changsha, Hunan 410000, P.R. China

Received April 27, 2020; Accepted January 20, 2021

DOI: 10.3892/etm.2021.10055

Abstract. Audio-visual (AV) or music distraction may be used to reduce pain during several healthcare procedures. The present manuscript is a systematic review and meta-analysis to assess the effectiveness of media distraction in reducing pain and anxiety in extracorporeal shock wave lithotripsy (ESWL) patients. The PubMed, Embase, Scopus, BioMed Central, Ovoid and CENTRAL (Cochrane Central Register of Controlled Trials) databases were screened for studies assessing the role of media distraction (music/AV media) in reducing pain and anxiety of ESWL patients. Data were summarized using the mean difference (MD) with 95% confidence intervals (CI). A total of 11 randomized controlled trials were included. Pooled analysis indicated a statistically significant difference in pain outcomes with media distraction [mean difference (MD): -1.18; 95% CI: -2.35, -0.01; $I^2=96.8\%$]. Subgroup analysis indicated that both AV media (MD: -2.94; 95% CI: -4.70, -1.17; $I^2=79.2\%$) and music (MD: -0.86; 95% CI: -1.37, -0.35; $I^2=62.5\%$), led to significant reduction in pain outcomes. Pooled analysis indicated a statistically significant reduction of anxiety scores with the use of media distraction (MD: -3.91; 95% CI: -6.44, -1.38; $I^2=77.7\%$). To conclude, the present review suggests that media distraction in the form of AV media or music may be beneficial in reducing the pain and anxiety of patients undergoing ESWL. Evidence is, however, weak considering the small effect size, confidence intervals being close to zero, and instability of the results on sensitivity analysis. In clinical practice, media distraction may be used during ESWL as a nursing intervention, but a clinically important reduction of pain and anxiety may not be expected.

Introduction

Urinary stones are a significant health problem affecting ~12% of the population in North America, 5-9% in Europe and 1-5% in Asia (1). Analysis of global trends indicates that due to climate change, obesity and dietary modifications, the worldwide prevalence of urinary stones is gradually increasing (2). Since the 1980s, extracorporeal shock wave lithotripsy (ESWL) has been widely used in the management of urinary tract calculi, due to its high efficacy and low morbidity (3). After the manufacture of the first ESWL device in 1984, several modifications have made subsequent devices smaller and less powerful, allowing the management of urinary stones without the use of anesthetic (4,5). Despite technological development, ~30% of patients still complain of severe pain (6). Pain during ESWL can affect the patients' tolerance of the procedure as well as the effectiveness of treatment (7). Clinicians have reportedly used a number of pain control agents, including opioids, non-steroidal anti-inflammatory drugs (NSAIDs), nitrous oxide, local anesthetic infiltration and dermal anesthetics such as Emla, to manage the analgesic requirements of ESWL (6).

Nursing personnel play an important role in patient preparation and management of the patients' pain and anxiety (8). They are not only responsible for prescription drug administration but can also support pain management by a variety of complementary therapies (9,10). In the world of modern media, audio-visual (AV) or music distraction has been successfully used to reduce procedural pain for many non-urological procedures (11-14). Considering the widespread use of the ESWL procedure, a number of researchers have also studied the efficacy of such media distraction techniques for pain management in urinary stone patients (15,16). A recent meta-analysis from Kyriakides *et al* (15) evaluated the role of music in reducing pain in patients undergoing urological procedures. Their study, updated to 2017, included only six trials on patients undergoing ESWL. Their review was focused on only music therapy and did not present results of studies only on ESWL. Considering this lacuna in the literature, the purpose of the present study was to conduct a systematic review and meta-analysis to assess the effectiveness of any media distraction in reducing pain and anxiety of ESWL patients.

Correspondence to: Mr. Xu Lu, Department of Laboratory, Hunan Provincial People's Hospital, The First Affiliated Hospital of Hunan Normal University, Room 202, East Building 15-Luqiao Group, Shaoshan South Road, Yuhua, Changsha, Hunan 410000, P.R. China
E-mail: luxusky@126.com

Key words: music, audiovisual, distraction, complementary therapy, analgesia, urology, renal stones

Materials and methods

Inclusion criteria. The Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines (17) and guidelines from the Cochrane collaboration (18) were followed during the conduct of this systematic review. The Population, Intervention, Comparison, Outcome and Study design (PICOS) model was used to select studies for inclusion in the review (17). Studies conducted on patients undergoing ESWL (population) were considered for inclusion. All randomized controlled trials (RCTs) comparing any type of media distraction (music/AV media) used during the ESWL procedure (intervention), with control (comparison) and measuring pain and/or anxiety (outcomes) were included. Studies were included irrespective of sample size and language of publication. The following studies were excluded: i) Studies utilizing a combination of music/AV media with any other complementary therapy, including relaxation therapy, acupuncture and guided imagery, in the study group; ii) studies utilizing music/AV media for any procedures other than ESWL; iii) studies not reporting pain and/or anxiety as an outcome measure; iv) non-RCTs; v) studies not comparing music/AV media with a control and vi) review articles, cases series and case reports. In the case of duplication of data, the most updated version of the study was to be included.

Search strategy. An electronic search was performed independently by two reviewers. The PubMed, Embase, Scopus, BioMed Central, Ovid, and Cochrane Central Register of Controlled Trials (CENTRAL) databases were screened up to 15th February 2020. Both MeSH terms and free-text keywords were utilized for searching relevant articles. Search terms and search strategy along with the results of the PubMed database are presented in Table SI. The reviewers screened the search results initially by their titles and abstracts for every database. After identifying potentially pertinent articles, full texts of the articles were sourced. Both the reviewers assessed individual articles based on the inclusion and exclusion criteria. Any disagreements were resolved by discussion. Post-screening, the bibliography of included studies, as well as review articles on the subject, were hand searched for any additional references.

Data extraction and risk of bias assessment. The two reviewers independently extracted data from the included studies. Data regarding authors, publication year, study type, inclusion/exclusion criteria, sample size, demographic details, types of AV media, number of shock waves, ESWL power settings, use of analgesics and study outcomes were extracted. The authors were contacted by e-mail for any missing data. The primary outcome of the analysis was pain scores recorded after the ESWL procedure using the visual analog scale (VAS). The secondary outcome of interest was anxiety measured by the State and Trait Anxiety Inventory-State Anxiety (STAI-SA) test (19).

The Cochrane Collaboration risk assessment tool was used to assess the quality of included studies (18). Studies were assessed by two reviewers independently for: Random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting and other bias.

Statistical analysis. The software 'Open MetaAnalyst (Version 10.10)' was used for the meta-analysis (20). A meta-analysis was conducted only if at least three studies reported data on the same outcome variable. Predicting heterogeneity in the included studies, a random-effects model was used to calculate the pooled effect size for all analyses. Heterogeneity was assessed using the I^2 statistic. I^2 values of 25-50% represented low, values of 50-75% medium and >75% represented substantial heterogeneity. Continuous data were summarized using the mean difference (MD) with 95% confidence intervals (CI). Sub-group analysis was performed for the use of music or AV media and the use of noise-canceling headphones for the primary outcome. For studies not reporting mean and standard deviation scores of outcome variables, the same was calculated from median and interquartile range using methods reported by Wan *et al* (21). A sensitivity analysis was carried out to assess the influence of each study on the pooled effect size. In this analysis, data from the various included studies were excluded one at a time and the effect size was recalculated. Due to the inclusion of fewer than 10 studies in the review, funnel plots were not used to assess publication bias (18).

Results

Study characteristics. A total of 438 unique records were screened and 13 articles were selected for full-text review (Fig. 1). Finally, 11 articles were included in this systematic review and meta-analysis (22-32). Details of the included studies are presented in Table I. Amongst all RCTs, only one was a cross-over trial (27). The sample size per arm in the included studies varied from 21-200 patients. In the study of Karalar *et al* (28), the study sample was further subdivided into two groups, based on the use of noise cancellation headphones. Details of these sub-groups were pooled separately. El-Khoury *et al* (32) conducted a three-arm trial comparing AV media and music with control. The data of these study groups were pooled separately for the meta-analysis. Çift and Benlioglu (24) in a five-arm study, compared three different types of music with two control groups of which one was provided with headphones while the other was not. The results of the study and control groups were pooled using a random-effects meta-analysis model into a single group (18). While different types of music were used in the included studies, two studies used AV media during ESWL. In the study of Marsdin *et al* (22), patients were provided with DVDs and television sets, while in the trial of El-Khoury *et al* (32) an iPad with music videos was used. Media distraction was delivered via headphones in all trials. A total of three studies reported the use of noise-cancellation headphones (28,30,32). In studies reporting adequate data, there was no statistically significant difference in the number of shock waves and the power setting of ESWL between the study and control groups. Seven studies reported the use of analgesic/anxiolytics during the trial (22,23,25,26,28,30,32). In the trial of Yilmaz *et al* (26), midazolam was used only in the control group whereas in all remaining studies drugs were not restricted to any specific group.

The authors' assessment of the quality of included studies is presented in Table II. Appropriate methods of randomization were not used in majority studies. Due to the nature of




Figure 1. Study flow-chart.

the intervention, blinding of patients was not possible. The use of analgesics, which could have introduced bias in pain outcomes, was taken into consideration for ‘other bias’.

Outcomes. A total of nine studies reported pain outcomes on the VAS scale (22-24,26,28-32). Pooled analysis indicated a statistically significant difference in pain outcomes in favor of media distraction (MD: -1.18; 95% CI: -2.35, -0.01; $I^2=96.8\%$; Fig. 2). On subgroup analysis based on the type of media distraction, the use of both, AV media (MD: -2.94; 95% CI: -4.70, -1.17; $I^2=79.2\%$) and music (MD: -0.86; 95% CI: -1.37, -0.35; $I^2=62.5\%$), was indicated to lead to a significant reduction in pain outcomes (Fig. 2). Results were, however, not stable on sensitivity analysis. On the exclusion of the studies of Karalar *et al* (28) and El-Khoury *et al* (32), no significant effect of media distraction on pain outcomes was identified (Fig. 3). Data from study groups using noise-cancellation headphones were pooled separately in a sub-group analysis. The results indicated that there was a statistically significant reduction in pain scores with the use of noise-cancellation headphones (MD: -1.20; 95% CI: -2.17, -0.23; $I^2=69.8$) but no such effect

was seen with regular headphones (MD: -1.13; 95% CI: -2.66, 0.40; $I^2=97.2\%$) (Fig. 4).

There were two studies that did not report pain outcomes as mean \pm standard deviation. E-mail to the corresponding author for these missing data did not elicit a response. In the trial of Cepeda *et al* (25), alfentanil consumption was compared in the study and control groups. The authors reported no statistically significant difference in analgesic use and pain intensity during ESWL in the two groups. Akbas *et al* (27), on the other hand, in a cross-over RCT, reported a statistically significant reduction of pain with the use of music after both treatment sessions.

Anxiety outcomes using STAI-SA were reported by five studies (24,26,28,29,31). All studies had used only music in the study groups. Pooled analysis indicated a statistically significant reduction of anxiety scores with the use of media distraction (MD: -3.91; 95% CI: -6.44, -1.38; $I^2=77.7\%$; Fig. 5). The results were stable on sensitivity analysis as there was no change in significance on exclusion of one study at a time (Fig. 6). Exclusion of the study of Yilmaz *et al* (26) used midazolam in the control group and the ‘noise cancellation headphone’ group of Karalar *et al* (28) did not change the significance of results. Ordaz Jurado *et al* (30) compared anxiety

Table I. Characteristics of included studies.

Author/ Year	Inclusion criteria	Exclusion criteria	Sample size		Mean age		Male sex (%)		Media distraction	Number of shock waves		ESWL power settings		Analgesia during procedure
			Study	Control	Study	Control	Study	Control		Study	Control	Study	Control	
Cepeda <i>et al</i> (25), 1998	Age: 15-65 years First ESWL	Serum creatinine >1.5 mg/dl History of opioid use	97	96	40.7	41	49.4	48.9	Patient selected classical, Caribbean, soft rock, and jazz music via headphones started 10 mins before procedure and continued up to 10 mins post- procedure	NR	NR	All patients reached 17 V	Morphine 10 mg SC with ketor olac 30 mg IV + alfentanil PCA (0.25 mg dose with 1 min lockout interval)	
Koch <i>et al</i> (23), 1998	NR	NR	21	22	54	53	80	44	Any music pf patients choice via headphone during procedure	NR	NR	NR	Midazolam and Alfentanil 10 µg/kg PCA	
Yilmaz <i>et al</i> (26), 2003	Age: 19-68 years	NR	48	50	40.8	39.3	60.4	64	Relaxing, slow rhythm sedative music via headphones during procedure	3043± 110.42	3016± 64.22	21.96± 0.54 kV 21.03± 0.48 kV	15 g emla cream over flank, 25 mg fentanyl IV given to patients reporting pain. Midazolam 2 mg in control group only	

Table I. Continued.

Author/ Year	Inclusion criteria	Exclusion criteria	Sample size		Mean age		Male sex (%)		Media distraction	Number of shock waves		ESWL power settings		Analgesia during procedure
			Study	Control	Study	Control	Study	Control		Study	Control	Study	Control	
Marsdin <i>et al</i> (22), 2012	Elective outpatient ESWL patients	Ureteric stents Emergency patients	61	57	51.7	53.5	54	56	32-inch television with 30 different DVDs of 30-min programs provided with wireless headphones	3948± NR	3764± NR	NR	NR	Pethidine 100 mg IM and diclofenac 75 mg IM
Akbas <i>et al</i> (27), 2016	Age: >18 y No preoperative pain No analgesic use before the procedure No ureteral stent inserted Radio-opaque stone	Not completed procedure Development of complications Refusing to listen to music Not completing the second session	200	200	-	-	-	-	Turkish folk, classical, popular music during procedure	2724± 492	2710± 551	17.71± 2.59 J	17.04± 2.07 J	None
Karalar <i>et al</i> (28), 2016	Radio- opaque renal stone	Prior ESWL Ureteric stents Renal colic	S1: 28 S2: 29	32	S1: 47.8 S2: 47.7	48.2	S1: 61 S2: 62	59	S1: Turkish classical music by noise cancellation headphones during procedure S2: Turkish classical music via non noise cancellation headphones during procedure	2913± 444	2828± 272	20.1± 1.1 kV	19.9± 1.3 kV	Diclofenac 75 mg IM

Table I. Continued.

Author/ Year	Inclusion criteria	Exclusion criteria	Sample size		Mean age		Male sex (%)		Media distraction		Number of shock waves		ESWL power settings		Analgesia during procedure
			Study	Control	Study	Control	Study	Control	Study	Control	Study	Control	Study	Control	
Cakmak <i>et al</i> (29), 2017	Age: >18 y Radio-opaque renal stone Located in upper/middle calices, renal pelvis: <20 mm in diameter Proximal ureter: <10 mm in diameter	Prior ESWL Ureteric stents Hearing disorders Uncontrolled hypertension History of psychiatric disease and antidepressant drug use	95	105	NR	NR	NR	NR	Turkish folk, classical, popular, relaxing or slow rhythm music via headphones during procedure	2764.5± 699.1	2777.9± 724.9	17.4± 5.3 kV	17.5± 5.5 kV	NR	NR
Jurado <i>et al</i> (30), 2017	Age: 18-80 y diameter	Prior ESWL Hearing loss Analgesia prior to procedure Regularly taking painkillers	42	53	52.6	52	54.8	69.8	Patient selected classical, chill out, pop, jazz music via by noise cancellation headphones during procedure	NR ^a	NR	NR ^a	NR	Pethidine 20 mg IV and paracetamol 1 g IV. 10 cc (max 50 cc) of pethidine given to patients reporting pain.	NR
Gezginci <i>et al</i> (31), 2018	Age: >18 y First ESWL BMI: <30 kg/m ² No contradiction for ESWL	Analgesic before procedure Renal colic before procedure Ureteric stents	40	40	NR ^a	NR	33.3	35.2	Turkish (folk, art, arabesque, pop) music, foreign pop music, rock and classical music via headphones during procedure	Targeted 3,000 per session	NR	NR	NR	None	NR

Table I. Continued.

Author/ Year	Inclusion criteria	Exclusion criteria	Sample size		Mean age		Male sex (%)		Media distraction		Number of shock waves		ESWL power settings		Analgesia during procedure
			Study	Control	Study	Control	Study	Control	Study	Control	Study	Control	Study	Control	
EI-Khoury <i>et al</i> (32), 2020	Adult patients	NR	AV media: 33 Music: 30	31	AV media: 52.4 Music: 51.5	52.6	AV media: 70 Music: 90	71	AV media: music videos using Apple iPad Music: Any music/local radio Both delivered by noise cancellation headphones during procedure	AV media: 3205.9± 545.3 Music: 3175.1± 434.1	3150.4± 424.5	AV media: 74.3± 10 kV Music: 74.3± 9.1 kV	70.4±16 kV	NSAID suppository offered to all patients. Breakthrough pain treated by Fentanyl IM	
Çift <i>et al</i> (24), 2020	Age: >18 years Radiopaque stone 10-20 mm in diameter localized in the renal pelvis or ureteropelvic junction Absence of renal colic before the procedure	Previous ESWL history Ureteral stenosis Hearing deficit, Uncontrolled hypertension Antidepressant drug usage	S1: 30 S2: 30 S3: 30	C1: 30 C2: 30	S1: 37.7 S2: 36.3 S3: 36.4	C1: 38.4 C2: 39.2	S1: 83.3 S2: 70 S3: 73.3	C1: 53.3 C2: 56.7	S1: Turkish art music S2: Western classical music S3: Patient selected any type of music. All delivered via headphones during procedure C1: No headphones, no music C2: Headphones on, no music	S1: 2851.5 (2685.7- 3000) S2: 2842.5 (2437.2- 2958.7) S3: 2800 (2500- 2925)	C1: 2700 (2500- 2900) C2: 2725 (2500- 2900)	S1: 92 (82- 101.25) S2: 90 (64.25- 103.5) S3: 94 (79.75- 96.25)	C1: 80 (74.75- 92.5) C2: 85 (65.0- 93.25)	None	

Data presented as the mean ± SD. *No difference between the two groups for number of shock waves and energy, but data not reported as the mean ± SD. AV, audiovisual; ESWL, extracorporeal shock wave lithotripsy; PCA, patient controlled analgesia; SC, subcutaneous; IM, intramuscular; IV, intravenous; BMI, body mass index; S1, study group 1; S2, study group 2; S3, study group 3; C1, control group 1; C2, control group 2; NR, not reported; V, volts; kV, kilovolts; J, Joules.

Table II. Risk of bias in included studies.

Study	Random sequence generation	Allocation concealment	Blinding of participants and personnel	Blinding of outcome assessment	Incomplete outcome data	Selective reporting	Other bias
Cepeda <i>et al</i> (25)	Unclear risk	Unclear risk	High risk	High risk	Low risk	Unclear risk	Unclear risk
Koch <i>et al</i> (23)	Unclear risk	Unclear risk	High risk	High risk	Low risk	Unclear risk	Unclear risk
Yilmaz <i>et al</i> (26)	Unclear risk	Unclear risk	High risk	High risk	Low risk	Unclear risk	High risk
Marsdin <i>et al</i> (22)	Unclear risk	Unclear risk	High risk	High risk	Low risk	Unclear risk	Unclear risk
Akbas <i>et al</i> (27)	Unclear risk	Unclear risk	High risk	High risk	Low risk	Unclear risk	Low risk
Karalar <i>et al</i> (28)	High risk	Unclear risk	High risk	High risk	Low risk	Unclear risk	Unclear risk
Cakmak <i>et al</i> (29)	Unclear risk	Unclear risk	High risk	High risk	Low risk	Unclear risk	Low risk
Jurado <i>et al</i> (30)	Low risk	Unclear risk	High risk	Low risk	Low risk	Unclear risk	Unclear risk
Gezginci <i>et al</i> (31)	Low risk	Unclear risk	High risk	High risk	Low risk	Low risk	Low risk
El-Khoury <i>et al</i> (32)	Low risk	Unclear risk	High risk	High risk	Low risk	Unclear risk	Unclear risk
Çift <i>et al</i> (24)	Low risk	Low risk	High risk	Low risk	Low risk	Unclear risk	Low risk

in the two groups using a 10-point VAS scale and reported no statistically significant difference between the two groups.

Discussion

After a comprehensive review of the literature, the results of the present study indicated that media distraction in the form of AV media or music may lead to a statistically significant reduction of pain and anxiety in patients undergoing ESWL. Data from a limited number of studies also indicate that the use of noise-canceling headphones may lead to a significant reduction in pain and that such an effect may not be seen with regular headphones.

Several non-pharmacological methods of pain management, including massage, relaxation therapy, acupuncture, hypnosis, aromatherapy, music and AV distraction, have been used for a variety of healthcare-procedures worldwide (33,34). Music interventions for pain management can be in the form of music therapy, where a trained professional selects tempo-controlled melodies to have a calming effect, or in the form of random music listening which acts as a distraction tool similar to AV aids (35,36). Owing to its ease of application, a number of studies have assessed the efficacy of such music and/or AV distraction for different medical and surgical procedures. In a recent systematic review and meta-analysis of nine RCTs, Song *et al* (11) have demonstrated significantly reduced pain and anxiety with the use of music in patients undergoing biopsy. Another RCT has demonstrated that listening to music during labor significantly reduces pain and anxiety of mothers during all stages of labor (37). Similarly, AV aids are effective to reduce pain in patients with sickle cell disease (38), those undergoing colonoscopy (12), as well as those undergoing minor surgeries (39).

The application of media distraction, specifically music, during urological procedures has been evaluated by several trials with conflicting results. Chang *et al* (40) in a pilot RCT found music to be effective in reducing pain, discomfort, and dissatisfaction in patients undergoing transrectal ultrasound (TRUS)-guided prostate biopsy, however, no such effect was noted in a similar trial by Packiam *et al* (41). Music has also been used to reduce patient-reported pain and anxiety for other urological procedures, including percutaneous nephrostomy tube placement and urodynamic studies (42,43). As different urological procedures entail different degrees of pain, the current review was focused solely on ESWL to ensure reduced bias owing to non-homogenous procedures. Pain during ESWL can be attributed to several reasons, including trauma caused by shock waves passing through skin and muscles, stretching of the renal capsule where therapy is delivered, or due to an increase in the internal pressure of the kidney (7). Many factors can influence the degree of pain, including the dimensions and position of the urinary stone, frequency of shock waves, power settings, sex and the pain threshold of the patient and use of pre-procedural analgesic (44). As all included studies were RCTs, it is hypothesized that the effect of such baseline variables on outcomes is negated.

The results of the present analysis indicate that media distraction may significantly reduce pain and anxiety scores in patients undergoing ESWL. Similar results were obtained by Kyriakides *et al* (15) in their meta-analysis of the role




Figure 2. Forest plot of studies comparing media distraction vs. control for the outcome pain with sub-group analysis based on the type of media distraction (audiovisual or music). *Indicates different sub-group from the same study.




Figure 3. Sensitivity analysis of pain outcome presenting the resultant effect size after exclusion of one study at a time. Black squares in front of each study indicate the effect size and horizontal lines indicate the 95% confidence intervals, after exclusion of that study from the analysis. *Indicates different sub-group from the same study.

of music in reducing pain and anxiety for any urological procedure. Compared to the present review, they focused only on music and were able to include only six studies for ESWL. The present results also concur with a very recent systematic review of Saraogi *et al* (45), which indicates that complementary medicine strategies including music, acupressure, acupuncture, transcutaneous electrical nerve stimulation and audiovisual distractions can reduce pain and anxiety in patients undergoing ESWL.

In a sub-group analysis based on the type of media, both AV-media and music were found to significantly reduce pain. However, results should be interpreted with caution, as only two studies were available for the sub-group analysis of AV media. Also, relevant data on anxiety were reported by only five studies. While assessing the results of patient-reported outcomes, it is

important to bear in mind the concept of minimal clinically important difference (MCID). The MCID is defined as the smallest change in a measurement that signifies an important improvement in a symptom (46). Data from non-urological studies indicates that the MCID for the VAS scale is 1.4 while for STAI-SA is 10 (47,48). The pooled MD of VAS scores of all studies in the present analysis was -1.18 (95% CI: -2.35, -0.01), for studies on music was -0.86 (95% CI: -1.37, -0.35) and for STAI-SA scores was -3.91 (95% CI: -6.44, -1.38). Interpreted in the context of the reported MCID, it can be noted that the effect size of the intervention was small and the upper limits of 95% CI were close to zero for both pain and anxiety outcomes. This can have important clinical implications as, even though studies suggest a significant difference in pain and anxiety




Figure 4. Forest plot of studies comparing media distraction vs. control for the outcome pain with sub-group analysis based on the use of noise-cancelling headphones (NCH). *Indicates different sub-group from the same study.




Figure 5. Forest plot of studies comparing media distraction vs. control for the outcome anxiety.




Figure 6. Sensitivity analysis of anxiety outcome presenting the resultant effect size after exclusion of one study at a time.*Indicates different sub-group from the same study.

outcomes with media distraction, the resultant effect may not produce a clinically significant improvement in symptoms.

In another sub-group analysis, the present results indicated a significant reduction in pain scores with the use of noise cancellation headphones, but no such effect was noted with

regular headphones. Since ESWL is a noisy procedure, the continuous unpleasant sound of shockwaves may affect patient comfort and anxiety (28). Therefore, active noise control employed by such noise-cancellation headphones may lend a more comfortable environment for ESWL thereby influencing

pain outcomes. However, since only three studies utilized this technology, results should be interpreted with caution.

The limitations of the present review should be mentioned. Firstly, the quality of the included studies in the present review was not high. The absence of adequate methods of randomization and lack of blinding, due to the nature of the intervention, may have influenced outcomes. Secondly, there were methodological variations amongst included studies for the type of music/AV media, choice of music, use of pre-procedural analgesics and patient-controlled analgesia. This may have contributed to the high heterogeneity in the present meta-analyses. Thirdly, data from two studies were not available for the primary outcome and only five studies were pooled for the secondary outcome, which has limited the power of the present analyses. Further, all studies assessing the role of media distraction on anxiety utilized only music as the intervention. The role of AV media on anxiety scores could not be assessed. Fourthly, due to a limited number of studies in the analysis and limitations of the meta-analysis software, publication bias could not be assessed in the present review. Lastly, the results for pain were not stable on sensitivity analysis, as the exclusion of two studies resulted in non-significant results.

The present study has its strengths in being the first meta-analysis focused on the role of media distraction during ESWL. Only RCTs were analyzed to provide level-I evidence. Sensitivity and subgroup analysis were performed to assess the influence of each study on the effect size and to account for inter-study variations.

To conclude, the present study indicates that media distraction in the form of AV media or music may be beneficial in reducing the pain and anxiety of patients undergoing ESWL. The use of noise-canceling headphones to deliver the intervention may have some benefit over regular headphones. Evidence is, however, weak considering the small effect size, confidence intervals being close to zero and instability of the results on sensitivity analysis. In clinical practice, media distraction may be used during ESWL as a nursing intervention, but a clinically important reduction of pain and anxiety may not be expected.

Acknowledgements

Not applicable.

Funding

No funding was received.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

WH conceived and designed the study. KY, LZ and XL collected the data, performed the literature search and were responsible for confirming the authenticity of all the raw data. WH was involved in the writing of the manuscript. All authors have read and approved the final manuscript.

Ethical approval

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Amato M, Lusini ML and Nelli F: Epidemiology of nephrolithiasis today. *Urol Int* 72 (Suppl 1): S1-S5, 2004.
- Raheem OA, Khandwala YS, Sur RL, Ghani KR and Denstedt JD: Burden of urolithiasis: Trends in prevalence, treatments, and costs. *Eur Urol Focus* 3: 18-26, 2017.
- Kim CH, Shin DS, Kim TB and Jung H: The efficacy of early extracorporeal shockwave lithotripsy for the treatment of ureteral stones. *Urol J* 16: 331-336, 2019.
- Gupta NP and Kumar A: Analgesia for pain control during extracorporeal shock wave lithotripsy: Current status. *Indian J Urol* 24: 155-158, 2008.
- Chaussy C, Schmiedt E, Jocham D, Brendel W, Forssmann B and Walther V: First clinical experience with extracorporeally induced destruction of kidney stones by shock waves. *J Urol* 127: 417-420, 1982.
- Bach C, Zaman F, Kachrilas S, Kumar P, Buchholz N and Masood J: Drugs for pain management in shock wave lithotripsy. *Pain Res Treat* 2011: 259426, 2011.
- Ozsaker E and Dirmali A: The effect of transcutaneous electrical nerve stimulation for pain relief during extracorporeal shock-wave lithotripsy procedure. *Pain Manag Nurs* 15: 59-68, 2014.
- Brown MA: The role of nurses in pain and palliative care. *J Pain Palliat Care Pharmacother* 27: 300-302, 2013.
- Zins S, Gross C and Hooke MC: Complementary therapies for pain among individuals receiving hemodialysis: A systematic review. *Nephrol Nurs J* 45: 13-23, 2018.
- Aydin D, Sahiner NC and Ciftci EK: Non-pharmacological strategies used to reduce procedural pain in infants by nurses at family health centres. *J Pak Med Assoc* 67: 889-894, 2017.
- Song M, Li N, Zhang X, Shang Y, Yan L, Chu J, Sun R and Xu Y: Music for reducing the anxiety and pain of patients undergoing a biopsy: A meta-analysis. *J Adv Nurs* 74: 1016-1029, 2018.
- Xiaolian J, Xiaolin L and Lan ZH: Effects of visual and audiovisual distraction on pain and anxiety among patients undergoing colonoscopy. *Gastroenterol Nurs* 38: 55-61, 2015.
- Drahota A, Galloway E, Stores R, Ward D, Severs M and Dean T: Audiovisual distraction as an adjunct to pain and anxiety relief during minor surgery. *Foot (Edinb)* 18: 211-219, 2008.
- Sahiner NC and Bal MD: The effects of three different distraction methods on pain and anxiety in children. *J Child Health Care* 20: 277-285, 2016.
- Kyriakides R, Jones P, Geraghty R, Skolarikos A, Liatsikos E, Traxer O, Pietropaolo A and Somani BK: Effect of music on outpatient urological procedures: A systematic review and meta-analysis from the European Association of Urology section of uro-technology. *J Urol* 199: 1319-1327, 2018.
- Goh NM, Drake T, Rai BP and Somani BK: Complementary approaches to decreasing discomfort during shockwave lithotripsy (SWL). *Urolithiasis* 42: 189-193, 2014.
- Moher D, Liberati A, Tetzlaff J and Altman DG; PRISMA Group: Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med* 6: e1000097, 2009.
- Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page M and Welch V (eds): *Cochrane handbook for systematic reviews of interventions*. Version 6. Cochrane, 2019.
- Elwood LS, Wolitzky-Taylor K and Olatunji BO: Measurement of anxious traits: A contemporary review and synthesis. *Anxiety Stress Coping* 25: 647-666, 2012.
- Wallace BC, Schmid CH, Lau J and Trikalinos TA: Meta-analyst: Software for meta-analysis of binary, continuous and diagnostic data. *BMC Med Res Methodol* 9: 80, 2009.

21. Wan X, Wang W, Liu J and Tong T: Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 14: 135, 2014.
22. Marsdin E, Noble JG, Reynard JM and Turney BW: Audiovisual distraction reduces pain perception during shockwave lithotripsy. *J Endourol* 26: 531-534, 2012.
23. Koch ME, Kain ZN, Ayoub C and Rosenbaum SH: The sedative and analgesic sparing effect of music. *Anesthesiology* 89: 300-306, 1998.
24. Çift A and Benlioglu C: Effect of different musical types on patient's relaxation, anxiety and pain perception during shock wave lithotripsy: A randomized controlled study. *Urol J* 17: 19-23, 2020.
25. Cepeda MS, Diaz JE, Hernandez V, Daza E and Carr DB: Music does not reduce alfentanil requirement during patient-controlled analgesia (PCA) use in extracorporeal shock wave lithotripsy for renal stones. *J Pain Symptom Manage* 16: 382-387, 1998.
26. Yilmaz E, Ozcan S, Basar M, Basar H, Batislam E and Ferhat M: Music decreases anxiety and provides sedation in extracorporeal shock wave lithotripsy. *Urology* 61: 282-286, 2003.
27. Akbas A, Gulpinar MT, Sancak EB, Karakan T, Demirbas A, Utangaz MM, Dede O, Sancaktutar AA, Simsek T, Sahin B and Resorlu B: The effect of music therapy during shockwave lithotripsy on patient relaxation, anxiety, and pain perception. *Ren Fail* 38: 46-49, 2016.
28. Karalar M, Keles I, Doğanekin E, Kahveci OK and Sarici H: Reduced pain and anxiety with music and noise-canceling headphones during shockwave lithotripsy. *J Endourol* 30: 674-677, 2016.
29. Cakmak O, Cimen S, Tarhan H, Ekin RG, Akarken I, Ulker V, Celik O, Yucel C, Kisa E, Ergani B, *et al*: Listening to music during shock wave lithotripsy decreases anxiety, pain, and dissatisfaction: A randomized controlled study. *Wien Klin Wochenschr* 129: 687-691, 2017.
30. Ordaz Jurado DG, Budia Alba A, Bahilo Mateu P, Trassierra Villa M, López-Acón D and Boronat Tormo F: Litotricia por ondas de choque con música: Un tratamiento menos doloroso y más satisfactorio/Shockwave lithotripsy with music: Less painful and more satisfactory treatment. *Actas Urol Esp* 41: 584-589, 2017.
31. Gezginç E, Iyigün E, Yalçın S, Bedir S and Özgök İY: Comparison of two different distraction methods affecting the level of pain and anxiety during extracorporeal shock wave lithotripsy: A randomized controlled trial. *Pain Manag Nurs* 19: 295-302, 2018.
32. El-Khoury HJ, Brookes JDL, Tan B, Shahbaz S and McCahy PJ: Impact of modern media on analgesic requirements during extracorporeal shockwave lithotripsy. *ANZ J Surg* 90: 53-56, 2020.
33. Smith CA, Levett KM, Collins CT, Armour M, Dahlen HG and Sukanuma M: Relaxation techniques for pain management in labour. *Cochrane Database Syst Rev* 3: CD009514, 2018.
34. Lin YC, Wan L and Jamison RN: Using integrative medicine in pain management: An evaluation of current evidence. *Anesth Analg* 125: 2081-2093, 2017.
35. Bradt J, Potvin N, Kesslick A, Shim M, Radl D, Schriver E, Gracely EJ and Komarnicky-Kocher LT: The impact of music therapy versus music medicine on psychological outcomes and pain in cancer patients: A mixed methods study. *Support Care Cancer* 23: 1261-1271, 2015.
36. Gold C, Erkkilä J, Bonde LO, Trondalen G, Maratos A and Crawford MJ: Music therapy or music medicine? *Psychother Psychosom* 80: 304-305, 2011.
37. Simavli S, Gumus I, Kaygusuz I, Yildirim M, Usluogullari B and Kafali H: Effect of music on labor pain relief, anxiety level and postpartum analgesic requirement: A randomized controlled clinical trial. *Gynecol Obstet Invest* 78: 244-250, 2014.
38. Ezenwa MO, Yao Y, Engeland CG, Molokie RE, Wang ZJ, Suarez ML and Wilkie DJ: A randomized controlled pilot study feasibility of a tablet-based guided audio-visual relaxation intervention for reducing stress and pain in adults with sickle cell disease. *J Adv Nurs* 72: 1452-1463, 2016.
39. Hudson BF and Ogden J: Exploring the impact of intraoperative interventions for pain and anxiety management during local anesthetic surgery—a systematic review and meta-analysis. *J Perianesth Nurs* 31: 118-133, 2016.
40. Chang YH, Oh TH, Lee JW, Park SC, Seo IY, Jeong HJ and Kwon WA: Listening to music during transrectal ultrasound-guided prostate biopsy decreases anxiety, pain and dissatisfaction in patients: A pilot randomized controlled trial. *Urol Int* 94: 337-341, 2015.
41. Packiam VT, Nottingham CU, Cohen AJ, Eggner SE and Gerber GS: No effect of music on anxiety and pain during transrectal prostate biopsies: A randomized trial. *Urology* 117: 31-35, 2018.
42. Hamidi N and Ozturk E: The effect of listening to music during percutaneous nephrostomy tube placement on pain, anxiety, and success rate of procedure: A randomized prospective study. *J Endourol* 31: 457-460, 2017.
43. Diri MA, Çetinkaya F and Gül M: The effects of listening to music on anxiety, pain, and satisfaction during urodynamic study: A randomized controlled trial. *Urol Int* 103: 444-449, 2019.
44. Kumar A, Gupta NP, Hemal AK and Wadhwa P: Comparison of three analgesic regimens for pain control during shockwave lithotripsy using dornier delta compact lithotripter: A randomized clinical trial. *J Endourol* 21: 578-582, 2007.
45. Saraogi M, Geraghty RM, Hameed B, Rob S, Pietropaolo A, Sarica K, Gozen A, Liatsikos E and Somani BK: Role of complementary medicine (music, acupuncture, cupressure, TENS and audio-visual distraction) in shockwave lithotripsy (SWL): A systematic review from EAU sections of urolithiasis (EULIS) and uro-technology (ESUT). *Urology* 145: 38-51, 2020.
46. Tashjian RZ, Deloach J, Porucznik CA and Powell AP: Minimal clinically important differences (MCID) and patient acceptable symptomatic state (PASS) for visual analog scales (VAS) measuring pain in patients treated for rotator cuff disease. *J Shoulder Elbow Surg* 18: 927-932, 2009.
47. Delgado DA, Lambert BS, Boutris N, McCulloch PC, Robbins AB, Moreno MR and Harris JD: Validation of digital visual analog scale pain scoring with a traditional paper-based visual analog scale in adults. *J Am Acad Orthop Surg Glob Res Rev* 2: e088, 2018.
48. Corsaletti BF, Proença M-DGL, Bisca GKW, Leite JC, Bellinetti LM and Pitta F: Minimal important difference for anxiety and depression surveys after intervention to increase daily physical activity in smokers. *Fisioter E Pesqui* 21: 359-364, 2014.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.