

Case Report

Intraventricular meningioma admixed with choroid plexus papilloma: a rare case report with review of literature

So-Woon Kim¹, Jiwook Ryu², Kiyong Na¹

¹Department of Pathology, Kyung Hee University Hospital, College of Medicine, Kyung Hee University, Seoul, Republic of Korea; ²Department of Neurosurgery, Kyung Hee University Hospital, College of Medicine, Kyung Hee University, Seoul, Republic of Korea

Received November 23, 2020; Accepted February 4, 2021; Epub April 15, 2021; Published April 30, 2021

Abstract: A 42-year-old male presented with a history of headaches for the previous 2 weeks. Magnetic resonance imaging of the brain showed a 3 cm-sized well-defined, enhancing mass in the atrium of the right lateral ventricle. The tumor comprised two heterogeneous components: approximately one-third of the tumor exhibited complex and delicate papillary fibrovascular cores lined with uniform cuboidal-to-columnar epithelial cells, whereas the remaining part was seen as a solid sheet comprising ovoid-to-spindle cells with plump cytoplasm, which occasionally had a whorling pattern. Further, immunohistochemical staining with cytokeratin 7 (CK7) and epithelial membrane antigen (EMA) clearly demarcated each component: the CK7+/EMA- choroid plexus papilloma and CK7-/EMA+ meningioma. This report provides a description of an unusual case of concomitant choroid plexus papilloma and ventricular meningioma presenting as a single mass, along with a review of relevant literature.

Keywords: Choroid plexus papilloma, meningioma, intraventricular tumor

Introduction

Intraventricular tumors account for 2%-7% of all intracranial neoplasms [1]. They are classified according to their pathologic origins, which may include the walls of the ventricular system, the septum pellucidum, and the choroid plexus [2]. Based on the composition and patient's age, differential diagnosis primarily reveals choroid plexus papilloma (CPP) in patients under the age of 10 years, low-grade glioma in patients aged 10-40 years, and metastases, lymphoma, and meningioma (MG) after the fourth decade of life [3].

MG is the most common intracranial tumor, accounting for approximately 20% of all primary brain neoplasms [4]. They typically arise in the supratentorial compartment, and intraventricular MG constitutes approximately 0.5%-2% of all intracranial MGs [4, 5]. CPP is a rare benign neoplasm of the central nervous system, representing 0.3%-0.8% of all brain neoplasms and <1% of all intracranial neoplasms in adults [6]. CPP is commonly found surrounded by normal

choroid plexus structures such as the lateral ventricle [1, 6]. In childhood, CPP usually develops in the lateral ventricle with its atrial level as the most common site [7]. In adults, it frequently occurs in the fourth ventricle and/or lateral recess [8].

The occurrence of brain tumors with different pathologies in a single mass is rare [9]. In particular, the simultaneous presentation of a CPP and an MG as a single mass in the intraventricular system has never been reported. Herein, we describe an unusual case of concomitant CPP and MG in the lateral ventricle, supplemented with a review of pertinent literature.

Case presentation

Clinical features

A 42-year-old male presented with a history of headache for the previous 2 weeks. The pain was described as dull, intermittent, and not localized to one side. Neurological examination showed no signs of sensory, motor, or cranial

nerve abnormalities. He had a history of hepatitis B virus-associated liver cirrhosis, but no history of surgery, stroke, or a familial history of stroke or cancer. Magnetic resonance imaging (MRI) of the brain showed a well-defined and enhancing mass in the atrium of the right lateral ventricle, measuring 3.3 × 2.8 × 2.6 cm (**Figure 1A**). Obstructive dilatation of the occipital and temporal horn of the right lateral ventricle was observed. Based on radiologic examination, possible differential diagnoses included intraventricular MG, CPP, and exophytic intra-axial tumors. Cerebral angiography revealed that the mass was supplied by the right anterior and posterior choroidal arteries. Tumor embolization was technically difficult because of the thin wall of the supplying blood vessel. Therefore, the mass was surgically resected.

Pathologic findings

Microscopic examination of the mass revealed two different tumors: CPP and MG (**Figure 1B**). The CPP was composed of complex and delicate papillary fibrovascular cores covered by uniform cuboidal-to-columnar epithelial cells. Tumor cells were round to oval with monotonous, basally located nuclei and plump eosinophilic pale cytoplasm (**Figure 1C**). The border of the CPP was partially intermingled with the MG component, which was arranged in lobules with occasional whorling patterns. Tumor cells had ovoid-to-spindle shaped nuclei with pink cytoplasm and indistinct cell membranes (**Figure 1D**). In both the CPP and MG, cytologic atypia was minimal and mitotic figures were absent.

Immunohistochemical staining

Immunohistochemical staining of the CPP was strong and diffuse positive for cytokeratin 7 (CK7, 1:200, clone OV-TL, Dako, Glostrup, Denmark) and was negative for the epithelial membrane antigen (EMA, 1:100, clone GP1.4, Leica Biosystems: Novocastra, UK) (**Figure 1E**). In contrast, tumor cells in the MG were diffusely positive for EMA and negative for CK7 (**Figure 1F**). The proliferative labeling index measured by Ki-67 expression (1:100, clone MIB-1, Dako, Glostrup, Denmark) was less than 2% for both tumors.

Clinical course

The patient was discharged on postoperative day 13 without complications. Follow-up brain

imaging revealed residual linear or nodular enhancement along the right choroid plexus and right ventricular wall with hemorrhage, indicative of a residual tumor at the surgical site. Three months later, gamma knife radiosurgery was performed on the suspected residual tumor. The patient was alive without tumor recurrence 5 months after surgery.

Discussion

Here, we report an unusual case of a CPP and an MG constituting a single mass in the lateral ventricle. To date, only two cases with coincidental CPP and MG have been reported in the English literature (**Table 1**) [10, 11]. In Case 1 (**Table 1**), a 45-year-old female experienced headache for 2 months, accompanied by sensory and motor impairments due to extensive involvement of the lower cranial nerves and tumor adherence to the medullary pia. The CPP and MG were separate but both tumors had developed in the extraventricular area. In Case 2 (**Table 1**), a 45-year-old female complained of progressive confusion and altered sensory and motor functions. The CPP was filling the fourth ventricle, whereas the MG was distinctly located at the craniocervical junction. Unlike these cases, the present case showed an admixture of CPP and MG within a single mass in the intraventricular space. As the tumor was located in the ventricle, clinical presentation was mainly characterized by obstructive hydrocephalus and functional deficits in the neurological systems were absent.

Patients with two distinct primary intracranial neoplasms are very rare [12, 13]. In particular, it is extremely rare for multiple primary intracranial neoplasms to develop concurrently in patients without prior intracranial radiotherapy or underlying phacomatoses such as neurofibromatosis-2 or tuberous sclerosis [14]. Approximately, fewer than 100 cases of multiple concurrent primary intracranial neoplasms have been reported, and the majority of coincidental intracranial neoplasms are MGs and gliomas [14, 15]. Of all primary intracranial neoplasms, MGs are well-known to occur concurrently with other types of tumors [16, 17]. This may be due to the indolent course of such tumors, which allows its prolonged clinical evolution before diagnosis. Further, detection of MGs is primarily incidental [13, 18].

Choroid plexus papilloma and meningioma




Figure 1. (A) Brain MRI shows a well-defined and intense enhancing mass in the atrium of the right lateral ventricle. (B) The ventricular tumor comprises two distinct components: a choroid plexus papilloma (white arrow) and meningioma (black arrow). (C) In the choroid plexus papilloma, complex and delicate papillary fibrovascular fronds are covered by a single layer of uniform cuboidal-to-columnar epithelial cells. (D) Meningioma cells are arranged in lobules and are packed together in short fascicles, forming whorls and syncytial structures. (E) In the choroid plexus papilloma, tumor cells are negative for immunohistochemical staining of the epithelial membrane antigen (EMA, 1:100, clone GP1.4, Leica Biosystems: Novocastra, UK). (F) In contrast, tumor cells are diffusely and strongly positive for EMA in the meningioma component. (B-D, Hematoxylin and eosin; B, Scan view; C, D, $\times 40$; inset, $\times 400$; E, F, EMA, $\times 200$).

An admixture of CPP and MG within a single mass may complicate diagnosis during patho-

logic examinations. The main histologic differential diagnoses for such tumor samples in-

Choroid plexus papilloma and meningioma

Table 1. Summary of cases with coincidental choroid plexus papilloma and meningioma

Case (Ref.)	Age/Sex	Clinical presentation	Tumor location and size	Tumor continuity
1 (10)	45/F	Headache for 2 months Sensory and motor deficit	CPP: Petrotentorial junction, 2.5 cm MG: Cervicomedullary junction, 5 cm	Noncontiguous
2 (11)	45/F	Progressive confusion for 2 months Sensory and motor deficit	CPP: Fourth ventricle, 4.4 cm MG: Craniocervical junction, 2.3 cm	Noncontiguous
Present case	42/M	Headache for 2 weeks No sensory and motor deficit	CPP and MG: Lateral ventricle, 3 cm	Contiguous

Abbreviations: CPP, choroid plexus papilloma; MG, meningioma.

clude papillary MG, papillary ependymoma, and metastatic carcinoma with papillary features [1]. Papillary MG is an MG in which the perivascular pseudopapillary pattern involves most of the tumor [20]. The detection of high-grade histologic features with aggressive presentation is sufficient for a conclusive diagnosis of a papillary MG [1]. Papillary ependymomas exhibit finger-shaped processes lined with a single layer of cuboidal cell with smooth surfaces as well as glial fibrillary acidic protein-positive tumor cell processes. CPP can be distinguished from a papillary ependymoma by bumpy and hobnail cellular surfaces that are not characterized by extensive glial fibrillary acidic protein immunoreactivity [1].

The pathophysiology of the simultaneous development of multiple primary intracranial neoplasms at the same location remains unclear, especially the occurrence of intermingled components of CPP and MG. Various mechanisms have been proposed to explain concurrent primary intracranial tumors [20-28]. First, the concurrent occurrence of various tumor types at the same anatomical site may be due to the collision of two discrete tumors [20-22]. Second, the first tumor may act as a stimulant and cause excessive growth or metaplasia, leading to the development of a second tumor [23]. The slow growth of benign early tumors is generally considered to be a stimulus that leads to the production of microenvironmental factors, which may induce the growth of other *de novo* tumors [24-26]. Third, development of the mixed tumor may be associated with the independent development of two elements or bidirectional differentiation from common progenitor cells [27, 28]. Fourth, in rare situations, commonly occurring tumors may accidentally present at the same site [24-26]. Despite some evidence for the aforementioned theories, plausible underlying mechanisms

have not been conclusively proven. Further research and additional case studies are needed to develop a comprehensive understanding of the mechanisms that drive the simultaneous occurrence of multiple intracranial neoplasms.

In our patient, both CPP and MG were present in the right ventricle and could be removed using one surgical approach. A small remnant of the CPP remained attached to the pia of the medulla due to adherence. This tumor remnant was observed at the 3-month follow-up MRI scan. Equivocal growth was seen in serial imaging, following which gamma knife radiosurgery was successfully applied.

In conclusion, this is the first case report of coincident CPP and MG at the same locus of the lateral ventricle. This rare presentation of intraventricular tumors suggests that an admixture of tumors with papillary and lobular growth patterns in a single intraventricular mass may be due to the coincidental development of a CPP and MG at the same site.

Acknowledgements

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF), which is funded by the Ministry of Science, ICT, and Future Planning (NRF-2020R1G1A10036-92).

Informed consent was obtained from the patient described in this study.

Disclosure of conflict of interest

None.

Address correspondence to: Kiyong Na, Department of Pathology, Kyung Hee University Hospital, Kyung Hee University College of Medicine, 26 Kyungheedaero Dongdaemun-gu, Seoul 02447, Republic

Choroid plexus papilloma and meningioma

of Korea. Tel: 82-2-958-8740; Fax: 82-2-958-8744;
E-mail: rari papa@gmail.com

References

- [1] Louis DN, Perry A, Reifenberger G, von Deimling A, Figarella-Branger D, Cavenee WK, Ohgaki H, Wiestler OD, Kleihues P and Ellison DW. The 2016 world health organization classification of tumors of the central nervous system: a summary. *Acta Neuropathologica* 2016; 131: 803-20.
- [2] Koeller KK and Sandberg GD. From the archives of the AFIP. Cerebral intraventricular neoplasms: radiologic-pathologic correlation. *Radiographics* 2002; 22: 1473-505.
- [3] de Castro FD, Reis F and Guerra JG. Intraventricular mass lesions at magnetic resonance imaging: iconographic essay - part 1. *Radiologia Brasileira* 2014; 47: 176-81.
- [4] Buerki RA, Horbinski CM, Kruser T, Horowitz PM, James CD and Lukas RV. An overview of meningiomas. *Future Oncol* 2018; 14: 2161-77.
- [5] Marosi C, Hassler M, Roessler K, Reni M, Sant M, Mazza E and Vecht C. Meningioma. *Crit Rev Oncol Hematol* 2008; 67: 153-71.
- [6] Sethi D, Arora R, Garg K and Tanwar P. Choroid plexus papilloma. *Asian J Neurosurg* 2017; 12: 139-41.
- [7] Muñoz Montoya JE, Maldonado Moran MA, Santamaria Rodriguez P, Toro Lopez S, Perez Cataño CS and Luque Suarez JC. Choroid plexus papilloma of the fourth ventricle: a pediatric patient. *Asian J Neurosurg* 2019; 14: 585-8.
- [8] Kimura M, Takayasu M, Suzuki Y, Negoro M, Nagasaka T, Nakashima N and Sugita K. Primary choroid plexus papilloma located in the suprasellar region: case report. *Neurosurgery* 1992; 31: 563-6.
- [9] Lee Y, Kim SI, Kim SK, Kim IO and Park SH. A mixed choroid plexus papilloma and ependymoma. *Brain Tumor Pathol* 2016; 33: 147-50.
- [10] McIver JI, Link MJ, Giannini C, Cohen-Gadol AA and Driscoll C. Choroid plexus papilloma and meningioma: coincidental posterior fossa tumors: case report and review of the literature. *Surg Neurol* 2003; 60: 360-5.
- [11] Khayal HB, Abograra A and Lashhab M. Fourth ventricle papilloma and cranio-cervical junction meningioma: coincidental tumors: a case report and review of the literature. *Curr Opin Neurol Sci* 2018; 2: 463-7.
- [12] Lee EJ, Chang CH, Wang LC, Hung YC and Chen HH. Two primary brain tumors, meningioma and glioblastoma multiforme, in opposite hemispheres of the same patient. *J Clin Neurosci* 2002; 9: 589-91.
- [13] Maiuri F, Cappabianca P, Iaconetta G, Esposito F and Messina A. Simultaneous presentation of meningiomas with other intracranial tumours. *Br J Neurosurg* 2005; 19: 368-75.
- [14] Zhang Z, Yang Y, Zhang K, Zhuang J, Shao F, Liu H, Xing Y and Xu S. Collision tumor of glioblastoma and meningioma: case report and literature review. *World Neurosurg* 2018; 117: 137-41.
- [15] Gordon AS, Fallon KE and Riley KO. Meningioma interdigitated with primary central nervous system B-cell lymphoma: a case report and literature review. *Surg Neurol Int* 2011; 2: 181.
- [16] Honegger J, Buchfelder M, Schrell U, Adams EF and Fahlbusch R. The coexistence of pituitary adenomas and meningiomas: three case reports and a review of the literature. *Br J Neurosurg* 1989; 3: 59-69.
- [17] Suzuki K, Momota H, Tonooka A, Noguchi H, Yamamoto K, Wanibuchi M, Minamida Y, Hasegawa T and Houkin K. Glioblastoma simultaneously present with adjacent meningioma: case report and review of the literature. *J Neurooncol* 2010; 99: 147-53.
- [18] Slowik F and Jellinger K. Association of primary cerebral lymphoma with meningioma: report of two cases. *Clin Neuropathol* 1990; 9: 69-73.
- [19] Wang DJ, Zheng MZ, Gong Y, Xie Q, Wang Y, Cheng HX, Mao Y, Zhong P, Che XM, Jiang CC, Huang FP, Zheng K, Li SQ, Gu YX, Bao WM, Yang BJ, Wu JS, Xie LQ, Tang HL, Zhu HD, Chen XC and Zhou LF. Papillary meningioma: clinical and histopathological observations. *Int J Clin Exp Pathol* 2013; 6: 878-88.
- [20] Dohrmann GJ and Collias JC. Choroid plexus carcinoma. *J Neurosurg* 1975; 43: 225-32.
- [21] Karami KJ, Poulik J, Rabah R, Krass J and Sood S. Simultaneous choroid plexus carcinoma and pilocytic astrocytoma in a pediatric patient. *J Neurosurg Pediatr* 2010; 5: 104-12.
- [22] Iyer V, Sanghvi D, Shenoy A and Goel A. Three distinct co-existent primary brain tumors in a patient. *J Can Res Ther* 2009; 5: 293-6.
- [23] Spallone A, Santoro A, Palatinsky E and Giunta F. Intracranial meningiomas associated with glial tumours: a review based on 54 selected literature cases from the literature and 3 additional personal cases. *Acta Neurochir (Wien)* 1991; 110: 133-9.
- [24] Hokari M, Hida K, Ishii N, Seki T, Iwasaki Y and Nakamura N. Associated meningioma and neurofibroma at the same cervical level without clinical signs of neurofibromatosis: case report. *No Shinkei Geka* 2002; 30: 953-7.
- [25] Nakamizo A, Suzuki SO, Shimogawa T, Amano T, Mizoguchi M, Yoshimoto K and Sasaki T. Concurrent spinal nerve root schwannoma and meningioma mimicking single-component schwannoma. *Neuropathology* 2012; 32: 190-5.
- [26] Oichi T, Chikuda H, Morikawa T, Mori H, Kitamura D, Higuchi J, Taniguchi Y, Matsubayashi

Choroid plexus papilloma and meningioma

- Y, Oshima Y and Tanaka S. Concurrent spinal schwannoma and meningioma mimicking a single cervical dumbbell-shaped tumor: case report. *J Neurosurg Spine* 2015; 23: 784-7.
- [27] Tunthanathip T, Kanjanapradit K, Ratanalert S, Phuenpathom N, Oearsakul T and Kaewborisutsakul A. Multiple, primary brain tumors with diverse origins and different localizations: case series and review of the literature. *J Neurosci Rural Pract* 2018; 9: 593-607.
- [28] Deen HG and Laws ER. Multiple primary brain tumors of different cell types. *Neurosurgery* 1981; 8: 20-5.