

Analysis

Exemplary medical care or Trojan horse?

An analysis of the 'lifestyle medicine' movement

BACKGROUND

The rise of non-communicable diseases, many of which share common risk factors of smoking, alcohol, poor diet, and physical inactivity, has resulted in calls to develop and expand lifestyle medicine, giving 'hope to those suffering from chronic illness' (<https://bslm.org.uk/>). It has been argued that lifestyle medicine should be recognised as a new medical specialty,¹ with primary care leading.

There are numerous drivers for lifestyle medicine (Box 1). Our analysis does not aim to argue against the importance of these drivers as many of them are well informed. Instead we seek to balance existing discussions with aspects that, in our opinion, have been less well considered. With this in mind, we focus on the unintended consequences of uncritical endorsement and application of lifestyle medicine including the infiltration of pseudoscience, profiteering, and the potential for widening health inequalities by a continued focus on the 'individual'. We stress the need for greater attention to public health and community-level interventions and a more critical approach to current practice.

WHAT IS 'LIFESTYLE MEDICINE'?

Medical practice guidelines often advise on 'lifestyle factors'. These are usually in the form of individual behaviours that impact on health, framed as modifiable, often related to smoking, alcohol intake, physical activity, diet, and, to a lesser degree, sleep quality, stress, and social factors. However, translation of guidelines into achievable real-world benefits outside clinical trials is challenging.

The British Society of Lifestyle Medicine (BSLM) describes lifestyle medicine as 'an established approach that focuses on improving the health and wellbeing of individuals and populations ... It requires an understanding and acknowledgement of the physical, emotional, environmental and social determinants of disease.' (<https://bslm.org.uk/>). Society membership is open to registered health professionals, who can take a diploma, and associate membership is available to others, such as reflexologists, homeopaths, herbalists, and naturopaths.

Some have called for greater inclusion of lifestyle medicine education in professional training, including medical curricula,² based on evidence that knowledge of

"A key issue with the growth in prominence of lifestyle medicine ... is its use as a 'Trojan horse' to carry in non- or poorly evidenced practices under the auspices of 'evidence based'."

lifestyle medicine interventions is lacking.^{3,4} In the US, board certification in lifestyle medicine involves the ability to issue 'lifestyle prescriptions' with the definition 'the systematic practice of assisting individuals and families to adopt and sustain behaviours that can improve health and quality of life'.⁵ Current UK training in lifestyle medicine aimed at GPs is offered by the BSLM, the British Association for Nutrition and Lifestyle Medicine (BANT), and Prescribing Lifestyle Medicine (run by Clinical Education). Full BANT membership to practising UK therapists is available only to registrants on the Complementary and Natural Healthcare Council, which includes colon hydrotherapists, naturopaths, and reflexologists.

TROJAN HORSE AND CONFLICTS OF INTEREST

A key issue with the growth in prominence of lifestyle medicine, as broadly defined, is its use as a 'Trojan horse' to carry in non- or poorly evidenced practices under the auspices of 'evidence based'. We have two main concerns here: first, the opportunity taken by some to link alternative medicine practices to lifestyle medicine, and,

second, the association of some 'lifestyle medicine practitioners' with commercial opportunities.

For example, it doesn't take much of a search online to find multiple self-described, medically qualified practitioners advertising lifestyle medicine in the private sector, whose clinics offer discredited immunoglobulin G (IgG) tests for food intolerance, herbal remedies, mistletoe injections, intravenous vitamin infusions, bioidentical HRT (advised against by the British Menopause Society), and private health screening not recommended by the UK National Screening Committee. Conflicts may include ownership or sales of supplement or vitamin companies, or subscriptions to newsletters, books, or online lifestyle coaching.

Others have written extensively about the 'Trojan horse' of 'integrative medicine' (also termed 'functional medicine'),⁶ which claims that, by *integrating* complementary and alternative medicine (CAM) with conventional medicine, patients receive a complete, 'holistic' perspective. Integrative medicine often appeals to nature ('natural treatments'), antiquity ('ancient wisdom'), authority ('renowned universities run

Box 1. Key drivers for lifestyle medicine

Prevention is better than cure.

The patient is an active partner, not a passive recipient.

Lifestyle medicine treats the root cause (behaviours) of chronic disease, which medicine often overlooks.

Lifestyle risk factors are the primary cause of non-communicable disease and must therefore be addressed.

It is better to treat with lifestyle changes than to use potentially unnecessary drugs with the risk of side effects.

Lifestyle medicine is cost-effective compared with conventional medicine.

Professional satisfaction is greater, especially compared with usual approaches to chronic disease management (for example, the 'tick box' approach of the Quality and Outcomes Framework in England).

Because of weakening of public health through defunding within local government, individual approaches are necessary.

Public Health England has been perceived as not being sufficiently independent of government or industry.

“‘Lifestyle medicine’ needs a clear consensus on what constitutes evidence-based practice, with organisational standards and leadership commitment to the removal of bad science, and of financial and ideological conflicts.”

courses’), and popularity (‘demand is high’). The broad umbrella of CAM means the inclusion of poorly evidenced interventions including supplements, acupuncture, homeopathy, reiki, and reflexology. It also includes largely unproven physiological tests (for example, thermography, bioenergetic health scans, IgG food panels) to identify ‘root causes’ of symptoms (framed as ‘systems biology’), each at a financial cost to the patient (<https://lifestyleprescriptions.org/>).

Under this ecosystem, uncritical adoption of lifestyle medicine may result in inclusion of non-evidence-based ‘integrative medicine’ practices, thus providing a Trojan horse within which pseudoscience

can flourish. Disclosure of conflicts is not enough: indeed, this may be seen as giving ‘moral licence’ where transparency is equated with trustworthiness, regardless of content.

Notwithstanding the wide scope for the meaning of the word ‘holistic’, and variation in individual practice, systematic review evidence indicates that GPs have a strong understanding of whole-person care, the therapeutic value of an enduring patient–GP relationship, and an attentive, supportive, and collaborative approach.⁷ However, healthcare systems may not support, or may even be hostile to, these values.⁸ It is understandable that systems offering more time with a continuous practitioner are

popular, but this may be offset by a variable offering of evidence-based practice.

‘Lifestyle medicine’ needs a clear consensus on what constitutes evidence-based practice, with organisational standards and leadership commitment to the removal of bad science, and of financial and ideological conflicts. Movement towards this is welcome.⁹

LIFESTYLE ‘CHOICES’ AND HEALTH INEQUALITIES

The concept of individual ‘choices’ and ‘changes’, and by proxy ‘control’, plays a central role in many working definitions of lifestyle medicine. We are concerned about the potential for widening health inequalities when conceptualised and delivered in this way.

Potentially modifiable unhealthy behaviours are not evenly distributed across populations. They often coexist within individuals and communities, alongside multiple health (multimorbidity) and social problems, and are more concentrated in areas of socioeconomic deprivation.¹⁰ The clustering of fast food, tobacco, and alcohol outlets in deprived areas highlights the influence of environmental context on health-related behaviours.¹¹ While the BSLM definition of lifestyle medicine acknowledges ‘*environmental and social determinants of disease*’, the growth in private services are unlikely to yield benefits for people at highest risk of premature mortality from non-communicable diseases. Those in greatest need of support with health behaviour change are least likely to receive it.

The BSLM also states that ‘*Lifestyle Medicine has a wider responsibility to recognise upstream determinants of disease and to promote population health, to protect ecological health and to reduce health inequity*’ (<https://bslm.org.uk/>). However, we have seen limited evidence of advocacy for action on ‘upstream determinants’ among lifestyle medicine proponents. Rather, the focus is, and has historically been, on downstream individual-based interventions. There is emerging evidence that such approaches increase health inequalities¹² and the continuation of this as the dominant approach in policy draws attention away from the need for wider environmental or structural (public health) interventions.¹³

Ultimately, the biggest drivers of health, and associated risk factors, are the social determinants of health (‘*the conditions in which we are born, grow, live, work and age*’),¹⁴ shaped by the distribution of

Box 2. Individual and population approaches to health risk factors

Intervention for	Individual-level interventions	Public health interventions	
		Community level	National policy level
More modifiable risk factors			
Physical inactivity	Advice and support, exercise prescription, gym or home exercise resources	Parks, cycling infrastructure, bike hire schemes	Active travel policies, working hours
Poor diet	Advice and support, referral to weight management services	Community gardens, cooking classes.	Sugar tax, pack sizes, food labelling, restrictions on promotions and marketing of unhealthy foods
Excess alcohol intake	Screening, individual advice and support	Local alcohol licensing, recovery communities	Minimum Unit Price legislation, restrictions on advertising
Smoking	Advice from health professional at routine appointment	Open access to pharmacy quit services	Mass advertising for telephone advice, bans on smoking in public places
Less modifiable risk factors			
Poverty	Community links worker, financial inclusion advice	Anti-poverty community groups, food pantries, credit unions.	Active labour market policies, welfare system reform to improve ease of applications, minimum income for healthy living
Environmental	Access to private outdoor spaces, referral to social prescribing schemes	Green spaces, parks	Regulation to limit air pollution

“Those in greatest need of support with health behaviour change are least likely to receive it.”

money, power, and resources at global, national, and local levels. Efforts to change health-related behaviours among the most deprived members of society are unlikely to succeed unless they are supported by measures designed to improve the material circumstances and the drivers of those behaviours within communities.

INTEGRATING PERSONAL HEALTH AND PUBLIC HEALTH

We suggest that individual-level interventions are most likely to succeed when integrated with public health interventions, which focus on populations, either within communities or with higher-level (regional/national) policies (Box 2). Public health aims to subvert the ‘healthy attender paradox’ and avoid the need for healthcare professional guidance or action.

Taking alcohol as an example, individual interventions may have a role, but only population interventions can reach everyone. A 2018 Cochrane review found moderate evidence that a brief alcohol intervention could reduce alcohol consumption by around a pint of beer or a third of a bottle of wine per week, albeit with little impact on binges per week or alcohol-free days.¹⁵ The introduction of minimum unit pricing on alcohol in Scotland seems to have had a greater effect on the highest alcohol consumers.¹⁶ Similarly, brief advice to stop smoking can increase quit rates by 1–3%,¹⁷ but reduced cigarette smoking across the population, and decreased youth take-up, is most likely due to legal interventions to limit smoking and advertising.¹⁸

It is critical to know whether success in trials can be replicated in real-world practice. For example, a popular claim is that ‘brief physical activity advice interventions’ in primary care have a number needed to treat of 12 for increasing self-reported physical activity.¹⁹ This has led efforts to research and increase professional knowledge of physical activity guidelines. However, ‘brief advice’ in trials consisted of frequent face-to-face and telephone call support delivered by professionals from different disciplines, subsidised gym membership, postal support, and personalised reports.²⁰ A core problem with such trials is the dilution of any effect on patients who elected not to take part in the trial at the start. Weight

loss interventions described as successful have uptake in less than a third of at-risk patients and require systematic support.²¹ Even evidence-based interventions may not be effective in the real-world, higher-risk community.

REFERENCES

1. Sayburn A. Lifestyle medicine: a new medical specialty? *BMJ* 2018; **363**: k4442. <https://www.bmj.com/content/363/bmj.k4442> [accessed 22 Mar 2021].
2. Meldrum H, Katz D, Egger G. Lifestyle medicine in Expert Review of Cardiovascular Therapy. *Expert Rev Cardiovasc Ther* 2016; **14**(3): 263.
3. Crowley J, Ball L, Hiddink GJ. Nutrition in medical education: a systematic review. *Lancet Planet Health* 2019; **3**(9): e379–e389. DOI: 10.1016/s2542-5196(19)30171-8.
4. Chatterjee R, Chapman T, Brannan MGT, Varney J. GPs’ knowledge, use, and confidence in national physical activity and health guidelines and tools: a questionnaire-based survey of general practice in England. *Br J Gen Pract* 2017; DOI: 10.3399/bjgp17X692513.
5. Clarke CA, Hauser ME. Lifestyle medicine: a primary care perspective. *J Grad Med Educ* 2016; **8**(5): 665–667. DOI: 10.4300/JGME-D-15-00804.1.
6. Gorski D. Functional medicine: the ultimate misnomer in the world of integrative medicine. *Science-Based Medicine* 2020; **11 Apr**: <https://sciencebasedmedicine.org/functional-medicine-the-ultimate-misnomer-in-the-world-of-integrative-medicine/> [accessed 22 Mar 2021].
7. Thomas H, Mitchell G, Rich J, Best M. Definition of whole person care in general practice in the English language literature: a systematic review. *BMJ Open* 2018; **8**(12): e023758. DOI: 10.1136/bmjopen-2018-023758.
8. Hasegawa H, Reilly D, Mercer SW, Bikker AP. Holism in primary care: the views of Scotland’s general practitioners. *Primary Health Care Research & Development* 2005; **6**(4): 320–328. DOI: 10.1191/1463423605pc2480a
9. Lianov L, Johnson M. Physician competencies for prescribing lifestyle medicine. *JAMA* 2010; **304**(2): 202–203. DOI: 10.1001/jama.2010.903.
10. Meader N, King K, Moe-Byrne T, et al. A systematic review on the clustering and co-occurrence of multiple risk behaviours. *BMC Public Health* 2016; **16**: 657. DOI: 10.1186/s12889-016-3373-6.
11. Macdonald L, Olsen JR, Shortt NK, Ellaway A. Do ‘environmental bads’ such as alcohol, fast food, tobacco, and gambling outlets cluster and co-locate in more deprived areas in Glasgow City, Scotland? *Health Place* 2018; **51**: 224–231
12. McGill R, Anwar E, Orton L, et al. Are interventions to promote healthy eating equally effective for all? Systematic review of socioeconomic inequalities in impact. *BMC Public Health* 2015; **15**: 457. <https://doi.org/10.1186/s12889-015-1781-7> [accessed 22 Mar 2021].
13. Kriznik NM, Kinmonth A-L, Ling T, Kelly MP. Moving beyond individual choice in policies to reduce health inequalities: the integration of dynamic with individual explanations. *J Public Health (Oxf)* 2018; **40**(4): 764–775. DOI: 10.1093/pubmed/tdy045.
14. World Health Organization. Social determinants of health. 2020. <https://www.who.int/teams/social-determinants-of-health> [accessed 22 Mar 2021].
15. Kaner EF, Beyer FR, Muirhead C, et al. Effectiveness of brief alcohol interventions in primary care populations. *Cochrane Database Syst Rev* 2018; **2**(2): CD004148. DOI: 10.1002/14651858.cd004148.pub4.
16. O’Donnell A, Anderson P, Jané-Llopis E, et al. Immediate impact of minimum unit pricing on alcohol purchases in Scotland: controlled interrupted time series analysis for 2015–18. *BMJ* 2019; **366**: i5274. DOI: 10.1136/bmj.i5274.
17. Stead LF, Buitrago D, Preciado N, et al. Physician advice for smoking cessation. *Cochrane Database Syst Rev* 2013; **5**(5): CD000165. DOI: 10.1002/14651858.cd000165.pub4.
18. Frazer K, Callinan JE, McHugh J, et al. Legislative smoking bans for reducing harms from secondhand smoke exposure, smoking prevalence and tobacco consumption. *Cochrane Database Syst Rev* 2016; **2**(2): CD005992. DOI: 10.1002/14651858.cd005992.pub3.
19. Brooks J, Ahmad I, Easton G. Promoting physical activity: the general practice agenda. *Br J Gen Pract* 2016; DOI: 10.3399/bjgp16x686689.
20. Orrow G, Kinmonth A-L, Sanderson S, Sutton S. Effectiveness of physical activity promotion based in primary care: systematic review and meta-analysis of randomised controlled trials. *BMJ* 2012; **344**: e1389. DOI: 10.1136/bmj.e1389.
21. Aveyard P, Lewis A, Tearne S, et al. Screening and brief intervention for obesity in primary care: a parallel, two-arm, randomised trial. *Lancet* 2016; **388**(10059): 2492–2500. DOI: 10.1016/s0140-6736(16)31893-1.

CONCLUSION

Optimally, ‘lifestyle medicine’ is concerned with preventive or therapeutic targeting of potentially modifiable risk factors using evidence-based strategies to support favourable change in health behaviours. Yet usual, good medical practice should encompass all of this, embedded within considerate relationships between patients and professionals. The affiliation between lifestyle medicine and non-evidence-based, fringe, and alternative tests, diagnoses, and interventions in many areas risks

“Effective care requires not simply calls to education, but resources where they are needed most, assessment of opportunity cost, and critical evaluation of interventions.”

ADDRESS FOR CORRESPONDENCE

David Nunan

Nuffield Department of Primary Care Health Sciences, Radcliffe Primary Care Building, Radcliffe Observatory Quarter, University of Oxford, Woodstock Road, Oxford OX2 6GG, UK.

Email: david.nunan@phc.ox.ac.uk

disrepute, conflict, and confusion for patients. We recommend that evidence-based interventions should be part of routine training for clinical staff, with limitations explained, and critical thinking and reflection employed.

Effective care requires not simply calls to education, but resources where they are needed most, assessment of opportunity cost, and critical evaluation of interventions. Lifestyle medicine's continued emphasis on the individual as the change agent most likely results in the people at lowest risk having the greatest amount of intervention, while people carrying the greatest risk are not receiving the support they need. Understanding the environmental drivers of unhealthy behaviours requires primary care practitioners to work more closely with public health colleagues to develop local community approaches, particularly in disadvantaged areas.

We support public health colleagues working with the government to research and implement evidence-based population

interventions targeting the drivers of potentially modifiable risks, as these have the highest chance of benefiting populations already at greatest risk of premature mortality and morbidity.

David Nunan,

Senior Research Fellow, Nuffield Department of Primary Care Health Sciences, University of Oxford, Oxford.

David N Blane,

Clinical Research Fellow in General Practice and Primary Care, Institute of Health and Wellbeing, University of Glasgow, Glasgow.

Margaret McCartney,

Honorary Senior Lecturer, CSO Fellow, University of St Andrews, St Andrews, Fife, Scotland.

Provenance

Freely submitted; externally peer reviewed.

Competing interests

David Nunan has received expenses and fees

for his media work and teaching workshops on evidence-based medicine for the Centre for Evidence-Based Medicine (CEBM). He has held grant funding from the National Institute for Health Research (NIHR), the NIHR School for Primary Care Research, and the Royal College of General Practitioners (RCGP). David Blane has received research funding from the Chief Scientist Office (CSO) Scotland and the RCGP. Margaret McCartney is a CSO career fellow, Hon. Fellow CEBM Oxford, and nationally elected RCGP council member and trustee. She has donated money to Keep Our NHS Public and has campaigned for mandatory disclosures of interest. She receives royalties for books and payments for journalism and broadcasting, and occasionally for lectures to health professionals.

The authors alone are responsible for the views expressed in this article, which do not necessarily represent the views, decisions, or policies of the institutions with which the authors are affiliated.

Acknowledgements

We thank Jeffrey Aronson for his comments on early drafts of this article.

DOI: <https://doi.org/10.3399/bjgp21X715721>