




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Letter to the Editors-in-Chief

Therapeutic anticoagulation in COVID-19 patients



ARTICLE INFO

Keywords

COVID-19

Anticoagulant treatment

Coagulopathy

Our pilot investigation [1] was performed in an early phase of this pandemic when the first SARS-CoV-2 infected patients emerged in our region.

We thank your interest in our research, and we clarify some questions raised. First, about the prophylactic anticoagulation, our institutional protocol indicated these two treatment regimens (enoxaparin or unfractionated heparin); thus, enoxaparin was not used for all patients. There were no pre-defined criteria to choose between the two regimens, and we respected the doctors' option.

Second, all the patients were evaluated for enrollment in the emergency department immediately after admission. Most of the patients were randomized in the first 24 h after admission, and they received the first dose anticoagulant during this period.

Third, the control and intervention group had similar baseline characteristics. The tidal volume (TV) per Kilograms of predicted body weight and the positive end-expiratory pressure (PEEP) were not statistically different between these two groups. They exhibited the same average value for the TV per Kg of predicted body weight and only 1 cm of water of difference in the PEEP relation, which is not clinically significant to distinguish different severities between them. The number of patients submitted to prone positioning was similar between the groups (80% vs. 70%; $p = 0.606$); however, the prophylactic group required a higher number of prone positioning sessions because they did not improve the gas exchange over time.

We want to highlight that early endotracheal intubation and mechanical ventilation were preconized in the initial phase of this pandemic. The experience acquired during the pandemic course has shown that several patients could be maintained in the high-flow nasal cannula or non-invasive ventilation. Therefore, some patients included in our investigation would currently not meet the criteria for mechanical ventilation.

Fourth, we excluded seven patients before randomization, and the two patients with circulatory shock who required high doses of vasopressors were excluded before the enrollment. Despite this, four patients in the prophylactic group and two patients in the therapeutic group developed late severe circulatory shock after randomization; however, they were not excluded from this intention-to-treat analysis.

Fifth, the bleeding was evaluated through the Thrombolysis in Myocardial Infarction (TIMI) bleeding criteria [2]. The bleeding

requiring medical attention was defined as any sign of hemorrhage necessitating intervention, prolonged hospitalization, or need for prompt evaluation and did not meet major or minor bleeding criteria. A high proportion of patients presented an isolated drop in hemoglobin levels, without any clinically evident sign of hemorrhage; consequently, these events were not classified as minor or major bleeding. There is significant heterogeneity among different bleeding definitions. For example, the International Society of Thrombosis and Haemostasis (ISTH) considers an isolated fall in hemoglobin level of 2 g/dl as major bleeding [3]. If we had considered this last definition, we would have had two major bleeding in the prophylactic group and four major bleeding in the therapeutic group. Two patients required blood transfusion in the therapeutic anticoagulation group and one patient in the prophylactic anticoagulation group. We think that the most important thing is that the occurrence of symptomatic bleeding in a critical organ is very low in COVID-19 patients.

Lastly, everyone is anxiously waiting for publication of the results of these clinical trials about therapeutic and intermediate dose of anticoagulation in COVID-19 patients. The interim analysis of the three multicenter (ATTACC, REMAP-CAP, ACTIV-4a) randomized, open-label trials showed that therapeutic dose anticoagulation decreased the need for organ support and mortality in moderately ill COVID-19 patients, on the other hand, therapeutic-dose anticoagulation did not decrease organ support and mortality in critically ill COVID-19 patients. [4]

Despite the uncertainties, we cannot forget that all severely ill patients were previous moderately ill; it is only a time question. We think an intermediate dose of anticoagulation is insufficient to mitigate this hypercoagulable state, as demonstrated in the INSPIRATION clinical trial. [5] Furthermore, the antithrombotic intervention could be too late in the severe COVID-19 patients admitted to the intensive care unit to improve their clinical condition. From our point of view, early hospitalization of COVID-19 patients with dyspnea and mild reduction in the arterial oxygen saturation (moderately ill patients) with prescription of anti-inflammatory drugs (corticosteroids, tocilizumab, etc.) [6,7], anti-viral drugs, and therapeutic anticoagulation in selected patients with low risk of bleeding are the available strategies to prevent the worsening of this disease.

<https://doi.org/10.1016/j.thromres.2021.04.027>

Received 3 March 2021; Received in revised form 22 April 2021; Accepted 26 April 2021

Available online 3 May 2021

0049-3848/© 2021 Elsevier Ltd. All rights reserved.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] A.C.B. Lemos, D.A. do Espírito Santo, M.C. Salvetti, R.N. Gilio, L.B. Agra, A. Pazin-Filho, C.H. Miranda, Therapeutic versus prophylactic anticoagulation for severe COVID-19: a randomized phase II clinical trial (HESACOVID), *Thromb. Res.* 196 (2020) 359–366.
- [2] R. Mehran, S.V. Rao, D.L. Bhatt, C.M. Gibson, A. Caixeta, J. Eikelboom, S. Kaul, S. D. Wiviott, V. Menon, E. Nikolsky, V. Serebruany, M. Valgimigli, P. Vranckx, D. Taggart, J.F. Sabik, D.E. Cutlip, M.W. Krucoff, E.M. Ohman, P.G. Steg, H. White, Standardized bleeding definitions for cardiovascular clinical trials: a consensus report from the Bleeding Academic Research Consortium, *Circulation* 123 (23) (2011) 2736–2747.
- [3] S. Schulman, C. Kearon, S. Subcommittee on Control of Anticoagulation of the T. Standardization Committee of the International Society on, Haemostasis, Definition of major bleeding in clinical investigations of antihemostatic medicinal products in non-surgical patients, *J Thromb Haemost* 3(4) (2005) 692–4.
- [4] Full-dose blood thinners decreased need for life support and improved outcomes in hospitalized COVID-19 patients, National Institutes of Health (NIH), 2021. <https://www.nih.gov/news-events/new-releases/full-dose-blood-thinners-decrease-need-life-support-improved-outcome-hospitalized-covid-19-patients>.
- [5] I. Investigators, P. Sadeghipour, A.H. Talasaz, F. Rashidi, B. Sharif-Kashani, M.T. Beigmohammadi, M. Farrokhpour, S.H. Sezavar, P. Payandemehr, A. Dabbagh, K.G. Moghadam, S. Jamalkhani, H. Khalili, M. Yadollahzadeh, T. Riahi, P. Rezaeifar, O. Tahamtan, S. Matin, A. Abedini, S. Lookzadeh, H. Rahmani, E. Zoghi, K. Mohammadi, P. Sadeghipour, H. Abri, S. Tabrizi, S.M. Mousavian, S. Shahmirzaei, H. Bakhshandeh, A. Amin, F. Rafiee, E. Baghizadeh, B. Mohebbi, S.E. Parhizgar, R. Aliannejad, V. Eslami, A. Kashefzadeh, H. Kakavand, S.H. Hosseini, S. Shafaghi, S.F. Ghazi, A. Najafi, D. Jimenez, A. Gupta, M.V. Madhavan, S.S. Sethi, S.A. Parikh, M. Monreal, N. Hadavand, A. Hajighasemi, M. Maleki, S. Sadeghian, G. Piazza, A.J. Kirtane, B.W. Van Tassel, P.P. Dobesh, G.W. Stone, G.Y.H. Lip, H.M. Krumholz, S.Z. Goldhaber, B. Bikdeli, Effect of intermediate-dose vs standard-dose prophylactic anticoagulation on thrombotic events, extracorporeal membrane oxygenation treatment, or mortality among patients with COVID-19 admitted to the intensive care unit: the INSPIRATION Randomized clinical trial, *JAMA* (2021).
- [6] R.C. Group, P. Horby, W.S. Lim, J.R. Emberson, M. Mafham, J.L. Bell, L. Linsell, N. Staplin, C. Brightling, A. Ustianowski, E. Elmahi, B. Prudon, C. Green, T. Felton, D. Chadwick, K. Rege, C. Fegan, L.C. Chappell, S.N. Faust, T. Jaki, K. Jeffery, A. Montgomery, K. Rowan, E. Juszczak, J.K. Baillie, R. Haynes, M.J. Landray, Dexamethasone in hospitalized patients with Covid-19, *N. Engl. J. Med.* 384 (8) (2021) 693–704.
- [7] H.J. Salama C, Yau L, Reiss WG, Kramer B, Neidhart JD, Criner GJ, Kaplan-Lewis E, Baden R, Pandit L, Cameron ML, Garcia-Diaz J, Chávez V, Mekebeb-Reuter M, Lima de Menezes F, Shah R, González-Lara MF, Assman B, Freedman J, Mohan SV, Tocilizumab in patients hospitalized with Covid-19 pneumonia, *N. Engl. J. Med.* 384 (2021) 20–30.

Anna Cristina Bertoldi Lemos, Douglas Alexandre do Espírito Santo,
 Carlos Henrique Miranda*
 Division of Emergency Medicine, Department of Internal Medicine, Ribeirão
 Preto School of Medicine, São Paulo University, Rua Bernardino de Campos,
 1000 Ribeirão Preto, São Paulo, Brazil

* Corresponding author.

E-mail address: chmiranda@fmrp.usp.br (C.H. Miranda).