

## Research Article

# Effects of Berberine on Circular RNA Expression Profiles in Human Gastric Cancer Cells

Meng Wang <sup>1,2</sup>, Letao Sun,<sup>3</sup> Li Wang,<sup>4</sup> and Yongning Sun <sup>1,4</sup>

<sup>1</sup>Shanghai Jiao Tong University Affiliated Sixth People's Hospital, Shanghai 200233, China

<sup>2</sup>Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China

<sup>3</sup>Gordon F. Derner School of Psychology, Adelphi University, New York 11530, USA

<sup>4</sup>Shanghai Municipal Hospital of Traditional Chinese Medicine, Shanghai University of Traditional Chinese Medicine, Shanghai 200071, China

Correspondence should be addressed to Yongning Sun; ynsun2002@126.com

Received 2 January 2021; Revised 23 March 2021; Accepted 29 March 2021; Published 4 May 2021

Academic Editor: Jianping Chen

Copyright © 2021 Meng Wang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Background.** Berberine has been demonstrated to have anticancer effects against gastric cancer (GC), but the mechanism of these actions is unclear. **Objectives.** To explore the impact of berberine on circular RNA (circRNA) expression profiles in GC and investigate the potential molecular mechanisms associated with circRNAs in GC. **Methods.** AGS and HGC27 GC cells were treated with various concentrations of berberine. Cell viability was measured using a Cell Counting Kit-8 assay. Cell proliferation was measured using a cell colony formation assay. Cell apoptosis was measured using flow cytometry. The mitochondrial membrane potential ( $\Delta\psi_m$ ) was determined using a JC-1 probe. RNA-seq was performed to identify circRNA expression profiles in AGS cells after berberine treatment. Selected differentially expressed (DE) circRNAs were verified using RT-qPCR. Bioinformatics analysis was performed to predict target miRNAs and mRNAs and construct a circRNA-miRNA-mRNA network. Pathway and process enrichment analyses were performed to explore the potential biological roles of DE circRNAs. **Results.** Berberine decreased GC cell viability, cell proliferation, and  $\Delta\psi_m$  and induced cell apoptosis. Thirty-one DE circRNAs were identified in the berberine-treated group compared to the control group, among which circRNA2499, hsa\_circ\_0003423, and hsa\_circ\_0006702 were validated using RT-qPCR. Enrichment analyses, based on the host genes of these 31 DE circRNAs and putative target mRNAs in the circRNA-miRNA-mRNA network of the validated circRNAs, indicated that berberine exerts anti-GC effects in multiple pathways including the Notch, MAPK, and NF- $\kappa$ B signaling pathways via specific circRNAs. **Conclusion.** This study elucidated the expression profile of circRNAs in human GC cells after berberine treatment. Our results demonstrate that berberine has the potential to influence cancer-related pathways by regulating circRNA expression and their corresponding target genes in GC cells.

## 1. Introduction

Despite advancements in early diagnosis and therapeutics, such as surgery and chemoradiotherapy, the prognosis of gastric cancer (GC) remains relatively poor [1, 2]. Multiple factors including genetics and epigenetics are involved in the development of GC [3, 4]. Circular RNAs (circRNAs) are noncoding RNAs with a covalently closed continuous loop [5]. circRNAs can regulate gene expression by acting as nuclear transcriptional regulators, miRNA sponges, and RNA-binding protein sponges [6]. Many researchers have highlighted the important functions of circRNAs in the

development of cancer, including hepatocellular carcinoma [7], GC [8], and colon cancer [9].

Natural products represent an important source for discovering anticancer agents. Several plant-derived agents have been successfully used in cancer treatment, such as vinca alkaloid, etoposide, and paclitaxel, and some others are currently under investigation [10, 11]. Some studies reported that natural products such as nitidine chloride [12] and quercetin [13] exert anticancer effects or improve the prognosis of patients by influencing the expression of circRNAs. Several studies have demonstrated the important anticancer roles of berberine against malignant tumors,

including GC [14]. However, the mechanisms of berberine against GC via circRNAs remain unclear.

In this study, RNA-seq analysis was performed to identify and analyze changes in circRNAs in GC cells in response to berberine. These results improve the understanding of the circRNAs targeted by berberine, which may be useful in developing treatments for GC.

## 2. Materials and Methods

**2.1. Cell and Drug Preparation.** Human AGS and HGC27 GC cells were obtained from the Cell Bank of the Chinese Academy of Sciences (Shanghai, China) and maintained in F-12K (Gibco, Thermo Fisher Scientific; Waltham, USA) and RPMI 1640 (Gibco) media, respectively. Both media were supplemented with 10% fetal bovine serum (Gibco). The cells were maintained in a humidified atmosphere of 5% CO<sub>2</sub> at 37°C. Berberine (MedChemExpress, Shanghai, China) was dissolved in dimethyl sulfoxide (Solarbio, Beijing, China) and diluted to the working concentration with culture media.

**2.2. Cell Viability Assay.** AGS and HGC27 cells were seeded into 96-well plates and incubated with berberine (0, 20, 50, and 80 μM) for 24, 48, and 72 h. After incubation, cell viability was examined using a Cell Counting Kit-8 (CCK-8) kit (Dojindo, Kumamoto, Japan) by measuring the optical density values.

**2.3. Cell Colony Formation Assay.** AGS and HGC27 cells were seeded into 6-well plates and incubated with berberine (0 and 50 μM). After 7 days in culture, cells were stained with crystal violet (Beyotime, Shanghai, China) and photographed with a digital camera. The colony formation rate was calculated, which can account for cell population dependence and proliferation ability.

**2.4. Flow Cytometry Evaluation of Cell Apoptosis.** AGS and HGC27 cells treated with berberine were harvested and cell apoptosis induced due to berberine was identified using the annexin V-FITC/PI (Beyotime, Shanghai, China) staining according to the instructions. The percentage of apoptotic cells was assessed using a FACSCalibur flow cytometer (BD Biosciences, San Jose, CA, USA).

**2.5. Measurement of Mitochondrial Membrane Potential.** AGS cells were cultured and treated with different concentrations of berberine in 24-well plates. Thereafter, the mitochondrial membrane potential ( $\Delta\psi_m$ ) was determined using a JC-1 probe kit (MedChemExpress) according to the instructions. Red and green fluorescence in the cells was examined using a fluorescence microscope (Leica, Wetzlar, Germany). The ratio of red to green fluorescence intensity was calculated using ImageJ software (National Institutes of Health, Bethesda, MD, USA).

**2.6. RNA Library Construction and circRNA Sequencing.** Three random samples from each group were subjected to RNA-seq. Total RNA was isolated using TRIzol reagent (Invitrogen, Carlsbad, CA, USA) according to the instructions. RNA concentration and purity were quantified with a NanoDrop ND-1000 (Thermo Fisher Scientific). RNA integrity was assessed using an Agilent 2100 (Agilent Technologies, Santa Clara, CA, USA). Ribosomal RNA was depleted from total RNA according to the Ribo-Zero rRNA Removal Kit instructions (Illumina, San Diego, CA, USA). Preparation of RNA libraries and sequencing were conducted by LC Bio Co., Ltd. (Hangzhou, China). Sequencing was performed on a NovaSeq 6000 (Illumina) according to the instructions. Differentially expressed (DE) circRNAs were identified as those showing a  $|\log_2(\text{fold-change})| \geq 1$  and statistical significance ( $P$  value  $< 0.05$ ) according to R package edgeR [15].

**2.7. Pathway and Process Enrichment Analyses.** The effects of circRNAs on their host genes were predicted using GO (<http://www.geneontology.org>) and KEGG (<http://www.kegg.jp>) enrichment analyses to explore meaningful gene annotations. The  $P$  values represent enrichment scores. GO analysis was performed to evaluate biological processes, cellular components, and molecular functions.

**2.8. RT-qPCR Validation.** Total RNA was isolated from the cell lines using TRIzol reagent (Invitrogen) and then reverse-transcribed into cDNA using the PrimerScript RT reagent kit with gDNA Eraser (Takara Bio, Shiga, Japan). RT-qPCR was performed using SYBR Premix Ex Taq (Takara), and *GAPDH* was used as an internal control. The expression of circRNAs was defined based on the threshold cycle (Ct), and relative expression levels were calculated using the  $2^{-\Delta\Delta C_t}$  method. PCR amplification was performed as follows: 95°C for 30 s, 40 cycles of 95°C for 5 s, and 60°C for 34 s.

**2.9. Construction of circRNA-miRNA-mRNA Network.** Interactions of circRNA-miRNA and miRNA-mRNA, which were predicted using TargetScan (<http://www.targetscan.org/>) and miRanda (<http://www.miranda.org/>), were combined to construct a circRNA-miRNA-mRNA network. Cytoscape software v3.7.2 [16] was used to visualize the network. The Metascape database [17] provides reliable and productive bioinformatics analyses of gene or protein lists, which can help users to make better data-driven decisions. The predicted mRNAs were entered into the Metascape database for annotation, functional analysis, and MCODE algorithm [18] analysis.

**2.10. Statistical Analyses.** Experimental data are presented as the means  $\pm$  SEM. Student's two-tailed unpaired  $t$ -test was used to evaluate the differences between the two groups.  $P$  values  $< 0.05$  represent statistical significance. GraphPad Prism 8.0 (<http://www.graphpad.com>) was used for statistical analyses.

### 3. Results

**3.1. Berberine Decreased Cell Viability, Cell Proliferation, and  $\Delta\psi_m$  and Induced Cell Apoptosis.** The CCK-8 analysis showed that AGS and HGC27 cell viability decreased in berberine concentration- and treatment time-dependent manners (0–80  $\mu\text{M}$  and 0–72 h, respectively) (Figure 1(a)). After 7 days of treatment, 50  $\mu\text{M}$  berberine significantly decreased AGS and HGC27 cell colony formation rate (Figure 1(b)). Treatment with different concentrations of berberine for 48 h resulted in an increased number of apoptotic AGS and HGC27 cells (Figure 1(c)), as well as a reduction in  $\Delta\psi_m$  in AGS cells (Figure 1(d)).

**3.2. Overview of circRNA Profiles.** After 48 h treatment, 50  $\mu\text{M}$  berberine induced changes in the circRNA expression profiles. A total of 12,256 circRNAs were detected using RNA-seq analysis between berberine-treated and control AGS cells. Among these, 5561 potential novel circRNAs have not been identified previously. Figure 2(a) represents the chromosome distribution of the identified circRNAs. No circRNAs were distributed in chromosome Y. The length distribution of all identified circRNAs is visualized in Figure 2(b). The lengths of the circRNAs ranged from <1000 nucleotides (nt) to >10,000 nt, with the majority having <1000 nt. The genomic origins of the identified circRNAs are presented in Figure 2(c). Based on their location in the genome, most candidate circRNAs' host genes were obviously derived from exonic regions, suggesting the importance of these circRNAs.

**3.3. DE circRNAs Affected by Berberine.** Figure 3(a) shows a clustered heatmap of the DE circRNAs in the berberine-treated group and control group determined using the described cutoff. The volcano plot shows significant DE circRNAs (Figure 3(b)). We identified 31 DE circRNAs, among which 19 were upregulated and 12 were downregulated in the berberine-treated group (Figures 3(b) and 3(c)).

**3.4. DE circRNA-Hosting Gene GO and KEGG Analyses.** To annotate the DE circRNAs, enrichment analysis was performed on their host genes. GO analysis (Figures 4(a) and 4(b)) revealed that the altered circRNAs were associated with terms such as negative regulation of DNA binding (GO: 0043392), NF- $\kappa\text{B}$  binding (GO:0051059), phospholipid binding (GO:0005543), regulation of interleukin-1 production (GO:0032732), and phosphorylase kinase activity (GO:0004689). KEGG analysis (Figure 4(c)) yielded enrichment in terms such as nucleotide excision repair (ko03420), Notch signaling pathway (ko04330), insulin signaling pathway (ko04910), and cell cycle (ko04110).

**3.5. Validation of DE circRNAs.** To verify the RNA-seq results, four DE circRNAs were manually selected at random for further validation by RT-qPCR in AGS and HGC27 cells. The primer sequences for the candidate circRNAs are shown

in Table S1. The expression levels of circRNA2499 (1.33-fold;  $P < 0.01$ ), hsa\_circ\_0003423 (1.76-fold;  $P < 0.001$ ), hsa\_circ\_0006702 (3.25-fold;  $P < 0.0001$ ), and hsa\_circ\_0070562 (1.76-fold;  $P < 0.01$ ) were significantly upregulated in the berberine-treated group of AGS cells. The expression levels of circRNA2499 (1.47-fold;  $P < 0.01$ ), hsa\_circ\_0003423 (1.31-fold;  $P < 0.01$ ), hsa\_circ\_0006702 (1.69-fold;  $P < 0.0001$ ), and hsa\_circ\_0070562 (1.09-fold;  $P < 0.05$ ) were also significantly upregulated in the berberine-treated group of HGC27 cells. The RT-qPCR results of circRNA2499, hsa\_circ\_0003423, and hsa\_circ\_0006702 in AGS and HGC27 cells agreed with those of RNA-seq (Figure 5).



**3.6. circRNA-miRNA-mRNA Network and Biological Function Prediction for Validated circRNAs.** To estimate the functions of circRNA2499, hsa\_circ\_0003423, and hsa\_circ\_0006702, the circRNAs were assumed to act as miRNA sponges and take part in a circRNA-miRNA-mRNA network. The top five miRNAs predicted to bind to the circRNAs as well as each miRNA's top ten target mRNAs are shown in Figures 6(a)–6(c). The target mRNAs represented by their transcript IDs in Figures 6(a)–6(c) are listed in Supplementary Table S2.

To further evaluate the circRNAs, enrichment analysis based on putative target mRNAs was conducted and the results were visualized using the Metascape database. Analysis of each circRNA (Figure 6(d) and Table S3) indicated that the target genes of circRNA2499 were associated with terms such as mitotic prometaphase (R-HSA-68877); target genes of hsa\_circ\_0003423 were associated with terms such as MAPK family signaling cascades (R-HSA-5683057) and response to endoplasmic reticulum stress (GO: 0034976); and target genes of hsa\_circ\_0006702 were associated with terms such as NIK/NF- $\kappa\text{B}$  signaling (GO: 0038061), hormone metabolic process (GO:0042445), and G2/M transition of the mitotic cell cycle (GO:0000086). In addition, the merged target genes of these three circRNAs (Figure 6(e) and Table S4) were associated with terms such as cell-substrate adhesion (GO:0031589), apoptosis (R-HSA-109581), and PTEN regulation (R-HSA-680707). MCODE module analysis using the Metascape database showed that the target mRNAs were mainly enriched in the biological process of cell cycle (Table 1).


### 4. Discussion

The use of berberine, a compound isolated from medicinal plants such as *Coptis chinensis*, has been reported for preventing and treating GC via targeting the AMPK/HNF4 $\alpha$ /WNT5A pathway [19], Akt/mTOR/p70S6/S6 pathway [20], interleukin-8 [21], and survivin and STAT3 [22]. Our study focused on the effects of berberine on circRNA expression profiles in human GC cells.

After observing berberine-induced changes in GC cell viability, cell colony formation rate, cell apoptosis, and  $\Delta\psi_m$ , the circRNA expression alterations in AGS cells treated with berberine were compared with those in the control group. The length of most circRNAs was 0–1000 nt, which is




(a)



(b)

FIGURE 1: Continued.



(c)

FIGURE 1: Continued.




FIGURE 1: Anticancer effects of berberine (BBR) in gastric cancer cells. (a) AGS and HGC27 cell viability following berberine treatment for 24, 48, and 72 h determined using CCK-8 assay. (b) Colony formation assay showed the clone number in AGS and HGC27 cells treated with 50 μM BBR or control. (c) AGS and HGC27 cell apoptosis after treatment with BBR for 48 h determined using flow cytometry. (d) Mitochondrial membrane potential of AGS cells following treatment with BBR for 48 h and staining with JC-1 probe. Data are presented as the means ± SEM; *P* value significance: 0 ≤ \*\*\*\* < 0.0001 ≤ \*\*\* < 0.001 ≤ \*\* < 0.01 ≤ \* < 0.05 ≤ ns; ns: not significant.

consistent with the findings of a previous study [23]. CircRNAs are mainly derived from exons or introns of their host linear transcripts and are involved in regulating their host gene expression [24, 25]. Accordingly, after screening

31 DE circRNAs induced by berberine, the annotations of their host genes were estimated by GO and KEGG analyses. Numerous pathways have been demonstrated to exert important effects in the mechanisms of GC onset and





FIGURE 2: Continued.




FIGURE 2: Overview of circular RNA (circRNA) profiles based on RNA-seq analysis. (a) Distribution of identified circRNAs on the human chromosome. (b) Length distribution of the circRNAs. (c) Genomic origin of the identified circRNAs (circRNA: exonic origin; ciRNA: intronic origin).




FIGURE 3: Continued.







FIGURE 3: Differentially expressed (DE) circRNAs in AGS cells after treatment with berberine (BBR). (a) Clustered heatmap showing all DE circRNAs between the BBR-treated group and control group. (b) Volcano plot filtering identified DE circRNAs selected with  $|\log_2(\text{fold change})| \geq 1$  and  $P$  value  $< 0.05$ . (c) Total number of up- and downregulated DE circRNAs.

development, including interleukin-1 [26], NF- $\kappa$ B [27], and the Notch signaling pathway [28].


Analyses of host genes cannot completely reveal the roles of circRNAs. circRNAs can also function as miRNA sponges [29]. In this study, such circRNAs (circRNA2499, hsa\_circ\_0003423, and hsa\_circ\_0006702) were identified and further analyzed. circRNA2499 is a newly predicted circRNA in our current RNA-seq analysis. circRNA-miRNA-mRNA networks with these validated circRNAs were constructed to predict their biological functions. Several functions were reported to be closely associated with GC, including cell

cycle, cell-substrate adhesion, apoptosis, and the NF- $\kappa$ B and MAPK signaling pathways.

In enrichment analyses, cell apoptosis and mitochondrion alterations induced by berberine were validated in our study. Several crucial regulators of apoptosis maintain or destroy the integrity of the mitochondrial membrane [30]. Our results indicate that berberine can reduce  $\Delta\psi_m$  in AGS cells. NF- $\kappa$ B is one of the most important transcription factors linking chronic inflammation and cancer and is activated in cancer cells and the tumor microenvironments of most cancers [31]. Previous reports demonstrated that




(a)



(b)

FIGURE 4: Continued.



(c)

FIGURE 4: GO and KEGG pathway analyses of DE circRNA-hosting genes. (a) Representative GO terms of biological processes, cellular components, and molecular functions, (b) top 20 GO analyses, and (c) top 20 KEGG analyses.






FIGURE 5: Continued.



(c)

FIGURE 5: Validation of DE circRNAs. (a) Four circRNAs with significantly different expression levels between the berberine-treated group and the control group in AGS cells, which were assessed via the RNA-seq approach. (b, c) RT-qPCR validation of DE circRNAs in AGS and HGC27 cells. Data are presented as means  $\pm$  SEM. *P* value significance:  $0 \leq **** < 0.0001 \leq *** < 0.001 \leq ** < 0.01 \leq * < 0.05$ .



(a)

FIGURE 6: Continued.






FIGURE 6: circRNA-miRNA-mRNA network and biological function prediction of validated circRNAs. circRNA-miRNA-mRNA regulation networks of (a) circRNA2499, (b) hsa\_circ\_0003423, and (c) hsa\_circ\_0006702 visualized using Cytoscape v3.7.2. Enrichment analysis of (d) target genes of each circRNA and (e) merged target genes of the three circRNAs using Metascape database.

TABLE 1: MCODE module enrichment analysis of merged target genes in three validated circRNAs in the circRNA-miRNA-mRNA network based on the Metascape database.

| Term          | Category           | Description          | Log10 (P) |
|---------------|--------------------|----------------------|-----------|
| R-HSA-69278   | Reactome gene sets | Cell cycle, mitotic  | -6.1      |
| R-HSA-68877   | Reactome gene sets | Mitotic prometaphase | -5.8      |
| R-HSA-1640170 | Reactome gene sets | Cell cycle           | -5.6      |

berberine can suppress NF- $\kappa$ B expression in gastric [32], colon [33], lung [34], breast [35], and prostate [36] cancer cells to exert anticancer effects. The MAPK family signaling cascade, which includes JNK, p38 MAPK, and ERK, is one of the main intracellular pathways for apoptosis [37]. It has

been reported that berberine can induce colon cancer cell apoptosis through continuous phosphorylation of JNK and p38 MAPK [38], whereas another study reported that berberine inhibits GC cell growth by inactivating the p38/JNK pathway [32]. Our findings indicate that circRNA-

miRNA-mRNA regulatory interactions play important roles in the treatment of GC with berberine.

Our study provides novel insight into the treatment of GC using berberine. Nevertheless, the animal and clinical assays are needed to investigate the regulatory relationships between DE circRNAs and their target miRNAs and mRNAs.

## 5. Conclusions

To summarize, circRNA sequencing analysis was conducted on berberine-treated and untreated AGS cells after investigating the appropriate concentrations and treatment durations for berberine to exert its anti-GC effects. The results, for the first time, demonstrated that berberine may influence cancer-related pathways by regulating circRNA expression and provided novel understandings of the mechanisms of berberine treatment for GC. The identified circRNAs, such as circRNA2499, hsa\_circ\_0003423, and hsa\_circ\_0006702, could be potential targets for GC treatment. Further studies are needed to evaluate the precise functions of these DE circRNAs in GC.

## Data Availability

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## Authors' Contributions

Meng Wang and Letao Sun contributed equally to this work.

## Acknowledgments

This work was supported by the National Natural Science Foundation of China (grant number 81874377), Shanghai Leading Talents Program (grant number 2020068), and internal fund of the Shanghai Jiao Tong University Affiliated Sixth People's Hospital (grant number LYZY-0297).

## Supplementary Materials

Table S1: primers used for RT-qPCR assay. Table S2: mRNA names corresponding to transcript IDs. Table S3: enrichment analysis of target genes of each validated circRNA in the circRNA-miRNA-mRNA network based on the Metascape database. Table S4: enrichment analysis of the merged target genes of three validated circRNAs in the circRNA-miRNA-mRNA network based on the Metascape database (*Supplementary Materials*)

## References

- [1] E. Van Cutsem, X. Sagaert, B. Topal, K. Haustermans, and H. Prenen, "Gastric cancer," *The Lancet*, vol. 388, no. 10060, pp. 2654–2664, 2016.
- [2] F. Bray, J. Ferlay, I. Soerjomataram, R. L. Siegel, L. A. Torre, and A. Jemal, "Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries," *CA: A Cancer Journal for Clinicians*, vol. 68, no. 6, pp. 394–424, 2018.
- [3] Z. Zhou, Z. Lin, Y. He et al., "The long noncoding RNA D63785 regulates chemotherapy sensitivity in human gastric cancer by targeting miR-422a," *Molecular Therapy-Nucleic Acids*, vol. 12, pp. 405–419, 2018.
- [4] P. Hohenberger and S. Gretschel, "Gastric cancer," *The Lancet*, vol. 362, no. 9380, pp. 305–315, 2003.
- [5] S. Meng, H. Zhou, Z. Feng et al., "CircRNA: functions and properties of a novel potential biomarker for cancer," *Molecular Cancer*, vol. 16, no. 1, p. 94, 2017.
- [6] J. Salzman, "Circular RNA expression: its potential regulation and function," *Trends in Genetics*, vol. 32, no. 5, pp. 309–316, 2016.
- [7] D. Han, J. Li, H. Wang et al., "Circular RNA circMTO1 acts as the sponge of microRNA-9 to suppress hepatocellular carcinoma progression," *Hepatology*, vol. 66, no. 4, pp. 1151–1164, 2017.
- [8] W. Tang, K. Fu, H. Sun, D. Rong, H. Wang, and H. Cao, "CircRNA microarray profiling identifies a novel circulating biomarker for detection of gastric cancer," *Molecular Cancer*, vol. 17, no. 1, p. 137, 2018.
- [9] K.-Y. Hsiao, Y.-C. Lin, S. K. Gupta et al., "Noncoding effects of circular RNA CCDC66 promote colon cancer growth and metastasis," *Cancer Research*, vol. 77, no. 9, pp. 2339–2350, 2017.
- [10] A. Bishayee and G. Sethi, "Bioactive natural products in cancer prevention and therapy: progress and promise," *Seminars in Cancer Biology*, vol. 40, pp. 1–3, 2016.
- [11] K. Changmin and K. Bonglee, "Anti-cancer natural products and their bioactive compounds inducing er stress-mediated apoptosis: a review," *Nutrients*, vol. 10, no. 8, p. 1021, 2018.
- [12] D. D. Xiong, Z. B. Feng, Z. F. Lai et al., "High throughput circRNA sequencing analysis reveals novel insights into the mechanism of nitidine chloride against hepatocellular carcinoma," *Cell Death & Disease*, vol. 10, no. 9, pp. 616–658, 2019.
- [13] Z. Zhang, B. Li, P. Xu, and B. Yang, "Integrated whole transcriptome profiling and bioinformatics analysis for revealing regulatory pathways associated with quercetin-induced apoptosis in HCT-116 cells," *Frontiers in Pharmacology*, vol. 10, p. 798, 2019.
- [14] C. Zhang, J. Sheng, G. Li et al., "Effects of berberine and its derivatives on cancer: a systems pharmacology review," *Frontiers in Pharmacology*, vol. 10, p. 1461, 2020.
- [15] M. D. Robinson, D. J. McCarthy, and G. K. Smyth, "edgeR: a Bioconductor package for differential expression analysis of digital gene expression data," *Bioinformatics*, vol. 26, no. 1, pp. 139–140, 2010.
- [16] P. Shannon, A. Markiel, O. Ozier et al., "Cytoscape: a software environment for integrated models of biomolecular interaction networks," *Genome Research*, vol. 13, no. 11, pp. 2498–2504, 2003.
- [17] Y. Zhou, B. Zhou, L. Pache et al., "Metascape provides a biologist-oriented resource for the analysis of systems-level datasets," *Nature Communications*, vol. 10, no. 1, p. 1523, 2019.
- [18] G. D. Bader and C. W. V. Hogue, "An automated method for finding molecular complexes in large protein interaction networks," *BMC Bioinformatics*, vol. 4, no. 2, 2003.

- [19] Q. Hu, L. Li, X. Zou et al., "Berberine attenuated proliferation, invasion and migration by targeting the AMPK/HNF4 $\alpha$ /WNT5A pathway in gastric carcinoma," *Frontiers in Pharmacology*, vol. 9, p. 1150, 2018.
- [20] T. Yi, L. Zhuang, G. Song, B. Zhang, G. Li, and T. Hu, "Akt signaling is associated with the berberine-induced apoptosis of human gastric cancer cells," *Nutrition and Cancer*, vol. 67, no. 3, pp. 523–531, 2015.
- [21] H.-L. Shi, X.-J. Wu, Y. Liu, and J.-Q. Xie, "Berberine counteracts enhanced IL-8 expression of AGS cells induced by evodiamine," *Life Sciences*, vol. 93, no. 22, pp. 830–839, 2013.
- [22] A. Pandey, K. Vishnoi, S. Mahata et al., "Berberine and curcumin target survivin and STAT3 in gastric cancer cells and synergize actions of standard chemotherapeutic 5-fluorouracil," *Nutrition and Cancer*, vol. 67, no. 8, pp. 1295–1306, 2015.
- [23] Q. Zheng, C. Bao, W. Guo et al., "Circular RNA profiling reveals an abundant circHIPK3 that regulates cell growth by sponging multiple miRNAs," *Nature Communications*, vol. 7, p. 11215, 2016.
- [24] Y. Zhang, X.-O. Zhang, T. Chen et al., "Circular intronic long noncoding RNAs," *Molecular Cell*, vol. 51, no. 6, pp. 792–806, 2013.
- [25] Z. Li, C. Huang, C. Bao et al., "Exon-intron circular RNAs regulate transcription in the nucleus," *Nature Structural & Molecular Biology*, vol. 22, no. 3, pp. 256–264, 2015.
- [26] Z. Gong, J. Ma, H. Su et al., "Interleukin-1 receptor antagonist inhibits angiogenesis in gastric cancer," *International Journal of Clinical Oncology*, vol. 23, no. 4, pp. 659–670, 2018.
- [27] O. Sokolova and M. Naumann, "NF- $\kappa$ B signaling in gastric cancer," *Toxins*, vol. 9, no. 4, p. 119, 2017.
- [28] E. S. Hibdon, N. Razumilava, T. M. Keeley et al., "Notch and mTOR signaling pathways promote human gastric cancer cell proliferation," *Neoplasia*, vol. 21, no. 7, pp. 702–712, 2019.
- [29] B. Han, J. Chao, and H. Yao, "Circular RNA and its mechanisms in disease: from the bench to the clinic," *Pharmacology & Therapeutics*, vol. 187, pp. 31–44, 2018.
- [30] C. Adrain and S. J. Martin, "The mitochondrial apoptosome: a killer unleashed by the cytochrome seas," *Trends in Biochemical Sciences*, vol. 26, no. 6, pp. 390–397, 2001.
- [31] K. Taniguchi and M. Karin, "NF- $\kappa$ B, inflammation, immunity and cancer: coming of age," *Nature Reviews Immunology*, vol. 18, no. 5, pp. 309–324, 2018.
- [32] Y. Wang, M. Zhou, and D. Shang, "Berberine inhibits human gastric cancer cell growth via deactivation of p38/JNK pathway, induction of mitochondrial-mediated apoptosis, caspase activation and NF-kappa B inhibition," *Journal of the Balkan Union of Oncology*, vol. 25, no. 1, pp. 314–318, 2020.
- [33] K. N. C. Murthy, G. K. Jayaprakasha, and B. S. Patil, "The natural alkaloid berberine targets multiple pathways to induce cell death in cultured human colon cancer cells," *European Journal of Pharmacology*, vol. 688, no. 1, pp. 14–21, 2012.
- [34] L. Fu, W. Chen, and W. Guo, "Berberine targets AP-2/hTERT, NF- $\kappa$ B/COX-2, HIF-1 $\alpha$ /VEGF and cytochrome-c/caspase signaling to suppress human cancer cell growth," *PLoS One*, vol. 8, no. 7, Article ID e69240, 2013.
- [35] H.-P. Kuo, T.-C. Chuang, S.-C. Tsai et al., "Berberine, an isoquinoline alkaloid, inhibits the metastatic potential of breast cancer cells via Akt pathway modulation," *Journal of Agricultural and Food Chemistry*, vol. 60, no. 38, pp. 9649–9658, 2012.
- [36] S. B. Muralimanoharan, A. B. Kunnumakkara, B. Shylesh et al., "Butanol fraction containing berberine or related compound from nexrutine inhibits NF $\kappa$ B signaling and induces apoptosis in prostate cancer cells," *The Prostate*, vol. 69, no. 5, pp. 494–504, 2009.
- [37] E. K. Kim and E.-J. Choi, "Compromised MAPK signaling in human diseases: an update," *Archives of Toxicology*, vol. 89, no. 6, pp. 867–882, 2015.
- [38] W.-H. Hsu, Y.-S. Hsieh, H.-C. Kuo et al., "Berberine induces apoptosis in SW620 human colonic carcinoma cells through generation of reactive oxygen species and activation of JNK/p38 MAPK and FasL," *Archives of Toxicology*, vol. 81, no. 10, pp. 719–728, 2007.