



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Short communication

COVID-19 vaccine hesitancy among persons living in homeless shelters in France



C. Longchamps^a, S. Ducarroz^b, L. Crouzet^b, N. Vignier^{a,c,d}, L. Pourtau^e, C. Allaire^f, A.C. Colleville^f, T. El Aarbaoui^a, M. Melchior^{a,c}, the ECHO study group¹

^a Sorbonne Université, INSERM, Institut Pierre Louis d'Épidémiologie et de Santé Publique, IPLESP, Équipe de Recherche en Épidémiologie Sociale, Paris F75012, France

^b Health Services and Performance Research lab (HeSPeR, EA 7425), Université Claude Bernard Lyon 1, Lyon F6900, France

^c French Collaborative Institute on Migration, CNRS, Aubervilliers F93300, France

^d Inserm CIC 1424 & Corevih Guyane, Centre Hospitalier de Cayenne, France

^e Université Paul Valéry Montpellier 3, Laboratoire d'Etudes et de Recherche en Sociologie et en Ethnologie de Montpellier, (EA 4584), Montpellier F34080, France

^f Santé Publique France, French National Public Health Agency, Saint-Maurice F94415, France

ARTICLE INFO

Article history:

Received 5 December 2020

Received in revised form 2 March 2021

Accepted 4 May 2021

Available online 12 May 2021

ABSTRACT

COVID-19 vaccine hesitancy is frequent and can constitute a barrier to the dissemination of vaccines once they are available. Unequal access to vaccines may also contribute to socioeconomic inequalities with regard to COVID-19. We studied vaccine hesitancy among persons living in homeless shelters in France between May and June 2020 (n = 235). Overall, 40.9% of study participants reported vaccine hesitancy, which is comparable to general population trends in France. In multivariate regression models, factors associated with vaccine hesitancy are: being a woman (OR = 2.55; 95% CI 1.40–4.74), living with a partner (OR = 2.48, 95% CI 1.17–5.41), no legal residence in France (OR = 0.51, 95% CI 0.27–0.92), and health literacy (OR = 0.38, 95% CI 0.21, 0.68). Our results suggest that trends in vaccine hesitancy and associated factors are similar among homeless persons as in the general population. Dissemination of information on vaccine risks and benefits needs to be adapted to persons who experience severe disadvantage.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

As the COVID-19 pandemic spreads in many parts of the world, access to an effective vaccine offers hope of a new much-needed preventive measure in upcoming months. However, in the area of COVID-19 infection as well as with regard to other infectious diseases, there are concerns that vaccine hesitancy may be a barrier to rapid roll-out of a new vaccine. Indeed, in Europe, it is estimated that approximately one person out of four would not want to be vaccinated [1–3], young persons, women, and those not feeling at risk with COVID-19 being most likely to be reluctant. Among reasons cited to explain vaccine hesitancy, fear of side effects is the most frequent one (55% of vaccine hesitant individuals), particularly among women [3]. In recent months, and particularly since

the approval of several COVID-19 vaccines, opinion polls suggest that vaccine hesitancy has decreased. However, inequalities with regard to vaccination intentions remain [4].

Members of disadvantaged populations, appear disproportionately impacted by the COVID-19 epidemic. In particular, persons who live in shared facilities such as homeless shelters have been shown to be at high risk of getting infected [5]. Moreover, migrants from low or middle-income countries have also been shown to have high levels of COVID-19 morbidity and mortality [6], and for those who live in homeless shelters the risks may be compounded. Elevated levels of COVID-19 morbidity and mortality among persons who experience socioeconomic disadvantage and migrants likely reflect a high prevalence of risk factors of severe COVID-19 infection (ex. diabetes, hypertension) and insufficient information access to the health care system [7]. However there is also suggestion of lower levels of adherence to preventive measures, although data on this topic are lacking [8].

Persons residing in homeless shelters and migrants may constitute relevant target groups for COVID-19 vaccination, however vaccine acceptability in this population is not known. We studied

¹ The ECHO study group includes Pierre Anquetil, Léa Balage, Estelle Dussert, Betty Girard, Laure Luyinga Nzuzi, Hermine Metias, Nathalie Oprescu, Philippe Rebouffat-Roux (Habitat et Humanisme), Anahaid Armenian, Marianne Auffret, Perrine Leclerc (Association Aurore), Faouzi Bertrand (Groupe SOS), Mourad Bouderbail (Croix Rouge Française), Antoine Denis (SIAO 67), Christelle Witczak (Empreintes Sud 77), François Fortin (La Rose des Vents).

the intention to be vaccinated against COVID-19 among persons living in homeless shelters in France, a majority of whom are recent migrants, integrating risk factors identified by the working group on vaccine hesitancy established by the WHO [9]. The study was conducted in the Spring 2020, before anti-COVID-19 vaccines were approved for use.

2. Methods

2.1. Study population

Our investigation is based upon data collected in ECHO, a cross-sectional study conducted among residents of 18 short and long-term homeless shelters in the Paris ($n = 12$), Lyon ($n = 5$) and in Strasbourg ($n = 1$) regions. The study interview, aiming to measure participants' level of information regarding COVID-19, their knowledge, implementation and acceptability of various preventive measures, was conducted between May 2nd and June 28, 2020, corresponding to the 1st COVID-19 lockdown period (March 16–May 11, 2020) and its immediate aftermath, by trained interviewers, in person or by telephone. The questionnaire was administered in French, in English or in the participant's language: 25% of questionnaires were completed with the assistance of a trained translator contacted by telephone (most frequently used languages: Arabic, Pashto, Dari, Tigrinya, and Amharic). In total, 929 residents were invited to take part in the survey, 131 refused participation, 263 were unavailable, and 535 (57.6%) completed the study questionnaire. An item on the intention to be vaccinated against COVID-19 was added to the ECHO questionnaire from the end of May 2020, and 235 study participants provided information.

The ECHO study received approval of the Ethical Research Committee of the University of Paris (CER-2020–41).

2.2. Measures

The intention to be vaccinated was ascertained by the following item: "If a vaccine existed would you be willing to get vaccinated"? Participants who responded 'no' or 'I don't know' were considered probably vaccine hesitant.

Factors potentially associated with 'vaccine hesitancy', which we studied, based on the SAGE consortium, included sociodemographic characteristics (sex, age, household composition, presence of children, WHO region of birth, administrative status, educational level, French language level, employment prior to lockdown, health insurance, duration of residence in the homeless shelter, social support), health (depression ascertained with the PHQ-9 using a cut-off of 10 to identify persons with a significant level of symptoms [10], and self-reported chronic health problems, fear of getting infected by COVID-19), as well as health-seeking information (trust in official information about COVID-19, health literacy ascertained using the 'Appraisal of health information' subscale ($n = 5$ items) of the Health Literacy Questionnaire (HLQ) [11], primary sources of COVID-19 information).

2.3. Statistical analysis

To examine factors associated with the intention to be vaccinated against COVID-19, we conducted bivariate analyses with the chi-square statistic. Next all variables associated with the study outcome with a $p \leq 0.25$ were included in a step-wise logistic regression model.

All statistical analyses were conducted using the R software version 1.3.1093.

3. Results

As shown in Table 1, ECHO study participants were mainly male (66.3%), young (61.1% below 35 years of age), lived alone and without children (respectively 81.5 and 65.5%), were migrant (92.5% born outside of France), did not have legal residence in France (60.8%), had low educational attainment (62.9% with secondary school education or less), low or intermediate level knowledge of the French language (80.7%), no employment prior to the COVID-19 lockdown (70.5%), and resided in the shelter where they were interviewed for a short period of time (60.4% at most one year). In terms of health, 27.9% of participants had symptoms of clinical depression, 25.5% reported another chronic disease, 67.0% feared getting infected by COVID-19, 76.4% expressed trust in official information on COVID-19, 49.3% had low health literacy, and the leading sources for information about COVID-19 were the TV (63.0%) and social media (84.0%).

Overall, 40.9% reported not being willing to get vaccinated against SARS-CoV-2 – among those 71.2% indicated they would not want to be vaccinated and 28.8% did not know what to answer. In bivariate analyses factors associated with 'vaccine hesitancy' included sex (55.9% in women vs. 32.5% in men, $p = 0.0004$), household composition (living with a partner: 56.1 vs. living alone: 36.9%, $p = 0.0232$), living with children (yes: 55.7 vs. no: 33.1%, $p = 0.0009$), region of birth (Europe: 52.7%, Africa: 43.6%, Eastern Mediterranean: 30.0%, Other: 12.5%, $p = 0.05$), legal status (French/legal residence: 48.9% vs. no legal residence: 35.9%, $p = 0.05$), and health literacy (low: 50.9% vs. intermediate/high: 31.3%, $p = 0.0026$). In multivariate logistic regression models, factors which remained associated with 'vaccine hesitancy' were sex (female vs. male: OR = 2.55; 95% CI 1.40–4.74), household composition (participants living with a partner vs. living alone: OR = 2.48, 95% CI 1.17–5.41), administrative status (no legal residence vs. French/with legal residence: OR = 0.51, 95% CI 0.27–0.92), and health literacy (intermediate/high vs. low: OR = 0.38, 95% CI 0.21, 0.68).

4. Discussion

Our study conducted among persons living in short and long-term homeless shelters in France, among whom a majority were migrant, shows approximately 41% of COVID-19 hesitancy in the Spring of 2020. Factors associated with low intention to be vaccinated against COVID-19 include female sex, living with a partner, as well as French citizenship/legal residence, and low health literacy. Our data suggest that factors associated with 'vaccine hesitancy' appear similar in this highly disadvantaged group to those reported in the general population, however their absolute contribution may be higher.

First, we need to address limitations and strengths of this study, which can influence data interpretation. First, ECHO is not representative of persons living in homeless shelters in France. However, we collaborated with several different organizations and surveyed 18 homeless shelters hosting different types of populations. Importantly, homeless shelters and participants were recruited independently of the risk of COVID-19 infection, which rules out the possibility of systematic bias. Moreover, our sample's characteristics (a majority of young men who are migrant) is similar to that observed across homeless facilities across France [12]. Second, we had no information regarding participants' COVID-19 infection, which could influence perceptions of preventive measures and particularly vaccine acceptance. However at the time we conducted our study (May–June 2020), neither virological nor serological COVID-19 tests were easily accessible, making it difficult to ascertain the actual prevalence of infection. This lead us

Table 1
Factors associated with COVID-19 vaccine hesitancy (ECHO study, France, n = 240, May–June 2020, %, chi-square, multivariate logistic regression, OR; 95% CI).

	% in the ECHO study	% vaccine hesitant	p-value	Multivariate OR (95% CI)
Socio-demographic characteristics				
Sex: Male	63.3	32.5	0.0004	1.0
Female	36.7	55.5		2.55 (1.40, 4.74)
Age: 18–24 years	19.3	38.6	0.86	
25–34 years	41.8	43.4		
35–49 years	25.9	41.6		
>= 50 years	13.0	35.4		
Household composition: Living alone	81.5	36.9	0.02	1.0
Living with a partner	18.5	56.1		2.48 (1.17, 5.41)
Living with children: No	65.5	33.1	0.0009	
Yes	34.5	55.7		
Country of birth (WHO region):	15.6	52.7	0.05	
Europe	59.5	43.6		
Africa	21.5	30.0		
Eastern Mediterranean	3.4	12.5		
Other				
Administrative status: French/legal residence	39.2	48.9	0.05	1.0
No legal residence	60.8	35.9		0.51 (0.27, 0.92)
Educational level: No schooling	24.1	37.7	0.66	
Secondary school	38.8	39.3		
≥ High school degree	37.1	44.7		
French language level: Low	49.2	36.2	0.24	
Intermediate	31.5	48.6		
High	19.3	42.2		
Work status prior to lockdown: Every day	19.2	55.6	0.06	
From time to time	10.3	45.8		
No	70.5	36.2		
Health insurance: No	17.4	21.9	0.01	
Yes	82.6	44.6		
Duration of residence in the shelter: ≤1 month	11.3	26.9	0.09	
1 month – 1 year	49.1	39.3		
> 1 year	39.6	49.4		
Social support outside the shelter: No	5.5	23.1	0.18	
Yes	94.5	41.9		
Health				
Depression (PHQ-9): No	72.1	42.6	0.61	
Yes	27.9	39.1		
Chronic disease: No	74.5	43.9	0.15	
Yes	25.5	33.3		
Fear of getting infected by COVID-19: No	30.9	51.4	0.07	
Yes	67.0	36.7		
Doesn't know	2.1	20.0		
Trust in official information about COVID-19: No	23.6	52.7	0.04	
Yes	76.4	37.4		
Health literacy: Low	49.3	50.9	0.003	1.0
Intermediate/high	50.7	31.3		0.38 (0.21, 0.68)
Source of COVID-19 related information				
Television: No	37.0	37.5	0.42	
Yes	63.0	42.8		
Social media: No	16.0	38.9	0.78	
Yes	84.0	41.4		

to focus on perceptions of the illness and its impact. Main strengths of ECHO are: a) the inclusion of a large sample of individuals living in homeless shelters in the two largest metropolitan areas of France (Paris and Lyon); b) access to translation services which made it possible to interview persons who did not speak French.

Our findings are generally in line with observations from general population surveys. Previous studies had suggested that women are more likely to express COVID-19 vaccine hesitancy than men [2,3]. In particular, they seem especially concerned about possible side effects, which may reflect overall higher levels of concerns about health [3]. Additionally, women are more likely than men to seek information about health on the Internet and social media [13], which may put them in contact with negative information regarding vaccines. This suggests that it is necessary to better control information that is disseminated via the Internet, as well as provide persons with accurate information in these much accessed media.

Our study found that persons who live with a partner and are French or have legal residence in France are especially likely to report a low intention to be vaccinated against COVID-19. This finding may be related to a higher dissemination of anti-vaccine messages in media used by this group. It may also be that these persons have come across vaccine-critical content (in mainstream news media for instance) in the years prior to the COVID-19 epidemic, as controversies in this area have emerged in France as early as in 2009. Additionally, health care is not always adapted to migrants' needs, resulting in inadequate or negative experiences, which they are likely to encounter with time [14]. Prior studies reported an influence of political views on vaccine hesitancy [1], which recent migrants may be less exposed to. Moreover, we found a relationship between low health literacy, which indicates individuals' limited capacity to seek and evaluate health information, and vaccine hesitancy. This is consistent with research on vaccine hesitancy in other areas [15] and indicates that

all messages on measures which can prevent COVID-19 should be adapted to be easily understood by all.

In conclusion, our study shows elevated levels of low intention to be vaccinated against COVID-19 among persons living in homeless shelters, who may be at higher risk of COVID-19 infection. As this population is likely to be a relevant target group for future COVID-19 vaccine deployment [16], it is important to ensure that the information regarding the COVID-19 infection and vaccine is disseminated in an appropriate way and all barriers lifted, in order to inform properly and increase vaccine adherence.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement

The funders of our study are: The French Collaborative Institute on Migration, France's Public Health Agency, and the French National Research Agency (Flash COVID-19).

References

- [1] Ward JK, Alleaume C, Peretti-Watel P. The French public's attitudes to a future COVID-19 vaccine: the politicization of a public health issue. *Soc Sci Med* 2020;265:113414.
- [2] Detoc M, Bruel S, Frappe P, Tardy B, Botelho-Nevers E, Gagneux-Brunon A. Intention to participate in a COVID-19 vaccine clinical trial and to get vaccinated against COVID-19 in France during the pandemic. *Vaccine*. 2020;38(45):7002–6.
- [3] Neumann-Böhme S, Varghese NE, Sabat I, Barros PP, Brouwer W, van Exel J, et al. Once we have it, will we use it? A European survey on willingness to be vaccinated against COVID-19. *Eur J Health Econ*. 2020;21(7):977–82.
- [4] Burger AE, Reither EN, Mamelund SE, Lim S. Black-white disparities in 2009 H1N1 vaccination among adults in the United States: a cautionary tale for the COVID-19 pandemic. *Vaccine*. 2021;39(6):943–51.
- [5] Baggett TP, Keyes H, Sporn N, Gaeta JM. Prevalence of SARS-CoV-2 infection in residents of a large homeless shelter in Boston. *J Am Med Assoc* 2020;323(21):2191–2.
- [6] Drefahl S, Wallace M, Mussino E, Aradhya S, Kolk M, Brandén M, et al. A population-based cohort study of socio-demographic risk factors for COVID-19 deaths in Sweden. *Nat Commun* 2020;11(1):5097.
- [7] Clark E, Fredricks K, Woc-Colburn L, Bottazzi ME, Weatherhead J. Disproportionate impact of the COVID-19 pandemic on immigrant communities in the United States. *PLoS Negl Trop Dis* 2020;14(7):e0008484.
- [8] Ralli M, Cedola C, Urbano S, Morrone A, Ercoli L. Homeless persons and migrants in precarious housing conditions and COVID-19 pandemic: peculiarities and prevention strategies. *Eur Rev Med Pharmacol Sci*. 2020;24(18):9765–7.
- [9] World Health Organization. SAGE working group dealing with vaccine hesitancy; 2014. [https://www.who.int/immunization/sage/sage_wg_vaccine_hesitancy_apr12/en/].
- [10] Costantini L, Pasquarella C, Odone A, Colucci ME, Costanza A, Serafini G, et al. Screening for depression in primary care with Patient Health Questionnaire-9 (PHQ-9): a systematic review. *J Affect Disord* 2020;279:473–83.
- [11] Debussche X, Lenclume V, Balcou-Debussche M, Alakian D, Sokolowsky C, Ballet D, et al. Characterisation of health literacy strengths and weaknesses among people at metabolic and cardiovascular risk: Validity testing of the Health Literacy Questionnaire. *SAGE Open Med*. 2018;6:2050312118801250.
- [12] Cour des Comptes. L'hébergement des personnes sans domicile: des résultats en progrès, une stratégie à préciser; 2017. [<https://www.ccomptes.fr/sites/default/files/EzPublish/06-hebergement-personnes-sans-domicile-Tome-2.pdf2017>].
- [13] Calixte R, Rivera A, Oridota O, Beauchamp W, Camacho-Rivera M. Social and demographic patterns of health-related Internet use among adults in the United States: a secondary data analysis of the Health Information National Trends Survey. *Int J Environ Res Public Health* 2020;17(18).
- [14] Priebe S, Sandhu S, Dias S, Gaddini A, Greacen T, Ioannidis E, et al. Good practice in health care for migrants: views and experiences of care professionals in 16 European countries. *BMC Public Health*. 2011;11:187.
- [15] Wilson L, Rubens-Augustson T, Murphy M, Jardine C, Crowcroft N, Hui C, et al. Barriers to immunization among newcomers: a systematic review. *Vaccine*. 2018;36(8):1055–62.
- [16] Haute Autorité de Santé. Stratégie de vaccination contre le Sars-Cov-2 - Recommandations préliminaires sur la stratégie de priorisation des populations à vacciner; 2020. [https://www.has-sante.fr/jcms/p_3221338/fr/strategie-de-vaccination-contre-le-sars-cov-2-recommandations-preliminaires-sur-la-strategie-de-priorisation-des-populations-a-vacciner].