



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



# Digital contact tracing, community uptake, and proximity awareness technology to fight COVID-19: a systematic review

George Grekousis<sup>a,b,\*</sup>, Ye Liu<sup>a,b</sup>

<sup>a</sup> School of Geography and Planning, Department of Urban and Regional Planning, No 135, Xingang Xi Road, Guangzhou, Haizhu, 510275, China

<sup>b</sup> Guangdong Key Laboratory for Urbanization and Geo-simulation, Sun Yat-sen University, No 135, Xingang Xi Road, Guangzhou, Haizhu, 510275, China

## ARTICLE INFO

### Keywords:

Digital contact tracing  
manual contact tracing  
SARS-CoV-2  
location-awareness  
proximity awareness  
smartphone

## ABSTRACT

Digital contact tracing provides an expeditious and comprehensive way to collect and analyze data on people's proximity, location, movement, and health status. However, this technique raises concerns about data privacy and its overall effectiveness. This paper contributes to this debate as it provides a systematic review of digital contact tracing studies between January 1, 2020, and March 31, 2021. Following the PRISMA protocol for systematic reviews and the CHEERS statement for quality assessment, 580 papers were initially screened, and 19 papers were included in a qualitative synthesis. We add to the current literature in three ways. First, we evaluate whether digital contact tracing can mitigate COVID-19 by either reducing the effective reproductive number or the infected cases. Second, we study whether digital is more effective than manual contact tracing. Third, we analyze how proximity/location awareness technologies affect data privacy and population participation. We also discuss proximity/location accuracy problems arising when these technologies are applied in different built environments (i.e., home, transport, mall, park). This review provides a strong rationale for using digital contact tracing under specific requirements. Outcomes may inform current digital contact tracing implementation efforts worldwide regarding the potential benefits, technical limitations, and trade-offs between effectiveness and privacy.

## 1. Introduction

The coronavirus disease 2019 (COVID-19) is probably the most severe public health crisis our world has faced in the last decades (Maiti et al., 2021; Velraj & Haghighat, 2020). It has not only posed a serious threat against human life, but it has significantly impacted every aspect of social and economic activity (Ge et al., 2020; Rahmani and Mirmahaleh, 2020). Considering its unprecedented scale, the United Nations declared the pandemic as a social, human, and economic crisis that threatens human wellbeing and society's sustainability (United Nations, 2020). The COVID-19 pandemic created the worst recession since the Great Depression, lasting from 1929 to 1939 (International Monetary Fund, 2020). As a consequence, people are likely to become poorer in the mid-to-long term (Batty, 2020), with an anticipated increase in suicide rates, substance abuse, domestic violence, homelessness, and food insecurity (DeLuca, Papageorge, Mitchel, & Kalish, 2020). Moreover, the psychological impact of social distancing and other non-pharmaceutical interventions (i.e., lockdowns) is likely to worsen

due to the post-traumatic disorder effect (Fokas, Cuevas-Maraver, & Kevrekidis, 2020).

The COVID-19 pandemic with a multiple hit to health, economy, and society directly threatens the achievement of various Sustainable Development Goals (SDGs) (Zhou et al., 2020). The United Nations SDGs aim to achieve a better and more sustainable future for all societies by addressing global challenges (United Nations Development Programme, 2015). For example, SGD-3 (one of the 17 goals of the United Nations program) aims to ensure good health and wellbeing for everyone (United Nations Development Programme, 2015). Specifically, target SDG 3.3 aims to end epidemics (i.e., AIDS, malaria) or other communicable diseases. However, in the absence of medicines, or widespread vaccination, controlling an epidemic needs careful management and targeted interventions to alleviate its severe impact on society (Rahmani and Mirmahaleh, 2020). Many strategies have been introduced to mitigate the effects of COVID-19 pandemic, based on the concept of sustainable cities and environments (Jadidi et al., 2021). These include city management (Elavarasan, Pugazhendhi, Shafiullah, Irfan, &

\* Corresponding author at: School of Geography and Planning, Department of Urban and Regional Planning, No 135, Xingang Xi Road, Guangzhou, Haizhu, 510275, China.

E-mail addresses: [geograik@gmail.com](mailto:geograik@gmail.com), [graikousis@mail.sysu.edu.cn](mailto:graikousis@mail.sysu.edu.cn) (G. Grekousis), [liuye25@mail.sysu.edu.cn](mailto:liuye25@mail.sysu.edu.cn) (Y. Liu).

<https://doi.org/10.1016/j.scs.2021.102995>

Received 20 January 2021; Received in revised form 2 April 2021; Accepted 2 May 2021

Available online 12 May 2021

2210-6707/© 2021 Elsevier Ltd. All rights reserved.

Anvari-Moghaddam, 2021; Zhou et al., 2020), lockdown management (Rahmani and Mirmahaleh, 2020), and digital contact tracing through the use of proximity awareness technologies, geospatial technologies, and wireless communication infrastructure (Jadidi et al., 2021).

### 1.1. Digital contact tracing

Given the life-saving importance of rapid action against a novel pathogen, tracing the contacts of infected individuals is critical for controlling its spread (Rahmani and Mirmahaleh, 2020). Contact tracing is a well-established method to interrupt chains of infection transmission by locating and isolating those in close contact with an infected individual (Braithwaite, Callender, Bullock, & Aldridge, 2020). Successful contact tracing is a powerful tool to keep the virus spread at manageable levels. However, a precondition should be met; finding active cases as soon as possible (World Health Organization, 2020). Failing to trace asymptomatic infected individuals or newly infected patients rapidly results in silent spread of the virus in the community for days before a significant number of patients have been diagnosed and health services become aware. In particular, for viruses such as the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) where more than half of the transmissions (70%) occur before someone is symptomatic (Mizumoto, Kagaya, Zarebski, & Chowell, 2020), contact tracing should be as fast as possible. For a 3-day delay in notification, models show that the COVID-19 epidemic cannot be controlled (Ferretti, Wymant, & Kendall, 2020; Kretzschmar & Rozhnova, 2020). In such cases, manual contact tracing cannot significantly assist in containing the virus, especially when the epidemic is accelerating exponentially (Hellewell, Abbott, & Gimma, 2020; Peak Corey, Childs Lauren, Grad Yonatan, & Buckee Caroline, 2017). This emphasizes the need for rapid and effective contact tracing. By the same token, the World Health Organization (WHO) urges to “shift from reliance on existing surveillance networks to system of rapid, population-level active surveillance” (World Health Organization, 2020).

Digital contact tracing responds to this need and may overcome the problems mentioned above (Ferretti et al., 2020). Digital contact tracing uses a variety of technologies such as, location-based services, geospatial technologies, proximity awareness technology, machine learning algorithms, and automated decision making to scrutinize individual’s digital footprint and trace those who are potentially infected, locate their close contacts and enforce specific health protocols or social distancing (Calvo, Deterding, & Ryan, 2020). With digital contact tracing, many of the logistic challenges of massive contact tracing following the traditional approach are eased (Aleta, Martín-Corral, & Piontti, 2020). Through proximity awareness technology, mobile apps, and in some case GIS analysis and mapping (Geographical Information Systems), digital contact tracing decreases the response time compared to manual tracing, allowing for quicker tracing of individuals exposed to the virus. Therefore, a rapid assessment of close contacts through digital contact tracing can break the chain of infection earlier, curbing the virus spread (Ferretti et al., 2020). For a more thorough review of digital contact tracing and related apps against COVID-19, one may refer to Gasser, Ienca, Scheibner, Sleight, and Vayena (2020); Salehinejad, Niakan Kalhori, Hajesmael Gohari, Bahaadinbeigy, and Fatehi (2021), or Fagherazzi, Goetzinger, Rashid, Aguayo, and Huiart (2020).

### 1.2. Research gaps and contribution of this review

Despite the initial enthusiasm for novel technologies to contain COVID-19, especially in the spring of 2020, there is no widespread integration of digital contact tracing strategies within governmental response plans across the world. Current digital tracing endeavors portray heterogeneities on preferred system architecture (i.e., decentralized vs. centralized) (Gasser et al., 2020), proximity or location awareness technologies (i.e., Bluetooth vs. GPS) (Rosenkrantz, Schuurman, Bell, & Amram, 2020), and data privacy (privacy first vs. public

safety first) (Cohen, Gostin, & Weitzner, 2020), while their overall usefulness and effectiveness are debated (Braithwaite et al., 2020). Therefore, a systematic review is needed to summarize the key findings of digital contact tracing-related studies

Currently, there are two reviews on a similar subject (Anglemyer, Moore, & Parker, 2020; Braithwaite et al., 2020). The first review studied automated, and non-automated contact tracing approaches applied for COVID-19, SARS, MERS, influenza, and Ebola published up to April 14, 2020 (Braithwaite et al., 2020). The second review studied digital contact tracing in various epidemics (Anglemyer et al., 2020). Only four papers are related to COVID-19, out of which two report the impact of digital contact tracing on the  $R_{\text{eff}}$ . Hence, conclusions are difficult to be drawn.

The above two reviews offer a valuable understanding of multiple aspects of digital contact tracing. Yet, they fail to address some critical questions in the context of a systematic literature review, which can be attributed to the early analysis of a limited number of studies published up to May 5, 2020. First, they do not study whether digital contact tracing reduces the effective reproductive number ( $R_{\text{eff}}$ ). The  $R_{\text{eff}}$  represents the average number of secondary cases generated by a single infection case and is one of the most important metrics to monitor an epidemic (Flaxman et al., 2020). A value of  $R_{\text{eff}}$  under one would indicate that the epidemic is controlled. As  $R_{\text{eff}}$  is a measure of virus transmission in the community, it is widely used to assess the effectiveness of pharmaceutical and non-pharmaceutical interventions over time (Flaxman et al., 2020). On this account, interventions like digital contact tracing, are commonly evaluated based on their ability to drop the  $R_{\text{eff}}$  metric preferable below one. Current reviews neglect to study this aspect.

Second, there is a gap in comparing manual contact tracing to digital contact tracing. Only one study has been reported in previous reviews that compared digital to manual contact tracing effectiveness for COVID-19 (Kucharski, Klepac, & Edmunds, 2020). Third, as underlined by Braithwaite et al. (2020), there is a gap in studying how technology affects data privacy, community uptake, and the overall effectiveness of digital contact tracing. Although the above questions have been studied from individual studies, a systematic literature review that summarizes the findings is missing.

This work aims to fill these gaps by systematically reviewing digital contact tracing papers for controlling COVID-19 from January 1, 2020, to March 31, 2021. The contributions of this paper are threefold:

- First, opposed to existing studies that analyze the number of contacts identified through digital contact tracing (Braithwaite et al., 2020), we focus on the actual effect of digital contact tracing on reducing the effective reproductive number, or the number of infected individuals, an outcome extremely important for drawing related health policies. Hence, our review includes studies that numerically present the reduction in the  $R_{\text{eff}}$  or the number of infections. We also report this finding in relation to contact tracing app uptake (percentage of people using the app) and we search for functional relations between app uptake and the effectiveness of digital contact tracing. Braithwaite et al. (2020) reported one study that compared digital to manual contact. Here, we report eight studies that compare digital to manual tracing. This provides a better understanding of which strategy outperforms the other.
- Second, manual contact tracing is compared to digital contact tracing regarding their ability to reduce the  $R_{\text{eff}}$  or the number of infected cases. At least to our knowledge, this is the first time of such comparison in a systematic review for COVID-19.
- Third, we respond to another research gap as underlined by Braithwaite et al. (2020): *how technology affects digital contact tracing effectiveness*. Specifically, we also analyze how the system architecture (i.e., centralized vs. decentralized) may affect data privacy and subsequently the community uptake. We also examine the suitability of proximity awareness sensors, from the perspective of proximity

and location accuracy, across different spaces as this has a direct impact on the effectiveness of digital contact tracing. Low proximity/location accuracy may lead to false alerts and exposure notifications. Such an aspect of the problem has not been considered in previous reviews.

In conclusion, early reviews evaluating digital contact tracing did not consider the quantitative impact of digital contact tracing in reducing either the  $R_{\text{eff}}$  or the number of cases infected. Other essential features were neglected, such as how the proximity awareness technology infers inaccuracies in contact tracing and potential infringement of privacy. By evaluating the necessity, effectiveness, proximity/location accuracy, and personal data privacy concerns of such systems, this review offers vital information that can support decision making and guide future contact tracing endeavors.

## 2. Methods

### 2.1. Research objectives

This review attempts to assess the effect of digital contact tracing and balance the potential technological and ethical concerns in containing COVID-19. Specifically, the review analyzes a set of studies to provide evidence on:

- The effect of digital contact tracing in reducing the effective reproductive number ( $R_{\text{eff}}$ ) or the number of COVID-19 infections
- The necessary contact-tracing app adoption rate by the population to ensure control of the epidemic ( $R_{\text{eff}} < 1$ )
- The functional relation, if any, between the contact-tracing app uptake by the public and the effectiveness of digital contact tracing
- Whether digital contact tracing is more effective (decreasing  $R_{\text{eff}}$  or the number of infections more) than manual contact tracing
- How technical aspects such as the system architecture or the proximity/location accuracy affect data privacy, community uptake, and the overall effectiveness of digital contact tracing.

### 2.2. Eligibility criteria

We considered studies that reported a direct effect on mitigating the epidemic. Specifically, we only included studies that reported either the reduction in the effective reproductive number  $R_{\text{eff}}$  (average number of secondary cases generated by a single infection case) or the reduction in the number of infections.

Studies that did not report any numerical effect of digital contact tracing in mitigating the epidemic or were purely qualitative, were excluded. We also excluded papers that focus on the ethical and privacy concerns of digital contact tracing from a conceptual perspective only. We did not compare digital contact tracing to all potential combinations with other interventions, such as isolation, social distancing, and quarantine. Currently, many countries have developed various types of digital contact-tracing apps. We do not review such apps. For a more thorough presentation of such systems, one may refer to [Salehinejad et al. \(2021\)](#) and [Skoll, Miller, and Saxon \(2020\)](#). This paper is not a review of the limitations of digital or manual contact tracing strategies, although we discuss some of them. Further evaluation of security and privacy considerations related to hardware (i.e., data servers, firewalls) and software used (i.e., operating system) are beyond the scope of this review, but readers may refer to [Nguyen et al. \(2020\)](#) for a recent review. All reviewed manuscripts were in English.

### 2.3. Search strategy and study selection

Similar to others ([Braithwaite et al., 2020](#)), we followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) protocol. The PRISMA protocol has been the standard way

of reporting systematic reviews and meta-analysis of studies evaluating the effects of interventions ([Moher, Liberati, Tetzlaff, & Altman, 2009](#); refer to Supplementary Table S1 for the PRISMA statement). Further reading on other protocols used for systematic reviews and their comparisons is provided by [Booth et al. \(2020\)](#).

We searched PubMed, bioRxiv, medRxiv, and arXiv for articles published between January 1, 2020, and March 31, 2021. Search terms included “SARS-CoV-2” OR “COVID-19” AND “digital contact tracing” OR “automated contact tracing” OR “smartphone.” The complete query (with additional keywords) used to search the databases is presented in Supplementary Text S1. A total of 580 papers were retrieved ([Fig. 1](#)). The screening process was conducted independently by two authors and any disagreements were discussed. The initial screening of the titles and abstracts, and exclusion of duplicates, yielded 96 papers. The full text of these papers was meticulously reviewed. Those not meeting the primary criteria (i.e., reporting reduction either in  $R_{\text{eff}}$  or the number of infected cases resulting from digital contact tracing) were excluded. A final set of 19 papers was used for qualitative synthesis. Due to the high heterogeneity in the studies design, variations in technologies, and outputs, no meta-analysis has been conducted. For this reason, we synthesized study findings narratively.

### 2.4. Data extraction

Data were extracted manually using a Microsoft Excel spreadsheet. The following data were collected: year, authors, study sample, system architecture, proximity/location awareness technology used, proximity/location accuracy, data privacy risks, app uptake,  $R_{\text{eff}}$  change for digital contact tracing,  $R_{\text{eff}}$  change for manual contact tracing, infections change for digital contact tracing, infections change for manual contact tracing, description of the intervention.

### 2.5. Risk of bias

The Consolidated Health Economic Evaluation Reporting Standards (CHEERS) statement was used to assess the quality of the studies included in this review and reduce the risk of bias, an additional checkpoint of the PRISMA protocol ([Husereau et al., 2020](#)). The CHEERS statement aims to evaluate the quality of modelling studies related to health interventions and has been used in other systematic reviews ([Braithwaite et al., 2020](#)). Questions 1,6-7,9-17 and 22-25 from this statement were omitted here as they refer to the economic output of a study, and were not relevant to the modeling studies included in this review (refer to Supplementary Table S2 for the CHEERS summary). The assessment of risk of bias was conducted by the authors independently, and any conflicting opinions were resolved through discussion.

## 3. Results

We identified 19 papers on digital contact tracing, all of which were modeling studies ([Table 1](#)). Similar to others ([Braithwaite et al., 2020](#)), and due to this research topic’s high importance and timeliness, we included five preprints, while the rest fourteen papers have been peer-reviewed. The results vary significantly across studies, as different modeling and scenario settings are applied.

### 3.1. Digital contact tracing effect on the number of infections and $R_{\text{eff}}$

Nine studies reported the effect of digital contact tracing on the number of infections. [Abueg et al. \(2021\)](#) reported that if a digital tracing app is used by 75% of the population, the number of infections is reduced by 73–79%. Another study showed that an 80% uptake leads to 89% reduction in cases at the peak of the epidemic ([Almagor & Picascia, 2020](#)), while [Barrat, Cattuto, Kivela, Lehmann, & Saramaki \(2020\)](#) revealed that a 60% uptake produces a 36% reduction in epidemic size. Further, COVIDSafe, an existing tracing app created by the Australian




Fig. 1. PRISMA flowchart.

government, was assessed for its effectiveness (Currie, Peng, Jameson, Frommer, & Lyle, 2020). The model indicated that a 61% uptake of the app would reduce the total number of new cases by almost 50%. Another study estimated that a 50% adoption rate of digital contact tracing results in a 90% decrease in the peak number of infections (Nuzzo et al., 2020). López et al. (2021) calibrated an agent-based model on the French population integrating demographic and social-contact data simulated how digital contact tracing impacted COVID-19 transmission. Assuming  $R_{\text{eff}} = 2.6$ , digital tracing at a 60% adoption rate would produce a 67% reduction in peak incidence. Finally, Wilmlink, Summer, and Marsyla (2020) reported 12% fewer infections for 100% uptake. This study refers to nursing homes and long-term care facilities, and population uptake refers only to personnel, residents, and visitors. Meanwhile, the former studies used different metrics to assess the epidemic containment due to digital contact tracing (i.e., total infections, new infections, peak infections), and the results are not comparable. However, they provide evidence that a high uptake leads to the mitigation of the epidemic.

Another four studies reported a decrease in  $R_{\text{eff}}$  at various uptake levels (Bradshaw, Alley, Huggins, Lloyd, & Esvelt, 2021; Kretzschmar & Rozhnova, 2020; Kucharski et al., 2020; Plank et al., 2020). With an uptake value ranging from 20% to 90%, a reduction in  $R_{\text{eff}}$  was estimated between 17.6% and 47%. An additional set of seven papers also reported that digital contact tracing could reduce  $R_{\text{eff}}$  below one under various uptake values and are presented in the following section.

### 3.2. Minimum uptake to control the epidemic

A critical question from both a policy and a feasibility perspective is how many people should use a digital contact tracing app to control the epidemic. For this reason, we searched if a specific threshold value in uptake would be necessary to reduce  $R_{\text{eff}}$  to less than one, that is, to control the epidemic (excluding manual tracing, but including isolation of contacts and quarantine of infected people). One study called this as “digital herd immunity” to emphasize the transition to the smartphone era (Bulchandani, Shivam, Moudgalya, & Sondhi, 2020). Digital herd immunity is defined as a population that “can be immune to epidemics

even if not a single one of its members is immune to the disease.” According to this work, digital immunity is possible due to smartphone capabilities, regardless of the share of non-symptomatic transmission. To achieve digital herd immunity, the fraction of the population that needs to use a contact tracing app ranges between 75% and 95% for  $R_0 = 3$  (basic reproductive number). For smaller  $R_0$  values resulting from social distancing or other interventions, this fraction further decreases. Five additional studies also converged to a nearly 90% uptake needed to control the epidemic (Bradshaw et al., 2021; Kucharski et al., 2020; Nakamoto, Jiang, & Zhang, 2020; Pollmann, Pollmann, & Wiesinger, 2020; Xia & Lee, 2020). Finally, Hinch, Probert, and Nurta (2020) estimated that an epidemic could be suppressed if 80% of smartphone owners (corresponding to 56% of the overall population) use the app.

### 3.3. Relation between app uptake level and effectiveness

Four studies quantified the relation between the effectiveness of digital contact and mitigating COVID-19. Kim and Paul (2021) defined the effectiveness of digital contact tracing as the ratio of the number of individuals notified to the minimum number that should be notified to control the epidemic. Under specific assumptions, digital contact tracing effectiveness drops drastically, having a quadratic relation to the population uptake rate. For example, a 70% uptake will result in 49% effectiveness, while 50% uptake will result in a 25% effectiveness. Consequently, for digital contact tracing to succeed, most of the population should enroll. A similar finding was reported in another study, which identified that the reduction in the epidemic size grows quadratically with the app adoption uptake (Barrat et al., 2020). This may be attributed to the fact that the infected case and the contacts need to use the app. Based on this fact, Kucharski et al. (2020) found a quadratic relation between uptake and  $R_{\text{eff}}$  reduction. However, Hinch et al. (2020) reported a nearly linear relation between the cumulative deaths and app usage. This nearly linear dependence is explained by the combined effect of two non-linear effects in opposite directions: the quadratic proportion of contacts that use the app and the non-linear relation of epidemic size to  $R_0$  (Hinch et al., 2020).

**Table 1**  
Papers studied and outcomes of interest

Study	Data	Arch/re	Tech/ogy	Privacy risks	App Uptake	$R_{eff}$ / Infections		Key finding
						Digital	Manual	
Abueg et al. (2021)	USA	Not reported	Not reported	Not examined	75%	73-79% reduction in infections	30% more infections compared to digital	50% fewer infections for digital and manual combined
Almagor and Picascia (2020)	103,000 agents from 2011 UK Census	Not reported	Not reported	Not examined	80%	89% reduction in cases at the peak of the epidemic	Not reported	Digital contact tracing can contribute to reducing infection rates when accompanied by a sufficient testing capacity
Barrat et al. (2020)	Copenhagen Networks Study	Decentralized	BLE	Not examined	60%	36% reduction in epidemic size	60% reduction in epidemic size only with manual tracing	Digital and manual combined leads to an 80% reduction in the epidemic size
Bradshaw et al. (2021)	Hypothetical population	Decentralized	Not reported	Not examined	90%	$R_{eff}$ reduction close to 1	$R_{eff}$ reduction close to 1	Digital exposure notification alone is unlikely to control the epidemic
Bulchandani et al. (2020)	Hypothetical population	Not reported	Not reported	Not examined	75%-95%	$R_{eff} < 1$	Not reported	Digital immunity is possible with uptake of 75%-95%
Currie et al. (2020)	Australia COVIDSafe app	Centralised	BLE	Examined	61%	50% less infected	Not reported	COVIDSafe app an important tool adjunct to testing and social distancing
Ferretti et al. (2020)	40 source-recipient pairs	Decentralized	BLE	Not examined	High	$R_{eff} < 1$	$R_{eff}$ cannot get below 1 for a 3-day delay	A three-day delay assumed in manual tracing leads to an outbreak of control epidemic
Hinch et al. (2020)	1 million/UK	Decentralized	BLE	Not examined	56%	$R_{eff} < 1$	Not reported	High rates of app uptake lead to epidemic containment.
Kim and Paul (2021)	Hypothetical population	Not reported	Not reported	Not examined	High	Able to reduce infections if uptake is high	Not reported	Uptake rate has a quadratic relationship with digital contact tracing effectiveness
Kretzschmar and Rozhnova (2020)	Polymod study for the Netherlands	Not reported	Not reported	Not examined	20%	17.6% reduction in $R_{eff}$	2.5% reduction in $R_{eff}$	Digital more effective than manual tracing even with low uptake
Kucharski et al. (2020)	40,162 individuals/UK	Not reported	Not reported	Not examined	53%	47% reduction in $R_{eff}$	64% reduction in $R_{eff}$	66% reduction for manual and digital combined
López et al. (2021)	Demographic social-contact data/France	Not reported	Not reported	Not examined	60%	67% decrease at peak incidents	Not reported	For $R_0 > 2$ , digital contact tracing alone can not control the epidemic
Nakamoto et al. (2020)	Japan/ COCOA app	Decentralized	BLE	Examined	90%	$R_{eff} < 1$	Not reported	Data privacy first
Nuzzo et al. (2020)	Hypothetical population	Not reported	Not reported	Not examined	50%	90% decrease in peak number of infections	Not reported	Digital contact tracing successfully mitigates infection spread
Plank et al. (2020)	Hypothetical population	Centralized	BLE	Not examined	80%	$R_{eff}$ reduction from 2.4 to 1.46	Manual contact tracing alone reduction from 2.4 to 1.5	$R_{eff}$ reduced from 2.4 to 1.12 for digital and manual combined
b)	Hypothetical population	Decentralized	BLE	Not examined	80%	Not reported	Not reported	$R_{eff}$ reduced from 2.4 to 1.40 for digital and manual combined
c)	Hypothetical population	Not reported	QR	Not examined	80%	Not reported	Not reported	$R_{eff}$ reduced from 2.4 to 1.41 for digital and manual combined
Pollmann et al. (2020)	Hypothetical population	Not reported	BLE	Not examined	90%	$R_{eff} < 1$	Not reported	Random testing and social distancing necessary to push $R_{eff}$ below 1
Wilmink et al. (2020)	Hypothetical population in nursing homes	Centralized	Wearable device/ BLE Beacons	Not examined	100%	12% fewer infections compared to manual	Not reported	Digital contact tracing essential for nursing homes and long-term care facilities
Xia and Lee (2020)	Hypothetical population	Decentralized	Wearable BLE	Examined	>90%	$R_{eff} < 1$	Not reported	Uptake between 90%-95% to return to full normalcy
Yasaka et al. (2020)	Hypothetical population	Centralized and peer to peer	QR	Examined	25%	25% fewer infections compared to zero uptake	Not reported	Even a low adoption of 25% contributes to lower transmissions

**Note:** BLE = Bluetooth, QR = Quick Response,  $R_{eff}$  = effective reproductive number, APP = mobile application

### 3.4. Comparing digital and manual contact tracing

The studies in the previous section presented convincing evidence that digital contact tracing can reduce the epidemic spread at various

levels depending on the uptake rate and even control it, if the uptake surpasses the 90% threshold level. Nonetheless, this conclusion is based on the modeling assumption that no manual contact tracing is conducted concurrently. Hence, there is no direct comparison between the

effectiveness of digital-to-manual contact tracing. We identified eight studies that compared digital contact tracing to manual contact tracing (a detailed description of these studies is given in Supplementary Table S3). However, they draw contradictory outcomes on whether digital contact tracing is better than manual tracing.

Of the eight papers, four reported that digital contact tracing alone (i.e., without manual tracing) reduces significantly infections or  $R_{\text{eff}}$  more than manual tracing (Abueg et al., 2021; Ferretti et al., 2020; Kretzschmar & Rozhnova, 2020; Wilmink et al., 2020), and two reported only a marginal improvement of  $R_{\text{eff}}$  (Bradshaw et al., 2021; Plank et al., 2020). For example, without manual contact tracing, digital tracing reduces  $R_{\text{eff}}$  from 2.4 to 1.46, while manual contact tracing (in the absence of digital) reduces  $R_{\text{eff}}$  to 1.5 (Plank et al., 2020). On the other hand, two out of eight papers reported that manual contact tracing reduces  $R_{\text{eff}}$  or infections more than digital tracing alone (Barrat et al., 2020; Kucharski et al., 2020). For instance, Barrat et al. (2020) estimated a nearly double reduction in the epidemic size with manual contact tracing (60%) compared to digital (36%), while Kucharski et al. (2020) reported a 64% reduction in  $R_{\text{eff}}$  attributed only to manual tracing and 47% reduction if only digital tracing was applied. The above heterogeneous outcomes indicate that more evidence is needed to decide on the effectiveness of each strategy when applied alone.

Furthermore, we searched for studies that assessed the combined effect of manual and digital contact tracing on either  $R_{\text{eff}}$  or the number of infections. One study showed that both policies combined reduced new infections by 50% (Abueg et al., 2021). Another study showed that manual and digital tracing combined, at the same 60% coverage/uptake level, would result in a nearly 80% reduction in the epidemic size (Barrat et al., 2020). This is a significant improvement compared to a 60% reduction if only manual tracing is applied, or 36% if digital contact tracing is the only tracing policy. One study showed that, with manual and digital tracing combined (uptake of 80%),  $R_{\text{eff}}$  decreased from 2.4 to 1.12 (Plank et al., 2020). Bradshaw et al. (2021) showed that the hybrid use of manual contact tracing with bidirectional digital tracing, assuming 80% population uptake, led to an additional 0.26 lower  $R_{\text{eff}}$ . Finally, Kucharski et al. (2020) estimated only a slight improvement (from 64% to 66%  $R_{\text{eff}}$  reduction) of manual tracing once digital contact tracing was added.

### 3.5. System architecture, proximity awareness technology, and privacy

Ten out of the 19 papers reported information about the system architecture and the proximity/location awareness technology used. There are two main system architecture approaches in digital contact tracing: centralized and decentralized (Cohen et al., 2020). The centralized approach, reported in four papers in this review (Currie et al., 2020; Plank et al., 2020; Wilmink et al., 2020; Yasaka, Lehigh, & Sahyouni, 2020), is a top-down architecture where data are collected from the smartphone through an app and are stored at a central remote server. Therefore, the centralized approach has been criticized for potential privacy infringement (Cohen et al., 2020). Data are analyzed, and in the case of a confirmed case, all close contacts are notified to take specific actions (e.g., self-isolate). The decentralized approach is a bottom-up architecture and was reported in six papers (Barrat et al., 2020; Currie et al., 2020; Ferretti et al., 2020; Hinch et al., 2020; Nakamoto et al., 2020; Plank et al., 2020). The user has control over the data and decides whether they will be uploaded to a central server. Although this approach is less privacy-invasive, it struggles to ensure that sufficient data will be collected for successful contact tracing (Gasser et al., 2020).

In both architectures, proximity and location awareness technologies offer a set of tools to survey an individual's movement. Global Navigation Satellite Systems (GNSS), is a location awareness technology as it provides the geographic location of the user. WiFi, Bluetooth Low Energy (BLE), Beacons, or Quick Response (QR) codes collect proximity data. In other words, proximity-based approaches directly detect nearby

smartphones (or other devices) and not the exact geographic location (Abueg et al., 2021). However, under specific system architectural settings and calculations, proximity awareness technologies can also provide the location of an individual (see Table 2). Data are then analyzed in anticipation of rapid tracing of secondary virus cases, thus enforcing specific health protocols to mitigate the spread of the virus (see Supplementary Table S4 for a description of proximity awareness technologies).

This review found that BLE is the most widely used technology matched to a decentralized architecture (Barrat et al., 2020; Currie et al., 2020; Ferretti et al., 2020; Hinch et al., 2020; Nakamoto et al., 2020; Plank et al., 2020; Pollmann et al., 2020; Xia & Lee, 2020). Wilmink et al. (2020) coupled beacons with wearables. Following a centralized architecture, real-time location data are uploaded to cloud-based software to visualize egocentric contact networks. This system is suitable for indoor environments, such as nursing homes. The results show that this technology could reduce infections by more than 12% compared to manual tracing. Another work emphasizes privacy issues, and therefore the proposed app does not use personal or location data (Yasaka et al., 2020). Users should create "checkpoints" by scanning QR codes whenever they meet other people, either in public or private. In the case of infection, users should anonymously self-report their health status to the network.

It is worth mentioning that only one paper modelled different combinations of alternative digital contact tracing system architectures with varying proximity awareness technologies (Plank et al., 2020). A centralized BLE approach was compared to a decentralized BLE approach, and a QR code exposure notification system. Only the effect of the combined use of manual and digital contact tracing at an 80% uptake was assessed. The centralized BLE approach reduced  $R_{\text{eff}}$  to 1.1, nearly succeeding in containing the epidemic. The other two approaches (i.e., decentralized BLE approach, and a QR code exposure notification system) reduced  $R_{\text{eff}}$  to 1.4, a significant difference when compared to the centralized BLE approach.

Privacy considerations were examined in four studies (Currie et al., 2020; Nakamoto et al., 2020; Xia & Lee, 2020; Yasaka et al., 2020). For the COVIDSafe app, the Australian government used a centralized architecture. However, it maximized data security with legislation restricting data transfer, storage, use, and disposal (Currie et al., 2020). Only one study prioritized the privacy protection of users (Nakamoto et al., 2020). In this study, a peer-to-peer framework was developed for the COCOA app to provide individuals with reliable updates of COVID-19 without exposing their private data. Xia and Lee (2020) proposed a stand-alone device, called "contact recorder", specifically designed for contact tracing. A contact recorder, equipped with BLE technology, would be easier to preserve data privacy as it would store only data related to proximity and health status, minimizing thus privacy infringement. Lastly, Yasaka et al. (2020) used an anonymized graph of interpersonal interactions through a specifically designed privacy-preserving smartphone app that implements peer-to-peer contact tracing through the use of QR codes.

### 3.6. Quality assessment

The quality assessment of the 19 papers of this review varied considerably and is presented in Supplementary Table S2. All studies clearly defined their objectives, summarized the key findings, and discussed their limitations. Most of the studies sufficiently presented the applied models, and the related assumptions and parameters. However, some studies reported briefly or not at all the choice of the model (Nakamoto et al., 2020; Nuzzo et al., 2020), the model assumptions (Currie et al., 2020; Nakamoto et al., 2020; Nuzzo et al., 2020), the study parameters (Barrat et al., 2020; Currie et al., 2020; Kim & Paul, 2021; Nakamoto et al., 2020; Nuzzo et al., 2020; Yasaka et al., 2020). Some studies did not describe the characteristics of the base-case population (Bradshaw et al., 2021; Bulchandani et al., 2020; Currie et al., 2020;

**Table 2**  
Proximity/Location awareness technologies for COVID-19 digital contact tracing

Technology	Location/ Proximity Accuracy		COVID-19 tracing		Privacy Concerns
	Outdoors	Indoors	Suitable for	Unsuitable for	
GNSS	10 m GPS only, 5 m GPS + WiFi	Most likely not operating	Outdoors / Tracking overlapping routes / Detection of hotspots	Indoors	High
BLE	<2 m	<2 m	Tracing individuals within 2 meters	Spaces with airborne transmission of SARS-CoV-2	Low
Beacons	Building level	Room/floor level	Same room/ floor/building	Assessing the distance between individuals	Low
QR	Building level	Room/floor level	Same room/ floor/building	Assessing the distance between individuals	Moderate to high
WiFi	Depending on Access Points	<1m	Indoors	Outdoors	Low
UWB	Depending on UWB transmitters	<0.5 m	Indoors	Currently, few smartphones have this technology	Low

Ferretti et al., 2020; Kim & Paul, 2021; Kretzschmar & Rozhnova, 2020; Nakamoto et al., 2020; Nuzzo et al., 2020; Plank et al., 2020; Pollmann et al., 2020; Wilmlink et al., 2020; Xia & Lee, 2020; Yasaka et al., 2020), while other studies described in a limited way the analytical methods that supported the evaluation of the results (Currie et al., 2020; Kim & Paul, 2021; Kretzschmar & Rozhnova, 2020; Nuzzo et al., 2020; Plank et al., 2020; Wilmlink et al., 2020; Yasaka et al., 2020).

#### 4. Discussion

Here we discuss the lessons learned from the 19 papers presented in the previous section. Identified gaps and related suggestions are highlighted to guide future contact tracing research and implementation.

##### 4.1. Lesson 1. Digital contact tracing can mitigate COVID-19 if population uptake is substantial

All studies (19/19) in this review show that digital contact tracing reduces the  $R_{eff}$  and the number of infected cases. However, its effectiveness exhibits a quadratic dependence from the uptake rate (Barrat et al., 2020; Kim & Paul, 2021). This implies that a considerable value in app adoption is needed for digital contact tracing to be effective on its own. Most studies of this review converged to the same uptake rate of at least 90% (in the absence of manual contact tracing) to control the epidemic (drop  $R_{eff}$  below one). This finding is in agreement with other works that consider a broad uptake in the population as the key for digital contact tracing success (Munzert, Selb, Gohdes, Stoetzer, & Lowe, 2021).

In practice, it is tough to achieve such a high level of community uptake (Watts, 2020). However, a low adoption rate may still have a positive epidemiological impact. As shown in this review, even with low adoption rates the  $R_{eff}$  and the number of infections decrease. Other empirical studies have confirmed this finding. For example, a study in Spain showed that a 33% app uptake, could significantly reduce infections (Rodríguez et al., 2021). The drawback is that individuals may gain a false sense of security at a low uptake level if they do not receive exposure notifications, which may encourage a more relaxed behaviour (i.e., not following social distancing measures) (Kim & Paul, 2021). However, the notifications' absence could be attributed to the low uptake, not that individuals with COVID-19 have not crossed their pathways. For this reason, statistics should be summarized at small geographical units, creating GIS COVID-19 risk assessment maps that would be readily available to policymakers and users (Grekousis, 2020). This, underscores that accurate and updated content through the app and the subsequent spatial analysis of the identified patterns is crucial for people to align their behaviour (i.e., adhere to social distancing) with the actual epidemic severity.

The benefits of digital contact tracing can be applied to sub-populations even if the uptake rate is low at the general population. For

example, digital contact tracing apps would have greater epidemiological importance for older age groups. Yet, older people are less inclined to technology due to the phenomenon of technological exclusion (Watts, 2020). As such, they are less likely to download or systematically use a digital contact tracing app through their smartphone, something that has been already observed in many parts of the world, as for example, the UK and Singapore (Huang, Guo, Lim, & Chow, 2021; Watts, 2020). In this case, wearable devices could be used instead of smartphones to make digital contact tracing approachable to older persons. In nursing homes or long-term care facilities, with a COVID-19 mortality rate of nearly 25% among the elders, the need for specifically designed digital contact tracing systems is even higher (Wilmlink et al., 2020). We argue that in closed structures with controlled access and high surveillance, such as nursery homes, or health care facilities digital contact tracing based on high location accuracy technologies (i.e., beacons, UWB) integrated with wearable devices should be prioritized (Fig. 2C).


##### 4.2. Lesson 2. Digital contact tracing should be used in combination with manual contact tracing

This review examines another critical question: Is digital contact tracing more effective than manual contact tracing if used as the single tracing strategy? There is no consensus among this review's studies on whether digital is more effective than manual contact tracing. Of the eight papers that compared manual to digital contact tracing, four reported that digital is better than manual, two reported only marginal gains of digital tracing, and two papers that manual is more effective than digital contact tracing. These contradictory conclusions suggest that more research is needed to establish the superiority, if any, of one tracing strategy over the other (Abueg et al., 2021; Ferretti et al., 2020). Given that digital contact tracing achieves a satisfactory reduction in  $R_{eff}$  only at high uptake rates, we argue that excluding manual tracing is not rational for the time being. Other empirical studies also support this conclusion. For example, Lai, Tang, Kurup, and Thevendran (2021) showed that although various digital contact tracing apps (i.e., Safe-Entry and TraceTogether) helped in contact tracing in Singapore, they were not sufficient to replace manual contact tracing conducted by experienced personnel. Similar to others, we suggest the combined use of these strategies to maximize the benefits (Barrat et al., 2020; Kucharski et al., 2020). This is also highlighted in four studies of this review that assessed that the joint effect of these strategies could reduce infections by 50% and reduce  $R_{eff}$  down to 1.1, very close to the critical value below which the epidemic enters a controlled phase.

##### 4.3. Lesson 3. Proximity/location accuracy of digital contact tracing app varies across different built environments and facilities

Proximity/location awareness technologies are not suitable for every space or facility, something rarely mentioned in the modeling studies of





**Fig. 2.** Proximity/location awareness technologies for digital contact tracing in various environments. **A)** GNSS (belongs to location awareness technologies) tracks the location in outdoor environments (i.e. park). BLE can be used for proximity sensing but not for identifying the exact location. **B)** Beacons, WiFi, or QR code can be used on subway, buses, or other modes of transport. **C)** Wearables, coupled with beacons, are efficient for nursing homes and other closed structures. **D)** BLE may provide a false exposure notification for individuals separated by a wall. GNSS is hard to operate indoors. **E)** UWB can identify crossing pathways in indoor environments, while beacons and QR codes can provide proximity accuracy at the room level. **F)** Depending on smartphone's placement and orientation, BLE can identify as close contacts those sitting back-to-back and miss those at the same table. For indoor airborne transmission of the virus, WiFi, beacons, and QR are sufficient.

this review. Almost half (9/19) of the studies did not report any technical characteristics, a severe shortcoming to their modeling outcomes. Moreover, no study reported how the proximity/location accuracy may affect the effectiveness of digital contact tracing, a significant literature gap. However, the digital contact tracing efficiency is directly linked to how the proximity is determined and how data are retrieved, processed, and stored (Silva et al., 2021). On this account, more focus should be placed on the joint analysis of the technological and epidemiological components of a digital contact tracing system. Here, we extend the discussion to describe how the proximity/location accuracy of proximity/location awareness technologies varies on different environments and facilities to guide the development of future digital contact tracing systems (Fig. 2 and Table 2).

BLE is the preferred proximity awareness technology applied in 80% (8/10) of the studies that reported information about the system architecture and the proximity awareness technology used. Nevertheless, it comes with accuracy problems when applied in different environments (Table 2) (Nguyen et al., 2020). BLE cannot identify physical barriers between two individuals (i.e., separated by a wall or anti-droplet plexiglass) (Fig. 2D). If one of these individuals is infected, then the second one is likely to be notified and isolated, even when he or

she should not. BLE signal strength may vary not only with smartphone orientation but also with its placement (i.e., in the pocket or the hand if its owner) (Zastrow, 2020). Consequently, people sitting at the same table in a restaurant may seem to have less contact than those sitting back to back on a nearby table (Fig. 2F).

Digital contact tracing based on BLE traces contacts within 2 m (Table 2) (Rosenkrantz et al., 2020). However, SARS-CoV-2 also spreads through aerosols' airborne transmission up to tens of meters (Morawska & Cao, 2020). Hence, contacts within the same room or space cannot be notified if the app is based only on BLE (Xia & Lee, 2020). A centralized architecture could probably handle tracing under such conditions and would easily learn from the collected data. In this manner, the system could modify the criteria to label a contact as exposed, reflecting the new findings of a novel virus and making alerts more precise.

Another approach could combine BLE with Beacons, WiFi access point, or UWB to offer higher proximity accuracy in indoor environments (Fig. 2E,F) (Wilmink et al., 2020). In addition, the use of specifically designed devices, instead of smartphones, may increase the usability and the accuracy of contact tracing. Studies have shown that smart watches with digital contact tracing apps increase the chances that the two devices are in line of sight, while a smartphone usually remains

in a pocket or bag (Maccari & Cagno, 2021).

Finally, the GNSS of smartphone malfunctions in indoor environments with a location accuracy of approximately 10 m outdoors (Nguyen et al., 2020). Therefore, GNSS can assist in tracking rather than doing proximity evaluation in the digital contact tracing context (Fig. 2A).

In conclusion, the system architecture and proximity awareness technology are critical and should be considered in the modeling process to assess a digital contact tracing system's effectiveness.

#### 4.4. Lesson 4. The decentralized architecture offers more privacy but it is not as efficient as the centralized architecture

The centralized app-based BLE architecture outperformed the decentralized app-based BLE system and the QR-notification system (Plank et al., 2020). Probably due to governmental advice/order (Fagherazzi et al., 2020), the app adoption rate is generally higher in the centralized architecture than in the decentralized one, but at a privacy cost (Nakamoto et al., 2020). On the other hand, a decentralized approach with voluntary app usage suffers from lower adoption rates potential due to fears of personal data privacy infringement (Ferretti et al., 2020). More similar studies are needed to help policymakers better decide on the most effective system architecture.

#### 4.5. Lesson 5. Personal data privacy is the key for digital contact tracing success

With digital contact tracing critically depending on wide adoption for success, as shown in Lesson 1, governments opting for such a strategy should build a legal and technical framework, ensuring personal data privacy. The MIT Covid Tracing Tracker project reported that by July 30, 2020, 20 out of 31 countries that had officially adopted/supported the use of apps for digital contact tracing had an uptake of less than 8% (O'Neill, Ryan-Mosley, & Johnson, 2020). Surprisingly, an online survey of 5995 participants across five countries (i.e., France, Germany, Italy, the UK, and the US) found a high willingness to download a digital contact tracing app, with an overall acceptance rate of 74.8% (Altmann, Milsom, & Zillesen, 2020). The large gap between the intention to use such an app and the actual uptake is attributed to severe concerns about a) how governments will handle data after the end of the pandemic (42%), and b) cybersecurity (35%) (Altmann et al., 2020).

Balancing the need for epidemiological information with legitimate data privacy is critical (Altmann et al., 2020; Park, Choi, & Ko, 2020). As many concerns are raised regarding technology intervention for an individual's privacy, a trade-off between protecting public health and retaining personal data control should be found. Most publications in this review emphasized the importance of data privacy and suggested a decentralized architecture (Barrat et al., 2020; Ferretti et al., 2020; Hinch et al., 2020; Nakamoto et al., 2020; Xia & Lee, 2020). This architecture offers more tools to preserve anonymity and overall data privacy than the centralized approach (Rosenkrantz et al., 2020). However, studies have shown that, even with anonymized data, there are techniques to identify the actual individuals (Rocher, Hendrickx, & de Montjoye, 2019). Consequently, privacy standards should be the highest possible, while data should be kept only for a limited time and automatically deleted (Maccari & Cagno, 2021). Due to ethical and cybersecurity concerns, oversight from an inclusive advisory board is suggested (Ferretti et al., 2020). Studies show that the usefulness of digital tracing is leveraged only in combination with manual tracing, and other interventions such as testing, isolation, and social distancing (Rahmani and Mirmahaleh, 2020). On this account, a privacy infringement is not justified, as digital tracing can be observed only as supplementary to controlling an epidemic. However, as digital contact tracing is a strategy that can be highly effective under specific settings (i.e., closed structures), it involves technological, legal, and ethical issues that should be on top of their architectural designs. Key lessons, research

gaps, and suggestions are summarized in Table 3.

This review has certain limitations. Due to the limited number of available papers that report comparable numerical outputs in the effective reproductive number  $R_{eff}$ , or the reduction in infections, no meta-analysis has been conducted although initially planned. No empirical studies are included as none exists. Finally, we limited our search to English literature only.

## 5. Conclusions

Learning lessons from the current pandemic will be an essential step towards a stronger and more resilient society. This paper contributes towards informing policymakers on some lessons learned regarding digital contact tracing, community uptake, and proximity awareness technology to fight COVID-19 on the following topics/debates:

- the effectiveness (and thus its potential necessity as a policy against COVID-19) of digital contact tracing in reducing  $R_{eff}$ ,
- the necessary adoption rate (if any) by the public to ensure that digital contact tracing can assist in controlling the pandemic
- the superiority (if any) of the digital contact tracing over the manual contact tracing
- the proximity/location accuracy and privacy concerns raised with digital contact tracing

Policy-wise, the take-home message of this review is that, the success of digital contact tracing depends on a complex interplay of app uptake in the community, proximity awareness technologies, and public's trust (see also Table 3). If governments consider the barriers that keep away people from adopting tracing apps, and scientists address technical limitations, digital contact tracing could be a successful strategy (Ferretti et al., 2020; Nguyen et al., 2020). It would offer a powerful toolkit for decision makers and the public that could be an essential part of long-term response not only to COVID-19 but also to future epidemics. Even when the COVID-19 pandemic is over, other epidemics will strike (Xia & Lee, 2020). In the wake of a novel pathogen, effective treatments and vaccines will lag behind the virus spread, and other means should be readily available to contain the new disease. Although digital contact

**Table 3**  
Key lessons learned from this review along with suggestions

	Lessons learned	Suggestions/research gaps
Effectiveness	Digital contact tracing can control COVID-19 if the population uptake surpasses 90%.	As a 90% uptake is difficult to be achieved, digital contact tracing should be combined to manual contact tracing.
Digital vs. manual	There is no clear evidence that digital contact tracing can substitute manual.	Further research is needed with empirical data.
Proximity/location accuracy	Proximity/location accuracy highly varies on the technology used and the indoor or outdoor setting	To avoid false alerts or exposure notifications, the choice of proximity awareness technology should be central when designing a digital contact tracing system.
Proximity awareness technology	BLE is preferred.	As proximity accuracy is low with BLE, alternative technologies such as UWB should be promoted.
Architecture	Decentralized architecture allows for higher personal data privacy	Architecture should ensure the highest data privacy standards.
Privacy	Most studies raise privacy and ethical concerns related to personal data.	The need for epidemiological information should not lead to personal data privacy infringement. Governments should build a legal framework ensuring personal data privacy to gain people's trust.

tracing does not guarantee epidemic control, it provides technologies that respond quickly, have high location and proximity accuracy, and preserve anonymity.

Digital contact tracing should not only be linked to smartphones as their penetration varies significantly across countries and age groups, with a remarkably low usage among the elderly, the most susceptible to COVID-19 (López et al., 2021). Wearables or other stand-alone devices would ensure high uptake from elders and groups in need (Wilmink et al., 2020; Xia & Lee, 2020). For example, over a billion of the world's population (15% of the total) lives with some form of disability (World Health Organization, 2018). This number is expected to double by 2050, and the WHO calls for the development of assistive technologies disability (World Health Organization, 2018). For this reason, future innovation should focus on creating stand-alone, smaller, and cheaper wearable devices with low energy consumption. In the case of new outbreaks, wearable devices such as watches or pins could be freely distributed among populations, especially in low-income countries. Along with mass testing, isolation, social distancing, and personal hygiene, a new virus is likely to be contained quickly, minimizing loss of lives, economic and societal costs that threaten sustainability, and allowing for faster and safer return to normality.

## Funding

This work was supported the National Natural Science Foundation of China, grant number [41871140]. The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

## Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.scs.2021.102995>.

## References

- Abueg, M., Hinch, R., Wu, N., Liu, L., Probert, W., Wu, A., et al. (2021). Modeling the effect of exposure notification and non-pharmaceutical interventions on COVID-19 transmission in Washington state. *npj Digital Medicine*, 4(1), 1–10.
- Aleta, A., Martín-Corral, D., Piontti, A. P., et al. (2020). Modelling the impact of testing, contact tracing and household quarantine on second waves of COVID-19. *Nature Human Behaviour*, 4(9), 964–971.
- Almagor, J., & Picascia, S. (2020). Exploring the effectiveness of a COVID-19 contact tracing app using an agent-based model. *Scientific reports*, 10(1), Dec 17, 1–1.
- Altmann, S., Milsom, L., Zillesen, H., et al. (2020). Acceptability of App-Based Contact Tracing for COVID-19: Cross-Country Survey Study. *JMIR mHealth and uHealth*, 8(8), Article e19857.
- Anglemyer, A., Moore, T., Parker, L., et al. (2020). Digital contact tracing technologies in epidemics: a rapid review. *Cochrane Database of Systematic Reviews*, 8. <https://doi.org/10.1002/14651858.CD013699>
- Barrat, A., Cattuto, C., Kivela, M., Lehmann, S., & Saramaki, J. (2020). Effect of manual and digital contact tracing on COVID-19 outbreaks: a study on empirical contact data. *medRxiv*. Published on line July 25. <https://www.medrxiv.org/content/10.1101/2020.07.24.20159947v1> (preprint).
- Batty, M. (2020). The Coronavirus crisis: What will the post-pandemic city look like? *Environment and Planning B: Urban Analytics and City Science*, 47(4), 547–552. <https://doi.org/10.1177/2399808320926912>
- Booth, A., Mitchell, A. S., Mott, A., James, S., Cockayne, S., Gascoyne, S., et al. (2020). An assessment of the extent to which the contents of PROSPERO records meet the systematic review protocol reporting items in PRISMA-P. *F1000Research*, 9.
- Bradshaw, W. J., Alley, E. C., Huggins, J. H., Lloyd, A. L., & Esvelt, K. M. (2021). Bidirectional contact tracing could dramatically improve COVID-19 control. *Nature communications*, 12(1), 1–9.
- Braithwaite, I., Callender, T., Bullock, M., & Aldridge, R. W. (2020). Automated and partly automated contact tracing: a systematic review to inform the control of COVID-19. *The Lancet Digital Health*, 2(11), e607–e621.

- Bulchandani, V. B., Shivam, S., Moudgalya, S., & Sondhi, S. (2020). Digital herd immunity and COVID-19. *ArXiv*. published online April 15. <https://arxiv.org/abs/2004.07237> (preprint).
- Calvo, R. A., Deterding, S., & Ryan, R. M. (2020). Health surveillance during covid-19 pandemic. *BMJ*, 369. <https://doi.org/10.1136/bmj.m1373>
- Cohen, I. G., Gostin, L. O., & Weitzner, D. J. (2020). Digital Smartphone Tracking for COVID-19: Public Health and Civil Liberties in Tension. *JAMA*, 323(23), 2371–2372.
- Currie, D. J., Peng, C. Q., Jameson, B. A., Frommer, M. S., & Lyle, D. M. (2020). Stemming the flow: How much can the Australian smartphone app help to control COVID-19? *Public Health Research and Practice*, 30(2), Article e3022009.
- DeLuca, Papageorge, N., Mitchel, B., & Kalish, E. (2020). *The unequal cost of social distancing* Accessed in March 25, 2021, from <https://coronavirus.jhu.edu/from-our-experts/the-unequal-cost-of-social-distancing>.
- Elavarasam, R. M., Pugazhendhi, R., Shafullah, G. M., Irfan, M., & Anvari-Moghaddam, A. (2021). A hover view over effectual approaches on pandemic management for sustainable cities—The endowment of prospective technologies with revitalization strategies. *Sustainable Cities and Society*, 68, Article 102789.
- Fagherazzi, G., Goetzinger, C., Rashid, M. A., Aguayo, G. A., & Huiart, L. (2020). Digital health strategies to fight COVID-19 worldwide: challenges, recommendations, and a call for papers. *Journal of Medical Internet Research*, 22(6), Article e19284.
- Ferretti, L., Wymant, C., Kendall, M., et al. (2020). Quantifying SARS-CoV-2 transmission suggests epidemic control with digital contact tracing. *Science*, 368, Article eabb6936.
- Flaxman, S., Mishra, S., Gandy, A., Unwin, H. J. T., Mellan, T. A., Coupland, H., et al. (2020). Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe. *Nature*, 584(7820), 257–261.
- Fokas, A. S., Cuevas-Maraver, J., & Kevrekidis, P. G. (2020). A quantitative framework for exploring exit strategies from the COVID-19 lockdown. *Chaos, solitons, and fractals*, 140, Article 110244. <https://doi.org/10.1016/j.chaos.2020.110244>
- Gasser, U., Ienca, M., Scheibner, J., Sleight, J., & Vayena, E. (2020). Digital tools against COVID-19: taxonomy, ethical challenges, and navigation aid. *The Lancet Digital Health*, 2(8), e425–e434.
- Ge, X. Y., Pu, Y., Liao, C. H., Huang, W. F., Zeng, Q., Zhou, H., et al. (2020). Evaluation of the exposure risk of SARS-CoV-2 in different hospital environment. *Sustainable cities and society*, 61, Article 102413.
- Grekousis, G. (2020). *Spatial analysis methods and practice: Describe, Explore Explain, through GIS*. Cambridge, UK: Cambridge University Press.
- Hellewell, J., Abbott, S., Gimma, A., et al. (2020). Feasibility of controlling COVID-19 outbreaks by isolation of cases and contacts. *Lancet Glob Health*, 8, e488–96.
- Hinch, R., Probert, W., Nurtay, A., et al. (2020). *Effective configurations of a digital contact tracing app: a report to NHSX* (Accessed April 15, 2020) [https://github.com/BDI-pathogens/covid-19\\_instant\\_tracing](https://github.com/BDI-pathogens/covid-19_instant_tracing).
- Huang, Z., Guo, H., Lim, H. Y., & Chow, A. (2021). Awareness, acceptance, and adoption of the national digital contact tracing tool post COVID-19 lockdown among visitors to a public hospital in Singapore. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases*. <https://doi.org/10.1016/j.cmi.2021.01.007>. S1198-743X(21)00033-1. Advance online publication.
- Husereau, D., Drummond, M., Petrou, S., Carswell, C., Moher, D., Greenberg, D., et al. (2020). *CHEERS Task Force. Consolidated health economic evaluation reporting standards (CHEERS)*.
- International Monetary Fund. (2020). *A year like no other. IMF Annual report 2020* (Accessed at March 29, 2021) from <https://www.imf.org/external/pubs/ft/ar/2020/eng/downloads/imf-annual-report-2020.pdf>.
- Jadidi, M., Jamshidiha, S., Masroori, I., Moslemi, Mohammadi, A., & Pourahmadi, V. (2021). A two-step vaccination technique to limit COVID-19 spread using mobile data. *Sustainable Cities and Society*, 70, Article 102886. <https://doi.org/10.1016/j.scs.2021.102886>, 2021, ISSN 2210-6707.
- Kim, H., & Paul, A. (2021). Automated contact tracing: a game of big numbers in the time of COVID-19. *Journal of the Royal Society Interface*, 18(175), Article 20200954.
- Kretzschmar, M. E., & Rozhnova, G. (2020). Bootsma MCJ, van Boven M, van de Wijgert JHJM, Bonten MJM. Impact of delays on effectiveness of contact tracing strategies for COVID-19: a modelling study. *The Lancet Public Health*, 5(8), e452–e459.
- Kucharski, A. J., Klepac, P., Edmunds, W. J., et al. (2020). Effectiveness of isolation, testing, contact tracing, and physical distancing on reducing transmission of SARS-CoV-2 in different settings: a mathematical modelling study. *The Lancet Infectious Diseases*, 20(10), 1151–1160.
- Lai, S. H. S., Tang, C. Q. Y., Kurup, A., & Thevendran, G. (2021). The experience of contact tracing in Singapore in the control of COVID-19: highlighting the use of digital technology. *International orthopaedics*, 45(1), 65–69.
- López, J. A. M., García, B. A., Bentkowski, P., Bioglio, L., Pinotti, F., Boëlle, P. Y., et al. (2021). Anatomy of digital contact tracing: role of age, transmission setting, adoption and case detection. *Science advances*, Article eabd8750.
- Maccari, L., & Cagno, V. (2021). Do we need a contact tracing app? *Computer Communications*, 166, 9–18.
- Maiti, A., Zhang, Q., Sannigrahi, S., Pramanik, S., Chakraborti, S., Cerda, A., et al. (2021). Exploring spatiotemporal effects of the driving factors on COVID-19 incidences in the contiguous United States. *Sustainable cities and society*, 68, Article 102784.
- Mizumoto, K., Kagaya, K., Zarebski, A., & Chowell, G. (2020). Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Eurosurveillance*, 25(10), Article 2000180.
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*, 6, Article e1000097.

- Morawska, L., & Cao, J. (2020). Airborne transmission of SARS-CoV-2: The world should face the reality. *Environment International*, 139, Article 105730.
- Munzert, S., Selb, P., Gohdes, A., Stoetzer, L. F., & Lowe, W. (2021). Tracking and promoting the usage of a COVID-19 contact tracing app. *Nature Human Behaviour*, 5(2), 247–255.
- Nakamoto, I., Jiang, M., Zhang, J., et al. (2020). A Peer-To-Peer COVID-19 Contact Tracing Mobile App in Japan: Design and Implementation Evaluation. *JMIR mHealth and uHealth*. <https://doi.org/10.2196/22098>
- Nguyen, C. T., Saputra, Y. M., Van Huynh, N., Nguyen, N. T., Khoa, T. V., Tuan, B. M., et al. (2020). Enabling and emerging technologies for social distancing: A comprehensive survey. *arXiv preprint arXiv:2005.02816*.
- Nuzzo, A., Tan, C. O., Raskar, R., DeSimone, D. C., Kapa, S., & Gupta, R. (2020). Universal Shelter-in-Place Versus Advanced Automated Contact Tracing and Targeted Isolation: A Case for 21st-Century Technologies for SARS-CoV-2 and Future Pandemics. *Mayo Clinic Proceedings*, 95(9), 1898–1905.
- O'Neill, P. H., Ryan-Mosley, T., & Johnson, B. (2020). A flood of coronavirus apps are tracking us. Now it's time to keep track of them. *MIT Technology Review*, May 7 2020 (Accessed November 13, 2020) <https://www.technologyreview.com/2020/05/07/1000961/launching-mittr-covid-tracing-tracker/>.
- Park, S., Choi, G. J., & Ko, H. (2020). Information Technology-Based Tracing Strategy in Response to COVID-19 in South Korea - Privacy Controversies. *JAMA*, 323(21), 2129–2130.
- Peak Corey, M., Childs Lauren, M., Grad Yonatan, H., & Buckee Caroline, O. (2017). Comparing nonpharmaceutical interventions for containing emerging epidemics. *Proceedings of the National Academy of Sciences*, 114(15), 4023–4028.
- Plank, M., James, A., Lustig, A., Steyn, N., Binny, R., & Hendy, S. (2020). Potential reduction in transmission of COVID-19 by digital contact tracing systems. *medRxiv*. Published online September 1. <https://www.medrxiv.org/content/10.1101/2020.08.27.20068346v1> (preprint).
- Pollmann, T., Pollmann, J., Wiesinger, C., et al. (2020). The impact of digital contact tracing on the SARS-CoV-2 pandemic - a comprehensive modelling study. *medRxiv*. Published online September 14. <https://www.medrxiv.org/content/10.1101/2020.09.13.20192682v1> (preprint).
- Rahmani, A. M., & Mirmahaleh, S. Y. H. (2020). Coronavirus disease (COVID-19) prevention and treatment methods and effective parameters: A systematic literature review. *Sustainable cities and society*, Article 102568.
- Rocher, L., Hendrickx, J. M., & de Montjoye, Y. A. (2019). Estimating the success of re-identifications in incomplete datasets using generative models. *Nature Communications*, 10(1), 1–9.
- Rodríguez, P., Graña, S., Alvarez-León, E. E., Battagliani, M., Darias, F. J., Hernán, M. A., et al. (2021). A population-based controlled experiment assessing the epidemiological impact of digital contact tracing. *Nature communications*, 12(1), 1–6.
- Rosenkrantz, L., Schuurman, N., Bell, N., & Amram, O. (2020). The need for GIScience in mapping COVID-19. *Health and Place*. <https://doi.org/10.1016/j.healthplace.2020.102389>
- Salehinejad, S., Niakan Kalhori, S. R., Hajesmael Gohari, S., Bahaadinbeigy, K., & Fatehi, F. (2021). A review and content analysis of national apps for COVID-19 management using Mobile Application Rating Scale (MARS). *Informatics for Health and Social Care*, 46(1), 42–55. <https://doi.org/10.1080/17538157.2020.1837838>
- Silva, J. C. S., de Lima Silva, D. F., Neto, A. D. S. D., Ferraz, A., Melo, J. L., Júnior, N. R. F., et al. (2021). A city cluster risk-based approach for Sars-CoV-2 and isolation barriers based on anonymized mobile phone users' location data. *Sustainable cities and society*, 65, Article 102574.
- Skoll, D., Miller, J. C., & Saxon, L. A. (2020). COVID-19 testing and infection surveillance: Is a combined digital contact-tracing and mass-testing solution feasible in the United States? *Cardiovascular Digital Health Journal*. <https://doi.org/10.1016/j.cvdhj.2020.09.004>
- United Nations Development Programme. (2015). *What are the Sustainable Development Goals* Accessed January 26, 2020 from <https://www.undp.org/content/undp/en/home/sustainable-development-goals.html/>.
- United Nations. (2020). *Liquidity and Debt Solutions to Invest in the SDGs: The Time to Act is Now* Accessed March 29, 2021 from [https://www.un.org/sites/un2.un.org/files/sg\\_policy\\_brief\\_on\\_liquidity\\_and\\_debt\\_solutions\\_march\\_2021.pdf](https://www.un.org/sites/un2.un.org/files/sg_policy_brief_on_liquidity_and_debt_solutions_march_2021.pdf).
- Velraj, R., & Haghghat, F. (2020). The contribution of dry indoor built environment on the spread of Coronavirus: Data from various Indian states. *Sustainable cities and society*, 62, Article 102371.
- Watts, G. (2020). COVID-19 and the digital divide in the UK. *The Lancet. Digital health*, 2(8), e395–e396. [https://doi.org/10.1016/S2589-7500\(20\)30169-2](https://doi.org/10.1016/S2589-7500(20)30169-2)
- Wilmink, G., Summer, I., Marsyla, D., et al. (2020). Real-Time Digital Contact Tracing: Development of a System to Control COVID-19 Outbreaks in Nursing Homes and Long-Term Care Facilities. *JMIR public health and surveillance*, 6(3), Article e20828.
- World Health Organization. (2018). *Disability and health*. Geneva: World Health Organization (Accessed November 13, 2020) <https://www.who.int/news-room/fact-sheets/detail/disability-and-health>.
- World Health Organization. (2020). *COVID-19 strategy update—14 april 2020*. Geneva: World Health Organization (Accessed March 20, 2021) <https://www.who.int/publications/i/item/covid-19-strategy-update—14-april-2020>.
- Xia, Y., & Lee, G. (2020). How to return to normalcy: fast and comprehensive contact tracing of COVID-19 through proximity sensing using mobile devices. *ArXiv*. published online April 27. <https://arxiv.org/abs/2004.12576> (preprint).
- Yasaka, T. M., Lehrich, B. M., & Sahyouni, R. (2020). Peer-to-peer contact tracing: development of a privacy-preserving smartphone app. *JMIR mHealth and uHealth*, 8(4), Article e18936.
- Zastrow, M. (2020). Coronavirus contact-tracing apps: can they slow the spread of COVID-19? *Nature*. <https://doi.org/10.1038/d41586-020-01514-2>
- Zhou, C., Su, F., Pei, T., Zhang, A., Du, Y., Luo, B., et al. (2020). COVID-19: challenges to GIS with big data. *Geography and sustainability*, 1(1), 77–87.