

REVIEW

Brucellosis: Prevalence with reference to East African community countries – A rapid review

Juvenal Djangwani^{1,2}  | George Ooko Abong¹  | Lucy Gicuku Njue¹ | Dasel W. M. Kaindi¹

¹College of Agriculture and Veterinary Sciences, Department of Food Science, Nutrition and Technology, University of Nairobi, Kangemi, Kenya

²College of Agriculture, Animal Sciences and Veterinary Medicine, School of Agriculture and Food Sciences, University of Rwanda, Musanze, Rwanda

Correspondence

Juvenal Djangwani, College of Agriculture and Veterinary Sciences, Department Food Science, Nutrition and Technology, P. O. Box 29053, -00625, University of Nairobi, Kangemi, Nairobi, Kenya.
Email: djangwani@students.uonbi.ac.ke

Funding information

Michigan State University

Abstract

Brucellosis is a zoonotic disease which is endemic to certain regions of the world including Sub-Saharan Africa. The aim of this article is to provide a recent and rapid review on brucellosis prevalence in East African Community (EAC) countries. Literature was obtained using Google Scholar search engine and screened for relevancy and fulfilment of criteria to 1, 17, 4, 4, 30 and 29 articles retained for brucellosis prevalence in Burundi, Kenya, Rwanda, South Sudan, Tanzania and Uganda. Recent literature (published in the last decade 2010 to 2019) was considered for prevalence results in this review. In EAC, livestock had an animal-level prevalence of 0.2% to 43.8%, 0.0% to 20.0% and 0.0% to 13.8% for cattle, goats and sheep respectively. In humans, the prevalence varied mostly between 0.0% and 35.8%. In conclusion, brucellosis is quite prevalent in the region. The reported prevalence calls for plans or more efforts from individual member countries and from EAC, as a region, to control brucellosis.

KEYWORDS

Brucella, brucellosis, East Africa, prevalence

1 | INTRODUCTION

Brucellosis is a bacterial zoonotic infection which is among the most widespread and top neglected diseases in the world (World Health Organization, 2012).

Brucellosis is caused by bacteria belonging to the genus *Brucella* of which the most relevant species to livestock animal health and public health are *B. abortus*, *B. melitensis*, *B. ovis* and *B. suis* (Corbel, 2006; Godfroid et al., 2014). The infection in livestock animals occurs when the animals come in contact with infected animals, materials from infected animals (aborted materials, fetal membranes, vaginal discharges, milk, manure) or infected environment (Hamdy & Amin, 2002; Kaur et al., 2018; Langoni et al., 2000; Mugizi, Boqvist, et al., 2015; Mugizi, Muradrasoli, et al., 2015; Tekle et al., 2019). In livestock animals such as cattle, goats, sheep and pigs, brucellosis

causes abortions and other reproductive disorders such as stillbirths, weak calves at birth, retained placenta and longer calving intervals in female animals (Boukary et al., 2013; McDermott et al., 2013; Schmutz et al., 1996).

Being a zoonotic disease, brucellosis is transmitted from animals to humans. Humans contract brucellosis when their skin (especially skin with cuts) or nasopharyngeal mucous tissues/membrane come in contact with infected animals' materials such as abortion materials, fetuses, placental materials, vaginal discharges, urine and manure (Corbel, 2006; Pappas et al., 2006). Humans are also infected following the consumption of unpasteurized milk and milk products and improperly cooked meat from infected animals (Casalnuovo et al., 2016; Dadar et al., 2019). In humans, brucellosis causes a febrile illness with intermittent fevers, sweats, chills, weakness, malaise, headache, insomnia, anorexia, joint and muscle pain,

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors *Veterinary Medicine and Science* Published by John Wiley & Sons Ltd

constipation, sexual impotence, nervousness and depression (Acha & Szyfres, 2001; Corbel, 2006; Food & Drug Administration, 2012; Pappas et al., 2006). In human females, brucellosis may also be accompanied by abortion (Ali et al., 2016; Khan et al., 2001; Yang et al., 2018).

Although neglected, brucellosis results in animal health burdens for the farmers and is a public health concern. It is, therefore, important for countries to implement necessary measures to control and eradicate brucellosis. For any measures to be implemented, however, brucellosis prevalence and epidemiological data are needed to form the basis for planning and tackling effectively this zoonotic infection. In African countries, in general, brucellosis prevalence and epidemiological data have been presented in a number of reviews (Akakpo & Bornarel, 1987; Boukary et al., 2014; Craighead et al., 2017; Ducrotoy et al., 2015; McDermott & Arimi, 2002; McDermott et al., 2013) and showed the presence and prevalence of brucellosis on the continent in the recent past years. In EAC countries, in particular, a review is yet to be conducted to focus and provide updated data on brucellosis prevalence in the region.

The aim of this article was, therefore, to review and summarize recent literature (2010–2019) on the prevalence of brucellosis in animals and humans, with reference to EAC countries.

2 | METHODOLOGY

2.1 | Literature search

To write this review, a literature search was conducted using Google Scholar database. Literature search was conducted for all relevant and recent articles (published from 2010 to 2019) on brucellosis prevalence in the six EAC countries, namely, Burundi, Kenya, Rwanda, South Sudan, Tanzania and Uganda (Figure 1). The used search terms were “Brucellosis AND Prevalence AND Country”, “Animal Brucellosis AND country”, “Human Brucellosis AND country”, “Bovine Brucellosis AND country”, “Caprine brucellosis and country”, “Ovine brucellosis AND country” and “Porcine brucellosis AND country”. Following literature search and articles screening, 1, 17, 4, 4, 30 and 29 articles were retained for results presentation on brucellosis prevalence in Burundi, Kenya, Rwanda, South Sudan, Tanzania and Uganda respectively (Figure 2).

In addition to Google Scholar literature search, relevant reports on country's profiles were obtained and consulted from websites of governments' institutions such as National Bureaus of Statistics and Ministries of Agriculture and from international institutions such as Food and Agriculture Organization of the United Nations (FAO).

2.2 | Inclusion and exclusion criteria

To have a recent picture of brucellosis prevalence in EAC countries, articles were included if they reported on or included data on brucellosis (animal and/or human) prevalence in at least one of the




FIGURE 1 Map showing the location of the six EAC member countries (in grey)

EAC countries (Burundi, Kenya, Rwanda, South Sudan, Tanzania and Uganda), if they were published in the last decade (between 2010 and 2019) and if they were articles published in scientific journals. Articles were excluded if they did not report on brucellosis prevalence in EAC countries and if they reported on brucellosis prevalence in EAC countries but were published before 2010. Articles were also excluded if they were review articles or were other scientific work (theses, posters and reports) not published in scientific journals.

3 | RESULTS

In EAC and in the last decade (2010 to 2019), a number of studies have been conducted on both animal and human brucellosis prevalence. The number of brucellosis studies in the EAC countries is, however, different from more brucellosis studies conducted in Uganda, Tanzania and Kenya compared with Rwanda, South Sudan and Burundi.

3.1 | Brucellosis prevalence in Burundi

Burundi is the second smallest EAC country located at the West of the regional block (Figure 1) and sharing borders with Rwanda at the North, Tanzania at the East and South, Rwanda and Democratic republic of Congo at the West. Burundi has a total area of 27,834 square kilometres and a population estimated at 11.5 million (World Bank, 2019). Burundi's most popular livestock are small ruminants (Desiere et al., 2015) with 40% to 60% of rural households owning goats and/or sheep compared to 10% to 20% of the rural households owning cattle (Jeniček & Grofová, 2015). According to 2018 estimations by FAO, there were 3,249,827 goats, 1,110,936 cattle, 774,689 pigs and 548,608 sheep in Burundi (FAO Corporate Statistical Database, 2020). Cattle rearing is especially observed in the Imbo




FIGURE 2 Literature search and screening for articles to be included in results on brucellosis prevalence in EAC countries

plain at the most western part of the country and in the provinces of Bururi, Mwaro and Muramvya (Manirakiza et al., 2017; United States Agency for International Development, 2009). In Burundi, the traditional extensive agro-pastoral cattle production system is gradually being replaced by the mixed crops–livestock system in which cattle is kept in closed spaces and fed with cut and carried forage and crop residues. Around cities like Bujumbura where the demand in fresh milk is high, the intensive cattle production system is the most commonly practiced (Manirakiza et al., 2020).

Very few studies on brucellosis prevalence in Burundi exist in literature. Although the detection of the first cases of human brucellosis were reported in what used to be Rwanda–Urundi in 1930s (Pergher & Noel, 1936), a few further studies on brucellosis in Burundi were reported, one in the 1960s and the other in the 1980s (Merker & Schlichting, 1984; Thienpont et al., 1961). Recently and in the last decade, one study on brucellosis prevalence in Burundi was found in searched literature. The study, which was conducted on dairy herds in West and Central Africa by testing farm bulk milks with milk Enzyme Linked Immuno-Sorbent Assay, reported a sero-prevalence of 14.7% (95% CI: 9.4–20.8) among dairy cattle herds in peri-urban Bujumbura (Musallam et al., 2019). For the purpose of this review, recent (2010 to 2019) brucellosis studies on other

livestock and on humans in Burundi were not found or were not reported in searched literature.

3.2 | Brucellosis prevalence in Kenya

Kenya is the biggest economy in EAC. Kenya is located at the East of the regional block (Figure 1) and sharing borders with Ethiopia and South Sudan at the North, Somalia at the East, Tanzania at the South and Uganda at the West. Kenya is 580,370 square kilometres and has a population of 47.6 million (Kenya National Bureau of Statistics, 2019). Kenya's livestock census of 2009 indicated that cattle population was estimated at 17,467,774 heads and goats were 27,740,153 heads while the sheep population was 17,129,606 (Kenya National Bureau of Statistics, 2019). According to the same census, high populations of livestock were observed in the Rift Valley region. In Kenya, livestock is raised under different livestock production systems including intensive, semi-intensive and extensive production systems. Dairy cattle, which are an important part of livestock in Kenya, are raised under different production systems. It was estimated that 45% of dairy cattle farms practice the semi-intensive/semi-grazing system while 35% of the farms practice the intensive system but on a small scale. The

rest of dairy cattle farms practice controlled extensive system (10% of farms), uncontrolled extensive system (5% of farms) and intensive system on a large scale (5% of farms; Food and Agriculture Organization of United Nations, 2018a). Livestock, in general, depends on rangelands resources which support 70% of the livestock in the country (Kenya Ministry of Agriculture & Livestock, 2019).

Brucellosis has been studied in Kenya for a long time as shown by a recent review on the disease frequency in humans and animals and risk factors for human infection from studies conducted as early as 1916 (Njeru, Wareth, et al., 2016). In the last decade (2010–2019), a number of studies have also reported on brucellosis prevalence in Kenya (Table 1).

Using a variety of diagnostic tests most of which were serological (Table 1), prevalence in cattle varied between 0.2% and 21.9% (Chota et al., 2016; Enström et al., 2017; Fèvre et al., 2017; Gicheru et al., 2015; Kairu-wanyoike et al., 2019; Kosgei et al., 2014; Nakkell et al., 2016; Okumu et al., 2019; Osoro et al., 2015). For the small ruminants, brucellosis prevalence ranged from 0.0% to 20.0% in goats and 0.0% to 13.8% in sheep (Chota et al., 2016; Kairu-wanyoike et al., 2019; Kosgei et al., 2014; Nakkell et al., 2016).

Human brucellosis was studied especially for small livestock keeping communities and for patients attending hospitals with brucellosis symptoms. The reported prevalence ranged from 0.6% to 35.8% in humans (Chota et al., 2016; de Glanville et al., 2017; Fèvre et al., 2017; Kairu-wanyoike et al., 2019; Maiyo & Obey, 2016; Nakkell et al., 2016; Njeru, Melzer, et al., 2016; Osoro et al., 2015). However, a higher county-level prevalence of 46.5% among humans was reported in the county of Marsabit in a study conducted for a group of three counties (including Marsabit) to determine sero-prevalence and risk factors for brucellosis among humans and their livestock (Osoro et al., 2015).

Of the two recent studies which covered brucellosis in camels, one reported a prevalence of 11.1% in camels in Marsabit county (Osoro et al., 2015) while the other reported a prevalence of 0% in five camels tested in West Pokot county (Chota et al., 2016).

In the last decade, a few studies also covered or extended their investigations on the prevalence of *Brucella* antibodies in milk in Kenya. One recent study, which investigated the incidence and knowledge of brucellosis in Kahuro district, Murang'a county, used the Milk Ring Test and reported that 22% of the analysed 150 pooled milk samples were positive to brucellosis (Njuguna et al., 2017). In the same study, 230 individual farm bulk milks were collected from farmers and analysed and 24% were positive to brucellosis. Such prevalence in raw milks, although not directly indicating the individual animal prevalence in cattle, is an indication of brucellosis prevalence in cattle in Kenya and the risk of transmission to consumers when milk is not pasteurized prior to using.

3.3 | Brucellosis prevalence in Rwanda

Rwanda is the smallest country in EAC located at the West of the regional block (Figure 1) and sharing borders with Uganda at the

North, Tanzania at the East, Burundi at the South and Democratic Republic of Congo at the West. Rwanda has a total area of 26,338 square kilometres. The total population in the country is estimated at 12.0 million (National Institute of Statistics of Rwanda, 2019). Rwanda's important livestock are cattle, goats, sheep and pigs. The reported numbers of different livestock showed that the country had a total of 1,856,490 cattle, 2,283,445 goats, 499,316 sheep and 703,145 pigs. The Eastern province had the highest number of cattle (28.3%) followed closely by the southern province with 27.3% of cattle (National Institute of Statistics of Rwanda, 2018). In Rwanda, cattle is raised mainly under the small-scale zero grazing system which is practiced by 80% of cattle keeping households. Semi-intensive and extensive/open grazing systems are practiced by 3% and 17% of the cattle keeping households respectively (Land O' lakes, 2014).

Brucellosis studies have been conducted in Rwanda, although few and scattered over the years (Akayezu, 1984; Chatikoba et al., 2008; Gafirita et al., 2017; Manishimwe et al., 2015; Ndazigaruye et al., 2018; Rujeni & Mbanzamihiho, 2014; Thienpont et al., 1961). In the last decade (2010–2019), a total of four studies were published on animal and human brucellosis in the country (Table 2). For animal brucellosis, a study published in 2015 (Manishimwe et al., 2015) was focusing on comparing Rose Bengal Test (RBT) with competitive Enzyme Linked Immuno-Sorbent Assay (c-ELISA) in detecting *Brucella* antibodies in cattle serum. The study was conducted on a total of 2017 sera previously collected from 157 cattle farms in Kigali and reported a bovine brucellosis prevalence of 2.0% using RBT and 1.7% using c-ELISA. A second study, published in 2018, was conducted in Nyagatare district with the aim of analysing the risk factors associated with brucellosis in cattle in the district (Ndazigaruye et al., 2018). The overall reported bovine brucellosis prevalence in Nyagatare district was 18.9%. With these studies, brucellosis prevalence in the last decade varied between 1.7% and 18.9% among cattle in Rwanda. No studies on brucellosis in small ruminants were found in searched literature for Rwanda and in the last decade.

Reported human brucellosis in Rwanda in the last decade varied between 6.1% and 25% according to two studies (Gafirita et al., 2017; Rujeni & Mbanzamihiho, 2014). In the first study conducted on women presenting with abortion or stillbirth of unknown origin at Huye district hospital, a prevalence of 25% among those women was reported (Rujeni & Mbanzamihiho, 2014). The second study covered patients attending Nyagatare district hospital, willing to participate in the study and having any of the following symptoms: intermittent or persistent fever, headache, weakness, profuse sweating, chills, arthralgia, weight loss and joint pain. The study reported a prevalence of 6.1% (Gafirita et al., 2017).

3.4 | Brucellosis prevalence in South Sudan

South Sudan is the newest EAC member country located at the north of the regional block (Figure 1) and bordered by Sudan at the North, Ethiopia at the East, Kenya, Uganda and Democratic Republic of Congo at the South and Central African Republic at the West.

TABLE 1 Studies on livestock and human brucellosis prevalence in Kenya in the last decade (2010–2019)

Species and study scope	Sample size	Diagnostic test	Prevalence in % (95% confidence interval)	Reference
Cattle				
Subnational	356	CFT	0.9–4.6 ^a	Chota et al. (2016)
Subnational	225	i-ELISA	12.4 (7.7–15.4)	Enström et al. (2017)
Subnational	983	Rapid immuno-chromatographic flow assay	0.2 (0.0–0.5) ^c	Fèvre et al. (2017)
Subnational	441	c-ELISA	6.3 (4.0–8.6)	Kairu-wanyoike et al. (2019)
Subnational	149	RBT	10.7 ^a	Kosgei et al. (2014)
Subnational	208	MRT	7.7 ^a	Gicheru et al. (2015)
Subnational	250	RBT & c-ELISA	21.9 ^a	Nakkel et al. (2016)
Subnational	398	c-ELISA	16.8 (13.2–20.4)	Okumu et al. (2019)
Subnational	2,978	i-ELISA	4.1 (3.4–4.8)	Osoro et al. (2015)
Goats				
Subnational	123	CFT	0.0–20.0 ^a	Chota et al. (2016)
Subnational	961	c-ELISA	3.3 (2.1–4.4)	Kairu-wanyoike et al. (2019)
Subnational	92	RBT	13.0 ^a	Kosgei et al. (2014)
Subnational	167	RBT & c-ELISA	7.3 ^a	Nakkel et al. (2016)
Subnational	4,080	c-ELISA	10.7 (9.3–12.3)	Osoro et al. (2015)
Sheep				
Subnational	30	CFT	0.0 to 13.8 ^a	Chota et al. (2016)
Subnational	623	c-ELISA	1.4 (0.5– 2.3)	Kairu-wanyoike et al. (2019)
Subnational	73	RBT	8.2 ^a	Kosgei et al. (2014)
Subnational	167	RBPT & c-ELISA	8.6 ^a	Nakkel et al. (2016)
Subnational	3,088	c-ELISA	7.3 (6.1–8.8)	Osoro et al. (2015)
Humans				
Subnational	562	^b	17.1 ^a	Chota et al. (2016)
Subnational	2,113	Rapid immuno- chromatographic flow assay	0.6 (0.2–0.9)	Fèvre et al. (2017)
Subnational	1,022	IgG-ELISA	35.8 (32.8–38.8)	Kairu-wanyoike et al. (2019)
Subnational	317	RBT & c-ELISA	1.3 ^a	Nakkel et al. (2016)
Subnational	2,811	IgG-ELISA	16.4 (13.5–19.6)	Osoro et al. (2015)
Subnational	825	Rapid immuno-chromatography flow assay	3.4 ^a	de Glanville et al. (2017)
Subnational	1,043	^b	32.3 ^a	Maiyo and Obey (2016)
Subnational	1,067	RBT, IgM/IgG-ELISA & q-PCR	13.7 (11.7–15.9)	Njeru, Melzer, et al. (2016)

Abbreviations: c-ELISA, Competitive Enzyme-Linked Immuno-Sorbent Assay; CFT, Complement Fixation Test; i-ELISA, Indirect Enzyme-Linked Immuno-Sorbent Assay; IgG/IgM-ELISA, Immunoglobulins G and M Enzyme-Linked Immuno-Sorbent Assay; IgG-ELISA, Immunoglobulin G Enzyme-Linked Immuno-Sorbent Assay; MRT, Milk Ring Test; q-PCR, Quantitative Polymerase Chain Reaction; RBT, Rose Bengal Test.

^aPrevalence confidence interval not provided in the original article.

^bDiagnostic test not specified (Study based on health centre results' records).

^cThe computed confidence interval value was <0.0.

South Sudan has an area of 644,329 square kilometres. South Sudan population was estimated at 12.2 million (United Nations Economic Commission for Africa, 2017). Livestock, including cattle, goats and sheep, is a very important agricultural subsector in South Sudan where the livestock per capita holding is considered among the first in Africa. Cattle, goats, sheep, camel and pig populations

were estimated at 17,729,188; 12,307,686; 11,682,172; 23,583 and 14,406 respectively (Onyango et al., 2015). Livestock is distributed all over South Sudan but more livestock populations are observed in the upper half of the country in the states of Northern Bhar El Ghazal, Warrap, Jonglei, Lakes, Western Bhar El Ghazal and Unit (Onyango et al., 2015). In South Sudan, livestock is reared

under agro-pastoralism and pastoralism mainly with 85% of South Sudanese households involved in livestock being agro-pastoralists and the remaining 15% being pastoralists (Emmanuel et al., 2018; FAO, 2016).

As is the case for Burundi and Rwanda, studies on brucellosis in South Sudan are still few. Among the few studies undertaken and published in the period of 2010 to 2019 (Table 3) was a cross-sectional study conducted in peri-urban Juba and in Terekeka county on bovine brucellosis (Lita et al., 2016). In this study, an overall individual animal sero-prevalence of 31.9% by RBT and 29.3% by c-ELISA was reported. A series of three brucellosis sero-prevalence studies was also conducted by Madut et al. on febrile patients attending hospital, on cattle and their herders and on slaughter house workers in the region of Bahr el Ghazal (Madut et al., 2019; Madut, Muwonge, et al., 2018; Madut, Nasinyama, et al., 2018). The prevalence among 416 febrile patients attending Wau hospital in Bahr el Ghazal region was 23.3% after tests of blood samples by RBT and Serum Agglutination and confirmation of results by c-ELISA (Madut, Muwonge, et al., 2018; Madut, Nasinyama, et al., 2018). Of the 893 bovine sera and 87 herders' sera tested using RBT and confirmation by c-ELISA, the reported overall prevalence was 31.0% among cattle and 33.3% among cattle herders (Madut, Muwonge, et al., 2018; Madut, Nasinyama, et al., 2018). In the same region of Bahr el Ghazal, a total 234 slaughterhouse workers were screened for brucellosis infection using RBT and the overall prevalence was 32.1% following c-ELISA confirmation of results (Madut et al., 2019). With these few studies in the last decade, reported brucellosis prevalence in South Sudan varied from 29.3% to 31.9% among cattle and from 23.3% to 33.3% among humans.

3.5 | Brucellosis prevalence in Tanzania

Tanzania is an EAC country located at the South of the regional block (Figure 1) and sharing borders with Uganda at the North, Kenya at the North-East, Mozambique at the South, Malawi at the South-West, Zambia and Democratic Republic of Congo at the West and Burundi and Rwanda at the North-West. According to Tanzania National Bureau of Statistics, Tanzania has a total

area of 947,300 square kilometres and a population of 54.2 million (Tanzania National Bureau of Statistics, 2019). Tanzania's most important livestock is cattle at an estimated population of 30,672,001 heads followed by goats estimated at 19,055,651 heads and sheep estimated at 5,565,986 heads (Tanzania National Bureau of Statistics, 2017). In Tanzania, livestock is especially raised in Shinyanga, Mwanza and Tabora regions (Engida et al., 2015).

Many brucellosis studies have been conducted in Tanzania. Recent brucellosis studies in cattle in Tanzania used serological diagnostic tests, mostly the RBT to screen samples and c-ELISA to confirm the results (Table 4). They reported an individual cattle brucellosis prevalence varying mostly between 0.2% and 11.7% (Asakura et al., 2018a,2018b; Assenga et al., 2015; Chitupila et al., 2015; Chota et al., 2016; Kayombo et al., 2017; Mathew, 2017; Sagamiko et al., 2018; Shirima et al., 2010; Shirima & John, 2016; Swai & Schoonman, 2010, 2012).

However, in one study which was conducted on animals from a single farm with a total of 350 cattle, a higher individual cattle brucellosis prevalence of 48% was reported (Mathew et al., 2015). The dairy herd from which the high individual cattle brucellosis was reported is located in the southern highlands of Tanzania and had been experiencing abortions (Mathew et al., 2015). Another investigative study following an abortion storm on a research farm reported an individual cattle brucellosis prevalence of 28.9% (Shirima et al., 2014). Brucellosis in this farm was eventually controlled through culling, among other measures, and brought to 0.0% over a period of 5 years.

Human brucellosis studies in Tanzania included mostly hospital patients with symptoms, like fever and spontaneous abortions, and communities in pastoral and agro-pastoral areas. The reported human brucellosis prevalence in Tanzania from recent studies (2010–2019) varied mostly between 0% and 28.2% (Assenga et al., 2015; Bouley et al., 2012; Carugati et al., 2018; Cash-goldwasser et al., 2017; Chipwaza et al., 2015; Chota et al., 2016; Crump et al., 2013; Mngumi et al., 2016; Nonga & Mwakapeje, 2017; Shirima et al., 2010; Shirima & John, 2016). It should, however, be mentioned that a prevalence as high as 58.1% was reported from a study which investigated the association of *Brucella* seropositivity with abortion for a group of 148 women with spontaneous abortions and 250 women with full-term

Species and study scope	Sample size	Diagnostic test	Prevalence in % (95% Confidence Interval)	Reference
Cattle				
Subnational	2,017	RBT	2.0 ^a	Manishimwe et al. (2015)
		c-ELISA	1.7 ^a	
Subnational	604	RBT	18.9 ^a	Ndazigaruye et al. (2018)
Humans				
Subnational	198	RBT	6.1 (0.6–7.8)	Gafirita et al. (2017)
Subnational	60	RBT	25 ^a	Rujeni and Mbanzamihigo (2014)

TABLE 2 Studies on livestock and human brucellosis prevalence in Rwanda in the last decade (2010–2019)

Note: c-ELISA: Competitive Enzyme-Linked Immuno-Sorbent Assay; RBT: Rose Bengal Test.

^aPrevalence confidence interval not provided in the original article.

TABLE 3 Studies on livestock and human brucellosis prevalence in South Sudan in the last decade (2010–2019)

Species and study scope	Sample size	Diagnostic test	Prevalence in % (95% Confidence Interval)	Reference
Cattle				
Subnational	160	RBT	31.9 ^a	Lita et al. (2016)
	147	c-ELISA	29.3 ^a	
Subnational	893	RBT & c-ELISA	31.0 (28.0–34.2)	Madut, Muwonge, et al. (2018); Madut, Nasinyama, et al. (2018)
Humans				
Subnational	416	RBT, SAT & c-ELISA	23.3 ^a	Madut, Muwonge, et al. (2018); Madut, Nasinyama, et al. (2018)
Subnational	87	RBT & c-ELISA	33.3 (23.9–44.3)	Madut, Muwonge, et al. (2018); Madut, Nasinyama, et al. (2018)
Subnational	234	RBPT & c-ELISA	32.1 ^a	Madut et al. (2019)

Abbreviations: c-ELISA, Competitive Enzyme-Linked Immuno-Sorbent Assay; RBT, Rose Bengal Test; SAT, Serum Agglutination Test.

^aPrevalence confidence interval not provided in the original article.

deliveries. The group of women with spontaneous abortions had a prevalence of 58.1%, while the group with full-term deliveries had a prevalence of 26% (Mujuni et al., 2018). A different but also focused study on 250 abattoir workers and meat vendors in the city of Mwanza reported that 48.4% of them were *Brucella* seropositive (Mirambo et al., 2018).

Fewer studies were conducted on small ruminants in Tanzania in the last decade. Brucellosis prevalence in goats was between 0% and 2.0% while brucellosis prevalence in sheep was between 0% and 5.7% (Assenga et al., 2015; Mathew et al., 2015; Shirima & John, 2016).

Apart from common livestock animals, a few recently published studies (2010–2019) in Tanzania covered other animals. A study on the epidemiology of *Brucella* infection in human, livestock and wild-life interface in the Katavi-Rukwa ecosystem included 38 buffaloes, two lions and two zebras and reported a brucellosis prevalence of 7.9% in buffaloes while one of the two lions was seropositive and none of the zebras was (Assenga et al., 2015). In a study to investigate a farm which had been experiencing cattle abortions, 6 dogs were included and none of the dogs tested positive for brucellosis (Mathew et al., 2015). Dogs were also covered in another study with a sample of 100 dogs randomly selected in the region of Morogoro and no dog was positive to *Brucella canis* (Muhairwa et al., 2012). A study conducted on camels in agro-pastoral communities of Northern Tanzania reported a prevalence of 2.1% at an animal level for a sample of 193 camels selected from 14 traditional herds (Swai et al., 2011). A low prevalence of 0.7% was reported in pigs from five selected pig slaughter facilities in Dar-Es-Salam (Simon et al., 2015).

A few recent studies (published from 2010 to 2019) covered also the prevalence of *Brucella* antibodies in raw marketed milk in Tanzania. One such study which sought to evaluate the microbiological quality and associated health risks for raw milk marketed in

Tanga region reported that 56% of 59 raw milk samples collected from selling points and deliverers were *Brucella* positive (Swai & Schoonman, 2011). This high milk contamination may be due to milk bulking and pooling from different cattle and farms. It presents a risk for transmission of brucellosis to consumers and calls for milk boiling or pasteurization prior to using. Another study which, in addition to animal sera, investigated the milk from dairy animals reported a prevalence of *Brucella* in 3.7% and 0% of cattle milk and goat milk respectively (Assenga et al., 2015). Herd-level brucellosis prevalence was, also, investigated in a study using bulk farm milk. The prevalence of *Brucella* in milk at herd level was 44.4% of 124 agro-pastoral farms investigated in Morogoro region (Asakura et al., 2018a, 2018b).

In addition to determining brucellosis prevalence in animals and humans, a few studies in Tanzania identified and characterized some prevalent *Brucella* species. In their investigative study on a single farm, Mathew and colleagues (Mathew et al., 2015) cultured, identified and characterized *Brucella* isolates from cattle to be *Brucella abortus* biovar 3. In another study published about the same time, molecular methods were used and *Brucella abortus* biovar 1 was identified and characterized from cattle milk (Assenga et al., 2015).

3.6 | Brucellosis prevalence in Uganda

Uganda is an EAC country at the Central-West of the regional block (Figure 1). Uganda shares borders with South Sudan at the North, Kenya at the East, Tanzania at the South, Rwanda at the South-West and Democratic Republic of Congo at the West. As indicated by Uganda Bureau of Statistics, Uganda is 241,550.7 square kilometres in total area and has a human population of 41.0 million (Uganda Bureau of Statistics, 2018). Uganda's important livestock are

TABLE 4 Studies on livestock and human brucellosis prevalence in Tanzania in the last decade (2010–2019)

Species and study scope	Sample size	Diagnostic test	Prevalence in % (95% Confidence Interval)	Reference
Cattle				
Subnational	667	RBT & c-ELISA	0.2 (0.0–1.1) ^d	Asakura et al. (2018a)
	673	RBT & c-ELISA	7.0 (5.7–8.4)	
Subnational	1,103	RBT & c-ELISA	6.8 (5.4–8.5)	Assenga et al. (2015)
Subnational	410	RBT & c-ELISA	5.6 (3.8–8.3)	Chitupila et al. (2015)
Subnational	1,376	RBT & c-ELISA	1.0–11.4 ^a	Chota et al. (2016)
Subnational	192	RBT & c-ELISA	4.2 ^a	Kayombo et al. (2017)
Subnational	658	i-ELISA	5.4 ^a	Mathew, 2017)
Subnational	200	RBT	21.5 (16–27)	Mathew et al. (2015)
		i-ELISA	48.0 (41–55)	
Subnational	296	RBT & c-ELISA	7.8 ^a	Shirima and John (2016)
Subnational	929	RBT & c-ELISA	2.8 (1.4–5.6)	Sagamiko et al. (2018)
	282	RBT & c-ELISA	11.3 (9.4–13.5)	
Subnational	2,723	c-ELISA	4.9 ^a	Shirima et al. (2010)
Subnational	51	RBT	11.7 (9.1–14.9)	Swai and Schoonman (2012)
Subnational	655	RBT	5.3 (3.1–7.8)	Swai and Schoonman (2010)
Subnational	483	RBT & c-ELISA	28.9 ^a	Shirima et al. (2014)
Goats				
Subnational	248	RBT & c-ELISA	1.6 (0.4–4.1)	Assenga et al. (2015)
Subnational	50 goats	RBT	0	Mathew et al. (2015)
		i-ELISA	2.0 (0.0–7.0) ^d	
Subnational	75	RBT & c-ELISA	0	Shirima and John (2016)
Subnational	^c	RBT & c-ELISA	0	Shirima et al. (2014)
Sheep				
Subnational	35	RBT	0	Mathew et al. (2015)
		i-ELISA	5.7 (0.0–17.0) ^d	
Subnational	42	RBT & c-ELISA	0	Shirima and John (2016)
Humans				
Subnational	340	RBT, BAPA & Riv. T	0.6 (0.1–2.1)	Assenga et al. (2015)
Subnational	455	MAT & Blood culture	3.5 ^a	Bouley et al. (2012)
Subnational	1,095	MAT & Blood culture	2.9 ^a	Carugati et al. (2018)
Subnational	562	MAT	6.9 ^a	Cash-Goldwasser et al. (2017)
Subnational	370	IgM-ELISA, IgG-ELISA & MAT	7.0 ^a	Chipwaza et al. (2015)
Subnational	578	^b	28.2 ^a	Chota et al. (2016)
Subnational	118	MAT	13.6 ^a	Crump et al. (2013)
Subnational	250	SAT	48.4 (42–54)	Mirambo et al. (2018)
Subnational	382	Rapid <i>Brucella</i> serum agglutination	14.1 (10.6–17.5)	Mngumi et al. (2016)

(Continues)

TABLE 4 (Continued)

Species and study scope	Sample size	Diagnostic test	Prevalence in % (95% Confidence Interval)	Reference
Subnational	148	SAT	58.1 (50–66)	Mujuni et al. (2018)
Subnational	13,642	^b	5.8 ^a	Nonga and Mwakapeje (2017)
Subnational	82	RBT & c-ELISA	0	Shirima and John (2016)
Subnational	120	c-ELISA	10.0 ^a	Shirima et al. (2014)
Subnational	460	c-ELISA	8.3 ^a	Shirima et al. (2010)

Abbreviations: BAPA, Buffered Acidified Plate Antigen Test; c-ELISA, Competitive Enzyme-Linked Immuno-Sorbent Assay; i-ELISA, Indirect Enzyme-Linked Immuno-Sorbent Assay; IgG-ELISA, Immunoglobulin G Enzyme-Linked Immuno-Sorbent Assay; IgM-ELISA, Immunoglobulin M Enzyme-Linked Immuno-Sorbent Assay; MAT, Microscopic Agglutination Test; RBT, Rose Bengal Test; Riv. T, Rivanol precipitation Test; SAT, Slide Agglutination Test.

^aPrevalence confidence interval not provided in the original article.

^bDiagnostic test not specified (Study based on health centre results' records).

^cSample size not specified.

^dThe computed confidence interval value was <0.0.

cattle, goats, sheep, pigs and poultry. There were 14,189,000 cattle, 16,034,000 goats, 4,445,000 sheep and 4,109,000 pigs in 2017 (Uganda Bureau of Statistics, 2018). The most important livestock in Uganda is cattle. Cattle is especially found in the "cattle corridor" which extends from the northeast of the country, with the highest concentration of cattle, to the southwest (Egeru et al., 2014). Cattle in Uganda is raised under commercial ranching, pastoral, agro-pastoral and semi-intensive production systems, the agro-pastoral system being the most common one (FAO, 2019).

In a recent report, FAO estimated brucellosis prevalence in Uganda at a national level to be 10% in cattle and 5.5% in cattle keepers (FAO 2018b). In addition to FAO's national estimations, many other studies on brucellosis have been conducted in Uganda. According to most recent cattle brucellosis studies published from 2010 to 2019 (Table 5), brucellosis was diagnosed using mostly serological tests (Table 5). The reported brucellosis prevalence in cattle (animal level) varied a lot and was between 1.2% and 43.8% (Bugeza et al., 2019; Ezama et al., 2019; Kabi et al., 2015; Kashiwazaki et al., 2012; Lolli et al., 2016; Makita et al., 2011; Miller et al., 2016; Mugizi, Boqvist, et al., 2015; Mugizi, Muradrasoli, et al., 2015; Nanfuka, 2018; Nguna et al., 2019; Nina et al., 2017; Nizeyimana et al., 2013).

Small ruminants were also covered in recent brucellosis studies in Uganda. The reported brucellosis prevalence in goats was lower compared with cattle and varied between 0.3% and 9.8% (Dubad et al., 2015; Lolli et al., 2016; Miller et al., 2016; Nguna et al., 2019). The reported prevalence in sheep varied between 2.6% and 10.5% (Dubad et al., 2015; Lolli et al., 2016).

Among recent studies, one study investigated brucellosis in pigs from three districts in Uganda where pig keeping is commonly practiced. The reported prevalence in the districts varied from 0.0% to 0.2% (Erume et al., 2016).

Human brucellosis was also reported in recent brucellosis studies in Uganda published from 2010 to 2019. The reported human brucellosis prevalence varied between 4.0% and 33.0% (Ezama et al., 2018,

2019; Frank et al., 2017; Kansime et al., 2015; Majalija et al., 2018; Migisha et al., 2018; Miller et al., 2016; Muloki et al., 2018; Nabukenya et al., 2013; Nanfuka, 2018; Nasinyama et al., 2014; Nguna et al., 2019; Tumwine et al., 2015). These human brucellosis results should, however, be interpreted considering that most of them were conducted on suspected patients attending hospitals with febrile illness or having prolonged fevers and on exposed cattle keepers and farm attendants in pastoralist and agro-pastoralist communities. Human brucellosis studies used serological diagnosis tests with only one study adding blood culturing to confirm the presence of *Brucella* spp. (Migisha et al., 2018).

A few recent studies in Uganda were also conducted to determine the prevalence of *Brucella* antibodies in cattle milk. Raw milk samples collected from dairy farms, milk shops, street vendors, milk deliverers, boiling points, dairies and milk collection centres were tested using serological diagnosis tests and the prevalence varied between 6.5% and 40% (Hoffman et al., 2016; Kamwine et al., 2017; Makita et al., 2010; Rock et al., 2016). While these results are from bulk and pooled raw milks and cannot be related to individual cattle prevalence, they are still an indication of the presence of brucellosis in cattle.

4 | DISCUSSION

Brucellosis is considered endemic across the African continent (Franc et al., 2018). The aim of this article was to review recent literature (2010–2019) on the prevalence of brucellosis in animals and humans with reference to EAC countries. This review indicates that brucellosis is prevalent in EAC with prevalence ranges that are quite variable within individual EAC countries and between countries. This variation should, however, be looked at considering that the retained and reviewed studies used different serological diagnostic tests and different sampling techniques from populations of various sizes. The practiced livestock production systems were also different, from the

TABLE 5 Studies on livestock and human brucellosis prevalence in Uganda in the last decade (2010–2019)

Species and study scope	Sample size	Diagnostic test	Prevalence in % (95% Confidence Interval)	Reference
Cattle				
Subnational	728	RBT & i-ELISA	3.2 (1.9–4.5)	Bugeza et al. (2019)
Subnational	839	RBT	34.7 ^a	Ezama et al. (2019)
National	925	i-ELISA & c-ELISA	8.64 ^a	Kabi et al. (2015)
Subnational	1,237	RBT & i-ELISA	21.5 ^a	Kashiwazaki et al. (2012)
	1,033	RBT & i-ELISA	3.4 ^a	
Subnational	3,935	RBT	9.8 (8.9–10.7)	Lolli et al. (2016)
Subnational	423	c-ELISA	5.0 (2.7–9.3)	Makita et al. (2011)
Subnational	768	RBT (<i>Abortus</i> antigen)	12.9 (10.0–16.2)	Miller et al. (2016)
		RBT (<i>Melitensis</i> antigen)	15.7 (12.4–19.3)	
Subnational	1,007	i-ELISA & c-ELISA	7.5 (6.1–9.4)	Mugizi, Boqvist, et al. (2015); Mugizi, Muradrasoli, et al. (2015)
Subnational	1,503	SAT & i-ELISA	23.0 ^a	Nanfuka, 2018)
Subnational	345	i-ELISA	1.2 ^a	Nguna et al. (2019)
Subnational	1,749	RBT & i-ELISA	43.8 ^a	Nina et al. (2017)
Subnational	149	i-ELISA	3.3 ^a	Nizeyimana et al. (2013)
Goats				
Subnational	305	RBT	8.8 ^a	Dubad et al. (2015)
Subnational	729	RBT	8.8 (6.9–11.1)	Lolli et al. (2016)
Subnational	315	RBT (<i>Abortus</i> antigen)	1.1 (0–6.0) ^d	Miller et al. (2016)
		RBT (<i>Melitensis</i> antigen)	9.8 (3.8–15.7)	
Subnational	351	i-ELISA	0.3 ^a	Nguna et al. (2019)
Sheep				
Subnational	95	RBT	10.5 ^a	Dubad et al. (2015)
Subnational	306	RBT	2.6 (1.2–5.3)	Lolli et al. (2016)
Humans				
Subnational	216	RBT	33 (27–39)	Ezama et al. (2019)
Subnational	216	RBT & IgM-ELISA	13.4 ^a	Ezama et al. (2018)
Subnational	177	Rapid agglutination test	10.7 ^a	Frank et al. (2017)
Subnational	9,177	PAT	14.4 ^a	Kansiime et al. (2015)
Subnational	200	SAT and TAT	7.5 ^a	Majalija et al. (2018)
Subnational	235	RBT	14.9 (10.6–20.1)	Migisha et al. (2018)
		Blood culture	4.3 ^a	
Subnational	236	IgG/IgM-LFA	8.1 (3.2–13.0)	Miller et al. (2016)
Subnational	251	PAT	18.7 ^a	Muloki et al. (2018)
Subnational	232	MAT & TAT	10 (6–16)	Nabukenya et al. (2013)
Subnational	113	i-ELISA	4–12 ^a	Nanfuka (2018)
Subnational	161	Rapid Agglutination Test, TAT & c-ELISA	5.8 (3.3–8.3)	Nasinyama et al. (2014)
	168	Rapid Agglutination Test, TAT & c-ELISA	9.0 (4.7–13.3)	
Subnational	451	i-ELISA	4.4 ^a	Nguna et al. (2019)

(Continues)

TABLE 5 (Continued)

Species and study scope	Sample size	Diagnostic test	Prevalence in % (95% Confidence Interval)	Reference
Subnational	235	SAT & RBT	17.0 ^a	Tumwine et al. (2015)

Abbreviations: c-ELISA, Competitive Enzyme-Linked Immuno-Sorbent Assay; i-ELISA, Indirect Enzyme-Linked Immuno-Sorbent Assay; IgG/IgM-LFA, Immunoglobulins G and M lateral flow assay; IgM-ELISA, Immunoglobulin M Enzyme-Linked Immuno-Sorbent Assay; MAT, Microplate Agglutination Test; PAT, Plate Agglutination Test; RBT, Rose Bengal Test; SAT, Serum Agglutination Test; TAT, Tube Agglutination Test.

^aPrevalence confidence interval not provided in the original article.

^bDiagnostic test not specified (Study based on health centre results' records).

^cSample size not specified.

^dThe computed confidence interval value was <0.0.

traditional free open grazing system through mixed agro-pastoral system to zero grazing system.

The cattle brucellosis prevalence range of 0.2% to 43.8% observed over EAC is comparable but higher than the prevalence in Sub-Saharan Africa (SSA). A review of brucellosis sero-prevalence studies published from 2003 to 2015 from 12 different SSA countries showed that brucellosis prevalence was between 1.0% and 36.6% among cattle raised under a variety of livestock production systems (Ducrotoy et al., 2015). The cattle brucellosis prevalence ranges of 0.2% to 43.8% reported in the current review is, also, comparable but higher than previous cattle brucellosis prevalences reported for specific EAC countries (McDermott & Arimi, 2002) or estimated for EAC as a region (International Livestock Research Institute, 2012). In their review on cattle brucellosis in SSA including specific EAC countries (Tanzania, Kenya, Burundi and Uganda), McDermott and Arimi reported a brucellosis prevalence range varying mostly from 1.8% in Tanzania to 25.4% in Burundi (McDermott & Arimi, 2002). Grace and colleagues also estimated a lower prevalence (8.2%) among cattle in the East African region (International Livestock Research Institute, 2012). In the current review and across other reports (Ducrotoy et al., 2015; International Livestock Research Institute, 2012; McDermott & Arimi, 2002), higher cattle brucellosis prevalence is observed in cattle raised in larger pastoral and agro-pastoral herds and lower prevalence is observed in small holder farms. The pastoral and agro-pastoral livestock production systems, in which cattle closely interact within herds and between different herds, are practiced by at least some communities or subregions in each of the EAC countries. In such livestock production systems, cattle share grazing areas, watering sources and bulls for natural breeding. Brucellosis is transmitted from animal to animal and has been reported to spread and stay in herds through contaminated grazing areas, contaminated water and through natural breeding by brucellosis-positive bulls (Aparicio, 2013). In their efforts to control cattle brucellosis, EAC and EAC countries should, therefore, include and adapt brucellosis control strategies recommended for pastoral and agro-pastoral livestock production systems with high brucellosis prevalence.

Outside Africa, a number of developed countries including New Zealand, Sweden, Denmark, Norway, Finland, Germany, Switzerland, Canada, Japan, UK, Netherlands, Belgium, Luxembourg and Austria have controlled and eradicated brucellosis (Benkirane, 2006; Pappas et al., 2006). Other developing countries in Central and South America,

South and South-East Asia and Middle East are, like Africa, still facing bovine brucellosis, although the reported prevalence is lower. In Central American countries of Guatemala, Belize, Honduras, El Salvador, Nicaragua, Costa Rica and Panama, bovine brucellosis was estimated to be from 4% to 8% and bovine brucellosis was 3% to 4%, 2% to 2.5% and 0.04% to 0.28% among cattle in Paraguay, individual Dairy cattle in Argentina and across federal Brazil (Lopes et al., 2010). Brucellosis estimates from South Asia and South East Asia regions were 16.0% and 2.9% respectively (International Livestock Research Institute, 2012). Cattle individual sero-prevalence in Middle East was from 0.8% to 12.2% (Musallam et al., 2016).

As observed in the current review, studies on brucellosis in small ruminants are fewer compared with studies on cattle. Indeed, for three of the six covered EAC countries, brucellosis in small ruminants has not yet been studied or is yet to be reported. This is a similar situation across SSA where there is limited information on brucellosis studies on small ruminants (Ducrotoy et al., 2015). For the few reported studies on goats and sheep brucellosis in EAC, the prevalence range was 0% to 20.0% among goats and 0% to 13.8% among sheep. This EAC prevalence range is higher compared with the SSA prevalence which varied between 0% and 4.8% for sheep and between 0% and 5.5% for goats (Ducrotoy et al., 2015). Although studies on brucellosis in small ruminants are still few, there is a need to evaluate the significance of this infection among small ruminants in EAC and across Africa. Indeed, goats and sheep are natural hosts and mainly infected by *Brucella melitensis* (Bruce, 1887). Among *Brucella* species, *B. melitensis* is the most virulent to humans resulting in a more acute infection (Mantur et al., 2007; Moreno, 2014). Outside Africa, brucellosis in small ruminants is still prevalent and remains a major problem in the Mediterranean region, the Middle East, Central Asia, South Asia and South-East (McDermott et al., 2013; Musallam et al., 2016). The recent brucellosis studies in EAC indicate the prevalence of the infection among livestock and human in the region and call for plans in individual countries to control brucellosis. FAO has given general guidelines in controlling brucellosis, which can be very well adapted by the different individual EAC countries. In regions, like EAC, where brucellosis prevalence is higher than 10%, FAO recommends a mass brucellosis vaccination until the prevalence is reduced to below 2% (FAO, 2009). Once the prevalence is brought under control, further eradication strategies can, then, be considered. Prior to any brucellosis control strategy, epidemiological studies are needed to determine the prevalence taking into account

the different regions within the same country or different countries within a same region (Blasco, 2010).

Most of the studies considered in the current review were targeting livestock and humans in identified suspected agro-pastoral communities within individual EAC countries. In designing their approach to control brucellosis, EAC countries could take advantage of such studies to determine the epidemiological units of intervention and design-appropriate brucellosis control strategies. In designing any brucellosis control strategy, there are, however, important considerations for the strategy to be implemented successfully. In countries where brucellosis control in livestock was, for example, implemented and attained success, adequate veterinary infrastructure was in place or put in place, awareness campaigns conducted to concerned farmers and economic resources availed by government and government donors (Blasco, 2010; FAO, 2014). It is also critical that the country's or region's political will to control brucellosis is demonstrated and maintained throughout the implementation of any brucellosis control strategy. In developed countries where brucellosis was controlled and eradicated, a strong political support and a legal framework to enforce control measures were important to the success of the control and eradication programs (FAO, 2014).

With a brucellosis control strategy in place, the invested efforts and economic resources have to be maintained over the course of the strategy implementation which is usually long. In countries where brucellosis is high, FAO recommends to start by long-term mass vaccination to control the disease. This vaccination can take up to 10 years to sustainably bring brucellosis prevalence to low levels (FAO, 2014).

Once a brucellosis control strategy is designed and needed resources availed, the implementing country needs to also ensure that good farm management is practiced to enable the control strategy to be effective and successful. The control of livestock movement, screening of replacement livestock prior to their introduction to the farms, hygienic disposal of abortive materials are all good farm management practices that have been reported as additional elements to the success of any brucellosis control strategy (Avila-Granados et al., 2019; Bamaiyi et al., 2014; Perez-Sancho et al., 2015; Zamri-Saad & Kamarudin, 2016).

5 | CONCLUSION

The literature presented in this review shows that brucellosis is prevalent in livestock in EAC. For cattle, which was the most studied among livestock, the reported individual cattle brucellosis prevalence varied mostly from 0.2% to 43.8% in the region. In small ruminants, brucellosis prevalence studies were very few and reported a prevalence varying between 0.0% and 20.0% in goats and 0.0 and 13.8% in sheep at EAC level. With most animal studies in EAC having focused on cattle, establishing the prevalence of brucellosis in small ruminants is also needed.

At EAC level, the reported prevalence of human brucellosis among studied patients attending hospitals and exposed pastoral and agro-pastoral communities varied mostly from 0.0% to 35.8%.

The reviewed studies show the continued existence and prevalence of brucellosis in EAC. These studies are a basis for individual EAC countries and the region to plan or revive efforts towards the control and eradication of brucellosis. Any brucellosis control plan will need, however, a well-designed strategy considering the involved economic resources, adequate veterinary services, time and consistency over the implementation time.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The authors have adhered to ethical policies of the journal. No ethical approval or consent forms were necessary as this is a review article.

ACKNOWLEDGEMENTS

This work was funded by the Borlaug Higher Education for Agricultural Research and Development program based at Michigan State University.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest.

AUTHOR CONTRIBUTION

Juvenal Djangwani: Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Project administration; Resources; Visualization; Writing-original draft; Writing-review & editing. **George Abong:** Conceptualization; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Writing-review & editing. **Lucy Gicuku Njue:** Conceptualization; Methodology; Resources; Supervision; Validation; Visualization; Writing-review & editing. **Dasel Wambua Mulwa Kaindi:** Conceptualization; Methodology; Resources; Supervision; Validation; Visualization; Writing-review & editing.

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1002/vms3.425>.

DATA AVAILABILITY STATEMENT

This is a review article and no new data were created or analysed.

ORCID

Juvenal Djangwani  <https://orcid.org/0000-0002-6872-9151>

George Ooko Abong  <https://orcid.org/0000-0003-2524-7862>

REFERENCES

- Acha, P. N., & Szyfres, B. (2001). Bacterioses. In P. N. Acha, & B. Szyfres (Eds.), *Zoonoses and communicable diseases common to man and animals* (Vol. 1, Issue 3, p. 423). Pan American Health Organization. <https://doi.org/10.1590/s1135-57272005000300012>
- Akakpo, A. J., & Bornarel, P. (1987). Epidémiologie des brucelloses animales en Afrique tropicale: Enquêtes clinique, sérologique et

- bactériologique. *Revue Scientifique et Technique de l'OIE*, 6(4), 981–1027. <https://doi.org/10.20506/rst.6.4.313>
- Akayezu, J. M. V. (1984). *A propos d'une enquête sero-épidémiologique sur la brucellose bovine au Rwanda*. Ecole Inter-Etats des Sciences et Médecine Vétérinaires (EISMV)-Dakar-Sénégal.
- Ali, S., Akhter, S., Neubauer, H., Scherag, A., Kesselmeier, M., Melzer, F., Khan, I., El-Adawy, H., Azam, A., Qadeer, S., & Ali, Q. (2016). Brucellosis in pregnant women from Pakistan: An observational study. *BMC Infectious Diseases*, 16(1), 1–6. <https://doi.org/10.1186/s12879-016-1799-1>
- Aparicio, E. D. (2013). Epidemiology of brucellosis in domestic animals caused by *Brucella melitensis*, *Brucella suis* and *Brucella abortus*. *Revue Scientifique et Technique de l'OIE*, 32(1), 53–60. <https://doi.org/10.20506/rst.32.1.2187>
- Asakura, S., Makingi, G., Kazwala, R., & Makita, K. (2018a). Brucellosis risk in urban and agro-pastoral areas in Tanzania. *EcoHealth*, 15, 41–51. <https://doi.org/10.1007/s10393-017-1308-z>
- Asakura, S., Makingi, G., Kazwala, R., & Makita, K. (2018b). Herd-level risk factors associated with *Brucella* sero-positivity in cattle, and perception and behaviours on the disease control among agro-pastoralists in Tanzania. *Acta Tropica*, 187(August), 99–107. <https://doi.org/10.1016/j.actatropica.2018.07.010>
- Assenga, J. A., Matamba, L. E., Muller, S. K., Malakalinga, J. J., & Kazwala, R. R. (2015). Epidemiology of Brucella infection in the human, livestock and wildlife interface in the Katavi-Rukwa ecosystem. *Tanzania. BMC Veterinary Research*, 11(1), 189. <https://doi.org/10.1186/s12917-015-0504-8>
- Avila-Granados, L. M., Garcia-Gonzalez, D. G., Zambrano-Varon, J. L., & Arenas-Gamboa, A. M. (2019). Brucellosis in Colombia: Current status and challenges in the control of an endemic disease. *Frontiers in Veterinary Science*, 6(SEP), 1–12. <https://doi.org/10.3389/fvets.2019.00321>
- Bamaiyi, P. H., Hassan, L., & Zainal, M. A. (2014). Updates on brucellosis in Malaysia and Southeast Asia. *Malaysian Journal of Veterinary Research*, 71–82.
- Bank, W. (2019). *World bank-Burundi population*. World Bank. https://data.worldbank.org/indicator/SP.POP.TOTL?locations=BI&most_recent_value_desc=false&view=chart
- Benkirane, A. (2006). Ovine and caprine brucellosis: World distribution and control/eradication strategies in West Asia/North Africa region. *Small Ruminant Research*, 62(1-2), 19–25. <https://doi.org/10.1016/j.smallrumres.2005.07.032>
- Blasco, J. (2010). Control and eradication strategies for *Brucella melitensis* infection in sheep and goats. *Section of Biological and Medical Sciences of the Macedonian Academy of Sciences and Arts*, 165, 145–165.
- Boukary, A. R., Saegerman, C., Adehossi, E., Matthys, F., Vias, G. F., Yenikoye, A., & Thys, E. (2014). La brucellose en Afrique subsaharienne. *Annales de Médecine Vétérinaire*, 158(1), 283–289.
- Boukary, A. R., Saegerman, C., Abatih, E., Fretin, D., Bada, R. A., De Deken, R., Harouna, H. A., Yenikoye, A., & Thys, E. (2013). Seroprevalence and potential risk factors for *Brucella* Spp. infection in traditional cattle, sheep and goats reared in urban, periurban and rural areas of Niger. *PLoS ONE*, 8(12), 1–12. <https://doi.org/10.1371/journal.pone.0083175>
- Bouley, A. J., Biggs, H. M., Stoddard, R. A., Morrissey, A. B., Bartlett, J. A., Afwamba, I. A., Maro, V. P., Kinabo, G. D., Saganda, W., Cleaveland, S., & Crump, J. A. (2012). Brucellosis among hospitalized febrile patients in northern Tanzania. *American Journal of Tropical Medicine and Hygiene*, 87(6), 1105–1111. <https://doi.org/10.4269/ajtmh.2012.12-0327>
- Bruce, D. (1887). *Note on the discovery of a microorganism in malta fever*. John Brigg.
- Bugeza, J., Muwonge, A., Munyeme, M., Lasuba, P., Godfroid, J., & Kankya, C. (2019). Seroprevalence of bovine brucellosis and associated risk factors in Nakasongola district. *Uganda. Tropical Animal Health and Production*, 51(7), 2073–2076. <https://doi.org/10.1007/s11225-0-018-1631-6>
- Carugati, M., Biggs, H. M., Maze, M. J., Stoddard, R. A., Cash-Goldwasser, S., Hertz, J. T., Halliday, J. E. B., Saganda, W., Lwezaula, B. F., Kazwala, R. R., Cleaveland, S., Maro, V. P., Rubach, M. P., & Crump, J. A. (2018). Incidence of human brucellosis in the Kilimanjaro Region of Tanzania in the periods 2007–2008 and 2012–2014. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 112(3), 136–143. <https://doi.org/10.1093/trstmh/try033>
- Casalnuovo, F., Ciabrone, L., Cacia, A., & Ripa, P. (2016). Contamination of bovine, sheep and goat meat with *Brucella* spp. *Italian Journal of Food Safety*, 5(3). <https://doi.org/10.4081/ijfs.2016.5913>
- Cash-Goldwasser, S., Maze, M. J., Rubach, M. P., Biggs, H. M., Stoddard, R. A., Sharples, K. J., Halliday, J. E. B., Cleaveland, S., Shand, M. C., Mmbaga, B. T., Muiruri, C., Saganda, W., Lwezaula, B. F., Kazwala, R. R., Maro, V. P., & Crump, J. A. (2017). Risk factors for human brucellosis in Northern Tanzania. *American Journal of Tropical Medicine and Hygiene*, 98(2), 598–606. <https://doi.org/10.4269/ajtmh.17-0125>
- Chatikoba, P., Manzi, M., Kagarama, J., Rwaremika, J. D., & Umunezero, O. (2008). The prevalence of bovine brucellosis in milking dairy herds in nyagatare and its implications on dairy productivity and public health. *The 3rd International Conference on Appropriate Technology (3rd ICAT)*, 368–376. <http://www.howard.edu/library/scholarship@howard/books/2008/icat2008.pdf>
- Chipwaza, B., Mhamphi, G. G., Ngatunga, S. D., Selemani, M., Amuri, M., Mugasa, J. P., & Gwakisa, P. S. (2015). Prevalence of bacterial febrile illnesses in children in Kilosa district, Tanzania. *PLoS Neglected Tropical Diseases*, 9(5). <https://doi.org/10.1371/journal.pntd.0003750>
- Chitupila, G. Y., Komba, E. V. G., & Mtui-Malamsha, N. J. (2015). Epidemiological study of bovine brucellosis in indigenous cattle population in Kibondo and Kikonko districts, Western Tanzania. *Livestock Research for Rural Development*, 27(6).
- Chota, A., Magwisha, H., Stella, B., Bunuma, E., Shirima, G., Mugambi, J., Omwenga, S., Wesonga, H., Mbatha, P., & Gathogo, S. (2016). Prevalence of brucellosis in livestock and incidences in humans in east Africa. *African Crop Science Journal*, 24(1), 45. <https://doi.org/10.4314/acsj.v24i1.5s>
- Corbel, M. J. (2006). *Brucellosis in humans and animals*. World Health Organization, 1–88. <https://doi.org/10.2105/AJPH.30.3.299>
- Craighead, L., Meyer, A., Chengat Prakashbabu, B., Musallam, I., Akakpo, A. J., Kone, P., Guitian, J., & Häslar, B. (2017). Brucellosis in West and Central Africa: A review of the current situation in a changing landscape of dairy cattle systems. *Acta Tropica*, 179, 96–108. <https://doi.org/10.1016/j.actatropica.2017.12.026>
- Crump, J. A., Morrissey, A. B., Nicholson, W. L., Massung, R. F., Stoddard, R. A., Galloway, R. L., Ooi, E. E., Maro, V. P., Saganda, W., Kinabo, G. D., Muiruri, C., & Bartlett, J. A. (2013). Etiology of severe non-malaria febrile illness in northern Tanzania: A prospective cohort study. *PLoS Neglected Tropical Diseases*, 7(7). <https://doi.org/10.1371/journal.pntd.0002324>
- Dadar, M., Shahali, Y., & Whatmore, A. M. (2019). Human brucellosis caused by raw dairy products: A review on the occurrence, major risk factors and prevention. *International Journal of Food Microbiology*, 292, 39–47. <https://doi.org/10.1016/j.ijfoodmicro.2018.12.009>
- de Glanville, W. A., Conde-Álvarez, R., Moriyón, I., Njeru, J., Díaz, R., Cook, E. A. J., Morin, M., Bronsvort, B. M. D. C., Thomas, L. F., Kariuki, S., & Fèvre, E. M. (2017). Poor performance of the rapid test for human brucellosis in health facilities in Kenya. *PLoS Neglected Tropical Diseases*, 11(4), 1–15. <https://doi.org/10.1371/journal.pntd.0005508>
- Desiere, S., Niragira, S., & D'Haese, M. (2015). Cow or goat? Population pressure and livestock keeping in Burundi. *Agrekon*, 54(3), 23–42. <https://doi.org/10.1080/03031853.2015.1084941>
- Dubad, A. B., Baluka, S. A., & Kaneene, J. B. (2015). Sero-prevalence of brucellosis in small ruminants in Kiruhura District of Uganda. *Advance Tropical Medicine and Public Health International*, 5(3), 62–76.

- Ducrottoy, M., Bertu, W. J., Matope, G., Cadmus, S., Conde-Álvarez, R., Gusi, A. M., Welburn, S., Ocholi, R., Blasco, J. M., & Moriyón, I. (2015). Brucellosis in Sub-Saharan Africa: Current challenges for management, diagnosis and control. *Acta Tropica*, *165*, 179–193. <https://doi.org/10.1016/j.actatropica.2015.10.023>
- Egeru, A., Wasonga, O., Kyagulanyi, J., Majaliwa, G. M., MacOpiyo, L., & Mburu, J. (2014). Spatio-temporal dynamics of forage and land cover changes in Karamoja sub-region, Uganda. *Pastoralism: Research, Policy and Practice*, *4*(1), 1–21. <https://doi.org/10.1186/2041-7136-4-6>
- Emmanuel, T. A. J., Tijjani, K. I., & Çakır, A. (2018). Challenges and possible improvement of livestock sector in south Sudan: Review paper. *International Journal of Research-Granthaalayah*, *6*(2), 214–223. <https://doi.org/10.5281/zenodo.1194652>
- Engida, E., Guthiga, P., & Karugia, J. (2015). The Role of Livestock in the Tanzanian Economy: Policy Analysis Using a Dynamic Computable General Equilibrium Model for Tanzania) ** Senior Policy Analyst, The Regional Strategy Analysis and Knowledge Support System for Eastern and Central Africa ** *Coor. Agriculture in an Interconnected World*, 22.
- Enström, S., Nthiwa, D., Bett, B., Karlsson, A., Alonso, S., & Lindahl, J. F. (2017). Brucella seroprevalence in cattle near a wildlife reserve in Kenya. *BMC Research Notes*, *10*(1), 1–5. <https://doi.org/10.1186/s13104-017-2941-x>
- Erume, J., Roesel, K., Dione, M. M., Ejobi, F., Mboowa, G., Kungu, J. M., Akol, J., Pezo, D., El-Adawy, H., Melzer, F., Elschner, M., Neubauer, H., & Grace, D. (2016). Serological and molecular investigation for brucellosis in swine in selected districts of Uganda. *Tropical Animal Health and Production*, *48*(6), 1147–1155. <https://doi.org/10.1007/s11250-016-1067-9>
- Ezama, A., Gonzalez, J. P., Majaliwa, S., & Bajunirwe, F. (2018). Assessing short evolution brucellosis in a highly brucella endemic cattle keeping population of Western Uganda: A complementary use of Rose Bengal test and IgM rapid diagnostic test. *BMC Public Health*, *18*(1), 1–5. <https://doi.org/10.1186/s12889-018-5228-9>
- Ezama, A., Gonzalez, J., Sebhatu, T., Gabriel, T., & Majaliwa, S. (2019). Presumptive diagnosis of brucellosis and determination of risk factors for seropositivity among members of cattle keeping households in a high cattle traffic area in the South Western region of Uganda. *Global Journal of Infectious Diseases and Clinical Research*, *5*(1), 16–24. <https://doi.org/10.17352/2455-5363.000024>
- Fèvre, E. M., De Glanville, W. A., Thomas, L. F., Cook, E. A. J., Kariuki, S., & Wamae, C. N. (2017). An integrated study of human and animal infectious disease in the Lake Victoria crescent small-holder crop-livestock production system, Kenya. *BMC Infectious Diseases*, *17*(1), 1–14. <https://doi.org/10.1186/s12879-017-2559-6>
- Food and Agriculture Organization of United Nations. (n.d.). *FAOSTAT statistical database*. FAO, c1997-. <http://www.fao.org/faostat/en/#data/QA>
- Food and Agriculture Organization of United Nations. (2009). *Brucella melitensis* in Eurasia and the Middle East. In *Animal Production and Health* (Issue May). <http://www.fao.org/docrep/012/i1402e/i1402e00.pdf>
- Food and Agriculture Organization of United Nations. (2014). FAO works to curb the burden of brucellosis in endemic countries Case studies from Eurasia and the Near East. In *FAO Empres* (Issue 8).
- Food and Agriculture Organization of United Nations. (2016). *The Impact of conflict on the livestock sector in south Sudan*. <http://files/1298/Gebreyes-CTheImpactofConflictontheLivestockSector.pdf>
- Food and Agriculture Organization of United Nations (2018a). Livestock production systems spotlight-Kenya-Cattle and poultry sectors. FAO. [https://doi.org/10.1016/s0301-6226\(97\)89729-1](https://doi.org/10.1016/s0301-6226(97)89729-1)
- Food and Agriculture Organization of United Nations. (2018b). The monetary impact of zoonotic diseases on society: Uganda. In *African Sustainable Livestock 2050*.
- Food and Agriculture Organization of United Nations. (2019). *The future of livestock in Uganda-Opportunities and challenges in the face of uncertainty*. Food and Agriculture Organization of the United Nations. Retrieved from <http://www.fao.org/3/ca5420en/CA5420EN.pdf>
- Food and Drug Administration. (2012). Bad bug book: Handbook of foodborne pathogenic microorganisms and natural toxins. In *U.S. Department of Health and Human Services* (2nd ed.). [https://doi.org/10.1016/S1872-2040\(10\)60451-3](https://doi.org/10.1016/S1872-2040(10)60451-3)
- Franc, K. A., Krecek, R. C., Häsler, B. N., & Arenas-Gamboa, A. M. (2018). Brucellosis remains a neglected disease in the developing world: A call for interdisciplinary action. *BMC Public Health*, *18*(1), 1–9. <https://doi.org/10.1186/s12889-017-5016-y>
- Frank, S., Benson, O., & Ivan, T. (2017). Human Brucellosis in gazetted forest areas: A case of bwindi impenetrable. *Journal of Foodborne and Zoonotic Diseases*, *5*(1), 7–11.
- Gafirita, J., Kiiza, G., Murekatete, A., Ndahayo, L. L., Tuyisenge, J., Mashengesho, V., Ruhirwa, R., Nyandwi, T., Asiimwe-Kateera, B., Ndahindwa, V., & Njunwa, K. J. (2017). Seroprevalence of brucellosis among patients attending a District Hospital in Rwanda. *American Journal of Tropical Medicine and Hygiene*, *97*(3), 831–835. <https://doi.org/10.4269/ajtmh.16-0632>
- Gicheru, M. N., Mwangi, E., & Mbaire, M. R. (2015). Prevalence and knowledge of Brucellosis in dairy cattle in Makuyu Division, Murang'a County, Kenya. *International Journal of Scientific Engineering and Technology*, *4*(12), 549–555. <https://doi.org/10.17950/ijset/v4s12/1202>
- Godfroid, J., Debolle, X., Roop, R. M., O'callaghan, D., Tsolis, R. M., Baldwin, C. J., Santos, R. L., Mcgiven, J. A., Olsen, S. C., Nymo, I. H., Larsen, A., Al dahouk, S., & Letesson, J. J. (2014). The quest for a true One Health perspective of brucellosis. *OIE Revue Scientifique et Technique*, *33*(2), 521–538. <https://doi.org/10.20506/rst.33.2.2290>
- Hamdy, M. E. R., & Amin, A. S. (2002). Detection of Brucella species in the milk of infected cattle, sheep, goats and camels by PCR. *The Veterinary Journal*, *163*(3), 299–305. <https://doi.org/10.1053/tvjl.2001.0681>
- Hoffman, T., Rock, K., Mugizi, D. R., Murdrasoli, S., Lindahl-Rajala, E., Erume, J., Magnusson, U., Lundkvist, Å., & Boqvist, S. (2016). Molecular detection and characterization of Brucella species in raw informally marketed milk from Uganda. *Infection Ecology & Epidemiology*, *6*(1), 32442. <https://doi.org/10.3402/iee.v6.32442>
- International Livestock Research Institute. (2012). Mapping of poverty and likely zoonoses hotspots. *Zoonoses Report*, *4*, 1–119. <http://www.ilri.org/ilrinews/index.php/archives/9172>
- Jeníček, V., & Grofová, Š. (2015). Least developed countries – The case of Burundi. *Agricultural Economics*, *61*(5), 234–247. <https://doi.org/10.17221/48/2014-AGRICECON>
- Kabi, F., Muwanika, V., & Masembe, C. (2015). Spatial distribution of Brucella antibodies with reference to indigenous cattle populations among contrasting agro-ecological zones of Uganda. *Preventive Veterinary Medicine*, *121*(1–2), 56–63. <https://doi.org/10.1016/j.prevetmed.2015.06.007>
- Kairu-wanyoike, S., Nyamwaya, D., Id, M. W., Lindahl, J., Ontiri, E., Bukachi, S., Njeru, I., Karanja, J., Sang, R., Grace, D., & Id, B. B. (2019). Positive association between Brucella spp. seroprevalences in livestock and humans from a cross-sectional study in Garissa and Tana River Counties, Kenya. *PLoS Neglected Tropical Diseases*, 1–15.
- Kamwine, M., Orikiriza, P., Taseera, K., Iramiot, J. S., Ojuka, P., Ikiriza, S., Atwebembeire, J., Otieno, D., Tweshengyereze, S., Mwanga-Amumpaire, J., Bazira, J., & Boum, Y. (2017). Prevalence of antibodies to Brucella species in commercial raw bovine milk in Southwestern Uganda. *BMC Research Notes*, *10*(1), 1–5. <https://doi.org/10.1186/s13104-017-2537-5>
- Kansiime, C., Rutebemberwa, E., Asiimwe, B. B., Makumbi, F., Bazira, J., & Mugisha, A. (2015). Annual trends of human brucellosis in pastoralist communities of south-western Uganda: A retrospective

- ten-year study. *Infectious Diseases of Poverty*, 4(1), 1–8. <https://doi.org/10.1186/s40249-015-0072-y>
- Kashiwazaki, Y., Ecewu, E., Imaligat, J. O., Mawejje, R., Kirunda, M., Kato, M., Musoke, G. M., & Ademun, R. A. O. (2012). Epidemiology of bovine brucellosis by a combination of rose Bengal test and indirect ELISA in the five districts of Uganda. *Journal of Veterinary Medical Science*, 74(11), 1417–1422. <https://doi.org/10.1292/jvms.12-0164>
- Kaur, P., Sharma, N. S., Arora, A. K., & Deepti. (2018). Investigation of brucellosis in cattle and buffaloes by conventional and molecular assays. *Indian Journal of Animal Research*, 52(10), 1482–1487. <https://doi.org/10.18805/ijar.B-3375>
- Kayombo, G., Makingi, G., Nonga, H., Misinzo, G., & Kazwala, R. (2017). Studies of brucellosis in lactating cows in Babati district, Tanzania. *Tanzania Veterinary Journal*, 35(1), 90–101.
- Kenya Ministry of Agriculture, Livestock, F. and I. (2019). *Draft national livestock policy*. Kenya Ministry of Agriculture, Livestock, Fisheries and Irrigation - State Department for Livestock Draft. Retrieved from <https://www.kilimo.go.ke/wp-content/uploads/2019/02/Draftreviewed-National-Livestock-Policy-February-2019.pdf>
- Kenya National Bureau of Statistics. (2019). *2019 Kenya Population and Housing Census. Volume IV: Distribution of population by social-economic characteristics*. Kenya National Bureau of Statistics. Retrieved from <https://africacheck.org/wp-content/uploads/2020/06/VOLUME-IVKPHC-2019.pdf>
- Khan, M. Y., Mah, M. W., & Memish, Z. A. (2001). Brucellosis in Pregnant Women. *Clinical Infectious Diseases*, 32(8), 1172–1177. <https://doi.org/10.1086/319758>
- Kosgei, P. K., Beboru, L., Waiboci, L., Kitale, P., & Kiambi, S. (2014). Estimating prevalence of brucellosis in livestock and assessment of knowledge, attitudes and practices of respective communities in Baringo county, Kenya. *Fourth RUFORUM Biennial Regional Conference*, 21(July), 297–301. <http://www.ruforum.org/sites/default/files/Kosgei.pdf>
- Land O' lakes, I. (2014). *Baseline survey report for Rwanda Dairy Competitiveness Program (RDGP)*.
- Langoni, H., Ichihara, S. M., da Silva, A. V., Pardo, R. B., Tonin, F. B., Mendonca, L. J. P., & Machado, J. A. D. (2000). Isolation of brucella spp from milk of brucellosis positive cows in São Paulo and Minas Gerais states. *Brazilian Journal of Veterinary Research and Animal Science*, 37(6), <https://doi.org/10.1590/s1413-95962000000600004>
- Lita, E., Nasinyama, G. W., Ochi, E. B., James, B., & Erume, J. (2016). Seroprevalence and risk factors associated with bovine Brucellosis in Central. 3(7), 454–462. <https://doi.org/10.21276/sjavs.2016.3.7.3>
- Lolli, C., Marenzoni, M. L., Strona, P., Lappo, P. G., Etiang, P., & Diverio, S. (2016). Infections and risk factors for livestock with species of Anaplasma, Babesia and Brucella under semi-nomadic rearing in Karamoja Region, Uganda. *Tropical Animal Health and Production*, 48(3), 603–611. <https://doi.org/10.1007/s11250-016-1005-x>
- Lopes, B. L., Nicolino, R., & P.A. Haddad, J. (2010). Brucellosis – Risk factors and prevalence: A review. *The Open Veterinary Science Journal*, 4(1), 72–84. <https://doi.org/10.2174/1874318801004010072>
- Madut, N. A., Muleme, J., Kankya, C., Nasinyama, G. W., Muma, J. B., Godfroid, J., Jubara, A. S., & Muwonge, A. (2019). Sero-prevalence of brucellosis among slaughterhouse workers in Bahr el Ghazal region, South Sudan. *BMC Infectious Diseases*, 7(JUN), 1–7. <https://doi.org/10.3389/fpubh.2019.00156>
- Madut, N. A., Muwonge, A., Nasinyama, G. W., Muma, J. B., Godfroid, J., Jubara, A. S., Muleme, J., & Kankya, C. (2018). The sero-prevalence of brucellosis in cattle and their herders in Bahr el Ghazal region, South Sudan. *PLoS Neglected Tropical Diseases*, 12(6), 1–14. <https://doi.org/10.1371/journal.pntd.0006456>
- Madut, N. A., Nasinyama, G. W., Muma, J. B., Sube, K. L. L., Ocan, M., Muwonge, A., Godfroid, J., Jubara, A. S., & Kankya, C. (2018). Prevalence of brucellosis among patients attending Wau Hospital, South Sudan. *PLoS One*, 13(6), 1–12. <https://doi.org/10.1371/journal.pone.0199315>
- Maiyo, G., & Obey, J. K. (2016). Distribution and prevalence of human Brucellosis among patients reporting at Chemundu Dispensary, Nandi County, Kenya. *Baraton Interdisciplinary Research Journal*, 6, 73–82. <http://ueab.ac.ke/BIRJ/wp-content/uploads/2016/10/Article12.pdf>
- Majalija, S., Luyombo, P., & Tumwine, G. (2018). Sero-prevalence and associated risk factors of Brucellosis among Malaria negative febrile out-patients in Wakiso district, Central Uganda. *BMC Research Notes*, 11(1), 1–6. <https://doi.org/10.1186/s13104-018-3907-3>
- Makita, K., Fevre, E. M., Waiswa, C., Eisler, M. C., & Welburn, S. C. (2010). How human brucellosis incidence in urban Kampala can be reduced most efficiently? A stochastic risk assessment of informally-marketed milk. *PLoS One*, 5(12), <https://doi.org/10.1371/journal.pone.0014188>
- Makita, K., Waiswa, C., & Mthrusfieldedacuk, M. T. (2011). Herd prevalence of bovine brucellosis and analysis of risk factors in cattle in urban and peri-urban areas of. *BMC Veterinary Research*. <https://doi.org/10.1186/1746-6148-7-60>
- Manirakiza, J., Hatungumukama, G., Besbes, B., & Detilleux, J. (2020). Characteristics of smallholders' goat production systems and effect of Boer crossbreeding on body measurements of goats in Burundi. *Pastoralism*, 10(1), <https://doi.org/10.1186/s13570-019-0157-5>
- Manirakiza, J., Hatungumukama, G., Thévenon, S., Gautier, M., Besbes, B., Flori, L., & Detilleux, J. (2017). Effect of genetic European taurine ancestry on milk yield of Ankole-Holstein crossbred dairy cattle in mixed smallholders system of Burundi highlands. *Animal Genetics*, 48(5), 544–550. <https://doi.org/10.1111/age.12578>
- Manishimwe, R., Ntaganda, J., Habimana, R., Nishimwe, K., Byukusenge, M., Dutuze, F., Ayabagabo, J. D., And, U. L., & Rukundo, J. C. (2015). Comparison between rose Bengal plat test and competitive enzyme linked immunosorbent assay to detect bovine Brucellosis in Kigali City, Rwanda. *Journal of Veterinary Science & Technology*, 06(01), 2–5. <https://doi.org/10.4172/2157-7579.1000211>
- Mantur, B., Amarnath, S., & Shinde, R. (2007). Review of clinical and laboratory features of human brucellosis. *Indian Journal of Medical Microbiology*, 25(3), 188–202. <https://doi.org/10.4103/0255-0857.34758>
- Mathew, C. (2017). Reproductive Infections in Cattle in Tanzania – Lessons for Control Priorities. *SOJ Microbiology & Infectious Diseases*, 5(2), 1–9. <https://doi.org/10.15226/sojmid/5/2/00169>
- Mathew, C., Stokstad, M., Johansen, T. B., Klevar, S., Mdegela, R. H., Mwamengele, G., Michel, P., Escobar, L., Fretin, D., & Godfroid, J. (2015). First isolation, identification, phenotypic and genotypic characterization of Brucella abortus biovar 3 from dairy cattle in Tanzania. *BMC Veterinary Research*, 11(1), 1–9. <https://doi.org/10.1186/s12917-015-0476-8>
- McDermott, J. J., & Arimi, S. M. (2002). Brucellosis in sub-Saharan Africa: Epidemiology, control and impact. *Veterinary Microbiology*, 90(1–4), 111–134. [https://doi.org/10.1016/S0378-1135\(02\)00249-3](https://doi.org/10.1016/S0378-1135(02)00249-3)
- McDermott, J. J., Grace, D., & Zinsstag, J. (2013). Economics of brucellosis impact and control in low-income countries. *Scientific and Technical Review of the Office International Des Epizooties*, 32(1), 249–261. <https://doi.org/10.20506/rst.32.1.2197>
- Merker, M., & Schlichting, H. (1984). Note sur la brucellose au Burundi. *Revue D'elevage et de Medecine Veterinaire Des Pays Tropicaux*, 138–144.
- Migisha, R., Nyehangane, D., Boum, Y., Page, A. L., Zúñiga-Ripa, A., Conde-Álvarez, R., Bagenda, F., & Bonnet, M. (2018). Prevalence and risk factors of brucellosis among febrile patients attending a community hospital in south western Uganda. *Scientific Reports*, 8(1), 1–8. <https://doi.org/10.1038/s41598-018-33915-9>
- Miller, R., Nakavuma, J. L., Ssajjakambwe, P., Vudriko, P., Musisi, N., & Kaneene, J. B. (2016). The prevalence of Brucellosis in cattle, goats

- and humans in rural Uganda: A comparative study. *Transboundary and Emerging Diseases*, 63(6), e197–e210. <https://doi.org/10.1111/tbed.12332>
- Mirambo, M. M., Mgone, G. F., Malima, Z. O., John, M., Mngumi, E. B., Mhamphi, G. G., & Mshana, S. E. (2018). Seropositivity of *Brucella* spp. and *Leptospira* spp. antibodies among abattoir workers and meat vendors in the city of Mwanza, Tanzania: A call for one health approach control strategies. *PLoS Neglected Tropical Diseases*, 12(6), 39–52. <https://doi.org/10.1371/journal.pntd.0006600>
- Mngumi, E. B., Mirambo, M. M., Wilson, S., & Mshana, S. E. (2016). Predictors of specific anti-*Brucella* antibodies among humans in agro-pastoral communities in Sengerema district, Mwanza, Tanzania: The need for public awareness. *Tropical Medicine and Health*, 44(1), 5–10. <https://doi.org/10.1186/s41182-016-0034-5>
- Moreno, E. (2014). Retrospective and prospective perspectives on zoonotic brucellosis. *Frontiers in Microbiology*, 5(MAY), 1–18. <https://doi.org/10.3389/fmicb.2014.00213>
- Mugizi, D. R., Boqvist, S., Nasinyama, G. W., Waiswa, C., Ikwapa, K., Rock, K., Lindahl, E., Magnusson, U., & Erume, J. (2015). Prevalence of and factors associated with *Brucella* sero-positivity in cattle in urban and peri-urban Gulu and Soroti towns of Uganda. *Journal of Veterinary Medical Science*, 77(5), 557–564. <https://doi.org/10.1292/jvms.14-0452>
- Mugizi, D. R., Muradrasoli, S., Boqvist, S., Erume, J., Nasinyama, G. W., Waiswa, C., Mboowa, G., Klint, M., & Magnusson, U. (2015). Isolation and molecular characterization of *Brucella* isolates in cattle milk in Uganda. *BioMed Research International*, 2015, <https://doi.org/10.1155/2015/720413>
- Muhairwa, A. P., Mwakijungu, E. O., Msoffe, P. L. M., & Mtambo, M. M. A. (2012). Seroprevalence and factors affecting canine monocytic ehrlichiosis and canine brucellosis in Tanzania. *Research Opinions in Animal & Veterinary Sciences*, 0343.
- Mujuni, F., Andrew, V., Mngumi, E. B., Chibwe, E., Mshana, S. E., & Mirambo, M. M. (2018). Predominance of *Brucella abortus* antibodies among women with spontaneous abortion in the city of Mwanza: Unrecognized link or coincidence? *BMC Research Notes*, 1–5, <https://doi.org/10.1186/s13104-018-3906-4>
- Muloki, H. N., Erume, J., Owiny, D. O., Kungu, J. M., Nakavuma, J., Ogeng, D., & Nasinyama, G. W. (2018). Prevalence and risk factors for brucellosis in prolonged fever patients in post-conflict Northern Uganda. *African Health Sciences*, 18(1), 22–28. <https://doi.org/10.4314/ahs.v18i1.4>
- Musallam, I. I., Abo-Shehadeh, M. N., Hegazy, Y. M., Holt, H. R., & Guitian, F. J. (2016). Systematic review of brucellosis in the Middle East: Disease frequency in ruminants and humans and risk factors for human infection. *Epidemiology and Infection*, 144(4), 671–685. <https://doi.org/10.1017/S0950268815002575>
- Musallam, I., Ndour, A. P., Yempabou, D., Ngong, C.-A., Dzousse, M. F., Mouiche-Mouliom, M.-M., Feussom, J. M. K., Ntirandekura, J. B., Ntakirutimana, D., Fane, A., Dembele, E., Doumbia, A., Ayih-Akakpo, A.-H., Pato, P., Pali, M., Tapsoba, A. S. R., Compaore, G. M., Gagara, H., Garba, A. I., ... Guitian, J. (2019). Brucellosis in dairy herds: A public health concern in the milk supply chains of West and Central Africa. *Acta Tropica*, 197(March), 105042. <https://doi.org/10.1016/j.actatropica.2019.105042>
- Nabukenya, I., Kaddu-Mulindwa, D., & Nasinyama, G. W. (2013). Survey of *Brucella* infection and malaria among Abattoir workers in Kampala and Mbarara Districts, Uganda. *BMC Public Health*, 13(1), 2–7. <https://doi.org/10.1186/1471-2458-13-901>
- Nakkal, M. J., Arimi, S. M., Kitale, P. K., Nduhii, G., Njenga, J. M., & Wabacha, J. (2016). A sero-epidemiological survey of Brucellosis, Q-fever and Leptospirosis in livestock and humans and associated risk factors in Kajiado County- Kenya. *Journal of Tropical Diseases*, 4(3), <https://doi.org/10.4172/2329-891X.1000215>
- Nanfuka, M. (2018). Sero-prevalence of bovine and human brucellosis on selected farms in South-western Uganda. *Online Journal of Public Health Informatics*, 10(1), 2579. <https://doi.org/10.5210/ojphi.v10i1.8979>
- Nasinyama, G., Ssekawojwa, E., Opuda, J., Grimaud, P., Etter, E., & Bellinguez, A. (2014). *Brucella* sero-prevalence and modifiable risk factors among predisposed cattle keepers and consumers of un-pasteurized milk in Mbarara and Kampala districts, Uganda. *African Health Sciences*, 14(4), 790–796. <https://doi.org/10.4314/ahs.v14i4.3>
- National Institute of statistics of Rwanda. (2018). *Agricultural household survey 2016/2017*. <http://www.statistics.gov.rw/agriculture/Publications/AHS/AHS2016-17Report.pdf>
- National Institute of Statistics of Rwanda. (2019). *Statistical YearBook*.
- Ndazigaruye, G., Mushonga, B., Kandiwa, E., Samkange, A., Segwagwe, B. E., Segwagwe, B., & Province, E. (2018). Prevalence and risk factors for brucellosis seropositivity in cattle in Nyagatare District, Eastern Province, Rwanda. *Journal of the South African Veterinary Association*, 89, 1–8. <https://doi.org/10.4102/jsava.v89i0.1625>
- Nguna, J., Dione, M., Apamaku, M., Majalija, S., Mugizi, D. R., Odoch, T., Kato, C. D., Tumwine, G., Kabaasa, J. D., Curtis, K., Graham, M., Ejobi, F., & Graham, T. (2019). Seroprevalence of brucellosis and risk factors associated with its seropositivity in cattle, goats and humans in Iganga district, Uganda. *Pan African Medical Journal*, 33, 1–10. <https://doi.org/10.11604/pamj.2019.33.99.16960>
- Nina, P. M., Mugisha, S., Leirs, H., Basuta, G. I., & Van Damme, P. (2017). Brucellosis in cattle and micro-scale spatial variability of pastoral household income from dairy production in south western Uganda. *Acta Tropica*, 175, 130–137. <https://doi.org/10.1016/j.actatropica.2016.11.030>
- Nizeyimana, G., Mwiine, F. N., & Ayebazibwe, C. (2013). Comparative *Brucella abortus* antibody prevalence in cattle under contrasting husbandry practices in Uganda. *Journal of the South African Veterinary Association*, 84(1), 1–5. <https://doi.org/10.4102/jsava.v84i1.943>
- Njeru, J., Melzer, F., Wareth, G., El-Adawy, H., Henning, K., Pletz, M. W., Heller, R., Kariuki, S., Fèvre, E., & Neubauer, H. (2016). Human brucellosis in febrile patients seeking treatment at remote hospitals, north-eastern Kenya, 2014–2015. *Emerging Infectious Diseases*, 22(12), 2160–2164. <https://doi.org/10.3201/eid2212.160285>
- Njeru, J., Wareth, G., Melzer, F., Henning, K., Pletz, M. W., Heller, R., & Neubauer, H. (2016). Systematic review of brucellosis in Kenya: Disease frequency in humans and animals and risk factors for human infection. *BMC Public Health*, 16(1), 1–15. <https://doi.org/10.1186/s12889-016-3532-9>
- Njuguna, J. N., Gicheru, M. M., Kamau, L. M., & Mbatha, P. M. (2017). Incidence and knowledge of bovine brucellosis in Kahuro district, Murang'a County, Kenya. *Tropical Animal Health and Production*, 49(5), 1035–1040. <https://doi.org/10.1007/s11250-017-1296-6>
- Nonga, H. E., & Mwakapeje, E. R. (2017). Trends of human brucellosis in pastoralist communities based on hospital records during 2013–2016 in Ngorongoro District, Tanzania. *Tanzania Veterinary Association Proceedings*, 34, 34–40. <http://www.suaire.sua.ac.tz:8080/xmlui/bitstream/handle/123456789/2536/NongaandMwakapeje.pdf?sequence=1&isAllowed=y>
- Okumu, T. A., John, N. M., Wabacha, J. K., Tsuma, V., & Vanleeuwen, J. (2019). Seroprevalence of antibodies for bovine viral diarrhoea virus, *Brucella abortus* and *Neospora caninum*, and their roles in the incidence of abortion/foetal loss in dairy cattle herds in Nakuru District, Kenya. *BMC Veterinary Research*, 15(1), 1–6. <https://doi.org/10.1186/s12917-019-1842-8>
- Onyango, D., Oyoko, G., Too, R., & Masake, R. (2015). *The contribution of livestock to the south Sudan economy* (Issue December).
- Osoro, E. M., Munyua, P., Omulo, S., Ogola, E., Ade, F., Mbatha, P., Mbabu, M., Ng'ang'a, Z., Kairu, S., Maritim, M., Thumbi, S. M., Bitek, A., Gaichugi, S., Rubin, C., Njenga, K., & Guerra, M. (2015). Strong association between human and animal brucella seropositivity in a linked study in Kenya, 2012–2013. *American Journal of Tropical Medicine and Hygiene*, 93(2), 224–231. <https://doi.org/10.4269/ajtmh.15-0113>

- Pappas, G., Papadimitriou, P., Akritidis, N., Christou, L., & Tsianos, E. (2006). The new global map of human Brucellosis. *The Lancet Infectious Diseases*, 6, 91–99. [https://doi.org/10.1016/S1473-3099\(06\)70382-6](https://doi.org/10.1016/S1473-3099(06)70382-6)
- Perez-Sancho, M., Garci-Seco, T., Dominguez, L., & Alvarez, J. (2015). Control of animal Brucellosis – The most effective tool to prevent human Brucellosis. *Intech, Tourism*, 13. <https://doi.org/10.5772/57353>
- Pergher, G., & Noel, G. (1936). Note sur la Fievre Ondulante au Ruanda-Urundi. *Annales de La Societe Belge de Medecine Tropicale*.
- Rock, K. T., Mugizi, D. R., Ståhl, K., Magnusson, U., & Boqvist, S. (2016). The milk delivery chain and presence of *Brucella* spp. antibodies in bulk milk in Uganda. *Tropical Animal Health and Production*, 48(5), 985–994. <https://doi.org/10.1007/s11250-016-1052-3>
- Rujeni, N., & Mbanzamihiho, L. (2014). Prevalence of brucellosis among women presenting with abortion/stillbirth in Huye, Rwanda. *Journal of Tropical Medicine*, 2014, 3–5. <https://doi.org/10.1155/2014/740479>
- Sagamiko, F. D., Muma, J. B., Karimuribo, E. D., Mwanza, A. M., Sindato, C., & Hang'ombe, B. M. (2018). Sero-prevalence of Bovine Brucellosis and associated risk factors in mbeya region, Southern highlands of Tanzania. *Acta Tropica*, 178, 169–175. <https://doi.org/10.1016/j.actatropica.2017.11.022>
- Schmutz, S. M., Moker, J. S., Clark, E. G., & Orr, J. P. (1996). Chromosomal aneuploidy associated with spontaneous abortions and neonatal losses in cattle. *Journal of Veterinary Diagnostic Investigation*, 8(1), 91–95. <https://doi.org/10.1177/104063879600800114>
- Shirima, G., Fitzpatrick, J., Kunda, J., Mfinanga, S., Kazwala, R., Kambarage, D., & Cleaveland, S. (2010). The role of livestock keeping in human brucellosis trends in livestock keeping communities in Tanzania. *Tanzania Journal of Health Research*, 12. <https://doi.org/10.4314/thrb.v12i3.51261>
- Shirima, G., & John, S. K. (2016). Prevalence of brucellosis in the human, livestock and wildlife interface areas of Serengeti. *Onderstepoort Journal of Veterinary Research*, 83(1). <https://doi.org/10.4102/ojvr.v83i1.1032>
- Shirima, G. M., Masola, S. N., Malangu, O. N., & Schumaker, B. A. (2014). Outbreak investigation and control case report of brucellosis: Experience from livestock research centre, Mpwawwa, Tanzania. *The Onderstepoort Journal of Veterinary Research*, 81(1). <https://doi.org/10.4102/ojvr.v81i1.818>
- Simon, C., Karimuribo, E., & Kessy, B. M. (2015). Seroprevalence study of swine brucellosis and knowledge of pig traders and dressers at slaughter facilities in temeke municipality of dar es salaam, Tanzania. *Livestock Research for Rural Development*, 27.
- Swai, E. S., Moshy, W., Mbise, E., Lutatina, J., & Bwanga, S. (2011). Disease and health conditions affecting camel production in pastoral and agro-pastoral communities of northern Tanzania. *Research Opinions in Animal and Veterinary Sciences*, 1(2), 83–88.
- Swai, E. S., & Schoonman, L. (2010). The use of rose Bengal plate test to assess cattle exposure to *Brucella* infection in traditional and smallholder dairy production systems of Tanga Region of Tanzania. *Veterinary Medicine International*, 2010, <https://doi.org/10.4061/2010/837950>
- Swai, E. S., & Schoonman, L. (2011). Microbial quality and associated health risks of raw milk marketed in the Tanga region of Tanzania. *Asian Pacific Journal of Tropical Biomedicine*, 1(3), 217–222. [https://doi.org/10.1016/S2221-1691\(11\)60030-0](https://doi.org/10.1016/S2221-1691(11)60030-0)
- Swai, E. S., & Schoonman, L. (2012). A survey of zoonotic diseases in trade cattle slaughtered at Tanga city abattoir: A cause of public health concern. *Asian Pacific Journal of Tropical Biomedicine*, 2(1), 55–60. [https://doi.org/10.1016/S2221-1691\(11\)60190-1](https://doi.org/10.1016/S2221-1691(11)60190-1)
- Tanzania National Bureau of Statistics. (2017). *Annual agriculture sample survey crop and livestock report, 2016/17*.
- Tanzania National Bureau of Statistics. (2019). *Tanzania in figures 2018*.
- Tekle, M., Legesse, M., Edao, B. M., Ameni, G., & Mamo, G. (2019). Isolation and identification of *Brucella melitensis* using bacteriological and molecular tools from aborted goats in the Afar region of north-eastern Ethiopia. *BMC Microbiology*, 19(1), 1–6. <https://doi.org/10.1186/s12866-019-1474-y>
- Thienpont, D., Vandervelden, M., Fagard, P., & Mortelmans, J. (1961). L'hygroma brucellique : l'aspect clinique caractéristique de la brucellose bovine au Rwanda-Burundi. *Revue d'élevage et de Médecine Veterinaire Des Pays Tropicaux*, 1956.
- Tumwine, G., Matovu, E., Kabasa, J. D., Owiny, D. O., & Majalija, S. (2015). Human brucellosis: Sero-prevalence and associated risk factors in agro-pastoral communities of Kiboga District, Central Uganda. *BMC Public Health*, 15(1), 1–8. <https://doi.org/10.1186/s12889-015-2242-z>
- Uganda Bureau Of Statistics. (2018). *Statistical abstract*.
- United Nations Economic Commission for Africa. (2013). South Sudan-country profile 2017. *Africa yearbook* (Vol. 9). Author. https://doi.org/10.1007/978-3-319-01384-8_672
- United States Agency for International Development (USAID). (2009). Livelihoods Zoning “Plus” Activity in Burundi, A special Report by The Famine Early Warning System Network (FEWS NET), November 2009. https://fews.net/sites/default/files/documents/reports/bi_zonedescriptions_en.pdf
- World Health Organization. (2012). Research priorities for zoonoses and marginalized infections. *World Health Organization Technical Report Series*, 971.
- Yang, H. X., Feng, J. J., Zhang, Q. X., Hao, R. E., Yao, S. X., Zhao, R., Piao, D. R., Cui, B. Y., & Jiang, H. (2018). A case report of spontaneous abortion caused by *Brucella melitensis* biovar 3. *Infectious Diseases of Poverty*, 7(1), 1–4. <https://doi.org/10.1186/s40249-018-0411-x>
- Zamri-Saad, M., & Kamarudin, M. I. (2016). Control of animal brucellosis: The Malaysian experience. *Asian Pacific Journal of Tropical Medicine*, 9(12), 1136–1140. <https://doi.org/10.1016/j.apjtm.2016.11.007>

How to cite this article: Djangwani J, Ooko Abong' G, Gicuku Njue L, Kaindi DWM. Brucellosis: Prevalence with reference to East African community countries – A rapid review. *Vet Med Sci*. 2021;7:851–867. <https://doi.org/10.1002/vms3.425>