



Published in final edited form as:

*Neuropharmacology*. 2020 June 15; 170: 108053. doi:10.1016/j.neuropharm.2020.108053.

## **$\alpha$ 7 nicotinic acetylcholine receptors as therapeutic targets in schizophrenia: update on animal and clinical studies and strategies for the future**

**Alvin V. Terry Jr.<sup>1,2</sup>, Patrick M. Callahan<sup>1,2</sup>**

<sup>1</sup>Department of Pharmacology and Toxicology, Medical College of Georgia, Augusta University, Augusta, Georgia, 30912.

<sup>2</sup>Small Animal Behavior Core, Medical College of Georgia, Augusta University, Augusta, Georgia, 30912

### **Abstract**

Schizophrenia is a devastating mental illness and its effective treatment is among the most challenging issues in psychiatry. The symptoms of schizophrenia are heterogeneous ranging from positive symptoms (e.g., delusions, hallucinations) to negative symptoms (e.g., anhedonia, social withdrawal) to cognitive dysfunction. Antipsychotics are effective at ameliorating positive symptoms in some patients; however, they are not reliably effective at improving the negative symptoms or cognitive impairments. The inability to address the cognitive impairments is a particular concern since they have the greatest long-term impact on functional outcomes. While decades of research have been devoted to the development of pro-cognitive agents for schizophrenia, to date, no drug has been approved for clinical use. Converging behavioral, neurobiological, and genetic evidence led to the identification of the  $\alpha$ 7-nicotinic acetylcholine receptor ( $\alpha$ 7-nAChR) as a therapeutic target several years ago and there is now extensive preclinical evidence that  $\alpha$ 7-nAChR ligands have pro-cognitive effects and other properties that should be beneficial to schizophrenia patients. However, like the other pro-cognitive strategies, no  $\alpha$ 7-nAChR ligand has been approved for clinical use in schizophrenia thus far. In this review, several topics are discussed that may impact the success of  $\alpha$ 7-nAChR ligands as pro-cognitive agents for schizophrenia including the translational value of the animal models used, clinical trial design limitations, confounding effects of polypharmacy, dose-effect relationships, and chronic versus intermittent dosing considerations. Determining the most optimal pharmacologic strategy at  $\alpha$ 7-nAChRs: agonist, positive allosteric modulator, or potentially even receptor antagonist is also discussed.

---

**Corresponding Author:** Alvin V. Terry Jr., Ph.D. Department of Pharmacology & Toxicology, 1120 15th Street, CB-3545, Augusta University, Augusta, Georgia 30912, Phone: 706-721-9462, Fax: 706-721-2347, aterry@augusta.edu.  
Credit Author Statement

Alvin V. Terry Jr.: Conceptualization, Writing, Reviewing, and Editing

Patrick M. Callahan: Conceptualization, Writing, Reviewing, and Editing

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of interest: The authors do not declare any conflict of interest.

## Keywords

Cholinergic; pro-cognitive; psychosis; cognition; executive function; attention

---

## 1. Introduction

Schizophrenia is a debilitating mental illness characterized by positive symptoms (e.g., hallucinations, delusions), negative symptoms (e.g., depressed mood, anhedonia, social withdrawal) and cognitive impairments (e.g., deficits in information processing, attention, working memory, executive function, Green and Braff, 2001). Among these diverse symptoms, cognitive impairment is a core feature of schizophrenia that often appears prior to the onset of psychotic symptoms, it persists throughout the course of the illness, and it has the greatest long-term impact on functional outcomes (reviewed, Kahn and Keefe, 2013; Green and Harvey, 2014, Kahn, 2019). Unfortunately, the most commonly prescribed treatments for schizophrenia, the antipsychotics, while effective at improving positive symptoms in some patients, are not reliably effective at improving cognitive function. This unmet medical need has been an important focus of drug discovery efforts in both academia and the pharmaceutical industry for several decades; however, to date no pro-cognitive agent has been approved for clinical use in schizophrenia.

A key challenge to developing novel treatments for the cognitive dysfunction in schizophrenia is the complex and poorly understood etiology and pathophysiology of the illness. Multiple neurotransmitter systems have been implicated in the illness (Goff and Wine, 1997; Kapur and Mamo, 2003) including dopaminergic, serotonergic, glutamatergic, adrenergic, and cholinergic pathways and accordingly, new compounds designed to target these various systems have been developed and evaluated. Cholinergic targets, particularly nicotinic acetylcholine receptors (nAChRs) have been a focus of a number of drug discovery programs over the last 20–25 years based on multiple lines of behavioral, neurobiological, and genetic evidence. From the behavioral perspective, a remarkable observation in schizophrenic patients is their especially heavy abuse of tobacco products. According to the National Institute on Drug Abuse (NIDA, 2020) tobacco smoking rates in schizophrenia patients range as high as 70–85%, which is dramatically higher than the general population (~19–20%) and significantly higher than in any other mental illness (George and Krystal, 2007). Smokers with schizophrenia have also been documented to extract more nicotine per cigarette and to smoke a higher number of cigarettes per day compared to smokers in the general population (Olincy et al., 1997; Strand and Nybäck, 2005). It has been suggested that this high level of nicotine consumption may represent an attempt of schizophrenia patients to self-medicate the cognitive symptoms, particularly deficits of information processing and attention (Olincy et al., 1997; Leonard et al., 2007).

## 2. Focus on $\alpha 7$ -nAChRs

In drug discovery programs for neuropsychiatric illnesses, the heteromeric  $\alpha 4\beta 2$  and homomeric  $\alpha 7$ -nAChRs have been the most commonly targeted nAChR subtypes to date since they are the most predominant subtypes found in the mammalian brain. However, the

$\alpha 7$ -nAChR has been more commonly targeted for the cognitive deficits in schizophrenia. This is likely based, in part, on several factors including postmortem evidence of  $\alpha 7$ -nAChR deficits in the frontal cortex and hippocampus of schizophrenic patients (Guan et al., 1999) and linkage analysis implicating chromosome 15q14 (the region that includes the  $\alpha 7$ -nAChR gene). Polymorphisms in the core promoter of the  $\alpha 7$ -nAChR gene (CHRNA7; GeneBank accession no. **Z23141**) have been associated with reduced inhibition of the P50 evoked response to repeated auditory stimuli in schizophrenic patients, which is indicative of sensory gating abnormalities (reviewed, Freedman et al., 2003).  $\alpha 7$ -nAChR deficits may also contribute to abnormalities of smooth pursuit eye movements, sustained attention, and other domains of cognition in schizophrenia (reviewed Martin et al., 2004). In addition to schizophrenia, the CHRNA7 gene is also linked to multiple disorders where cognitive deficits are present including bipolar disorder, autism spectrum disorders, attention deficit hyperactivity disorder, Alzheimer disease, epilepsy, and sensory processing deficits (reviewed in Corradi and Bouzat, 2016).

The information in the paragraph above regarding the importance of  $\alpha 7$ -nAChR as a potential therapeutic target for the cognitive defects in schizophrenia is also supported by extensive preclinical evidence. From a neurobiological and neuropharmacological perspective,  $\alpha 7$ -nAChRs modulate multiple (cognition-related) processes in neurons that are calcium-dependent including neurotransmitter release (McGehee et al., 1995; Gray et al., 1996), postsynaptic signaling (Chang and Berg, 1999; Hefft et al., 1999) and neuronal survival (Messi et al., 1997; Berger et al., 1998). Moreover,  $\alpha 7$ -nAChRs are also abundant in regions of the brain that are important for learning and memory and executive function such as the hippocampus and prefrontal cortex (Gotti et al., 2007). In addition, agonists of  $\alpha 7$ -nAChRs have been shown to increase the phosphorylation of ERK and CREB (signaling pathways linked to long-term potentiation and memory formation) in the rodent brain (Bitner et al., 2007, 2010). There is also extensive evidence that  $\alpha 7$ -nAChR ligands improve behavioral processes that are relevant to schizophrenia (see Fig 1) such as auditory-evoked gating and prepulse inhibition in rodents, as well as multiple domains of cognition including attention, working memory, reference memory, social cognition, and executive function in rodent models as well as non-human primates (see reviews, Young and Geyer, 2013; Freedman, 2014; Bertrand and Terry, 2018). Table 1 provides a list of representative  $\alpha 7$ -nAChRs ligands that have been developed as pro-cognitive agents for potential use in schizophrenia and other disorders of cognition. Although not all-inclusive, Table 1 includes  $\alpha 7$ -nAChR agonists, partial agonists, and positive allosteric modulators (PAMs), a summary of some of the positive behavioral properties associated with each compound, and representative references.

### 3. Clinical Trial Failures

Despite extensive preclinical evidence to support the pro-cognitive effects of  $\alpha 7$ -nAChR ligands noted above (and summarized in Table 1) and positive results in some early (Phase I and II) clinical trials, to date, no compound has met the primary objective of cognitive improvement in schizophrenic patients in a large phase III, double blind, placebo controlled clinical trial or unanticipated side effects emerged (see review, Tregellas and Wylie, 2019). The failure of pro-cognitive agents in schizophrenia clinical trials have reduced the

enthusiasm of pharmaceutical companies and, unfortunately, many have abandoned this line of research. While it is possible that the lack of robust (pro-cognitive) efficacy or side effect burden may represent real limitations of  $\alpha 7$ -nAChR ligands, it should be noted, that given the number of compounds that have been developed, only one agent have been evaluated in a large scale Phase III clinical trial in schizophrenia patients (Encenicline-EVP-6124, see Table 2). Moreover, there is an increasing amount of discussion about possible limitations of both the preclinical and clinical studies conducted to date that may have resulted in the so-called “treatment failures” of pro-cognitive agents. Some of these potential limitations are listed in Table 3. See also several recent reviews (Bertrand and Terry, 2018, Terry and Callahan, 2019; Tregellas and Wylie, 2019) on this subject.

#### 4. Translational Gaps and Overreliance on Rodent Models

The unfavorable results of clinical trials for pro-cognitive agents in schizophrenia described above have led to questions about the “translational validity” of animal models used in preclinical studies (see Lewis et al., 2018). In drug discovery research for schizophrenia and other neuropsychiatric disorders, addressing the translational elements, face, construct, and predictive validity in animal models is particularly challenging. In this context, multiple challenges include the subjective nature of many of the human symptoms, the lack of biomarkers and objective diagnostic tests, and our relatively poor understanding of the neurobiology and genetics of neuropsychiatric disorders (see review Nestler and Hyman, 2010; Monteggia et al., 2018). One of the goals of the Research Domain Criteria (RDoC) paradigm launched in 2010 by the National Institute of Mental Health (NIMH) was to improve translation between animal experiments and clinical studies in psychiatry research. The basic concept in the preclinical realm was to encourage basic scientists to identify molecular or neural mechanisms (or neural circuitry) that contributes to specific domains of a mental function rather than creating animal models of diseases. In the clinical realm, researchers were encouraged to conceptualize normal human behavior, emotion, and cognition as dimensional, with mental illnesses as dimensional extremes as opposed to being restricted by DSM diagnostic categories (see Morris and, Cuthbert, 2012; Ross and Margolis, 2019). However, more than 10 years after the introduction of the RDoC paradigm, it is unclear if it has improved research progress in psychiatry especially in the drug discovery arena and its clinical relevance is increasingly being questioned (see Carpenter, 2016; Ross and Margolis, 2019).

Regarding translational validity, rodent models are undoubtedly important in basic research for testing disease-related hypotheses and the early evaluations of novel therapeutic agents, however, it is likely they have been relied upon too much in neuropsychiatric drug discovery especially at the later preclinical stages of drug development. Compared to humans, the behavioral repertoire of rodents is quite limited and, while debated (see, Laubach et al., 2018), there are major anatomical differences in their brains, most notably, the development of cortical regions of the forebrain, particularly the dorsolateral prefrontal cortex (DLPFC). This portion of the brain of humans and more advanced non-human primates (e.g., macaques) has been implicated specifically in the most complex cognitive processes such as working memory, sustained attention, decision taking, and executive function (reviewed, Barbey et al., 2013).

Regarding,  $\alpha 7$ -nAChRs, there are also significant differences in the genetics, pharmacology, biophysical properties, and neuronal localization of  $\alpha 7$ -nAChRs between rodents and humans that could underlie different outcomes in preclinical and clinical trial evaluations of  $\alpha 7$ -nAChRs ligands (reviewed Bertrand and Terry, 2018). From a genetics standpoint, the recent findings of Yin et al., 2017 are particularly notable. In humans with 15q13.3 microdeletion syndrome, caused by heterozygous deletions involving the CHRNA7 gene, behavioral abnormalities often observed in neuropsychiatric conditions such as schizophrenia and autism were observed, whereas Chrna7 knockout mice did not exhibit similar neurobehavioral phenotypes. From a pharmacological perspective, the effects of specific agonists in vitro differ between human and rodent sequences coding for  $\alpha 7$ -nAChRs. For example, the partial agonist DMXB-A (GTS-21) activates the rat  $\alpha 7$ -nAChR to a maximal response greater than twice that of the human  $\alpha 7$ -nAChR, and the  $K_i$  of GTS-21 at the rat receptor is roughly an order of magnitude less than at the human receptor (Meyer et al., 1998), suggesting that similar serum levels might have disparate effects between the species. Finally, differences in the synaptic receptor expression between rodents and humans may be relevant as suggested by a recent study in nonhuman primates where postsynaptic localization of  $\alpha 7$ -nAChRs on spines were demonstrated in glutamatergic synapses of layer III dorsolateral prefrontal cortex (Yang et al., 2013). Most physiological studies in rodent frontal cortex, in contrast, have demonstrated presynaptic  $\alpha 7$ -nAChR actions and it has been suggested that spine  $\alpha 7$ -nAChRs are not prevalent or only have subtle effects on neuronal physiology in rodents compared to primates. Collectively, the information discussed here would appear to justify a renewed interest in the use of nonhuman primate species (see Monteggia et al., 2018) in neuropsychiatry and drug discovery research, given their richer behavioral repertoire and more homologous brain anatomy with humans compared to rodents.

## 5. Polypharmacy and Drug Exposure History

In both preclinical and clinical evaluations of  $\alpha 7$ -nAChR ligands, it is uncommon for the subject of polypharmacy and chronic antipsychotic drug history to be adequately addressed. Clearly, there are practical limitations in clinical trials, but the patient's treatment history (which in many cases consists of multiple years of antipsychotic treatment) should be more carefully considered, not just concomitant antipsychotic treatment at the time of the clinical trial. Antipsychotics are well documented to have a variety of chronic effects on the mammalian brain including alterations of neurotransmitter receptor expression and neural plasticity (reviewed, Morrison and Murray, 2018), i.e., effects that could influence the response to a novel  $\alpha 7$ -nAChR ligand. Interestingly, guidelines related to polypharmacy and concomitant drug exposure have been developed for studies designed to evaluate potential pro-cognitive agents in schizophrenia trials. For example, a workshop on clinical trial design for evaluating cognitive enhancing drugs for schizophrenia was held in 2004 and it included experts from the FDA, NIMH, and scientists from academia and the pharmaceutical industry (see Buchanan et al., 2005). Among the various guidelines developed, it was recommended that polypharmacy (treatment with multiple antipsychotics) and combining a putative cognitive-enhancing agent with an antipsychotic with high affinity for the targeted receptor be avoided. However, it is unclear how closely such policies have been followed.

In preclinical evaluations of potential pro-cognitive agents for potential use in schizophrenia, the concomitant administration of antipsychotics has only rarely been done and when it has, the antipsychotic has most commonly been administered acutely (e.g., Marquis et al., 2011). We recently conducted a series of experiments in rats where the  $\alpha 7$ -nAChR partial agonist tropisetron was administered to rats that had been exposed to either risperidone or quetiapine for 30 or 90 days then tested them in a novel object recognition task (Poddar et al., 2018). Tropisetron markedly improved NOR performance in rats treated with either antipsychotic for 30 or 90 days indicating that in this particular case, the antipsychotic treatment history did not interfere with the pro-cognitive effect of tropisetron. Thus,  $\alpha 7$ -nAChR ligands like tropisetron may have potential as adjunctive medications in schizophrenia since the pro-cognitive effect was maintained in the presence of chronic antipsychotic treatment. However, in this study, tropisetron was administered acutely and future studies would need to be conducted to determine if this pro-cognitive effect of tropisetron is lasting.

Other factors related to polypharmacy and drug exposure history that have not been adequately addressed are how chronic nicotine exposure or the combination of chronic nicotine exposure and antipsychotic treatment might affect the efficacy of a pro-cognitive agent. Given the well-documented chronic effects of nicotine on nAChR expression (see Lewis and Picciotto 2013 for review), and the aforementioned high smoking rates in schizophrenia, this could certainly be an important consideration when evaluating an  $\alpha 7$ -nAChR ligand for pro-cognitive effects. Levin and colleagues (see review, Levin and Rezvani, 2007) have performed some experiments in rodents to investigate nicotinic interactions with antipsychotic drugs and cognitive function. For example, they have shown that nicotine and some nicotine agonists can reduce cognitive impairments caused by some antipsychotic drugs. In other studies, they have shown that nicotine-induced cognitive improvements were attenuated by the some antipsychotics (e.g., clozapine). However, the specific questions raised above (i.e., how chronic nicotine exposure or the combination of chronic nicotine exposure and antipsychotic treatment might affect the efficacy of a pro-cognitive agent including an  $\alpha 7$ -nAChR ligand) have not been rigorously investigated either in animal models or in clinical studies.

## 6. Pro-cognitive drug dose, frequency of administration, and duration of treatment

Another potential limitation of many clinical studies conducted to date to evaluate novel pro-cognitive agents was the choice of dose, the frequency of administration, and the duration of treatment. Most of the rodent studies (where robust cognitive effects were observed) have employed acute or sub-acute dosing of  $\alpha 7$ -nAChR ligands, which contrasts with most of the clinical trials that were conducted over several weeks or months. All neuronal nAChRs (including  $\alpha 7$ -nAChRs) become temporarily inactive after prolonged exposure to an agonist (Quick and Lester 2002), thus the repeated administration of  $\alpha 7$ -nAChR agonists in the clinical studies may have resulted in receptor desensitization, or potentially even functional antagonism. To support this argument are the disparate effects of the immediate and slow-release versions of DMXB-A (GTS-21) observed in clinical trials. Whereas, the immediate release formulation (which was rapidly absorbed, but quickly cleared) improved cognition

and P50 sensory gating in schizophrenic patients, the slow-release version was not effective (Olincy et al., 2006; Freedman et al., 2008; Kem et al., 2018). Another emerging hypothesis is that low concentrations of  $\alpha 7$ -nAChR ligands may be more effective than higher concentrations, as the latter will maintain receptors in a desensitized and unresponsive state (see Tregellas and Wylie, 2019). In many of the more recent animal studies, the greatest response to a nAChR agonist was observed with a low drug dose and increasing the dose often produced a decreasing effect, in an inverted U-shaped response curve, which is thought to be due to receptor desensitization. A particularly notable example was a non-human primate study where, a low dose of an  $\alpha 7$ -nAChR agonist (PHA543613) facilitated neuronal activity in the prefrontal cortex and improved performance of a spatial working memory task, whereas higher doses were not effective (Yang et al., 2013).

The ability to select the proper dose and frequency of administration of an  $\alpha 7$ -nAChR ligand for optimal target engagement could possibly be improved if functional biomarkers were identified for use in both preclinical and clinical studies. Noninvasive neuroimaging methods such as functional magnetic resonance imaging (fMRI) may be able to facilitate the identification of biomarkers since they can be used to investigate neural circuitry alterations that underlie symptoms of schizophrenia as well as how medications affect this neural circuitry. Here, a biomarker that can be linked to a disease mechanism is categorized as a type I biomarker and a biomarker that can be linked to a treatment response is categorized as a type II biomarker (see Wylie et al., 2016 for review). Interestingly, Tregellas and colleagues, using fMRI, linked hippocampal hyperactivity to smooth pursuit eye movement (SPEM) deficits in schizophrenia patients. The findings revealed a link between eye-tracking abnormalities and a hypothesized disease mechanism, thereby potentially qualifying hippocampal hyperactivity during SPEM as a type I biomarker. Additional studies in schizophrenia patients demonstrated that nicotine and the  $\alpha 7$ -nAChR partial agonist DMXB-A could normalize hippocampal hyperactivity during SPEM, suggesting that SPEM during fMRI could also represent a potential type II biomarker of treatment response (see Wylie et al., 2016; Tregellas and Wylie, 2019). In summary, while the use of fMRI for the development of biomarkers for schizophrenia and other neuropsychiatric disorders is in the early stages, it has the potential to facilitate drug development by improving the translation from animal models to the clinical realm as well to inform investigators as to the best dosing strategies to optimize target engagement.

## 7. Optimizing $\alpha 7$ -nAChR activity

A large-scale effort to overcome the challenges related to nAChR desensitization and the administration of orthosteric agonists and partial agonists has been the development of positive allosteric modulators (PAMs). PAMs are thought to bind to sites that are distinct from the well-conserved (orthosteric) agonist binding domains and they require the presence of the endogenous ligand to increase receptor activity. To date, two types of PAMs have been developed, designated Type I and Type II. Type I PAMs are defined as molecules that predominately affect the apparent peak current, agonist sensitivity, and Hill coefficient, but not the receptor desensitization profile. Type II PAMs possess the aforementioned properties described for Type I PAMs as well as the ability to modify the desensitization profile of agonist responses (see Bertrand and Gopalakrishnan, 2007). It has been argued that Type II

PAMs (compared to Type I PAMs) are less likely to induce tolerance, which may occur after the chronic administration of nAChR agonists, whereas, Type I PAMs, may have the advantage (compared to Type II PAMs) of minimizing the potential for calcium induced cytotoxicity (Ng et al., 2007, see also Nikiforuk et al., 2015). A large number of PAMs from both classes have been developed with pro-cognitive properties in animal models (see Table 1); however, to date only a few clinical trials have been conducted (or are underway) with  $\alpha$ 7-nAChR PAMs (i.e., AVL-3288, galantamine, JNJ-39393406).

Finally, the classical view that nAChR stimulation is the key action responsible for the pro-cognitive effects of  $\alpha$ 7-nAChRs ligands may require additional consideration given the observations that low doses of the selective  $\alpha$ 7-nAChR antagonist methyllycaconitine (MLA) can (in some cases) improve cognition in animal models (Hahn et al, 2011; Burke et al., 2014). and facilitate LTP induction in hippocampal region CA1 in rats (Fujii et al., 2000). More recently, low concentrations/doses of MLA exerted surprising (positive) effects in several model systems. Specifically, in electrophysiological experiments, low concentrations of MLA potentiated receptor responses to acetylcholine in human  $\alpha$ 7-nAChR-transfected oocytes, enhanced long term potentiation (LTP) in rat hippocampal slices, increased hippocampal glutamate efflux in microdialysis experiments in rats, and improved the acquisition of a novel object recognition task in rats (van Goethem et al., 2019). Interestingly, the nonselective nAChR antagonist mecamylamine (in some cases) has also been found to exert pro-cognitive effects in working memory tasks in both rodents and monkeys as well as a recognition memory task in humans with attention deficit hyperactivity disorder (ADHD). For details of these studies, see Buccafusco et al., 2009; Bertrand and Terry 2018.

## 8. Additional Clinical Trial Design Issues

There is growing evidence that schizophrenia patients whose cognitive performance is comparable to healthy controls (i.e., up to 25% of patients) may not benefit from pro-cognitive agents (Granger et al., 2018; DeTore et al., 2019). Therefore, the inclusion of such individuals in a clinical trial designed to determine a compound's pro-cognitive efficacy may limit the power of the study. Unfortunately, a recent analysis of 87 randomized, double-blind, placebo-controlled, clinical trials listed on [ClinicalTrials.gov](https://clinicaltrials.gov) indicated that the vast majority of such clinical trials may have been underpowered due to the inclusion of cognitively "normal" patients (Cotter et al., 2019).

The argument that pro-cognitive strategies (specifically  $\alpha$ 7-nAChR-based approaches) should target subgroups of individuals who exhibit lower levels of cognitive function is supported by recent clinical studies with the selective  $\alpha$ 7-nAChR agonist CDP-choline in both healthy subjects and patients with schizophrenia. In healthy study participants showing relatively lower cognitive and sensory gating scores at baseline, CDP-choline was found to enhance multiple domains of cognition (Knott et al., 2015a, 2015b) and to improve sensory gating (Knott et al., 2014b). The same laboratory also reported CDP-choline-mediated improvements of P50 sensory gating scores in schizophrenia patients who exhibited impaired gating (Aidelbaum et al., 2018). They also demonstrated that combining CDP-choline with galantamine (an acetylcholinesterase inhibitor and nAChR PAM) improved



sensory gating to speech stimuli in schizophrenia patients who expressed low baseline suppression (Choueiry et al., 2019).

Another factor that may be important to consider in the evaluation of pro-cognitive agents in schizophrenia is the chronicity of the illness. The duration of the disease and the efficacy of antipsychotics has been the subject of several reviews (e.g., Leucht et al., 2008), but the concept could also certainly apply to pro-cognitive drug evaluations. Most participants in antipsychotic clinical trials have been chronically ill having experienced multiple episodes and hospitalizations. While these patients may represent the “typical” cases of schizophrenia, there is increasing interest in the effects of medications on patients who have been recently diagnosed (e.g., first episode patients). As reviewed by Leucht et al., 2008, there are multiple differences between recent onset and chronic schizophrenic patients that could result in different outcomes in clinical trials. These differences include the level of cognitive impairment, the level of treatment compliance, the sensitivity to treatment side effects, and changes in brain morphology (Molina et al., 2005; Rabinowitz et al., 2006; Mori et al., 2007). Thus, studies explicitly recruiting recent onset or first episode patients should be conducted and in large clinical trials, these patients could be included along with chronically ill patients and the study outcomes stratified by group.

Another observation that has been commonly made in failed studies of pro-cognitive agents in neuropsychiatric patients is an improvement in symptoms across all treatment arms once patients are randomized, suggesting that being in a study itself may have a powerful therapeutic effect. It has been suggested that efforts to simplify the studies by reducing the number of interactions of patients with study staff prior to the drug evaluation phase be considered. Alternatively, efforts to ensure that all study procedures, staff interactions, and assessments are included in any run-in period prior to actual randomization of patients might also help to reduce this apparent practice or placebo effect (Marder et al., 2017).

From a meta-analytic review of placebo-controlled trials of antidepressant drugs, Undurraga and Baldessarini, 2012 argued that when drug evaluations progress to Phase III and additional testing sites are recruited to increase the number of study subjects, an unanticipated result is that the quality of the sites diminish (a factor that may contribute to the study failure). Thus, new policies should be developed for neuropsychiatric drug evaluations to focus more on the quality of the recruitment and site conduct rather than on the volume of patients enrolled, which may result in fewer but more productive study sites. Such policies would necessitate continual monitoring of the conduct of the study sites and contract research organizations and the termination of sites that do not perform properly (Marder et al., 2017).

## 9. Conclusions and Future Directions

There are now decades of preclinical evidence to support the argument that  $\alpha 7$ -nAChRs should be viable therapeutic targets for schizophrenia and other disorders of cognition. A wide variety of molecules developed to modulate  $\alpha 7$ -nAChRs exhibited pro-cognitive activity in animals and some have shown positive effects in early (Phase I and II) clinical trials. However, to date, there has been no large phase III clinical trial in schizophrenia

where an  $\alpha 7$ -nAChR ligand has shown clear efficacy as pro-cognitive agent without untoward side effects. A thorough review of the literature indicates that multiple factors can potentially affect the success of pro-cognitive agents in schizophrenia (including  $\alpha 7$ -nAChR ligands) such as the translational value of the animal models used, clinical trial design limitations, confounding effects of polypharmacy, and complex dose-effect and dose frequency considerations.

From the preclinical perspective, more studies should be conducted where the pro-cognitive agent is evaluated in animals that have chronically been treated with an antipsychotic drug. Moreover, the number of chronic dosing paradigms in animal studies should be increased to parallel chronic administration in clinical populations. In both the preclinical and clinical evaluations of  $\alpha 7$ -nAChR ligands, a wider dose range (to include low doses) and the frequency of administration (repeated versus intermittent administration) should be evaluated. In the later phases of preclinical drug discovery, a greater emphasis should be placed on non-human primates (as opposed to rodents) given their more complex behavioral repertoire and brain homology with humans. From the clinical perspective, patients whose cognitive performance is comparable to healthy controls should be excluded and efforts should be made to reduce apparent practice or placebo effects associated with multiple interactions of patients with study staff prior to the drug evaluation. Recent onset or first episode patients should be included along with chronically ill patients and the study outcomes stratified by group. Finally, new policies should also be developed to focus more on the quality of recruitment sites rather than on volume of patients enrolled.

## Acknowledgements

The authors would like to thank Ms. Ashley Davis for her administrative assistance in preparing this manuscript. The author's laboratories and/or salary are supported in part by the following funding sources, National Institutes of Health grants, MH097695, MH083317, and NS099455, and Prime Behavior Testing Laboratories, Evans, Georgia.

## References

- Aidelbaum R, Labelle A, Baddeley A, Knott V. Assessing the acute effects of CDP-choline on sensory gating in schizophrenia: A pilot study. *J Psychopharmacol.* 2018 5;32(5):541–551. doi: 10.1177/0269881117746903 [PubMed: 29338621]
- Barak S, Arad M, De Levie A, Black MD, Griebel G, Weiner I. Pro-cognitive and antipsychotic efficacy of the alpha7 nicotinic partial agonist SSR180711 in pharmacological and neurodevelopmental latent inhibition models of schizophrenia. *Neuropsychopharmacology.* 2009 6;34(7):1753–63. doi: 10.1038/npp.2008.232. [PubMed: 19158670]
- Barbey AK, Koenigs M, Grafman J. Dorsolateral prefrontal contributions to human working memory. *Cortex.* 2013;49(5):1195–1205. doi:10.1016/j.cortex.2012.05.022 [PubMed: 22789779]
- Barch DM, Marder SR, Harms MP, Jarskog LF, Buchanan RW, Cronenwett W, Chen LS, Weiss M, Maguire RP, Pezous N, Feuerbach D, Lopez-Lopez C, Johns DR, Behrje RB, Gomez-Mancilla B. Task-related fMRI responses to a nicotinic acetylcholine receptor partial agonist in schizophrenia: A randomized trial. *Prog Neuropsychopharmacol Biol Psychiatry.* 2016 11 3;71:66–75. doi: 10.1016/j.pnpbp.2016.06.013. [PubMed: 27371157]
- Berger F, Gage FH, Vijayaraghavan S. Nicotinic receptor-induced apoptotic cell death of hippocampal progenitor cells. *J Neurosci.* 1998 9 1;18(17):6871–81. [PubMed: 9712657]
- Bertrand D, Gopalakrishnan M. Allosteric modulation of nicotinic acetylcholine receptors. *Biochem Pharmacol.* 2007 10 15;74(8):1155–63. [PubMed: 17707779]

- Bertrand D, Terry AV Jr. The wonderland of neuronal nicotinic acetylcholine receptors. *Biochem Pharmacol.* 2018 5;151:214–225. doi: 10.1016/j.bcp.2017.12.008. Review. [PubMed: 29248596]
- Bitner RS, Bunnelle WH, Anderson DJ, Briggs CA, Buccafusco J, Curzon P, Decker MW, Frost JM, Gronlien JH, Gubbins E, Li J, Malysz J, Markosyan S, Marsh K, Meyer MD, Nikkel AL, Radek RJ, Robb HM, Timmermann D, Sullivan JP, Gopalakrishnan M. Broad-spectrum efficacy across cognitive domains by alpha7 nicotinic acetylcholine receptor agonism correlates with activation of ERK1/2 and CREB phosphorylation pathways. *J Neurosci.* 2007 9 26;27(39):10578–87 [PubMed: 17898229]
- Bitner RS, Bunnelle WH, Decker MW, Drescher KU, Kohlhaas KL, Markosyan S, Marsh KC, Nikkel AL, Browman K, Radek R, Anderson DJ, Buccafusco J, Gopalakrishnan M. In vivo pharmacological characterization of a novel selective alpha7 neuronal nicotinic acetylcholine receptor agonist ABT-107: preclinical considerations in Alzheimer's disease. *J Pharmacol Exp Ther.* 2010 9 1;334(3):875–86. doi: 10.1124/jpet.110.167213. [PubMed: 20504913]
- Boess FG, De Vry J, Erb C, Flessner T, Hendrix M, Luthle J, Methfessel C, Riedl B, Schnizler K, van der Staay FJ, Van Kampen M, Wiese WB, Koenig G. (2007). The novel alpha7 nicotinic acetylcholine receptor agonist N-(3R)-1-azabicyclo[2.2.2]oct-3-yl-(7-(2-methoxyphenyl)-1-benzofuran)-2-carboxamide improves working and recognition memory in rodents. *J Pharmacol Exp Ther* 321:716–725. [PubMed: 17308038]
- Boess FG, de Vry J, Erb C, Flessner T, Hendrix M, Luthle J, Methfessel C, Schnizler K, van der Staay FJ, van Kampen M, Wiese WB, König G. (2013). Pharmacological and behavioral profile of N-((3R)-1-azabicyclo(2.2.2)oct-3-yl)-6-chinolincarboxamide (EVP-5141), a novel  $\alpha$ 7 nicotinic acetylcholine receptor agonist/serotonin 5-HT<sub>3</sub> receptor antagonist. *Psychopharmacology* 227:1–17. doi: 10.1007/s00213-012-2933-4. [PubMed: 23241647]
- Bordia T, McGregor M, Papke RL, Decker MW, McIntosh JM, Quik M. The  $\alpha$ 7 nicotinic receptor agonist ABT-107 protects against nigrostriatal damage in rats with unilateral 6-hydroxydopamine lesions. *Exp Neurol.* 2015 1;263:277–84. doi: 10.1016/j.expneurol.2014.09.015. [PubMed: 25261754]
- Bristow LJ, Easton AE, Li YW, Sivarao DV et al. (2016). The novel, nicotinic alpha7 receptor partial agonist, BMS-933043, improves cognition and sensory processing in preclinical models of schizophrenia. *PLoS One* 11:1–29.
- Buccafusco JJ, Beach JW, Terry AV Jr. (2009) Desensitization of nicotinic acetylcholine receptors as a strategy for drug development. *J Pharmacol Exp Ther.* 2;328(2):364–70. doi: 10.1124/jpet.108.145292. Review. PMID: 19023041 [PubMed: 19023041]
- Buchanan RW, Davis M, Goff D, Green MF, Keefe RS, Leon AC, Nuechterlein KH, Laughren T, Levin R, Stover E, Fenton W, Marder SR. A summary of the FDA-NIMH-MATRICES workshop on clinical trial design for neurocognitive drugs for schizophrenia. *Schizophr Bull.* 2005 1;31(1):5–19 [PubMed: 15888422]
- Buchanan RW, Kelly DL, Weiner E, Gold JM, Strauss GP, Koola MM, McMahon RP, Carpenter WT. A Randomized Clinical Trial of Oxytocin or Galantamine for the Treatment of Negative Symptoms and Cognitive Impairments in People With Schizophrenia. *J Clin Psychopharmacol.* 2017 8;37(4):394–400. doi: 10.1097/JCP.0000000000000720 [PubMed: 28590362]
- Burke DA, Heshmati P, Kholdebarin E, Levin ED. Decreasing nicotinic receptor activity and the spatial learning impairment caused by the NMDA glutamate antagonist dizocilpine in rats. *Eur J Pharmacol.* 2014 10 15;741:132–9. doi: 10.1016/j.ejphar.2014.07.030. [PubMed: 25064338]
- Callahan PM, Terry AV Jr, Tehim A. Effects of the nicotinic  $\alpha$ 7 receptor partial agonist GTS-21 on NMDA-glutamatergic receptor related deficits in sensorimotor gating and recognition memory in rats. *Psychopharmacology (Berl).* 2014 9;231(18):3695–706. doi: 10.1007/s00213-014-3509-2. [PubMed: 24595504]
- Callahan PM, Bertrand D, Bertrand S, Plagenhoef MR, Terry AV Jr. (2017). Tropisetron sensitizes  $\alpha$ 7 containing nicotinic receptors to low levels of acetylcholine in vitro and improves memory-related task performance in young and aged animals. *Neuropharmacology*, 117:422–433. [PubMed: 28259598]
- Carpenter WT Jr. The RDoC Controversy: Alternate Paradigm or Dominant Paradigm? *Am J Psychiatry.* 2016 6 1;173(6):562–3. doi: 10.1176/appi.ajp.2016.16030347. [PubMed: 27245188]

- Chang KT, Berg DK. (1999) Nicotinic acetylcholine receptors containing alpha7 subunits are required for reliable synaptic transmission in situ. *J Neurosci.* 5 15;19(10):3701–10. [PubMed: 10234002]
- Chouairy J, Blais CM, Shah D, Smith D, Fisher D, Labelle A, Knott V. Combining CDP-choline and galantamine, an optimized  $\alpha 7$  nicotinic strategy, to ameliorate sensory gating to speech stimuli in schizophrenia. *Int J Psychophysiol.* 2019 11;145:70–82. doi: 10.1016/j.ijpsycho.2019.02.005. [PubMed: 30790597]
- Corradi J, Bouzat C. Understanding the Bases of Function and Modulation of  $\alpha 7$  Nicotinic Receptors: Implications for Drug Discovery. *Mol Pharmacol.* 2016 9;90(3):288–99. doi: 10.1124/mol.116.104240. Review. [PubMed: 27190210]
- Cotter J, Barnett JH, Granger K. The Use of Cognitive Screening in Pharmacotherapy Trials for Cognitive Impairment Associated With Schizophrenia. *Front Psychiatry.* 2019 9 6;10:648. doi: 10.3389/fpsy.2019.00648. Review. [PubMed: 31551837]
- DeTore NR, Mueser KT, Byrd JA, McGurk SR. Cognitive functioning as a predictor of response to comprehensive cognitive remediation. *J Psychiatr Res.* 2019 6;113:117–124. doi: 10.1016/j.jpsychires.2019.03.012. [PubMed: 30933830]
- Dunlop J, Lock T, Jow B, Sitzia F, Grauer S, Jow F, Kramer A, Bowlby MR, Randall A, Kowal D, Gilbert A, Comery TA, Larocque J, Soloveva V, Brown J, Roncarati R. Old and new pharmacology: positive allosteric modulation of the alpha7 nicotinic acetylcholine receptor by the 5-hydroxytryptamine(2B/C) receptor antagonist SB-206553 (3,5-dihydro-5-methyl-N-3-pyridinylbenzo[1,2-b:4,5-b']di pyrrole-1(2H)-carboxamide), *J. Pharmacol. Exp. Ther* 328 (2009) 766–776. [PubMed: 19050173]
- Eskildsen J, Redrobe JP, Sams AG, Dekermendjian K, Laursen M, Boll JB, Papke RL, Bundgaard C, Frederiksen K, Bastlund JF. (2014). Discovery and optimization of Lu AF58801, a novel, selective and brain penetrant positive allosteric modulator of alpha-7 nicotinic acetylcholine receptors: attenuation of subchronic phencyclidine (PCP)-induced cognitive deficits in rats following oral administration. *Bioorg Med Chem Lett* 24:288–293. [PubMed: 24291041]
- Feuerbach D, Pezous N, Weiss M, Shakeri-Nejad K, Lingenhoehl K, Hoyer D, Hurth K, Bilbe G, Pryce CR, McAllister K, Chaperon F, Kucher K, Johns D, Blaettler T, Lopez Lopez C. (2015). AQW051, a novel, potent and selective  $\alpha 7$  nicotinic ACh receptor partial agonist: pharmacological characterization and phase I evaluation. *Br J Pharmacol* 172:1292–1304. [PubMed: 25363835]
- Freedman R, Olincy A, Ross RG, Waldo MC, Stevens KE, Adler LE, Leonard S. The genetics of sensory gating deficits in schizophrenia. *Curr. Psychiatry Rep* 2003;5:155–161. [PubMed: 12685995]
- Freedman R, Olincy A, Buchanan RW, Harris JG, Gold JM, Johnson L, Allensworth D, Guzman-Bonilla A, Clement B, Ball MP, Kutnick J, Pender V, Martin LF, Stevens KE, Wagner BD, Zerbe GO, Soti F, Kem WR. Initial phase 2 trial of a nicotinic agonist in schizophrenia. *Am J Psychiatry.* 2008 8;165(8):1040–7. doi: 10.1176/appi.ajp.2008.07071135 [PubMed: 18381905]
- Freedman R. (2014)  $\alpha 7$ -nicotinic acetylcholine receptor agonists for cognitive enhancement in schizophrenia. *Annu. Rev. Med.* 65 (2014), pp. 245–261, 10.1146/annurev-med-092112-142937. Review [PubMed: 24111888]
- Fujii S, Jia Y, Yang A, Sumikawa K. Nicotine reverses GABAergic inhibition of long-term potentiation induction in the hippocampal CA1 region. *Brain Res.* 2000 4 28;863(1–2):259–65. [PubMed: 10773216]
- Gee KW, Olincy A, Kanner R, Johnson L, Hogenkamp D, Harris J, Tran M, Edmonds SA, Sauer W, Yoshimura R, Johnstone T, Freedman R. First in human trial of a type I positive allosteric modulator of alpha7-nicotinic acetylcholine receptors: Pharmacokinetics, safety, and evidence for neurocognitive effect of AVL-3288. *J Psychopharmacol.* 2017 4;31(4):434–441. doi: 10.1177/0269881117691590 [PubMed: 28196430]
- George TP, Krystal JH. Comorbidity of psychiatric and substance abuse disorders. *Curr Opin Psychiatry.* 2000;13(3):327–331.
- Goff DC, Wine L. (1997) Glutamate in schizophrenia: clinical and research implications. *Schizophr Res.* 10 30;27(2–3):157–68. doi: 10.1016/S0920-9964(97)00079-0 [PubMed: 9416645]
- Gotti C, Moretti M, Gaimarri A, Zanardi A, Clementi F, Zoli M. Heterogeneity and complexity of native brain nicotinic receptors. *Biochem Pharmacol.* 2007 10 15;74(8):1102–11. [PubMed: 17597586]

- Granger G, Cotter J, Baker E, Evenden J, Barnett J, Sand M. Exploring participant-level trajectories of cognitive performance among patients with schizophrenia in a multi-national trial. *Schizophr Bull* (2018) 44:S353. 10.1093/schbul/sby018.861
- Gray R, Rajan AS, Radcliffe KA, Yakehiro M, Dani JA. Hippocampal synaptic transmission enhanced by low concentrations of nicotine. *Nature*. 1996 10 24;383(6602):713–6. [PubMed: 8878480]
- Green MF, Braff DL. (2001) Translating the basic and clinical cognitive neuroscience of schizophrenia to drug development and clinical trials of antipsychotic medications. *Biol Psychiatry*. 2 15;49(4):374–84. [PubMed: 11239909]
- Green MF, Harvey PD. (2014) Cognition in schizophrenia: Past, present, and future. *Schizophr Res Cogn*. 3;1(1):e1–e9. [PubMed: 25254156]
- Guan ZZ, Zhang X, Blennow K, Nordberg A. (1999) Decreased protein level of nicotinic receptor alpha7 subunit in the frontal cortex from schizophrenic brain. *Neuroreport*. 6 3;10(8):1779–82. [PubMed: 10501574]
- Hahn B, Shoab M, Stolerman IP. Selective nicotinic receptor antagonists: effects on attention and nicotine-induced attentional enhancement. *Psychopharmacology (Berl)*. 2011 9;217(1):75–82. doi: 10.1007/s00213-011-2258-8. [PubMed: 21432025]
- Haig G, Wang D, Othman AA, Zhao J. The  $\alpha 7$  Nicotinic Agonist ABT-126 in the Treatment of Cognitive Impairment Associated with Schizophrenia in Nonsmokers: Results from a Randomized Controlled Phase 2b Study. *Neuropsychopharmacology*. 2016 11;41(12):2893–2902. doi: 10.1038/npp.2016.101. [PubMed: 27319970]
- Haig GM, Bain EE, Robieson WZ, Baker JD, Othman AA. A Randomized Trial to Assess the Efficacy and Safety of ABT-126, a Selective  $\alpha 7$  Nicotinic Acetylcholine Receptor Agonist, in the Treatment of Cognitive Impairment in Schizophrenia. *Am J Psychiatry*. 2016 8 1;173(8):827–35. doi: 10.1176/appi.ajp.2015.15010093 [PubMed: 26940805]
- Haig GM, Wang D, Zhao J, Othman AA, Bain EE. Efficacy and Safety of the  $\alpha 7$ -Nicotinic Acetylcholine Receptor Agonist ABT-126 in the Treatment of Cognitive Impairment Associated With Schizophrenia: Results From a Phase 2b Randomized Controlled Study in Smokers. *J Clin Psychiatry*. 2018 May-Jun;79(3). pii: 16m11162. doi: 10.4088/JCP.16m11162.
- Hashimoto K, Iyo M, Freedman R, Stevens KE. Tropicsetron improves deficient inhibitory auditory processing in DBA/2 mice: role of alpha 7 nicotinic acetylcholine receptors. *Psychopharmacology* 183 (2005) 13–19. [PubMed: 16136299]
- Hashimoto K, Fujita Y, Ishima T, Hagiwara H, Iyo M (2006). Phencyclidine-induced cognitive deficits in mice are improved by subsequent subchronic administration of tropisetron: role of  $\alpha 7$  nicotinic receptors. *Eur J Pharmacol* 553:191–195. [PubMed: 17094961]
- Hauser TA, Kucinski A, Jordan KG, Gatto GJ, Wersinger SR, Hesse RA, Stachowiak EK, Stachowiak MK, Papke RL, Lippiello PM, Bencherif M. TC-5619: an alpha7 neuronal nicotinic receptor-selective agonist that demonstrates efficacy in animal models of the positive and negative symptoms and cognitive dysfunction of schizophrenia. *Biochem. Pharmacol*. 78 (2009) 803–812.
- Hefft S, Hulo S, Bertrand D, Muller D. (1999) Synaptic transmission at nicotinic acetylcholine receptors in rat hippocampal organotypic cultures and slices. *J Physiol*. 3 15;515 (Pt 3):769–76. doi: 10.1111/j.1469-7793.1999.769abx [PubMed: 10066903]
- Hogenkamp DJ1, Ford-Hutchinson TA, Li WY, Whittemore ER, Yoshimura RF, Tran MB, Johnstone TB, Bascom GD, Rollins H, Lu L, Gee KW. Design, synthesis, and activity of a series of arylpyrid-3-ylmethanones as type I positive allosteric modulators of alpha7 nicotinic acetylcholine receptors. *J. Med. Chem* 56 (2013) 8352–8365. [PubMed: 24098954]
- Jones KM, McDonald IM, Bourin C, Olson RE, Bristow LJ, Easton A. Effect of alpha7 nicotinic acetylcholine receptor agonists on attentional set-shifting impairment in rats. *Psychopharmacology (Berl)*. 2014 2;231(4):673–83. doi: 10.1007/s00213-013-3275-6. [PubMed: 24057763]
- Kahn RS, Keefe RS. (2013) Schizophrenia is a cognitive illness: time for a change in focus. *JAMA Psychiatry*. 10;70(10):1107–12. doi: 10.1001/jamapsychiatry.2013.155. [PubMed: 23925787]
- Kahn RS. On the Specificity of Continuous Cognitive Decline in Schizophrenia. *Am J Psychiatry*. 2019 10 1;176(10):774–776. doi: 10.1176/appi.ajp.2019.19080794 [PubMed: 31569987]
- Kantrowitz JT, Javitt DC, Freedman R, Sehatpour P, Kegeles LS, Carlson M, Sobeih T, Wall MM, Choo TH, Vail B, Grinband J, Lieberman JA. Double blind, two dose, randomized, placebo-

controlled, cross-over clinical trial of the positive allosteric modulator at the alpha7 nicotinic cholinergic receptor AVL-3288 in schizophrenia patients. *Neuropsychopharmacology*. 2020 2 3. doi: 10.1038/s41386-020-0628-9.

- Kapur S, Mamo D. (2003) Half a century of antipsychotics and still a central role for dopamine D2 receptors. *Prog Neuropsychopharmacol Biol Psychiatry*. 10;27(7):1081–90. Review. [PubMed: 14642968]
- Keefe RS, Meltzer HA, Dgetluck N, Gawryl M, Koenig G, Moebius HJ, Lombardo I, Hilt DC. Randomized, Double-Blind, Placebo-Controlled Study of Encenicline, an  $\alpha 7$  Nicotinic Acetylcholine Receptor Agonist, as a Treatment for Cognitive Impairment in Schizophrenia. *Neuropsychopharmacology*. 2015 12;40(13):3053–60. doi: 10.1038/npp.2015.176. [PubMed: 26089183]
- Kem WR, Olincy A, Johnson L, Harris J, Wagner BD, Buchanan RW, Christians U, Freedman R. Pharmacokinetic Limitations on Effects of an Alpha7-Nicotinic Receptor Agonist in Schizophrenia: Randomized Trial with an Extended-Release Formulation. *Neuropsychopharmacology*. 2018 2;43(3):583–589. doi: 10.1038/npp.2017.182. [PubMed: 28825423]
- Knott V, Smith D, de la Salle S, Impey D, Choueiry J, Beaudry E, Smith M, Saghir S, Ilivitsky V, Labelle A. CDP-choline: effects of the procholine supplement on sensory gating and executive function in healthy volunteers stratified for low, medium and high P50 suppression. *J Psychopharmacol*. 2014 12;28(12):1095–108. doi: 10.1177/0269881114553254. [PubMed: 25315828]
- Knott V, de la Salle S, Smith D, Choueiry J, Impey D, Smith M, Beaudry E, Saghir S, Ilivitsky V, Labelle A. Effects of acute CDP-choline treatment on resting state brain oscillations in healthy volunteers. *Neurosci Lett*. 2015 3 30;591:121–5. doi: 10.1016/j.neulet.2015.02.032 [PubMed: 25700947]
- Knott V, de la Salle S, Choueiry J, Impey D, Smith D, Smith M, Beaudry E, Saghir S, Ilivitsky V, Labelle A. Neurocognitive effects of acute choline supplementation in low, medium and high performer healthy volunteers. *Pharmacol Biochem Behav*. 2015 4;131:119–29. doi: 10.1016/j.pbb.2015.02.004 [PubMed: 25681529]
- Kohnomi S, Suemaru K, Goda M, Choshi T, Hibino S, Kawasaki H, Araki H. Ameliorating effects of tropisetron on dopaminergic disruption of prepulse inhibition via the alpha(7) nicotinic acetylcholine receptor in Wistar rats, *Brain Res*. 1353 (2010) 152–158. [PubMed: 20673759]
- Laubach M, Amarante LM, Swanson K, White SR. What, If Anything, Is Rodent Prefrontal Cortex? *eNeuro*. 2018 10 25;5(5). doi: 10.1523/ENEURO.0315-18.2018. Review.
- Leonard S, Mexal S, Freedman R. (2007) Smoking, Genetics and Schizophrenia: Evidence for Self Medication. *J Dual Diagn*. 11 1;3(3–4):43–59. [PubMed: 19122786]
- Leucht S, Heres S, Hamann J, Kane JM. Methodological issues in current antipsychotic drug trials. *Schizophr Bull*. 2008;34(2):275–285. doi:10.1093/schbul/sbm159 [PubMed: 18234700]
- Levin ED, Rezvani AH. Nicotinic interactions with antipsychotic drugs, models of schizophrenia and impacts on cognitive function. *Biochem Pharmacol*. 2007 10 15;74(8):1182–91. [PubMed: 17714691]
- Lewis AS, Picciotto MR. High-affinity nicotinic acetylcholine receptor expression and trafficking abnormalities in psychiatric illness. *Psychopharmacology (Berl)*. 2013 10;229(3):477–85. doi: 10.1007/s00213-013-3126-5. [PubMed: 23624811]
- Lewis CC, Mettert KD, Dorsey CN, Martinez RG, Weiner BJ, Nolen E, Stanick C, Halko H, Powell BJ. An updated protocol for a systematic review of implementation-related measures. *Syst Rev*. 2018 4 25;7(1):66. doi: 10.1186/s13643-018-0728-3. [PubMed: 29695295]
- Lieberman JA, Dunbar G, Segreti AC, Girgis RR, Seoane F, Beaver JS, Duan N, Hosford DA. A randomized exploratory trial of an  $\alpha$ -7 nicotinic receptor agonist (TC-5619) for cognitive enhancement in schizophrenia. *Neuropsychopharmacology*. 2013 5;38(6):968–75. doi: 10.1038/npp.2012.259. [PubMed: 23303043]
- Marcus MM, Björkholm C, Malmerfelt A, Möller A, Pålsson N, Konradsson-Geuken Å, Feltmann K, Jardemark K, Schilström B, Svensson TH. Alpha7 nicotinic acetylcholine receptor agonists and PAMs as adjunctive treatment in schizophrenia. An experimental study. *Eur*

- Neuropsychopharmacol. 2016 9;26(9):1401–1411. doi: 10.1016/j.euroneuro.2016.07.004. [PubMed: 27474687]
- Marder SR, Laughren T, Romano SJ. Why are innovative drugs failing in phase III? *Am J Psychiatry*. 2017;174(9):829–831 [PubMed: 28859511]
- Marquis KL, Comery TA, Jow F, Navarra RL, Grauer SM, Pulicicchio C, Kelley C, Brennan JA, Roncarati R, Scali C, Haydar S, Ghiron C, Terstappen GC, Dunlop J. Preclinical assessment of an adjunctive treatment approach for cognitive impairment associated with schizophrenia using the alpha7 nicotinic acetylcholine receptor agonist WYE-103914/SEN34625. *Psychopharmacology (Berl)*. 2011 12;218(4):635–47. doi: 10.1007/s00213-011-2357-6. [PubMed: 21643676]
- Martin LF, Kem WR, Freedman R. Alpha-7 nicotinic receptor agonists: potential new candidates for the treatment of schizophrenia. *Psychopharmacology (Berl)*. 2004 6;174(1):54–64. [PubMed: 15205879]
- McGehee DS, Heath MJ, Gelber S, Devay P, Role LW. Nicotine enhancement of fast excitatory synaptic transmission in CNS by presynaptic receptors. *Science*. 1995 9 22;269(5231):1692–6. [PubMed: 7569895]
- McLean SL, Grayson B, Marsh S, Zarroug SH, Harte MK, Neill JC. Nicotinic  $\alpha 7$  and  $\alpha 4\beta 2$  agonists enhance the formation and retrieval of recognition memory: Potential mechanisms for cognitive performance enhancement in neurological and psychiatric disorders. *Behav Brain Res*. 2016 4 1;302:73–80. doi: 10.1016/j.bbr.2015.08.037. [PubMed: 26327238]
- Messi ML, Renganathan M, Grigorenko E, Delbono O. 1997 Activation of alpha7 nicotinic acetylcholine receptor promotes survival of spinal cord motoneurons. *FEBS Lett*. 1997 7 7;411(1):32–8. [PubMed: 9247137]
- Meyer EM, Tay ET, Papke RL, Meyers C, Huang GL, de Fiebre CM. 3-[2,4-Dimethoxybenzylidene]anabaseine (DMXB) selectively activates rat alpha7 receptors and improves memory-related behaviors in a mecamylamine-sensitive manner. *Brain Res*. 1997 9 12;768(1–2):49–56. [PubMed: 9369300]
- Meyer EM, Kuryatov A, Gerzanich V, Lindstrom J, Papke RL. Analysis of 3-(4-hydroxy, 2-Methoxybenzylidene)anabaseine selectivity and activity at human and rat alpha-7 nicotinic receptors. *J Pharmacol Exp Ther*. 1998 12;287(3):918–25 [PubMed: 9864273]
- Molina V, Sánchez J, Reig S, Sanz J, Benito C, Santamarta C, Pascau J, Sarramea F, Gispert JD, Misiego JM, Palomo T, Desco M. N-acetyl-aspartate levels in the dorsolateral prefrontal cortex in the early years of schizophrenia are inversely related to disease duration. *Schizophr Res*. 2005 3 1;73(2–3):209–19. [PubMed: 15653263]
- Monteggia LM, Heimer H, Nestler EJ. Meeting Report: Can We Make Animal Models of Human Mental Illness? *Biol Psychiatry*. 2018 10 1;84(7):542–545. doi: 10.1016/j.biopsych.2018.02.010. [PubMed: 29606372]
- Mori T, Ohnishi T, Hashimoto R, Nemoto K, Moriguchi Y, Noguchi H, Nakabayashi T, Hori H, Harada S, Saitoh O, Matsuda H, Kunugi H. Progressive changes of white matter integrity in schizophrenia revealed by diffusion tensor imaging. *Psychiatry Res*. 2007;154:133–145. [PubMed: 17276660]
- Morris SE, Cuthbert BN. Research Domain Criteria: cognitive systems, neural circuits, and dimensions of behavior. *Dialogues Clin Neurosci*. 2012 3;14(1):29–37. [PubMed: 22577302]
- Morrison PD, Murray RM. The antipsychotic landscape: dopamine and beyond. *Ther Adv Psychopharmacol*. 2018 4;8(4):127–135. doi: 10.1177/2045125317752915. Review. [PubMed: 29607005]
- Nestler EJ, Hyman SE. Animal models of neuropsychiatric disorders. *Nat Neurosci*. 2010 10;13(10):1161–9. doi: 10.1038/nn.2647. [PubMed: 20877280]
- Ng HJ, Whittemore ER, Tran MB, Hogenkamp DJ, Broide RS, Johnstone TB, Zheng L, Stevens KE, Gee KW. Nootropic alpha7 nicotinic receptor allosteric modulator derived from GABAA receptor modulators. *Proc Natl Acad Sci U S A*. 2007 5 8;104(19):8059–64 [PubMed: 17470817]
- Nikiforuk A, Kos T, Potasiewicz A, Popik P. Positive allosteric modulation of alpha 7 nicotinic acetylcholine receptors enhances recognition memory and cognitive flexibility in rats. *Eur Neuropsychopharmacol*. 2015 8;25(8):1300–13. doi: 10.1016/j.euroneuro.2015.04.018

- Noroozian M, Ghasemi S, Hosseini SM, Modabbernia A, Khodaie-Ardakani MR, Mirshafiee O, Farokhnia M, Tajdini M, Rezaei F, Salehi B, Ashrafi M, Yekehtaz H, Tabrizi M, Akhondzadeh S. A placebo-controlled study of tropisetron added to risperidone for the treatment of negative symptoms in chronic and stable schizophrenia. *Psychopharmacology (Berl)*. 2013 8;228(4):595–602. doi: 10.1007/s00213-013-3064-2. [PubMed: 23515583]
- Olinicy A, Young DA, Freedman R. (1997) Increased levels of the nicotine metabolite cotinine in schizophrenic smokers compared to other smokers. *Biol Psychiatry*. 7 1;42(1):1–5. [PubMed: 9193735]
- Olinicy A, Harris JG, Johnson LL, Pender V, Kongs S, Allensworth D, Ellis J, Zerbe GO, Leonard S, Stevens KE, Stevens JO, Martin L, Adler LE, Soti F, Kem WR, Freedman R. Proof-of-concept trial of an alpha7 nicotinic agonist in schizophrenia. *Arch Gen Psychiatry*. 2006 6;63(6):630–8. [PubMed: 16754836]
- Perkins KA, Roy Chengappa KN, Karelitz JL, Boldry MC, Michael V, Herb T, Gannon J, Brar J, Ford L, Rassnick S, Brunzell DH. Initial Cross-Over Test of A Positive Allosteric Modulator of Alpha-7 Nicotinic Receptors to Aid Cessation in Smokers With Or Without Schizophrenia. *Neuropsychopharmacology*. 2018 5;43(6):1334–1342. doi: 10.1038/npp.2017.292. [PubMed: 29185480]
- Pichat P, Bergis OE, Terranova JP, Urani A, Duarte C, Santucci V, Gueudet C, Voltz C, Steinberg R, Stemmelin J, Oury-Donat F, Avenet P, Griebel G, Scatton B. SSR180711, a novel selective alpha7 nicotinic receptor partial agonist: (II) efficacy in experimental models predictive of activity against cognitive symptoms of schizophrenia. *Neuropsychopharmacology*. 2007 1;32(1):17–34. [PubMed: 16936709]
- Pieschl RL, Miller R, Jones KM, Post-Munson DJ, Chen P, Newberry K, Benitex Y, Molski T, Morgan D, McDonald IM, Macor JE, Olson RE, Asaka Y, Digavalli S, Easton A, Herrington J, Westphal RS, Lodge NJ, Zaczek R, Bristow LJ, Li YW. (2017). Effects of BMS-902483, an  $\alpha 7$  nicotinic acetylcholine receptor partial agonist, on cognition and sensory gating in relation to receptor occupancy in rodents. *Eur J Pharmacol* 807:1–11 doi: 10.1016/j.ejphar.2017.04.024.1. [PubMed: 28438647]
- Pitsikas N, Borsini F. Different effects of tropisetron and ondansetron in learning and memory paradigms. *Pharmacol. Biochem. Behav* 56 (1997) 571–576.
- Poddar I, Callahan PM, Hernandez CM, Yang X, Bartlett MG, Terry AV Jr. Tropisetron enhances recognition memory in rats chronically treated with risperidone or quetiapine. *Biochem Pharmacol*. 2018 5;151:180–187. doi: 10.1016/j.bcp.2017.11.017 [PubMed: 29175423]
- Potasiewicz A, Holuj M, Kos T, Popik P, Arias HR, Nikforuk A (2017). 3-Furan-2-yl-N-p-tolyl-acrylamide, a positive allosteric modulator of the  $\alpha 7$  nicotinic receptor, reverses schizophrenia-like cognitive and social deficits in rats. *Neuropharmacology* 113:188–197. [PubMed: 27717880]
- Prickaerts J, van Goethem NP, Chesworth R, Shapiro G, Boess FG, Methfessel C, Reneerkens OA, Flood DG, Hilt D, Gawryl M, Bertrand S, Bertrand D, König G. EVP-6124, a novel and selective alpha7 nicotinic acetylcholine receptor partial agonist, improves memory performance by potentiating the acetylcholine response of alpha7 nicotinic acetylcholine receptors. *Neuropharmacology* 62 (2012) 1099–1110. [PubMed: 22085888]
- Quick MW, Lester RA. Desensitization of neuronal nicotinic receptors. *J Neurobiol*. 2002 12;53(4):457–78. Review. [PubMed: 12436413]
- Rabinowitz J, Harvey PD, Eerdeken M, Davidson M. Premorbid functioning and treatment response in recent-onset schizophrenia. *Br J Psychiatry*. 2006 7;189:31–5. [PubMed: 16816303]
- Radek RJ, Robb HM, Stevens KE, Gopalakrishnan M, Bitner RS. Effects of the novel alpha7 nicotinic acetylcholine receptor agonist ABT-107 on sensory gating in DBA/2 mice: pharmacodynamic characterization. *J. Pharmacol. Exp. Ther* 343 (2012) 736–745. doi: 10.1124/jpet.112.197970 [PubMed: 22988063]
- Roncarati R, Scali C, Comery TA, Grauer SM, Aschmi S, Bothmann H, Jow B, Kowal D, Gianfriddo M, Kelley C, Zanelli U, Ghiron C, Haydar S, Dunlop J, Terstappen GC. Procognitive and neuroprotective activity of a novel alpha7 nicotinic acetylcholine receptor agonist for treatment of neurodegenerative and cognitive disorders. *J. Pharmacol. Exp. Ther* 329 (2009) 459–468. [PubMed: 19223665]




- Ross CA, Margolis RL. Research Domain Criteria: Strengths, Weaknesses, and Potential Alternatives for Future Psychiatric Research. *Mol Neuropsychiatry*. 2019 10;5(4):218–236. doi: 10.1159/000501797 [PubMed: 31768375]
- Shiina A, Shirayama Y, Niitsu T, Hashimoto T, Yoshida T, Hasegawa T, Haraguchi T, Kanahara N, Shiraishi T, Fujisaki M, Fukami G, Nakazato M, Iyo M, Hashimoto K. A randomised, double-blind, placebo-controlled trial of tropisetron in patients with schizophrenia. *Ann Gen Psychiatry*. 2010 6 24;9:27. doi: 10.1186/1744-859X-9-27. [PubMed: 20573264]
- Stevens KE, Zheng L, Floyd KL, Stitzel JA. Maximizing the effect of an  $\alpha 7$  nicotinic receptor PAM in a mouse model of schizophrenia-like sensory inhibition deficits. *Brain Res*. 2015 6 22;1611:8–17. doi: 10.1016/j.brainres.2015.02.044. [PubMed: 25744161]
- Strand JE, Nybäck H. (2005) Tobacco use in schizophrenia: a study of cotinine concentrations in the saliva of patients and controls. *Eur Psychiatry*. 1;20(1):50–4. doi: 10.1016/j.eurpsy.2004.09.005 [PubMed: 15642444]
- Terry AV, Callahan PM. Nicotinic Acetylcholine Receptor Ligands, Cognitive Function, and Preclinical Approaches to Drug Discovery. *Nicotine Tob Res*. 2019 2 18;21(3):383–394. doi: 10.1093/ntr/nty166. Review. [PubMed: 30137518]
- Tietje KR, Anderson DJ, Bitner RS, Blomme EA, Brackemeyer PJ, Briggs CA, Browman KE, Bury D, Curzon P, Drescher KU, Frost JM, Fryer RM, Fox GB, Gronlien JH, Håkerud M, Gubbins EJ, Halm S, Harris R, Helfrich RJ, Kohlhaas KL, Law D, Malysz J, Marsh KC, Martin RL, Meyer MD, Molesky AL, Nikkel AL, Otte S, Pan L, Puttfarcken PS, Radek RJ, Robb HM, Spies E, Thorin-Hagene K, Waring JF, Ween H, Xu H, Gopalakrishnan M, Bunnelle WH. Preclinical characterization of A-582941: a novel  $\alpha 7$  neuronal nicotinic receptor agonist with broad spectrum cognition-enhancing properties. *CNS Neurosci. Ther* 2008;14:65–82. [PubMed: 18482100]
- Timmermann DB, Grønlien JH, Kohlhaas KL, Nielsen EØ, Dam E, Jørgensen TD, Ahring PK, Peters D, Holst D, Christensen JK, Malysz J, Briggs CA, Gopalakrishnan M, Olsen GM. An allosteric modulator of the alpha7 nicotinic acetylcholine receptor possessing cognition-enhancing properties in vivo. *J. Pharmacol. Exp. Ther* 323 (2007) 294–307. [PubMed: 17625074]
- Tregellas JR, Wylie KP. Alpha7 Nicotinic Receptors as Therapeutic Targets in Schizophrenia. *Nicotine Tob Res*. 2019 2 18;21(3):349–356. doi: 10.1093/ntr/nty034. Review. [PubMed: 30137618]
- Umbricht D, Keefe RS, Murray S, Lowe DA, Porter R, Garibaldi G, Santarelli L. A randomized, placebo-controlled study investigating the nicotinic  $\alpha 7$  agonist, RG3487, for cognitive deficits in schizophrenia. *Neuropsychopharmacology*. 2014 6;39(7):1568–77. doi: 10.1038/npp.2014.17. [PubMed: 24549101]
- Undurraga J, Baldessarini RJ. Randomized, placebo-controlled trials of antidepressants for acute major depression: thirty-year meta-analytic review. *Neuropsychopharmacology*. 2012 3;37(4):851–64. doi: 10.1038/npp.2011.306. Review. [PubMed: 22169941]
- van Goethem NP, Paes D, Puzzo D, Fedele E, Reboisio C, Gulisano W, Palmeri A, Wennogle LP, Peng Y, Bertrand D, Prickaerts J. Antagonizing  $\alpha 7$  nicotinic receptors with methyllycaconitine (MLA) potentiates receptor activity and memory acquisition. *Cell Signal*. 2019 10;62:109338. doi: 10.1016/j.cellsig.2019.06.003.
- Wallace TL, Callahan PM, Tehim A, Bertrand D, Tombaugh G, Wang S, Xie W, Rowe WB, Ong V, Graham E, Terry AV Jr, Rodefer JS, Herbert B, Murray M, Porter R, Santarelli L, Lowe DA. RG3487, a novel nicotinic alpha7 receptor partial agonist, improves cognition and sensorimotor gating in rodents. *J. Pharmacol. Exp. Ther* 336 (2011) 242–253. [PubMed: 20959364]
- Winterer G, Gallinat J, Brinkmeyer J, Musso F, Kornhuber J, Thuerauf N, Rujescu D, Favis R, Sun Y, Franc MA, Ouwerkerk-Mahadevan S, Janssens L, Timmers M, Streffer JR. Allosteric alpha-7 nicotinic receptor modulation and P50 sensory gating in schizophrenia: a proof-of-mechanism study. *Neuropharmacology* 64 (2013) 197–204. [PubMed: 22766391]
- Wylie KP, Smucny J, Legget KT, Tregellas JR. Targeting Functional Biomarkers in Schizophrenia with Neuroimaging. *Curr Pharm Des*. 2016;22(14):2117–23. [PubMed: 26818860]
- Yang Y, Paspalas CD, Jin LE, Picciotto MR, Arnsten AF, Wang M. Nicotinic  $\alpha 7$  receptors enhance NMDA cognitive circuits in dorsolateral prefrontal cortex. *Proc Natl Acad Sci U S A*. 2013 7 16;110(29):12078–83. doi: 10.1073/pnas.1307849110 [PubMed: 23818597]

- Yin J, Chen W, Yang H, Xue M, Schaaf CP. Chrna7 deficient mice manifest no consistent neuropsychiatric and behavioral phenotypes. *Sci Rep.* 2017 1 3;7:39941. doi: 10.1038/srep39941. [PubMed: 28045139]
- Young JW, Geyer MA. Evaluating the role of the alpha-7 nicotinic acetylcholine receptor in the pathophysiology and treatment of schizophrenia. *Biochem Pharmacol.* 2013 10 15;86(8):112–232. doi: 10.1016/j.bcp.2013.06.031. Review.
- Zhang XY, Liu L, Liu S, Hong X, Chen DC, Xiu MH, Yang FD, Zhang Z, Zhang X, Kosten TA, Kosten TR. Short-term tropisetron treatment and cognitive and P50 auditory gating deficits in schizophrenia. *Am J Psychiatry.* 2012 9;169(9):974–81. doi: 10.1176/appi.ajp.2012.11081289. [PubMed: 22952075]

### Highlights

- Cognitive impairment is a core feature of schizophrenia that is debilitating.
- Currently, there are no clinically effective treatments for these impairments.
- $\alpha 7$ -nAChRs are considered viable therapeutic targets for cognition in schizophrenia.
- However, to date no  $\alpha 7$ -nAChR ligand has been approved for schizophrenia.
- This review discusses the relevant  $\alpha 7$ -nAChR literature and future directions.



**Fig 1.** Diagram illustrating several domains of cognition and other behaviors often targeted in drug discovery programs for schizophrenia and other neuropsychiatric disorders. The homomeric, low affinity  $\alpha_7$ -nicotinic acetylcholine receptor ( $\alpha_7$ -nAChR) is abundant in regions of the brain (e.g., hippocampus, prefrontal cortex) that are important for cognitive function. The receptor consists of five subunits arranged around a central channel that opens when endogenous ligands such as acetylcholine or exogenous ligands (nicotine) bind at the orthosteric site allowing cations (e.g.,  $Ca^{++}$ ) to flow through the channel into the neuron causing depolarization. Allosteric sites are the target of positive allosteric modulators (PAMs) and they are located at a site which is distinct from the orthosteric where they serve to indirectly influence (modulate) the effects of the agonist.

**Table 1.***In vivo* Pharmacological Effects of  $\alpha 7$ -nAChR Ligands

Compound Name	$\alpha 7$ nAChR activity & other actions	Cognitive Domain Enhanced	Additional Behavioral Improvements	References
ABBF	$\alpha 7$ full agonist & 5-HT3 antagonist	object and social recognition, working memory		Boess et al., 2007
ABT-107	$\alpha 7$ full agonist	attention, working memory, social recognition	auditory gating	Bitner et al., 2010; Bordia et al., 2015; Radek et al., 2012
A-582941	$\alpha 7$ full agonist	working memory, social recognition, inhibitory avoidance	sensory gating	Tietje et al., 2008
AQW051	$\alpha 7$ partial agonist	object and social recognition, spatial reference memory		Feuerbach et al., 2015
AVL-3288 (Compound 6, CCM1 or XY4083)	$\alpha 7$ Type I PAM	object recognition, social recognition, working memory, executive function	auditory gating, conditioned avoidance responding	Ng et al., 2007; Nikiforuk et al., 2015
BMS-933043	$\alpha 7$ partial agonist	object recognition, working memory, executive function	auditory gating	Bristow et al., 2016
BMS-902483	$\alpha 7$ partial agonist & 5-HT3 antagonist	object recognition, executive function	auditory gating	Pieschl et al., 2017
Compound 7z	$\alpha 7$ Type I PAM	object recognition		Hogenkamp et al., 2013
DMXB-A (GTS-21)	$\alpha 7$ partial agonist & 5-HT3 antagonist	attention, object recognition, spatial reference memory, working memory	sensorimotor gating	Meyer et al, 1997; Callahan et al., 2014; Jones et al., 2014
EVP-6124 (encenicline)	$\alpha 7$ partial agonist & 5-HT3 antagonist	object recognition		Prickaerts et al., 2012
EVP-5141	$\alpha 7$ full agonist & 5-HT3 antagonist	object and social recognition, working memory		Boess et al., 2013
JNJ-39393406	$\alpha 7$ Type I, Type II PAM	object recognition, executive function	auditory gating	Winterer et al., 2013
Lu AF58801	$\alpha 7$ Type I PAM	object recognition		Eskildsen et al., 2014
NS1738	$\alpha 7$ Type I PAM	object and social recognition, spatial reference memory		Marcus et al., 2016; Timmermann et al., 2007
PAM-2	$\alpha 7$ Type II PAM	object recognition, executive function	social interaction model	Potasiewicz et al., 2017
PNU 120596	$\alpha 7$ Type II PAM	recognition memory, spatial learning, working memory, executive function	auditory gating	Dunlop et al., 2009; Nikiforuk et al., 2015; Stevens et al., 2015
PNU-282987	$\alpha 7$ full agonist	recognition memory	conditioned avoidance responding	McLean et al., 2016; Marcus et al., 2016
RG 3487	$\alpha 7$ partial agonist & 5-HT3 antagonist	attention, object recognition, spatial reference memory, executive function	sensorimotor gating	Wallace et al., 2011
SEN-12333	$\alpha 7$ full agonist & histamine H3 antagonist	object recognition	sensorimotor gating	Roncarati et al., 2009
SSR-180711	$\alpha 7$ partial agonist	recognition memory, working memory, novelty discrimination	latent inhibition	Pichat et al., 2007; Barak et al., 2009
TC-5619	$\alpha 7$ full agonist	object recognition	sensorimotor gating	Hauser et al., 2009

<b>Compound Name</b>	<b><math>\alpha 7</math> nAChR activity &amp; other actions</b>	<b>Cognitive Domain Enhanced</b>	<b>Additional Behavioral Improvements</b>	<b>References</b>
Tropisetron	$\alpha 7$ partial agonist & 5-HT <sub>3</sub> antagonist	attention, object recognition, working memory, spatial reference memory	sensorimotor gating	Callahan et al., 2017; Hashimoto et al., 2005, 2006; Kohnomi et al., 2010; Pitsikas et al., 1997

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 2.**

$\alpha$ 7-nAChR ligands that have been evaluated in human clinical trials for pro-cognitive effects

Compound Name	$\alpha$ 7 nAChR activity & other actions	Clinical Stage	Clinical Endpoints	References
ABT-126	Partial agonist	Phase II	Cognition, Negative Symptomology	Haig et al., 2016; Haig et al., 2016; Haig et al., 2018
AQW051	Partial agonist	Phase II	Cognition, fMRI Brain Activation	Barch et al., 2016
AVL-3288	PAM	Phase Ib	Cognition, Auditory Sensory Gating Negative Symptomology	Gee et al., 2017; Kantrowitz et al., 2020
Cytidine 5'-diphosphocholine (CDP-choline)	Full agonist	Phase II	Cognition, Auditory Sensory Gating	Knott et al., 2015; Aidelbaum et al., 2018; Choueiry et al., 2019
DMXB-A (GTS-21)	Partial agonist	Phase II	Cognition, Auditory Sensory Gating, Negative Symptomology	Olincy et al., 2006; Freedman et al., 2008
Encenicline (EVP-6124)	Partial agonist/5-HT3 antagonist	Phase III	Cognition, Negative Symptomology	Keefe et al., 2015
Galantamine	PAM/acetylcholinesterase inhibitor	Phase II	Cognition, Auditory Sensory Gating, Negative Symptomology	Choueiry et al., 2019; Buchanan et al., 2017
JNJ-39393406	PAM	Phase Ib	Cognition, Auditory Sensory Gating, Smoking Cessation	Winterer et al., 2013; Perkins et al., 2018
RG3487	Partial agonist/5-HT3 antagonist	Phase II	Cognition, Negative Symptomology	Umbricht et al., 2014
Tropisetron	Partial agonist/5-HT3 antagonist	Phase II	Cognition, Auditory Sensory Gating, Negative Symptomology	Shiina et al., 2010; Zhang et al., 2012; Noroozian et al., 2013
TC-5619	Full agonist	Phase II	Cognition, Negative Symptomology	Lieberman et al., 2013

**Table 3.**

Limitations of Preclinical and Clinical Studies of pro-cognitive agents to date

<b>Preclinical</b>	<b>Clinical</b>
Translational challenges due to the poor understanding of the etiology and pathophysiology of schizophrenia	Studies often underpowered due to the inclusion of cognitively "normal" patients
Overreliance on rodent models	Polypharmacy and drug exposure history not properly addressed
Overreliance on acute dose-effect analyses	Unanticipated practice effects masking a positive outcomes
Pro-cognitive agents not evaluated in test subjects that have been chronically treated with antipsychotic.	Chronicity of the illness not taken into account. Recently diagnosed and first episode patients should be evaluated.
Pro-cognitive agents not evaluated in test subjects that have been chronically treated with nicotine or nicotine plus antipsychotics	Inclusion of inexperienced trial sites and focus on volume of patients enrolled as opposed to the quality of the recruitment sites

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript