



Published in final edited form as:

J Am Chem Soc. 2021 January 20; 143(2): 687–692. doi:10.1021/jacs.0c12484.

Rapid Construction of Tetralin, Chromane, and Indane Motifs via Cyclative C–H/C–H Coupling: Four-Step Total Synthesis of (±)-Russujaponol F


Zhe Zhuang[†], Alastair N. Herron[†], Shuang Liu[†], Jin-Quan Yu^{*†}

[†] Department of Chemistry, The Scripps Research Institute, 10550 N. Torrey Pines Road, La Jolla, California 92037, United States

Abstract

The development of practical C–H/C–H coupling reactions remains a challenging yet appealing synthetic venture because it circumvents the need to prefunctionalize both coupling partners for the generation of C–C bonds. Herein, we report a cyclative C(sp³)-H/C(sp²)-H coupling reaction of free aliphatic acids enabled by a cyclopentane-based mono-*N*-protected β-amino acid ligand. This reaction uses inexpensive sodium percarbonate (Na₂CO₃·1.5H₂O₂) as the sole oxidant, generating water as the only byproduct. A range of biologically important scaffolds, including tetralins, chromanes, and indanes, could be easily prepared by this protocol. Finally, the synthetic application of this methodology is demonstrated by the concise total synthesis of (±)-russujaponol F in a four-step sequence starting from readily available phenylacetic acid and pivalic acid through the sequential functionalizations of four C–H bonds.

Graphical abstract



*Corresponding Author yu200@scripps.edu.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Full experimental details and characterization of new compounds (PDF)

The authors declare no competing financial interests.

Carbon-carbon (C–C) bond formation constitutes one of the most important classes of reactions in organic synthesis. Owing to its potential to shorten synthesis, the past two decades have witnessed rapid developments in using C–H activation strategies for the construction of C–C bonds.¹ While most coupling methods require prefunctionalized coupling partners (e.g. organoborons and organohalides), C–H/C–H coupling reactions offer a complementary strategy to construct a C–C bond directly from two simple C–H bonds (Scheme 1A).² Compared to traditional coupling methods, this green and atom-economical approach is highly attractive because water is potentially the sole stoichiometric byproduct of this process (Scheme 1A). To date, extensive studies have focused on the coupling of two relatively reactive C(sp²)–H bonds for biaryl synthesis,³ whereas only a few reactions have been reported for the construction of more challenging C(sp³)–C(sp²) bonds. Because these existing reaction protocols require exogenous directing groups (DGs) to promote cyclometallation, additional steps to install and remove the DG are necessary.^{5,6} Additionally, reported methods pose practical limitations, such as the stoichiometric use of precious silver salts^{4b,c,5,6b,c} and harsh conditions^{4b,c,5a,b,6} — with temperatures as high as 160 °C being reported. Moreover, current methods for C(sp³)–H/C(sp²)–H coupling initiated by C(sp³)–H activation are largely limited to more reactive heterocyclic C(sp²)–H bonds.^{5a,b,6} Despite the great value that C–H/C–H coupling reactions might have for organic synthesis, the development of C(sp³)–H/C(sp²)–H coupling reactions that use both a practical oxidant and native substrates remains a significant challenge.

Recent advances in C–H functionalization have provided chemists with creative and strategic retrosynthetic disconnections that are otherwise difficult to achieve using traditional methods.⁷ However, for C–H functionalization strategies to truly improve the overall efficiency of synthesis, three criteria should be met: (1) the ability to use a wide range of simple starting materials to enable the synthesis of diverse natural product families; (2) the use of native functionalities as the DG; (3) the site-selectivity of C–H functionalization reactions should be precisely controllable. Given the ubiquitous nature of C–H bonds in organic molecules, synthetic sequences that incorporate multiple C–H functionalizations are particularly attractive for the efficient synthesis of natural products. However, approaches that meet these aforementioned criteria are challenging to execute and so uncommon in literature.^{7a,8}

To address these challenges, we herein report a cyclative C(sp³)–H/C(sp²)–H coupling reaction using a native free carboxylic acid as the DG (Scheme 1B). The use of a cyclopentane-based mono-*N*-protected β-amino acid ligand and a practical and inexpensive oxidant sodium percarbonate (Na₂CO₃·1.5H₂O₂) proved crucial to the success of this reaction. Tetralins, chromanes, or indanes, common frameworks in natural products could be readily prepared by this protocol (Figure 1). The synthetic application of this methodology is further demonstrated by a concise total synthesis of (±)-russujaponol F (the shortest and highest yielding to date) via multiple C–H functionalizations in four steps from readily available phenylacetic acid and pivalic acid (Scheme 1C), demonstrating the potential of C–H activation disconnections to enhance the ideality of synthesis⁹.

Aliphatic carboxylic acids are ubiquitous and synthetically versatile motifs and are often inexpensive reagents in organic chemistry; as such, they are privileged substrates for C–H

activation reactions.¹⁰ Following our recent disclosure of the β -C(sp³)-H lactonization¹⁰ⁱ and acyloxylation^{10j} of free carboxylic acids using *tert*-butyl hydrogen peroxide (TBHP) as the sole oxidant, we initiated our investigation of cyclative C(sp³)-H/C(sp²)-H coupling reactions by selecting TBHP as the bystander oxidant and aliphatic acid **1a** as a model substrate. Under the optimal conditions of the aforementioned β -acyloxylation reaction^{10j}, we were delighted to observe a 50% ¹H NMR yield of the desired product **2a** without forming competing reductive elimination products, such as the β -lactone or β -hydroxy acid (see Supporting Information, Table S1). Further investigation of the bystander oxidants and bases revealed that the combination of Na₂CO₃·1.5H₂O₂ and LiOAc could further improve the yield to 57% (see Supporting Information, Table S1–S2). The yields using LiOAc are generally better than those using NaOAc as the metal additives under the optimized conditions (see Supporting Information, Table S4). The use of sodium percarbonate, one of the cheapest and most easily handled oxidants,¹¹ potentially renders this protocol practical and scalable. In light of recent advances in ligand-accelerated Pd(II)-catalyzed C–H activation,¹² we next searched for ligands that could substantially improve the reactivity of the catalyst. Guided by mono-*N*-protected amino acid (MPAA) ligand-enabled C(sp³)-H activation reactions of free carboxylic acids^{10c,d,g,i,j}, we tested a series of commercially available MPAA ligands (**L1–L4**): β -amino acid ligand **L4** showed superior reactivity over α -amino acid ligands **L1–L3** (57% vs. 19–45%), as was also observed in other C(sp³)-H functionalization reactions of free acids via Pd(II)/Pd(IV) catalytic cycles^{10d,i,j}. Through systematic modifications to the backbone of the β -amino acid ligand (**L5–L10**), we found that *cis*-cyclopentane-based ligand (\pm)-**L9** gave the optimal reactivity (78% isolated yield). The superior reactivity of (\pm)-**L9** might be attributed to the more rigid conformation enforced by the cyclopentane linkage. Control experiments showed that the yields were low in the absence of the ligand or in the presence of the γ -amino acid ligand (**L11**) (23% or 20%, respectively), indicating the importance of six-membered chelation by the ligand for reactivity.

With the optimal ligand and reaction conditions in hand, we evaluated the scope of the cyclative C(sp³)-H/C(sp²)-H coupling reaction (Table 2). A wide range of tertiary aliphatic acids bearing a single α -methyl group (**1a–1e** and **1h**) or an α -gem-dimethyl group (**1f** and **1g**) were all compatible, affording the tetralin products in moderate to good yields (52–78%). The reaction could also be conducted on a 2.0 mmol scale, delivering **2a** in 69% yield. The attempted desymmetrization of the α -gem-dimethyl group of **1f** using enantioenriched **L9** resulted in racemic product. Less reactive free carboxylic acids containing α -hydrogens (**1i–1l**) also reacted in synthetically useful yields (35–65%). Among these, a variety of functionalities on the aryl rings such as methyl (**2b**), methoxy (**2j** and **2k**), fluoro (**2c**, **2g**, and **2l**), and chloro (**2d**) as well as naphthyl (**2e**) were tolerated, with the halogen moiety (**2d**) serving as a useful synthetic handle for subsequent derivatization. This protocol could also be successfully extended to the synthesis of biologically important chromane products. β -Phenoxy carboxylic acids containing an α -gem-dimethyl group (**1m–1r**) or α -hydrogens (**1s**, from Roche ester) were all reactive substrates. While a range of electron-donating (methoxy, *tert*-butyl, cyclohexyl, and benzyl) (**2s** and **2n–2p**) groups on the aryl ring were well tolerated to afford the desired products in good yields (70–85%), aliphatic acids containing electron-withdrawing (bromo and trifluoromethyl) groups (**2q** and

2r) showed comparatively low reactivity (31% and 23%), likely due to the sluggish nature of C(sp²)-H activations of electron-deficient arenes. Although intermolecular KIE experiments of electron-rich **1m** and **1m-d₅** ($k_H/k_D = 1.1$) suggest that C(sp²)-H activation is not the rate-determining step (see Supporting Information, KIE experiments section), the possibility of C(sp²)-H activation with electron-deficient substrates as the rate-determining step cannot be ruled out. It is noteworthy that high regioselectivity was observed for the aliphatic acids containing *meta*-methoxy groups (**1j** and **1s**), while the substrate bearing a *tert*-butyl group (**1n**) afforded a mixture of regioisomers (**2n/2n'** = 3/1). Considering the previously observed high *para*-selectivity of Pd(IV)-mediated C-H coupling reactions of anisoles^{3g,5c,22}, it is likely that the alkyl-Pd(II) intermediate is oxidized to Pd(IV) prior to C-H activation. Under the present conditions, the carboxylic acid **1t** containing either NBoc or NTs groups failed to deliver tetrahydroisoquinoline (THIQ) product **2t**. This cyclative C-H/C-H coupling reaction was also amenable to the syntheses of indane scaffolds (**2u-2w**). Notably, an [F⁺] oxidant^{3g,13} (1-fluoro-2,4,6-trimethylpyridinium tetrafluoroborate) showed superior reactivity for tertiary aliphatic acids containing an α -gem-dimethyl group (**2v** and **2w**) (see Supporting Information, Table S5).

Illudalane sesquiterpenes comprise a large family of natural products, which typically feature an indane core (for which various oxidation states are possible) bearing a challenging all-carbon quaternary center (Scheme 2A).¹⁴ Owing to their promising biological activities, tremendous efforts have been devoted to the total syntheses of these targets.^{15,16} Given the power of this methodology for the construction of indane scaffolds, we embarked on the total synthesis of (\pm)-russujaponol F via multiple C-H functionalizations (Scheme 2B). Baudoin's group reported the first total synthesis of russujaponol F in racemic and enantioselective forms based on a C(sp³)-H arylation strategy in 13 steps (26% yield) and 15 steps (12% yield) respectively.¹⁵ Beginning with phenylacetic acid **3** that is commercially available or synthesized through *ortho*-C-H methylation¹⁷, we were able to prepare aryl iodide **4** by esterification and subsequent mono-iodination¹⁸ of **3** using I₂ and Selectfluor in 79% yield. Investigation of the C-H arylation of pivalic acid indicated that, with ligand **L12**^{10f,19}, the mono-arylated product **5** could be obtained in 62% yield, along with 12% of the cyclative C-H/C-H coupling product **6** (see Supporting Information, Table S6). The formation of **6** under these conditions might be attributed to a second arylation of **5** with additional aryl iodide serving as the bystander oxidant.²⁰ The cyclative C-H/C-H coupling was then performed under the standard conditions using an [F⁺] oxidant to give the desired product **6** in 41% yield. Finally, global reduction of **6** using LAH cleanly delivered (\pm)-russujaponol F in 96% yield, completing the total synthesis in four steps and 28% overall yield: the shortest and highest yielding total synthesis of russujaponol F to date.

On the basis of literature precedents³⁻⁵ and our recent work on the C-H activation of free acids^{10i,j}, we propose that this cyclative C(sp³)-H/C(sp²)-H coupling reaction proceeds via a Pd(II)/Pd(IV) catalytic cycle as outlined in Scheme 3. First, coordination of Pd(OAc)₂ to an MPAA ligand generates the active LPd(II) species. After coordination of the acid substrate **1** to Pd, both the counteranion Na⁺ or Li⁺ and the MPAA ligand accelerate the cyclopalladation of the β -C(sp³)-H bond to form **int-I**. Next, oxidative addition of the hydrogen peroxide occurs to produce **int-II**, a process established in previous studies on the

oxidation of Pd(II) to Pd(IV) by benzoyl peroxide^{21a}, *tert*-butyl peroxyacetate^{21b}, or hydrogen peroxide^{21c,d}. In the previously reported β -lactonization¹⁰ⁱ or acetoxylation^{10j} reactions, selective reductive elimination yields β -lactone or β -acetoxyated carboxylic acid. In this case, a reactive phenyl group on the side chain of the substrate undergoes a second C(sp²)-H activation of **int-II** to deliver **int-III** via a seven or six-membered palladacycle, enabled by the facile dissociation of the weakly coordinating free acid.²² However, it is also possible that the Pd(II) intermediate **int-I** performs the second C-H activation prior to the oxidative addition of hydrogen peroxide that generates **int-III**. Finally, reductive elimination of **int-III** generates the cyclative C-H/C-H coupling product **2** and regenerates the LPd(II) species.

In summary, we have realized a Pd(II)-catalyzed cyclative C(sp³)-H/C(sp²)-H coupling reaction enabled by a cyclopentane-based mono-*N*-protected β -amino acid ligand. The use of inexpensive sodium percarbonate as the sole oxidant and native free carboxylic acids as the directing group renders this reaction highly practical and potentially amenable to large-scale manufacturing. A range of biologically significant scaffolds, including tetralins, chromanes, and indanes, could be readily prepared by this protocol. The synthetic application of this methodology was demonstrated by a concise total synthesis of (\pm)-russujaponol F via multiple C-H functionalizations in four steps from readily available phenylacetic acid and pivalic acid.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

We gratefully acknowledge the NIH (NIGMS, R01GM084019), the NSF under the CCI Center for Selective C-H Functionalization (CHE-1700982), and The Scripps Research Institute for financial support. Z.Z. thanks Dr. Nelson Y. S. Lam for proofreading.




REFERENCES

- (1). For reviews on C-H activation/C-C bond-forming reactions, see:(a)Chen X.; Engle KM; Wang D-H; Yu J-Q Palladium(II)-catalyzed C-H activation/C-C cross-coupling reactions: versatility and practicality. *Angew. Chem., Int. Ed.* 2009, 48, 5094–5115.(b)Daugulis O.; Roane J.; Tran LD Bidentate, monoanionic auxiliary-directed functionalization of carbon-hydrogen bonds. *Acc. Chem. Res.* 2015, 48, 1053–1064. [PubMed: 25756616] (c)He G.; Wang B.; Nack WA; Chen G. Syntheses and transformations of α -amino acids via palladium-catalyzed auxiliary directed sp³ C-H functionalization. *Acc. Chem. Res.* 2016, 49, 635–645. [PubMed: 27015079]
- (2). For reviews on C-H/C-H coupling reactions, see:(a)Yeung CS; Dong VM Catalytic dehydrogenative cross-coupling: forming carbon-carbon bonds by oxidizing two carbon-hydrogen bonds. *Chem. Rev.* 2011, 111, 1215–1292. [PubMed: 21391561] (b)Girard SA; Knauber T.; Li C-J The cross-dehydrogenative coupling of C(sp³)-H bonds: a versatile strategy for C-C bond formations. *Angew. Chem., Int. Ed.* 2014, 53, 74–100.(c)Liu C.; Yuan J.; Gao M.; Tang S.; Li W.; Shi R.; Lei A. Oxidative coupling between two hydrocarbons: an update of recent C-H functionalizations. *Chem. Rev.* 2015, 115, 12138–12204. [PubMed: 26558751]
- (3). For early examples of C(sp²)-H/C(sp²)-H coupling reaction, see:(a)Stuart DR; Fagnou K. The catalytic cross-coupling of unactivated arenes. *Science* 2007, 316, 1172–1175. [PubMed: 17525334] (b)Xia J-B; You S-L Carbon-carbon bond formation through double sp² C-H activations: synthesis of ferrocenyl oxazoline derivatives. *Organometallics* 2007, 26, 4869–4871.



- (c)Hull KL; Sanford MS Catalytic and highly regioselective cross-coupling of aromatic C–H substrates. *J. Am. Chem. Soc.* 2007, 129, 11904–11905. [PubMed: 17845047] (d)Brasche G.; García-Fortanet J.; Buchwald SL Twofold C–H functionalization: palladium-catalyzed *ortho* arylation of anilides. *Org. Lett.* 2008, 10, 2207–2210. [PubMed: 18465866] (e)Cho SH; Hwang SJ; Chang S. Palladium-catalyzed C–H functionalization of pyridine *N*-oxides: highly selective alkenylation and direct arylation with unactivated arenes. *J. Am. Chem. Soc.* 2008, 130, 9254–9256. [PubMed: 18582040] (f)Zhao X.; Yeung CS; Dong VM Palladium-catalyzed *ortho*-arylation of *O*-phenylcarbamates with simple arenes and sodium persulfate. *J. Am. Chem. Soc.* 2010, 132, 5837–5844. [PubMed: 20359220] (g)Wang X.; Leow D.; Yu J-Q Pd(II)-catalyzed para-selective C–H arylation of monosubstituted arenes. *J. Am. Chem. Soc.* 2011, 133, 13864–13867. [PubMed: 21834576]
- (4). For Pd-catalyzed C(sp³)–H/C(sp²)–H coupling reactions initiated by C(sp²)–H activation, see: (a)Liésgault B.; Fagnou K. Palladium-catalyzed intramolecular coupling of arenes and unactivated alkanes in air. *Organometallics* 2008, 27, 4841–4843. (b)Pierre C.; Baudoin O. Intramolecular PdII-catalyzed dehydrogenative C(sp³)–C(sp²) coupling: an alternative to Pd⁰-catalyzed C(sp³)–H arylation from aryl halides? *Tetrahedron* 2013, 69, 4473–4478. (c)Shi J-L; Wang D.; Zhang X-S; Li X-L; Chen Y-Q; Li Y-X; Shi Z-J Oxidative coupling of sp² and sp³ carbon–hydrogen bonds to construct dihydrobenzofurans. *Nat. Commun.* 2017, 8, 238–244. [PubMed: 28798314]
- (5). For Pd-catalyzed C(sp³)–H/C(sp²)–H coupling reactions initiated by C(sp³)–H activation, see: (a)Jiang Y.; Deng G.; Zhang S.; Loh T-P Directing group participated benzylic C(sp³)–H/C(sp²)–H cross-dehydrogenative coupling (CDC): synthesis of azapolycycles. *Org. Lett.* 2018, 20, 652–655. [PubMed: 29328666] (b)Sun W-W; Liu J-K; Wu B. Practical synthesis of polysubstituted unsymmetric 1,10-phenanthrolines by palladium catalyzed intramolecular oxidative cross coupling of C(sp²)–H and C(sp³)–H bonds of carboxamides. *Org. Chem. Front.* 2019, 6, 544–550. (c)Hao H-Y; Mao Y-J; Xu Z-Y; Lou S-J; Xu D-Q Selective cross-dehydrogenative C(sp³)–H arylation with arenes. *Org. Lett.* 2020, 22, 2396–2402. [PubMed: 32124610]
- (6). For other metal-enabled C(sp³)–H/C(sp²)–H coupling reactions, see: (a)Wu X.; Zhao Y.; Ge H. Pyridine-enabled copper-promoted cross dehydrogenative coupling of C(sp²)–H and unactivated C(sp³)–H bonds. *Chem. Sci.* 2015, 6, 5978–5983. [PubMed: 29861919] (b)Tan G.; You J. Rhodium(III)-catalyzed oxidative cross-coupling of unreactive C(sp³)–H bonds with C(sp²)–H bonds. *Org. Lett.* 2017, 19, 4782–4785. [PubMed: 28853579] (c)Wang X.; Xie P.; Qiu R.; Zhu L.; Liu T.; Li Y.; Iwasaki T.; Au C-T; Xu X.; Xia Y.; Yin S-F; Kambe N. Nickel-catalysed direct alkylation of thiophenes via double C(sp³)–H/C(sp²)–H bond cleavage: the importance of KH₂PO₄. *Chem. Commun.* 2017, 53, 8316–8319. (d)Tan G.; Zhang L.; Liao X.; Shi Y.; Wu Y.; Yang Y.; You J. Copper- or nickel-enabled oxidative cross-coupling of unreactive C(sp³)–H bonds with azole C(sp²)–H bonds: rapid access to β-azolyl propanoic acid derivatives. *Org. Lett.* 2017, 19, 4830–4833. [PubMed: 28853578]
- (7). For reviews on C–H functionalization for natural product synthesis, see: (a)Baudoin O. Multiple catalytic C–H bond functionalization for natural product synthesis. *Angew. Chem., Int. Ed.* 2020, 59, 17798–17809. (b)Lam NYS; Wu K.; Yu J-Q Advancing the logic of chemical synthesis: C–H activation as strategic and tactical disconnections for C–C bond construction. *Angew. Chem., Int. Ed.* 2020, 59, DOI: 10.1002/anie.202011901. (c)Gutekunst WR; Baran PS C–H functionalization logic in total synthesis. *Chem. Soc. Rev.* 2011, 40, 1976–1991. [PubMed: 21298176] (d)Abrams DJ; Provencher PA; Sorensen EJ Recent applications of C–H functionalization in complex natural product synthesis. *Chem. Soc. Rev.* 2018, 47, 8925–8967. [PubMed: 30426998]
- (8). For selected examples of total synthesis using multiple C–H functionalizations, see: (a)Wang D-H; Yu J-Q Highly convergent total synthesis of (+)-lithospermic acid via a late-stage intermolecular C–H olefination. *J. Am. Chem. Soc.* 2011, 133, 5767–5769. [PubMed: 21443224] (b)Gutekunst WR; Baran PS Total synthesis and structural revision of the piperarborenines via sequential cyclobutane C–H arylation. *J. Am. Chem. Soc.* 2011, 133, 19076–19079. [PubMed: 22066860] (c)Rosen BR; Simke LR; Thuy-Boun PS; Dixon DD; Yu J-Q; Baran PS C–H functionalization logic enables synthesis of (+)-hongoquercin A and related compounds. *Angew. Chem., Int. Ed.* 2013, 52, 7317–7320. (d)Hong B.; Li C.; Wang Z.; Chen J.; Li H.; Lei X. Enantioselective total synthesis of (–)-incarviateone A. *J. Am. Chem. Soc.* 2015, 137, 11946–11949. [PubMed: 25811111]

- 26371964] (e)Dailler D.; Danoun G.; Ourri B.; Baudoin O. Divergent synthesis of aeruginosins based on a C(sp³)-H activation strategy. *Chem. Eur. J.* 2015, 21, 9370–9379. [PubMed: 26031222] (f)Wu F.; Zhang J.; Song F.; Wang S.; Guo H.; Wei Q.; Dai H.; Chen X.; Xia X.; Liu X.; Zhang L.; Yu J-Q; Lei X. Chrysomycin A derivatives for the treatment of multi-drug-resistant tuberculosis. *ACS Cent. Sci.* 2020, 6, 928–938. [PubMed: 32607440]
- (9). Gaich T.; Baran PS Aiming for the ideal synthesis. *J. Org. Chem.* 2010, 75, 4657–4673. [PubMed: 20540516]
- (10). (a)Giri R.; Maugel N.; Li J-J; Wang D-H; Breazzano SP; Saunder LB; Yu J-Q Palladium-catalyzed methylation and arylation of sp² and sp³ C–H bonds in simple carboxylic acids. *J. Am. Chem. Soc.* 2007, 129, 3510–3511. [PubMed: 17335217] (b)Chen G.; Zhuang Z.; Li G-C; Saint-Denis TG; Hsiao Y.; Joe CL; Yu J-Q Ligand-enabled β-C–H arylation of α-amino acids without installing exogenous directing groups. *Angew. Chem., Int. Ed.* 2017, 56, 1506–1509.(c)Zhu Y.; Chen X.; Yuan C.; Li G.; Zhang J.; Zhao Y. Pd-catalysed ligand-enabled carboxylate-directed highly regioselective arylation of aliphatic acids. *Nat. Commun.* 2017, 8, 14904. [PubMed: 28383026] (d)Ghosh KK; van Gemmeren M. Pd-catalyzed β-C(sp³)-H arylation of propionic acid and related aliphatic acids. *Chem. Eur. J.* 2017, 23, 17697–17700. [PubMed: 29148233] (e)Shen P-X; Hu L.; Shao Q.; Hong K.; Yu J-Q Pd(II)-catalyzed enantioselective C(sp³)-H arylation of free carboxylic acids. *J. Am. Chem. Soc.* 2018, 140, 6545–6549. [PubMed: 29741883] (f)Zhuang Z.; Yu C-B; Chen G.; Wu Q-F; Hsiao Y.; Joe CL; Qiao JX; Poss MA; Yu J-Q Ligand-enabled β-C(sp³)-H olefination of free carboxylic acids. *J. Am. Chem. Soc.* 2018, 140, 10363–10367. [PubMed: 30029574] (g)Hu L.; Shen P-X; Shao Q.; Hong K.; Qiao JX; Yu J-Q PdII-catalyzed enantioselective C(sp³)-H activation/cross-coupling reactions of free carboxylic acids. *Angew. Chem., Int. Ed.* 2019, 58, 2134–2138.(h)Ghosh KK; Utrry A.; Koldemir A.; Ong M.; van Gemmeren M. Direct β-C(sp³)-H acetoxylation of aliphatic carboxylic acids. *Org. Lett.* 2019, 21, 7154–7157. [PubMed: 31398049] (i)Zhuang Z.; Yu J-Q Lactonization as a general route to β-C(sp³)-H functionalization. *Nature* 2020, 577, 656–659. [PubMed: 31825951] (j)Zhuang Z.; Herron AN; Fan Z.; Yu J-Q Ligand-enabled monoselective β-C(sp³)-H acyloxylation of free carboxylic acids using a practical oxidant. *J. Am. Chem. Soc.* 2020, 142, 6769–6776. [PubMed: 32200639] (k)Ghiringhelli F.; Utrry A.; Ghosh KK; van Gemmeren M. Direct β- and γ-C(sp³)-H alkynylation of free carboxylic acids. *Angew. Chem., Int. Ed.* 2020, 59, 23127–23131.
- (11). (a)McKillop A.; Sanderson WR Sodium perborate and sodium percarbonate: Cheap, safe and versatile oxidising agents for organic synthesis. *Tetrahedron Lett.* 1995, 51, 6145.(b)Muzart J. Sodium perborate and sodium percarbonate in organic synthesis. *Synthesis* 1995, 1325.
- (12). For reviews, see:(a)He J.; Wasa M.; Chan KSL; Shao Q.; Yu J-Q Palladium-catalyzed transformations of alkyl C–H bonds. *Chem. Rev.* 2017, 117, 8754–8786. [PubMed: 28697604] (b)Shao Q.; Wu K.; Zhuang Z.; Qian S.; Yu J-Q From Pd(OAc)₂ to chiral catalysts: the discovery and development of bifunctional mono-N-protected amino acid ligands for diverse C–H activation reactions. *Acc. Chem. Res.* 2020, 53, 833–851. [PubMed: 32227915]
- (13). Engle KM; Mei T-S; Wang X.; Yu J-Q Bystanding F⁺ oxidants enable selective reductive elimination from high-valent metal centers in catalysis. *Angew. Chem., Int. Ed.* 2011, 50, 1478–1491.
- (14). (a)Yoshikawa K.; Kaneko A.; Matsumoto Y.; Hama H.; Arihara S. Russujaponols A–F, illudoid sesquiterpenes from the fruiting body of *Russula japonica*. *J. Nat. Prod.* 2006, 69, 1267–1270. [PubMed: 16989517] (b)Yoshikawa K.; Matsumoto Y.; Hama H.; Tanaka M.; Zhai H.; Fukuyama Y.; Arihara S.; Hashimoto T. Russujaponols G–L, illudoid sesquiterpenes, and their neurite outgrowth promoting activity from the fruit body of *Russula japonica*. *Chem. Pharm. Bull.* 2009, 57, 311–314.(c)Becker U.; Erkel G.; Anke T.; Sterner O. Puraquinonic acid, a novel inducer of differentiation of human HL-60 promyelocytic leukemia cells from *Mycena pura* (Pers. Ex Fr.). *Nat. Prod. Lett.* 1997, 9, 229–236.(d)Kuroyanagi M.; Fukuoka M.; Yoshihira K.; Natori S. The absolute configurations of pterosins, 1-indanone derivatives from bracken, *Pteridium aquilinum* var. *latiusculum*. *Chem. Pharm. Bull.* 1974, 22, 723–726.(e)Suzuki S.; Murayama T.; Shiono Y. Echinolactones C and D: two illudalane sesquiterpenoids isolated from the cultured mycelia of the fungus *Echinodontium japonicum*. *Z. Naturforsch., B* 2006, 61, 1295–1298.
- (15). (a)Melot R.; Craveiro M.; Bürgi T.; Baudoin O. Divergent enantioselective synthesis of (nor)illudalane sesquiterpenes via Pd⁰-catalyzed asymmetric C(sp³)-H activation. *Org. Lett.*


- 2019, 21, 812–815. [PubMed: 30633535] (b)Melot R.; Craveiro MV; Baudoin O. Total synthesis of (nor)illudalane sesquiterpenes based on a C(sp³)-H activation strategy. *J. Org. Chem.* 2019, 84, 12933–12945. [PubMed: 31322348]
- (16). For recent examples, see:(a)Tiong EA; Rivalti D.; Williams BM; Gleason JL A concise total synthesis of (R)-puraquinonic acid. *Angew. Chem., Int. Ed.* 2013, 52, 3442–3445.(b)Elmehriki AAH; Gleason JL A spiroalkylation method for the stereoselective construction of α -quaternary carbons and its application to the total synthesis of (R)-puraquinonic acid. *Org. Lett.* 2019, 21, 9729–9733. [PubMed: 31756114] (c)Zeng Z.; Zhao Y.; Zhang Y. Divergent total syntheses of five illudalane sesquiterpenes and assignment of the absolute configuration. *Chem. Commun.* 2019, 55, 4250–4253.(d)Xun MM; Bai Y.; Wang Y.; Hu Z.; Fu K.; Ma W.; Yuan C. Synthesis of four illudalane sesquiterpenes utilizing a one-pot Diels–Alder/oxidative aromatization sequence. *Org. Lett.* 2019, 21, 6879–6883. [PubMed: 31441309]
- (17). Thuy-Boun PS; Villa G.; Dang D.; Richardson P.; Su S.; Yu J-Q Ligand-accelerated *ortho*-C–H alkylation of arylcarboxylic acids using alkyl boron reagents. *J. Am. Chem. Soc.* 2013, 135, 17508–17513. [PubMed: 24124892]
- (18). Stavber S.; Kralj P.; Zupan M. Selective and effective iodination of alkyl-substituted benzenes with elemental iodine activated by SelectfluorTM F-TEDA-BF₄. *Synlett* 2002, 598–600.
- (19). For examples of C–H activation reactions using L12, see:(a)Le KKA; Nguyen H.; Daugulis O. 1-Aminopyridinium ylides as monodentate directing groups for sp³ C–H bond functionalization. *J. Am. Chem. Soc.* 2019, 141, 14728–14735. [PubMed: 31529954] (b)Zhuang Z.; Yu J-Q Pd(II)-catalyzed enantioselective γ -C(sp³)-H functionalizations of free cyclopropylmethylamines. *J. Am. Chem. Soc.* 2020, 142, 12015–12019. [PubMed: 32605367]
- (20). (a)Sun W-W; Cao P.; Mei R-Q; Li Y.; Ma Y-L; Wu B. Palladium-catalyzed unactivated C(sp³)-H bond activation and intramolecular amination of carboxamides: a new approach to β -lactams. *Org. Lett.* 2014, 16, 480–483. [PubMed: 24341538] (b)Zhang S-J; Sun W-W; Cao P.; Dong X-P; Liu J-K; Wu B. Stereoselective synthesis of diazabicyclic β -lactams through intramolecular amination of unactivated C(sp³)-H Bonds of carboxamides by palladium catalysis. *J. Org. Chem.* 2016, 81, 956–968. [PubMed: 26745308] (c)Tong H-R; Zheng W.; Lv X.; He G.; Liu P.; Chen G. Asymmetric synthesis of β -lactam via palladium-catalyzed enantioselective intramolecular C(sp³)-H amidation. *ACS Catal.* 2020, 10, 114–120.(d)Zhou T.; Jiang M-X; Yang X.; Yue Q.; Han Y-Q; Ding Y.; Shi B-F Synthesis of chiral β -lactams by Pd-catalyzed enantioselective amidation of methylene C(sp³)-H bonds. *Chin. J. Chem.* 2020, 38, 242–246.
- (21). (a)Canty AJ; Jin H.; Skelton BW; White AH Oxidation of complexes by (O₂CPh)₂ and (ER)₂ (E = S, Se), including structures of Pd(CH₂CH₂CH₂CH₂)(SePh)₂(bpy) (bpy = 2,2'-bipyridine) and MMe₂(SePh)₂(L₂) (M = Pd, Pt; L₂ = bpy, 1,10-phenanthroline) and C–O and C–E bond formation at palladium(IV). *Inorg. Chem.* 1998, 37, 3975–3981. [PubMed: 11670512] (b)Giri R.; Liang J.; Lei J-G; Li J-J; Wang D-H; Chen X.; Naggar IC; Guo C.; Foxman BM; Yu J-Q Pd-catalyzed stereoselective oxidation of methyl groups by inexpensive oxidants under mild conditions: a dual role for carboxylic anhydrides in catalytic C–H bond oxidation. *Angew. Chem., Int. Ed.* 2005, 44, 7420–7424.(c)Oloo W.; Zavalij PY; Zhang J.; Khaskin E.; Vedernikov AN Preparation and C–X reductive elimination reactivity of monoaryl Pd(IV)-X complexes in water (X = OH, OH₂, Cl, Br). *J. Am. Chem. Soc.* 2010, 132, 14400–14402. [PubMed: 20866056] (d)Abada E.; Zavalij PY; Vedernikov AN Reductive C(sp²)-N elimination from isolated Pd(IV) amido aryl complexes prepared using H₂O₂ as oxidant. *J. Am. Chem. Soc.* 2017, 139, 643–646. [PubMed: 28048934]
- (22). For examples of C–H activation of arenes at Pd(IV), see:(a)Rosewall CF; Sibbald PA; Liskin DV; Michael FE Palladium-catalyzed carboamination of alkenes promoted by *N*-fluorobenzenesulfonimide via C–H activation of arenes. *J. Am. Chem. Soc.* 2009, 131, 9488–9489. [PubMed: 19545153] (b)Sibbald PA; Rosewall CF; Swartz RD; Michael FE Mechanism of *N*-fluorobenzenesulfonimide promoted diamination and carboamination reactions: divergent reactivity of a Pd(IV) species. *J. Am. Chem. Soc.* 2009, 131, 15945–15951. [PubMed: 19824646]

A C–H activation/C–C bond-forming reactions**B Cyclative C(sp³)-H/C(sp²)-H coupling reaction of free aliphatic acids****C Concise synthesis of (±)-russujaponol F via multiple C–H functionalizations**

Scheme 1.
C–H Activation/C–C Bond-Forming Reactions

A Illudalane sesquiterpenes: an indane core containing a quaternary center**B Total synthesis of (±)-russujaponol F****Scheme 2.**Total Synthesis of (±)-Russujaponol F^a

^aConditions: (a) SOCl₂, EtOH, reflux, overnight; I₂ (0.5 equiv), Selectfluor (0.5 equiv), CH₃CN, 60 °C, 3 h. (b) Pd(OAc)₂ (10 mol%), **L12** (10 mol%), pivalic acid (3.0 equiv), CsOAc (1.0 equiv), Ag₂CO₃ (2.0 equiv), HFIP, 80 °C, 12 h. (c) Pd(CH₃CN)₄(BF₄)₂ (10 mol %), Ag₂CO₃ (1.0 equiv), 1-fluoro-2,4,6-trimethylpyridinium tetrafluoroborate (2.0 equiv), HFIP, 90 °C, 12 h. (d) LAH (3.0 equiv), THF, 0 °C to rt, overnight.



Scheme 3.
Proposed Mechanism for Cyclative C(sp³)-H/C(sp²)-H Coupling Reaction





Figure 1. Biologically significant natural products containing tetralin, chromane, or indane frameworks.

Table 1.


Ligand Investigation for the Cyclative C(sp³)-H/C(sp²)-H Coupling Reaction^{a,b}

^aConditions: **1a** (0.1 mmol), Pd(OAc)₂ (10 mol%), ligand (L) (10 mol%), LiOAc (1.0 equiv), Na₂CO₃·1.5H₂O (2.0 equiv), HFIP (1.0 mL), 60 °C, 12 h.

^bThe yields were determined by ¹H NMR analysis of the crude product using CH₂Br₂ as the internal standard.

^cIsolated yield.

Table 2.

Substrate Scope of the Cyclative C(sp³)-H/C(sp²)-H Coupling Reaction^{a,b}

^a Conditions A: **1** (0.1 mmol), Pd(OAc)₂ (10 mol%), (±)-**L9** (10 mol%), LiOAc (1.0 equiv), Na₂CO₃·1.5H₂O₂ (2.0 equiv), HFIP (1.0 mL), 60 °C, 12 h.

^b Isolated yields.

^cConditions B: **1** (0.1 mmol), Pd(CH₃CN)₄(BF₄)₂ (10 mol%), Ag₂CO₃ (1.0 equiv), 1-fluoro-2,4,6-trimethylpyridinium tetrafluoroborate (2.0 equiv), HFIP (1.0 mL), 90 °C, 12 h. ^dOn 2.0 mmol scale.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript