



HHS Public Access

Author manuscript

Nat Rev Neurosci. Author manuscript; available in PMC 2021 May 28.

Published in final edited form as:

Nat Rev Neurosci. 2016 December ; 17(12): 777–792. doi:10.1038/nrn.2016.141.

Network abnormalities and interneuron dysfunction in Alzheimer disease

Jorge J. Palop^{1,2}, Lennart Mucke^{1,2}

¹Gladstone Institute of Neurological Disease, 1650 Owens Street, San Francisco, CA 94158, USA.

²Department of Neurology, University of California, San Francisco, 1650 Owens Street, San Francisco, CA 94158, USA.

Abstract

The function of neural circuits and networks can be controlled, in part, by modulating the synchrony of their components' activities. Network hypersynchrony and altered oscillatory rhythmic activity may contribute to cognitive abnormalities in Alzheimer disease (AD). In this condition, network activities that support cognition are altered decades before clinical disease onset, and these alterations predict future pathology and brain atrophy. Although the precise causes and pathophysiological consequences of these network alterations remain to be defined, interneuron dysfunction and network abnormalities have emerged as potential mechanisms of cognitive dysfunction in AD and related disorders. Here, we explore the concept that modulating these mechanisms may help improve brain function in these conditions.

Certain activities of neurons and neuronal networks are associated with the successful encoding of memories and retention of new information, and thus may be necessary for learning and memory. In Alzheimer disease (AD), schizophrenia, epilepsy and other neurological and psychiatric diseases that cause cognitive impairment, network activities supporting cognition are altered, even during preclinical stages^{1–5}, that is, before symptoms are noticed by the patient or can be detected by neurocognitive exams. These network alterations include activation and deactivation deficits, abnormal oscillatory rhythmic activity and network hypersynchrony. In people at high risk of developing AD, for example, abnormal activation and deactivation of specific networks during memory encoding can be detected decades before the predicted onset of clinical disease^{6–9}. As these functional network alterations widely overlap with the brain regions that ultimately develop pathological hallmarks and atrophy in AD, they may be a harbinger and possibly even a cause of clinical disease manifestations.

Robust network alterations are associated with diverse cognitive disorders, but the mechanisms and pathophysiological consequences of the alterations are poorly understood. Do alterations in network activity contribute to cognitive impairment or are they incidental

Correspondence to: J.J.P. and L.M. jorge.palop@gladstone.ucsf.edu and lennart.mucke@gladstone.ucsf.edu.

Competing interests statement

The authors declare completing interests. See Web version for details.

by-products of disease-induced cellular dysfunction? Do alterations in network synchrony cause the dysfunction of microcircuits, larger distributed networks, or both? Most importantly, could cognitive alterations be prevented and even reversed by improving the function of cells that promote specific network activities?

Recent findings suggest that altered network activity can indeed contribute to cognitive impairment in AD and that network activities can be experimentally or behaviorally manipulated to improve cognitive functions in patients at risk of AD and related mouse models. In this Review, we explore two main concepts — that network abnormalities and interneuron dysfunction contribute to cognitive deficits in AD and related conditions, and that blocking or counteracting these mechanisms could be of both symptomatic and disease-modifying therapeutic value.

Neuronal synchrony and brain function

Cognitive functions depend on brain states that range from maximal focus and concentration to inattentiveness, drowsiness and sleep. These states probably do not reflect a continuous functional gradient of neuronal activities but rather represent distinct operating modes of brain activity that are closely linked to — and possibly determined by — changes in neuronal synchrony, the degree to which neuronal activities are correlated. Such synchrony fluctuates considerably with behavioural and brain states¹⁰. The degree of synchrony among neuronal populations (network synchrony) can be determined by measuring and correlating neuronal activities at multiple sites, for example, through electroencephalography (EEG) or the recording of local field potentials (LFPs) with multielectrode arrays. During non-active states, such as slow-wave sleep or quiet wakefulness, cortical neuronal activities at different sites tend to be synchronized and slowly fluctuate at high amplitudes (FIG. 1a). During active behaviours, such as paying attention or learning, this highly synchronized pattern of brain activity abruptly ceases, and neuronal activities at different sites become desynchronized and fluctuate at higher frequencies and smaller amplitudes¹¹ (FIG. 1a).

The fundamental relationship between the synchrony of neuronal activities and functional states is also evident at the circuit level (FIG. 1b). When mice are resting, adjacent layer 2 pyramidal cells in the somatosensory cortex show synchronous high-amplitude fluctuations in membrane potentials, which result in relatively low signal-to-noise ratios for evoking action potentials relative to ongoing fluctuations; however, with active whisker use, this high-amplitude synchrony of membrane potentials ceases abruptly, and signal-to-noise ratios increase^{12,13}. Thus, increasing synchrony among neurons and recruiting them into slow-frequency and high-amplitude fluctuations seem to be basic mechanisms for ‘switching off’ specific brain regions. By contrast, the suppression of high-amplitude fluctuations and desynchronization of neuronal activities increases the ability of neurons to respond to external information-rich inputs¹². Synchrony and functional states are closely related in many neocortical and hippocampal regions, suggesting that synchrony reflects a fundamental principle of nervous system design. Here, we propose that deficits in synchrony are an important pathogenic mechanism of AD-related cognitive dysfunction.

Amyloid- β and AD pathogenesis

The hypothesis that amyloid- β has a causal role in AD pathogenesis is supported by diverse lines of evidence, including the early accumulation of amyloid- β in the brain in people who go on to develop AD and the ability of mutations in the genes encoding the amyloid precursor protein (APP) or presenilins (PSENs), which alter amyloid- β production, to cause autosomal dominant, early-onset familial AD (FAD)¹⁴. Amyloid- β peptides are proteolytically released from APP by β -site APP-cleaving enzyme 1 (BACE1; also known as β -secretase) and γ -secretase, and exist in diverse assembly states, including monomers, oligomers, fibrils and amyloid plaques¹⁵. FAD is caused by *APP* duplications, mutations in *APP* located around the BACE 1 and γ -secretase cleavage sites or within the amyloid- β coding sequence, or by mutations in the genes that encode PSEN1 and PSEN2 (alternative catalytic subunits of the γ -secretase complex). Over 250 genetic alterations have been linked to FAD. Those alterations that have been tested for effects on APP metabolism increase the overall production of amyloid- β or the amyloid- β_{1-42} /amyloid- β_{1-40} ratio¹⁶⁻¹⁹, or promote the accumulation of pathogenic amyloid- β assemblies²⁰⁻²². Conversely, an *APP* mutation (A673T) that reduces the β -secretase-mediated cleavage of APP and, consequently, amyloid- β levels decreases the risk of late-onset sporadic AD^{23,24}.

Amyloid- β oligomers elicit abnormalities in synaptic and cognitive functions *in vivo* and *in vitro*^{15,25-30}. Neuronal expression of FAD-mutant human or humanized APP (with or without co-expression of FAD-mutant PSEN1 or PSEN2) in transgenic mice simulates key aspects of AD, including elevated levels of human amyloid- β , amyloid plaques, neuritic dystrophy, synaptodendritic impairments, astrogliosis, microgliosis, vasculopathy, network dysfunction, cognitive deficits and behavioural alterations³¹. We refer to these models collectively as FAD mice (Supplementary information S1 (table)). An example of such models is highlighted in Supplementary information S2 (table), which lists the AD-like alterations observed in the hAPP-J20 mouse model, which was generated in our laboratory. In combination, the above findings strongly support the amyloid- β hypothesis, which includes the notion that amyloid- β is a critical cause of cognitive decline in AD. Notably, several other factors, including tau accumulation, apolipoprotein E4 (APOE4) and inflammatory mediators, contribute to the complex pathogenesis of AD and can be introduced into FAD models through additional genetic modifications³²⁻³⁵.

A major unresolved question in the field is which forms of amyloid- β , tau and APOE4 are the most neurotoxic. Lead suspects include soluble amyloid- β_{1-42} oligomers^{15,30}, abnormal assemblies or post-translational modifications of non-fibrillar tau³⁵⁻³⁸ and APOE4 fragments³⁹. The importance of answering this question is illustrated by the following observations. Approximately 20% of healthy aged individuals, 60% of patients with mild cognitive impairment (MCI) and almost all patients with AD exhibit evidence of amyloid deposition in the cortex, as shown by positron emission tomography (PET) involving radiopharmaceutical ligands that bind to amyloid plaques such as Pittsburgh compound B (PiB)⁴⁰. In healthy aged individuals and patients with MCI, cortical amyloid load correlated inversely with episodic memory performance⁴⁰; this correlation weakened with amyloid accumulation and age and does not exist in AD^{40,41}. Discrepancies between amyloid deposition and cognitive dysfunction may be explained, at least in part, by the early plateau

of amyloid burden during disease progression⁴² and by dissociations between amyloid burden and the accumulation of soluble amyloid- β oligomers⁴³. The latter may be more bioactive than amyloid- β monomers, fibrils and amyloid plaques, and readily elicit neuronal and network dysfunction^{26,28,30,43–50}. Notably, patients with FAD who carry a mutation in the amyloid- β sequence that increases amyloid- β oligomerization⁴⁹ have prominent cognitive impairments without PiB-positive (PiB⁺) amyloid deposition²², suggesting that AD can be caused by pathogenic assemblies of amyloid- β (probably amyloid- β oligomers) that cannot be detected by PiB imaging.

Notably, amyloid- β oligomers cannot yet be accurately measured in the brain parenchyma of live patients, which has greatly hampered clinical testing of the amyloid- β hypothesis. For example, it is not clear whether amyloid- β oligomer levels were significantly lowered in any of the failed clinical trials of anti-amyloid- β compounds for the treatment of AD. Nor is it clear how much, or by which mechanism, they would have to be lowered to reduce amyloid- β -dependent synaptic and network dysfunction. Anti-amyloid- β treatments that failed to lower amyloid- β oligomer levels in brain also failed to reduce cognitive deficits in hAPP-J20 mice^{51,52}.

In conclusion, testing the amyloid- β hypothesis in humans will probably require therapeutics that effectively lower or counteract the most pathogenic amyloid- β assemblies as well as better methods to confirm their target engagement in the brain. The above evidence strongly supports the hypothesis that amyloid- β causally contributes to cognitive decline in AD. However, the mechanisms remain to be fully elucidated, and alternative interpretations have been offered^{53–56}.

Network dysfunction and preclinical AD

Brain activity can be monitored by functional MRI (fMRI), PET, single-photon emission computed tomography (SPECT), EEG or LFP recordings. The corresponding signals emerge from the complex relationships among electrical activity, cerebrovascular hemodynamics, oxygen consumption and the metabolism of neurons and glia⁵⁷. Attention-demanding tasks, such as sensory processing and memory encoding, require the coordinated regulation of many neuronal ensembles. In healthy people, attention-demanding cognitive tasks increase fMRI signals in specific brain regions (for example, the hippocampus during learning) but also cause a profound large-scale deactivation in brain regions that are collectively referred to as the default mode network (DMN)^{58,59} (FIG. 2a). This network is most active during inwardly oriented mental activity, such as introspection, daydreaming, mind wandering, wakeful rest, imagination and recalling, and it becomes deactivated during outwardly directed mental tasks such as acquisition and encoding of new information.

The DMN includes the precuneus, posterior cingulate cortex, lateral and inferior parietal cortex, and regions of the temporal and medial prefrontal cortex⁵⁹. A widely overlapping, but not identical, DMN that includes the hippocampus has been revealed by resting-state functional connectivity MRI⁶⁰. Task-induced activation and deactivation of networks correlate well with cognitive performance in young healthy people⁶¹. Notably, task-induced deactivation of regions of the DMN (for example, the precuneus) can be a better predictor of

good cognitive performance than the activation of regions that have been more extensively studied with regard to their involvement in cognitive functions (for example, the hippocampus)⁶¹. In fact, task-induced hippocampal activation without adequate DMN deactivation is associated with poor memory formation in healthy people (FIG. 2b), which highlights that proper execution of these complex functions requires well-coordinated regulation of widely distributed networks.

In AD, schizophrenia and other cognitive disorders, deactivation of DMN components is impaired during learning^{1–5}. Because synchrony regulates the functional states of networks (FIG. 1), the abnormal activation or deactivation of brain regions during specific tasks raises the possibility that cognitive dysfunction in these disorders results from synchrony deficits. Although AD is increasingly viewed as a heterogeneous, multicausal syndrome, its fMRI signature seems to be remarkably consistent, particularly during the early stages of the disease. Hippocampal hyperactivation and reduced deactivation of DMN components during memory-encoding tasks have been observed in cognitively normal individuals with cerebral amyloid deposits⁴ (a potential harbinger of AD), cognitively normal carriers of the *APOE* $\epsilon 4$ allele^{6,62–64} (the major genetic risk factor for AD), presymptomatic carriers of FAD-causing mutations^{65,66} and patients with MCI^{67–69}, which often develops into AD. In later stages of AD, the hippocampal formation is hypoactive during memory encoding, whereas the reduced deactivation of the DMN persists^{4,9,69,70}.

Early hippocampal hyperactivation has been interpreted as a mechanism that may compensate for emerging cognitive decline in early AD⁷¹ and *APOE* $\epsilon 4$ carriers⁶⁴. However, accumulating evidence suggests that this hyperactivation is primarily pathogenic and may impair learning and memory^{67,72,73}. In cognitively normal *APOE* $\epsilon 4$ carriers, increased hippocampal activation was associated with reduced grid cell-like representations in the entorhinal cortex during a virtual spatial-memory task⁶⁴, suggesting a potential link between entorhinal grid cell dysfunction and hippocampal hyperactivation.

In patients with MCI, hippocampal hyperactivation occurred during a pattern-separation task, which strongly depends on hippocampal functions; treatment with low doses of the antiepileptic drug levetiracetam for 2 weeks reversed the hyperactivation and improved performance on the task^{67,73} (FIG. 2c). In FAD mice (lines hAPP-J20 and 3xTg-AD (Supplementary information S1 (table))), levetiracetam suppressed network hyperexcitability, improved cognitive functions^{74,75} and reversed excessive neuronal DNA double-strand breaks⁷⁶, which may contribute to neurodegeneration^{77,78}. Thus, this drug may have both symptomatic and disease-modifying therapeutic potential. Collectively, these studies suggest that network hyperactivity is an early and detrimental alteration in the pathogenesis of AD-related cognitive dysfunction and that preventing or reversing this hyperactivity may be of therapeutic benefit.

Remarkably, there is strong evidence that functional brain alterations begin decades before the clinical onset of AD. The Dominantly Inherited Alzheimer Network (DIAN) consortium revealed that pathophysiological changes in the brain in people carrying autosomal dominant FAD mutations begin at least two decades before the predicted onset of clinical symptoms, including changes in cerebrospinal fluid (CSF) biomarkers, cerebral amyloid deposition, and

brain metabolism⁸. Increased brain activity precedes the diagnosis of AD by 30 years in *PSEN1* mutation carriers^{7,9}. Finally, 20–35-year-old asymptomatic carriers of *APOE ε4* show task-related hippocampal hyperactivation and altered DMN activity on fMRI⁶. Thus, FAD-causing mutations and APOE4 modulate brain function several decades before any clinical manifestations of the disease.

Network activity and amyloid deposits

Impaired deactivation of the DMN in older people with or without AD is closely linked to amyloid deposition in brain regions that make up this network^{4,9,70} (FIG. 2d, 3a). Whether the dysfunction of the DMN is a cause or a consequence of amyloid deposition is not known. When studying clinicopathological associations in humans, it is often impossible to conclusively discriminate between detrimental, beneficial and bystander alterations. Transgenic mouse or rat models expressing proteins that are suspected of contributing to the pathogenesis of AD can help to test related hypotheses *in vivo*^{31–33,79,80}.

In humans and mice, soluble amyloid- β levels in the interstitial fluid of different brain regions fluctuate with the brain state. Amyloid- β levels are higher during wakefulness than sleep^{81–83}. In mice, sleep promotes amyloid- β clearance⁸³, whereas sleep deprivation increases amyloid- β levels and amyloid deposition⁸², suggesting that brain states can modulate amyloid deposition. In young FAD mice (line Tg2576) (Supplementary information S1 (table)), interstitial levels of soluble amyloid- β vary markedly in different regions and relate closely to regional differences in metabolic activity⁸⁴. Higher levels of metabolic activity are associated with higher levels of soluble amyloid- β and predict the regional amyloid burden in older mice. Thus, networks with higher metabolic rates may be more prone to amyloid deposition. In addition, experimental increases in neuronal activity by chronic optogenetic activation promote amyloid deposition and trigger epileptiform activity in FAD mice (line APP-A7)⁸⁵ (FIG. 3b; see Supplementary information S1 (table)). Thus, the vulnerability of the DMN to amyloid deposition may be related to its relatively high baseline activity and to deactivation deficits that are associated with early stages of AD.

Synaptic functions and amyloid- β

Amyloid- β production and the amyloid- β_{1-42} /amyloid- β_{1-40} ratio, which influence the formation of pathogenic oligomers as well as amyloid deposition, are regulated by neuronal action potential firing and calcium-dependent conformational changes in PSEN1 at presynaptic terminals⁸⁶. Increasing the rates of regular or burst firing enhances amyloid- β_{1-40} and amyloid- β_{1-42} production in an activity-dependent manner (see also REFS 87,88) (FIG. 3c). Regular firing proportionally increases both amyloid- β_{1-40} and amyloid- β_{1-42} levels, whereas burst firing, which raises presynaptic calcium concentrations, selectively increases the level of amyloid- β_{1-40} , reducing the amyloid- β_{1-42} /amyloid- β_{1-40} ratio (FIG. 3c).

Synaptic plasticity is controlled by neurotransmitter release probability and calcium concentrations at presynaptic terminals. Excitatory synapses with a low initial probability of neurotransmitter release typically display greater facilitation during high-frequency

stimulation (potentiation) and act as high-pass filters. Thus, they tend to be unresponsive to infrequent or disorganized action potentials but potentiate their responses to coordinated firing (FIG. 3d). By contrast, excitatory synapses with a high initial release probability respond preferentially to single-spike firing patterns and act as low-pass filters^{89,90}. They tend to exhibit depression or reduced potentiation during sequential stimulation (FIG. 3d). In one study, the authors concluded that high-pass-filter synapses with higher calcium loading produce a lower amyloid- β_{1-42} /amyloid- β_{1-40} ratio, whereas low-pass-filter synapses with lower calcium loading produce a higher amyloid- β_{1-42} /amyloid- β_{1-40} ratio⁸⁶.

Amyloid- β and other APP metabolites seem to participate in pre- and postsynaptic homeostatic mechanisms that regulate synaptic activity^{86,88,91-95}. Amyloid- β may enhance presynaptic facilitation at low concentrations⁹³ and promote postsynaptic depression at high concentrations⁸⁸. Application of synthetic or recombinant amyloid- β_{1-42} oligomers or overexpression of FAD-mutant human APP reduces paired-pulse facilitation — a measure of increased release probability — and impairs long-term potentiation (LTP) at some, but not other, synapses in the mouse hippocampus^{91,93,96}. These alterations may shift the filtering properties of the affected synapses from high-pass to low-pass. This shift could enhance the transfer of single action potentials, increase synaptic depression and elevate the amyloid- β_{1-42} /amyloid- β_{1-40} ratio (FIG. 3d), promoting both hyperactivity in vulnerable networks and amyloid deposition. Reducing presynaptic release might counteract these processes. Indeed, levetiracetam, which has beneficial effects on network activity in FAD mice^{74,75} and humans with MCI^{67,73}, reduces presynaptic neurotransmitter release in a use-dependent manner^{90,97,98}. Thus, it may counteract amyloid- β -induced alterations in release probability.

In general, it is difficult to extrapolate from the effect that factors exert on specific synapses or neurons to the overall effect they have on microcircuits and complex networks. Indeed, this has been true for APP and amyloid- β . Although early studies suggested that increased neuronal amyloid- β production may be part of a homeostatic feedback loop that primarily reduces neuronal activity⁸⁸, we now know that amyloid- β accumulation can also cause neuronal hyperexcitability *in vitro* and *in vivo*^{32,96,99-111}. Acute amyloid- β application initially and transiently (10–20 minutes) increases the levels of surface AMPA-type glutamate receptors and GluN2B-containing NMDA-type glutamate receptors (NMDARs)^{100,103} and the frequency of spontaneous excitatory postsynaptic currents in primary neuronal cultures^{100,103}. In brain slices, amyloid- β application also acutely increases the rate of action potential firing by hippocampal pyramidal cells⁹⁹. FAD mice (lines hAPP-J20, hAPP-J9, APP23xPS45 and APP/PSEN1dE9) (Supplementary information S1 (table)) contain both hyperactive and hypoactive neurons — reflected in abnormally high or low rates of action potential-dependent calcium transients or levels of transcripts for the immediate early genes *Arc* and *Fos* — both before^{96,112,113} and after^{109,114,115} amyloid deposition becomes detectable.

Neurons exposed to pathologically elevated levels of amyloid- β *in vitro* or *in vivo* show signs of atrophy, including lower dendritic spine density and shorter dendrites^{116,117}, alterations that can increase intrinsic cellular excitability¹¹⁸ and may explain the hyperactivity of some excitatory neurons. These morphological and functional effects of amyloid- β may be mediated, at least in part, by neuronal depletion of the DNA repair factor

BRCA1 and the resulting accumulation of activity-induced DNA double-strand breaks⁷⁷. Compensatory increases in inhibitory interneuron activity may account for the hypoactivity of other excitatory neurons^{101,109}. Notably, hippocampal neuronal hyperactivity preceded amyloid plaque deposition and neuronal hypoactivity in APP23xPS45 mice and could be induced by local application of amyloid- β in wild-type mice¹⁰¹, suggesting that it is a very early step in the pathogenic cascade that is triggered by amyloid- β accumulation.

Hypersynchrony in AD and animal models

Besides network activation and deactivation deficits detectable by fMRI (FIG. 2), AD also results in network hypersynchrony (Table 1: see Supplementary information S3 (table)). Unlike normal fluctuations in network synchrony that are associated with physiological changes in behaviour and brain states (FIG. 1), network hypersynchrony is a pathological phenomenon in which aberrant synchronization of neuronal networks results in epileptiform discharges and seizures.

Neuronal expression or overexpression of proteins that accumulate in AD brains or are genetically linked to the disease, such as APP, amyloid- β , tau, APOE4 or α -synuclein, causes network hypersynchrony in transgenic mice^{96,119–121}. Spontaneous epileptiform activity has been documented in many FAD models, including hAPP-J20 (REF. 96), APP/PSEN1dE9 (REFS 102,103), Tg2576 (REF. 104), 5xFAD¹²², 3xTg-AD⁷⁵, APP/TTA-EC¹⁰⁵, APP/TTA-CaMKII α ¹⁰⁶, APP23 (REF. 107) and hAPPJ9/FYN¹⁰⁸ transgenic mice (FIG. 3e; see Supplementary information S1 (table)). Behavioural seizures and lower thresholds for chemically induced seizures have also been described in some of these models^{96,103,108,111,123,124}. Interestingly, network hypersynchrony in mouse models of AD and of epilepsy depends on the presence of endogenous, soluble wild-type tau, which seems to enable or promote aberrant synchronization^{32,35,110,111,125,126}.

In humans, epidemiological studies have consistently shown that AD is a risk factor for epileptic seizures^{127,128}. However, the reported incidence of seizures in patients with AD that were detected by observation has ranged from 0.5% to 64%^{129,130}. A recent review of 17 clinical studies that assessed epileptic activity in AD reported an average incidence of 15.1% (median: 9.0%)¹³¹. The incidence of seizures is roughly 7–8-fold higher in individuals with AD than in people without dementia^{132,133}. In a recent nationwide study of data from over 140 million hospitalizations, patients with AD were fourfold more likely to be hospitalized for a seizure than for non-seizure-related condition¹³⁴. Moreover, people with early-onset AD (50–65 years old) were more likely to be hospitalized for seizures or epilepsy than those with a later onset of AD (>81 years old)¹³⁴. These new findings are consistent with earlier reports indicating that early-onset AD (<65 years old) is a major risk factor for seizures^{129,133,135}. Conversely, patients with amnesic MCI or AD who had seizures or subclinical epileptic activity typically had an earlier onset and faster progression of cognitive decline than those without detectable epilepsy^{129,136,258}.

Although seizures in AD are widely thought to result from end-stage neurodegeneration, several reports indicate that they can occur early in the disease course and before a neurodegenerative disease diagnosis is made^{136,137}. Indeed, in patients who had MCI or

early AD with epilepsy, seizure onset clustered near the onset of cognitive decline¹³⁶ or preceded the MCI diagnosis by 4 to 7 years¹³⁸. At the time MCI was diagnosed, the latter group of patients had mild hippocampal atrophy¹³⁸. Seizures in MCI or early AD with epilepsy responded best to the antiepileptic drugs levetiracetam or lamotrigine and least well to phenytoin¹³⁶. The lower relative risk of seizures in patients with late-onset AD may reflect the fact that strokes and other ageing-related comorbidities increase seizure risk in people without dementia. It is also possible that old brains are less susceptible to amyloid- β -induced network hypersynchrony than younger brains and that amyloid- β has a less important role in the pathogenesis of late-onset AD, which may be more multifactorial than early-onset AD.

Epileptic activity is even more prominent in pedigrees of early-onset autosomal dominant FAD^{91,139,140}. Seizures have been observed in patients with FAD who carry any of 66 different *PSEN1* mutations^{141,142} (Supplementary information S3 (table)), in 31% of patients with AD carrying *PSEN2* mutations¹⁴³, and in 58% of patients with *APP* duplications^{144,145}. More recently, a multicenter study including 132 patients with FAD showed that 48% of patients harboring an FAD mutation had seizures, including 43% of *PSEN1* mutation carriers ($n = 94$), 43% of *PSEN2* mutation carriers ($n = 7$), 47% of *APP* mutation carriers ($n = 15$), and 81% of *APP* duplication carriers ($n = 16$)¹⁴⁶. Seizure frequency in patients with FAD seems to be even higher than that observed in most FAD mice (Supplementary information S1 (table)).

In addition, 84% of patients with Down's syndrome who progress to dementia also develop seizures¹⁴⁷. Interestingly, very early onset of AD (<40 years old) further increases the proportion of FAD pedigrees with epileptic phenotypes (>80%)¹⁴⁸. Some FAD mutations (for example, *PSEN1* L166P) and *APP* duplications even cause epileptic activity in childhood or adolescence, preceding cognitive decline by many years^{145,149}. Thus, network hypersynchrony can be a very early and prominent clinical feature of FAD. Indeed, many FAD-*PSEN1* pedigrees qualify as epileptic syndromes or disorders^{140,141}. This intriguing association between AD and epilepsy probably holds important clues about the mechanisms by which amyloid- β , tau, and APOE4 promote the development of AD and about potential entry points for therapeutic intervention.

The power of gamma oscillations

Whereas abnormal neuronal synchrony can result in epileptic activity, normal neuronal synchrony underlies the generation of oscillatory rhythmic activities, or brain rhythms, that promote cognitive functions. Neuronal ensembles coordinate their firing within a range of oscillatory frequencies (0–300 Hz). Within each frequency band, neuronal ensembles can increase or decrease their firing rate and synchrony and thus generate higher or lower oscillatory amplitudes (FIG. 4a), although it should be noted that only a small fraction of the cells contributes to each oscillatory cycle. The amplitude, phase and frequency in each oscillatory band modulate each other and neuronal firing rates in precise ways¹⁵⁰. For example, the firing of hippocampal pyramidal cells shows phase-coupling with low-frequency oscillations, such as theta (4–8 Hz). Thus, during periods of high theta, the action potential firing rate of pyramidal cells is increased and its timing is synchronized with this

rhythm¹⁵¹. By contrast, the firing of several types of interneurons shows phase-coupling with high-frequency oscillations, such as gamma (30–150 Hz)^{150–152}. Thus, during periods of high gamma power, action potential firing of interneurons is more prominent and synchronized with the phase of gamma¹⁵¹ (FIG. 4b,c).

The amplitude (or power) of gamma oscillatory activity increases during sensory processing or memory encoding, and such increases in gamma power predict successful memory formation in humans and mice^{153–158}. Encoding during sensory, motor or memory tasks is associated not only with increased power of high-frequency oscillations (gamma) but also with reduced power of lower-frequency oscillations (alpha, beta and theta) in task-related brain regions^{57,154,159,160} (FIG. 4d). However, the relationship between gamma power and attention can differ across cortical areas and tasks¹⁶¹. Increases in gamma power tend to be localized to networks that are directly involved in the task (task-related networks), whereas reductions in the power of lower-frequency oscillations typically involve multiple cortical regions^{162,163}. During sensory encoding, individuals with schizophrenia have reduced increases in gamma power in task-related networks and reduced decreases in the power of lower-frequency oscillations across multiple components of the DMN¹⁶⁴. It is tempting to speculate that these large-scale deficits of oscillatory activity may be related to the activation and deactivation deficits of fMRI signals in patients with schizophrenia or AD⁵ (FIG. 2). Patients with AD also have reductions in gamma power¹⁶⁵; however, the relationship between task-related deficits in fMRI signals, oscillatory frequencies and the observed prominent memory-encoding problems does not seem to have been directly studied in this disease.

In healthy individuals, the strength of fMRI blood-oxygen-level-dependent (BOLD) signals correlates positively with the power of gamma oscillations^{166,167} and negatively with the power of alpha oscillations^{168–170}. By contrast, decreases in DMN fMRI BOLD signals are associated with reductions in gamma power¹⁷¹. During visual stimulation in monkeys, an increase in oxygen level, which was associated with a rise in fMRI signals, coincided with increases in LFP gamma power in task-related networks (for example, visual area V3), whereas a decrease in oxygen level coincided with prominent decreases in LFP power in all oscillatory frequencies in DMN regions (for example, the posterior cingulate cortex)⁵⁷ (FIG. 4d). These findings suggest that fMRI signals may directly reflect changes in oscillatory activity. It should be informative to further explore the relationship between fMRI signals and oscillatory alterations in AD and other cognitive disorders.

Longitudinal video-EEG recordings revealed abnormal behaviour-dependent fluctuations in the power of gamma oscillations in hAPP-J20 mice (FIG. 5a). Notably, spontaneous epileptiform discharges emerged primarily during resting periods in these animals, when the intensity of gamma oscillations was low¹⁷². In two mouse models of absence epilepsy, pharmacologically induced increases or decreases of gamma power were associated with decreases or increases in epileptic activity, respectively¹⁷³. Thus, network hypersynchrony and reduced gamma power may be mechanistically linked across different conditions, and behaviour-induced increases in gamma oscillatory power may counteract AD-related epileptiform activity by activating task-related brain regions and reducing network synchrony.

These ideas are supported by findings in humans with epilepsy¹⁵⁴ (FIG. 5b). During EEG recordings, individuals were shown a set of images (encoding trials), and their memory of the images was probed 24 hours later (recognition trials). Encoding trials were deemed ‘correct’ or ‘incorrect’, depending on whether or not the old images were recognized during the recognition trial. Remarkably, gamma power increased and epileptiform activity reduced only during correct memory encoding.

Overall, these LFP, EEG and fMRI findings suggest that effective memory formation requires, or induces, a pattern of brain activity that increases the power of gamma oscillations and reduces the power of low-frequency oscillations, and that this ‘brain state’ reduces aberrant network hypersynchrony in humans and mice (FIG. 5c). Social interactions and mental activity may benefit some patients with AD¹⁷⁴, perhaps, at least in part, by promoting this state. Pharmacological interventions that more persistently improve brain rhythms could be of greater therapeutic benefit.

Interneuron impairments in AD

Network synchrony and oscillatory brain rhythms are promoted and controlled by the activity of inhibitory GABAergic interneurons¹⁷⁵. These cells are highly diverse, and different subtypes form electrically coupled functional networks¹⁷⁶. Their combined inhibitory synaptic input onto excitatory principal neurons and other interneurons generates precise oscillatory rhythms that in turn coordinate the timing of pyramidal cell firing (see above). Some types of inhibitory interneurons, such as somatostatin-positive (SOM⁺) or neuropeptide Y-positive (NPY⁺) cells, tend to fire tonically and independently of brain and behavioural states^{177,178}. Others, such as parvalbumin-positive (PV⁺) or vasoactive intestinal polypeptide-positive (VIP⁺) cells, fire predominantly during brain states that promote encoding^{177,178}. Interneurons and the oscillatory network activities that they regulate are altered in AD, epilepsy, schizophrenia and autism^{91,165,172,179–182}. Indeed, interneuron impairment is emerging as a potential common mechanism of brain dysrhythmias and cognitive dysfunction in many neurological and psychiatric disorders^{172,179,180,183,184}. What is more, recent findings support the hypothesis that modulating interneuron function may improve brain rhythms and cognitive functions in AD and related disorders^{172,182,185,186}.

Proper functioning of inhibitory interneurons is required for the generation of high-frequency gamma oscillatory activity, coordination among different oscillatory frequencies (cross-frequency interactions) and regulation of neuronal firing by brain rhythms (frequency-action potential interactions). Several lines of evidence suggest that fast-spiking PV⁺ cells are impaired in disorders associated with hypersynchronous network activity such as schizophrenia and epilepsy¹⁸⁷, and that other types of interneurons — for example, those containing cholecystokinin or NPY — are impaired in mood disorders and anxiety^{187–189}.

PV⁺ cells constitute ~40% of inhibitory interneurons and are the major source of perisomatic inhibition onto excitatory pyramidal cells. Because PV⁺ cells are electrically coupled by dendritic gap junctions and form an electrically coordinated interneuron network, they are particularly well suited to synchronously modulate the activity of many pyramidal cells¹⁷⁶. In optogenetic stimulation studies in wild-type mice, increasing the firing rate of PV⁺ cells

increased the power of gamma oscillations but not of other oscillations, whereas increasing the firing rate of pyramidal cells specifically increased the power of low-frequency oscillations^{179,190} (FIG. 4c), indicating a causal link between gamma and PV⁺ cell function. Increasing gamma power improved the signal-to-noise ratio of pyramidal cell firing, an effect that is likely to enhance neuronal processing^{179,190}.

In FAD mice (lines hAPP-J20, Tg2576, APP23, *APOE4*-KI, and APP/PSEN1dE9) (Supplementary information S1 (table)), gamma oscillatory activity is altered^{107,172,191–193}, suggesting that these animals have deficits in interneuron function. hAPP-J20 mice have brief peaks of increased gamma power and long periods of decreased gamma power^{172,192}. Similar abnormal fluctuations in gamma power occur in Tg2576 mice¹⁹¹. Increased gamma power is associated with behaviour-dependent brain activation, increased firing rate of PV⁺ cells^{177,194} and the suppression of epileptiform discharges in hAPP-J20 mice¹⁷² and humans with epilepsy¹⁵⁴. By contrast, gamma power is decreased during resting activity and is associated with reduced firing rate of PV⁺ cells^{177,194} and increased epileptiform discharges in hAPP-J20 mice¹⁷². These behaviour-related modulations of gamma oscillations probably differ mechanistically from aberrant increases in gamma power during seizures or hyper-synchronized network activity and from the overall increases in oscillatory power across multiple frequency bands in APP23 and APP/PSEN1dE9 mice^{107,193}. In APP23xPS45 mice, hyperactivity of cortical neurons was associated with decreased GABAergic inhibition rather than increased glutamatergic transmission, suggesting impaired inhibitory function¹⁰⁹. More recently, reduced inhibitory function has been linked to amyloid- β -induced deficits in slow-wave propagation^{195,196}. Aberrant increases in gamma power in the auditory cortex during evoked auditory stimulation in patients with AD have also been related to decreased inhibition^{197,198}. Overall, however, humans with AD typically show decreases in the power of higher-frequency oscillations and increases in the power of lower-frequency oscillations¹⁹⁹.

APOE4-KI mice (Supplementary information S1 (table)), in which the *ApoE* gene is replaced by knocking in the human *e4* allele, also have prominent GABAergic dysfunction and show reduced hippocampal power of slow gamma oscillations during sharp-wave ripple activity²⁰⁰. Amyloid- β application^{99,201} and APP overexpression^{202–204} markedly suppress the power of kainate-induced gamma oscillations in neuronal cultures. The mechanisms of gamma degradation by amyloid- β in brain slices are unclear but may involve changes in inhibition–excitation balance and cellular excitability⁹⁹. Amyloid- β treatment increased excitatory and decreased inhibitory postsynaptic currents in pyramidal cells and increased the firing rate of these excitatory neurons⁹⁹. Interestingly, although the mean phase of action potential firing did not change, the temporal firing window of excitatory neurons became wider, suggesting desynchronization of action potential firing⁹⁹. Perhaps, spike-theta phase dysregulation²⁰⁴ also contributes to the reduced gamma power and the epileptiform activity that are observed in FAD mice. Hippocampal injections of amyloid- β reduced firing rates in rhythmically bursting interneurons (probably PV⁺ cells) but not in tonically firing interneurons (probably cholinergic interneurons) in the septum²⁰³.

Across brain regions and behavioural tasks, effective encoding also seems to depend on cross-frequency interactions, particularly the coupling between the phase of theta

oscillations and the power of gamma oscillations (phase–amplitude coupling)^{205–209}, which is regulated by the synaptic activity of PV⁺ cells^{210,211}. Recordings from acute brain slices of young FAD mice and *in vivo* recordings from adult FAD mice (lines TgCRND8 and APP/PSEN1dE9, respectively) (Supplementary information S1 (table)) revealed deficits in theta-gamma coupling^{193,212}, which were associated with cognitive deficits²¹² and could result from the kind of PV⁺ cell dysfunction that we identified in hAPP-J20 mice. Indeed, by impairing PV⁺ cell functions, APP and amyloid- β could disrupt multiple aspects of neuronal coordination.

Single-unit recordings of inhibitory cells in behaving rodents have revealed that behavioural and brain states affect interneuron subtypes differentially^{177,178,189,213}. For example, PV⁺ and VIP⁺ cells drastically increase their action potential firing rate during specific behavioural states (for example, locomotor exploration), whereas NPY⁺ and SOM⁺ cells show persistent or tonic firing across multiple behavioural states (for example, locomotion, sleep and quiet wakefulness). These findings imply that certain behavioural states may be able to overcome, at least partly and for short periods, PV⁺ cell impairments that cause low gamma power, hypersynchronous network activity and cognitive deficits in FAD mice and humans with AD (FIG. 5c). Optogenetically suppressing the activity of interneurons that express the homeobox protein DLX1 in the dentate gyrus impaired learning and acutely suppressed memory retrieval in mice, further highlighting the crucial role of interneurons in cognitive tasks²¹⁴. It is tempting to speculate that fluctuations in interneuron activity may help to explain the fluctuations in cognitive performance of patients with AD that are frequently reported by care takers but have been difficult to document by clinical measurements.

The mechanisms by which interneuron deficits contribute to network and cognitive dysfunction in AD and related conditions remain to be fully elucidated, although much progress has recently been made on this front. Decreased synaptic inhibition and excitation–inhibition imbalances might explain deactivation deficits in the DMN of individuals with MCI or AD^{4,67,72}, but these hypotheses have not yet been tested experimentally. Impaired inhibition is another potential mechanism for plaque-related neuronal hyperactivity in APP23xPS45 mice¹⁰⁹. In hAPP-J20 mice, we found that PV⁺ cell impairments are caused by the depletion of the voltage-gated sodium channel subunit Nav1.1 and crucially contribute to network and cognitive dysfunctions¹⁷² (FIG. 6a). Voltage-gated sodium channels control intrinsic cellular excitability by modulating action potential firing in specific neuronal subtypes. Nav1.1 is expressed predominantly in interneurons, including PV⁺ cells, in both mice¹⁷² and humans²¹⁵, and its levels in the parietal cortex are reduced in hAPP-J20 mice and patients with AD¹⁷². Hypofunction of Nav1.1 has been also described in Tg2576 mice²¹⁶ and in transgenic mice overexpressing BACE1 (REF. 217). As indicated earlier, the activity of BACE1, which is expressed at high levels in the brains of patients with AD²¹⁸, is required for the release of amyloid- β from APP. Because this enzyme also cleaves the β -subunit of voltage-gated sodium channels, increased BACE1 levels lead to aberrant β -subunit cleavage and reduced transport of Nav1.1 to the membrane, resulting in Nav1.1 hypofunction^{216,217}. Neuroblastoma cells expressing the FAD-linked *PSEN1* E280A mutation also had reduced levels of Nav1.1 mRNA and protein levels²¹⁹.

APOE4-KI mice also develop spontaneous epileptic activity and seizures¹²⁰. In humans without dementia, hyperventilation induces network hypersynchronization (for example, sharp waves) more frequently in *APOE* e4 carriers than non-carriers²²⁰. Although network hypersynchrony has not been directly linked to interneuron dysfunction in *APOE4*-KI mice, these mice lose around 30% of SOM⁺ cells in the hilus of the dentate gyrus, and this reduction is associated with learning and memory deficits²²¹. Interestingly, chronic treatment with pentobarbital reversed the latter deficits but not the loss of hilar interneurons²²¹, suggesting that this positive allosteric modulator of GABA type A receptors can compensate for the partial loss of inhibitory input (FIG. 6b).

Synaptic depression and hypersynchrony

Synaptic loss is a pathological hallmark of AD and correlates well with cognitive decline^{222,223}. In experimental models, high levels of amyloid- β cause synaptic loss, reduce glutamatergic synaptic transmission and LTP, and increase long-term depression^{25,88,94,224–228}. Intriguingly, several proposed mechanisms of amyloid- β -induced synaptic depression might also contribute to network hypersynchrony²²⁹. For example, amyloid- β blocks neuronal glutamate uptake at synapses, which could result in glutamate spillover around the synaptic cleft²⁵. The rise in glutamate may desensitize synaptic NMDARs and aberrantly activate extra- or perisynaptic GluN2B-containing NMDARs and metabotropic glutamate receptors (mGluRs), and both GluN2B-containing NMDARs and mGluRs can promote long-term synaptic depression and spine retraction^{25,228,230}. Amyloid- β -induced NMDAR- and mGluR-dependent long-term depression can be prevented by lowering extracellular glutamate²⁵ and can be mimicked by application of the glutamate reuptake inhibitor thero- β -benzyloxyaspartate (TBOA), which can also trigger epileptiform discharges in wild-type brain slices²³¹. Thus, amyloid- β -induced network hypersynchrony may be directly linked to synaptic depression.

Future therapies

The effective therapy of AD will probably require combining treatments targeting the root causes of the disease with therapeutic strategies that can block or counteract the pathogenic processes that these causes trigger³³. From the evidence reviewed above, we think that brain dysrhythmias caused by aberrant interneuron activity and other mechanisms probably contribute to cognitive deficits and behavioural alterations in AD and related conditions. We further hypothesize that enhancing the function of interneurons in these conditions, particularly of PV⁺ and SOM⁺ cells, will improve network activities and cognitive performance.

In hAPP-J20 mice, restoring Nav1.1 levels enhanced PV⁺ cell-dependent gamma oscillatory power, reduced network hypersynchrony and improved cognitive performance (FIG. 6a), pinpointing Nav1.1- and PV⁺ cell-dependent oscillatory rhythms as potential therapeutic targets. Although it is never certain that therapeutic findings obtained in experimental models (or in early clinical trials in humans, for that matter) will hold up in large heterogeneous human populations, Nav1.1 depletion and various other molecular alterations identified in hAPP-J20 mice have been found in people with AD (Supplementary

information S2 (table)). These include depletions of calbindin in the dentate gyrus and of reelin in the entorhinal cortex, increased neuronal production of collagen VI, higher hippocampal levels of activated group IVA phospholipase A2 and increased expression of the adenosine A_{2A} receptor by hippocampal astrocytes^{100,172,232–236}. In addition, the anti-epileptic drug levetiracetam, which reduces synaptic, network and cognitive deficits in hAPP-J20 mice⁷⁴, also exerts beneficial effects in patients with amnesic MCI^{67,73}, which is widely viewed as an early stage of AD^{237–239}.

Manipulating interneuron function to improve brain rhythms is a promising therapeutic strategy for disorders involving interneuron and network dysfunctions, including AD, epilepsy, schizophrenia and autism. But how might these dysfunctions be approached therapeutically? Interneurons have highly specialized functions, electrophysiological properties and molecular profiles. Targeting molecules such as ion channels or receptors that are predominantly expressed in specific types of interneurons might make it possible to improve and harness the function of these cells without impairing other cell types. Indeed, Nav1.1 enhancers that preferentially activate Nav1.1-containing interneurons have been proposed for the treatment of epilepsy, schizophrenia and AD²⁴⁰. Because the firing rate of PV⁺ cells and gamma power strongly depend on behavioural activity^{154,177}, behavioural or sensorial interventions that promote brain states associated with increased power ratios in high- to low-frequency oscillations might also be of benefit.

Finally, embryonic interneuron precursors transplanted into adult brains can migrate and integrate into appropriate circuits and mature into fully functional interneurons¹⁸². Although cell therapy for cognitive disorders may seem to be rather daring and must be approached with appropriate caution, it has already been explored in a small clinical trial in patients with AD²⁴¹. In AD-related mouse models with prominent loss of hilar interneurons, including pilocarpine-treated wild-type mice¹⁸⁵ and untreated *APOE4*-KI mice¹⁸⁶, hippocampal transplants of wild-type interneuron precursors improved behavioural functions. However, it is not known how well such transplants would survive and function in the proteopathic, toxic microenvironment in AD brains^{172,203}. Genetic modifications may be required to render interneuron precursors resistant to such conditions. Potentially beneficial modifications include overexpression of proteins that protect against the disease, such as Nav1.1 (REF. 172) in PV⁺ cells, or reduction of proteins that promote or enable AD-related pathogenesis, such as tau¹¹⁰ and APOE4 fragments²⁴². These therapeutic opportunities also call for the development of reprogramming strategies to convert skin or blood cells into precursors of specific interneuron subtypes.

Conclusions

Network activities that support cognition are altered decades before the expected onset of clinical signs and symptoms in AD, and the affected networks predict future pathology and brain atrophy. Because neuronal synchrony regulates the functional state of brain circuits and networks, deficits in synchrony, including network hypersynchrony and altered oscillatory rhythmic activity, could contribute to AD-related cognitive dysfunction. Although the precise causes of these synchrony deficits remain to be defined, diverse lines of evidence suggest that interneuron dysfunction and network imbalance may be crucially

involved. Therapeutic strategies that improve the function of interneurons and counteract such abnormalities might improve cognitive functions in AD and related disorders. By preventing excitotoxic overstimulation of neurons and maladaptive compensatory processes, they may also help to prevent or stall disease progression.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors thank P. Sanchez, Y. Huang, A. Kreitzer and members of their laboratories for helpful discussions; S. Ordway for editorial review; and C. Dickerson and A. Cheung for administrative assistance. This work was supported by US National Institutes of Health grants AG011385 (L.M.), AG053981 (L.M.), and AG047313 (J.J.P.); an Alzheimer's Association grant IIRG-13-284779 (J.J.P.); an S.D. Bechtel, Jr. Young Investigator Award (J.J.P.); and a gift from the Dolby Family.

L. M. has received research funding from Bristol-Myers Squibb and consulting fees from AbbVie; he serves on the scientific advisory boards of Acumen Pharmaceuticals, Alkahest, Dolby Family Ventures, E-scape Bio, and Neuropore Therapies. J. P. declares no competing interests.

References

1. Seidman LJ et al. Medial temporal lobe default mode functioning and hippocampal structure as vulnerability indicators for schizophrenia: A MRI study of non-psychotic adolescent first-degree relatives. *Schizophr. Res* 159, 426–434 (2014). [PubMed: 25308834]
2. Fahoum F, Zelmann R, Tyvaert L, Dubeau F. & Gotman J. Epileptic discharges affect the default mode network--fMRI and intracerebral EEG evidence. *PLoS One* 8, e68038 (2013).
3. Oser N. et al. Default mode network alterations during language task performance in children with benign epilepsy with centrotemporal spikes (BECTS). *Epilepsy Behav.* 33, 12–17 (2014). [PubMed: 24583653]
4. Sperling RA et al. Amyloid deposition is associated with impaired default network function in older persons without dementia. *Neuron* 63, 178–188 (2009). [PubMed: 19640477] This study shows that PiB+ amyloid deposits are associated with deactivation deficits in a default network component in non-demented individuals and patients with MCI, linking aberrant neuronal activity and pathological changes.
5. Anticevic A. et al. The role of default network deactivation in cognition and disease. *Trends Cogn. Sci* 16, 584–592 (2012). [PubMed: 23142417]
6. Filippini N. et al. Distinct patterns of brain activity in young carriers of the APOE-ε4 allele. *Proc. Natl. Acad. Sci. USA* 106, 7209–7214 (2009). [PubMed: 19357304]
7. Mondadori CR et al. Enhanced brain activity may precede the diagnosis of Alzheimer's disease by 30 years. *Brain* 129, 2908–2922 (2006). [PubMed: 17012294]
8. Bateman RJ et al. Clinical and biomarker changes in dominantly inherited Alzheimer's disease. *N. Engl. J. Med* 367, 795–804 (2012). [PubMed: 22784036] This study demonstrates changes in CSF biomarkers, cerebral amyloid deposition, and brain metabolism in carriers of autosomal dominant FAD mutations at least two decades before the estimated onset of clinical symptoms.
9. Reiman EM et al. Brain imaging and fluid biomarker analysis in young adults at genetic risk for autosomal dominant Alzheimer's disease in the presenilin 1 E280A kindred: A case-control study. *Lancet. Neurol* 11, 1048–1056 (2012). [PubMed: 23137948] This study shows hippocampal hyperactivity and deactivation deficits in the DMN in 18–26-year-old asymptomatic PSEN1 mutation carriers.
10. Jasper HH Cortical excitatory state and variability in human brain rhythms. *Science* 83, 259–260 (1936). [PubMed: 17757098]

11. Destexhe A, Contreras D. & Steriade M. Spatiotemporal analysis of local field potentials and unit discharges in cat cerebral cortex during natural wake and sleep states. *J. Neurosci* 19, 4595–4608 (1999). [PubMed: 10341257]
12. Poulet JF & Petersen CC Internal brain state regulates membrane potential synchrony in barrel cortex of behaving mice. *Nature* 454, 881–885 (2008). [PubMed: 18633351]
13. Poulet JF, Fernandez LM, Crochet S. & Petersen CC Thalamic control of cortical states. *Nat. Neurosci* 15, 370–372 (2012). [PubMed: 22267163]
14. Karran E, Mercken M. & De Strooper B. The amyloid cascade hypothesis for Alzheimer’s disease: An appraisal for the development of therapeutics. *Nat. Rev. Drug Discov* 10, 698–712 (2011). [PubMed: 21852788]
15. Benilova I, Karran E. & De Strooper B. The toxic A β oligomer and Alzheimer’s disease: An emperor in need of clothes. *Nat. Neurosci* 15, 349–357 (2012). [PubMed: 22286176]
16. Karch CM, Cruchaga C. & Goate AM Alzheimer’s disease genetics: from the bench to the clinic. *Neuron* 83, 11–26 (2014). [PubMed: 24991952]
17. Gomez-Isla T. et al. The impact of different presenilin 1 and presenilin 2 mutations on amyloid deposition, neurofibrillary changes and neuronal loss in the familial Alzheimer’s disease brain: Evidence for other phenotype-modifying factors. *Brain*. 122, 1709–1719 (1999). [PubMed: 10468510]
18. Jankowsky JL et al. Mutant presenilins specifically elevate the levels of the 42 residue beta-amyloid peptide in vivo: evidence for augmentation of a 42-specific γ _secretase. *Hum. Mol. Genet* 13, 159–170 (2004). [PubMed: 14645205]
19. Steiner H. & Haass C. Intramembrane proteolysis by presenilins. *Nat. Rev. Mol. Cell Biol* 1, 217–224 (2000). [PubMed: 11252897]
20. Tsubuki S, Takaki Y. & Saido TC Dutch, Flemish, Italian, and Arctic mutations of APP and resistance of A β to physiologically relevant proteolytic degradation. *Lancet* 361, 1957–1958 (2003). [PubMed: 12801742]
21. Nilsberth C. et al. The ‘Arctic’ APP mutation (E693G) causes Alzheimer’s disease by enhanced A β protofibril formation. *Nat. Neurosci* 4, 887–893 (2001). [PubMed: 11528419]
22. Shimada H. et al. Clinical course of patients with familial early-onset Alzheimer’s disease potentially lacking senile plaques bearing the E693 mutation in amyloid precursor protein. *Dement. Geriatr. Cogn. Disord* 32, 45–54 (2011). [PubMed: 21846988]
23. Jonsson T. et al. A mutation in APP protects against Alzheimer’s disease and age-related cognitive decline. *Nature* 488, 96–99 (2012). [PubMed: 22801501]
24. Maloney JA et al. Molecular mechanisms of Alzheimer disease protection by the A673T allele of amyloid precursor protein. *J. Biol. Chem* 289 30990–31000, (2014). [PubMed: 25253696]
25. Li S. et al. Soluble oligomers of amyloid β -protein facilitate hippocampal long-term depression by disrupting neuronal glutamate uptake. *Neuron* 62, 788–801 (2009). [PubMed: 19555648] This study shows that amyloid- β blocks neuronal glutamate uptake at synapses, which results in glutamate spillover, aberrant activation of extra- or perisynaptic GluN2B-containing NMDARs, and enhanced long-term depression.
26. Shankar GM et al. Amyloid- β protein dimers isolated directly from Alzheimer’s brains impair synaptic plasticity and memory. *Nat. Med* 14, 837–842 (2008). [PubMed: 18568035]
27. Li S. et al. Soluble A β oligomers inhibit long-term potentiation through a mechanism involving excessive activation of extrasynaptic NR2B-containing NMDA receptors. *J. Neurosci* 31, 6627–6638 (2011). [PubMed: 21543591]
28. Cleary JP et al. Natural oligomers of the amyloid- β protein specifically disrupt cognitive function. *Nat. Neurosci* 8, 79–84 (2005). [PubMed: 15608634]
29. Musiek ES & Holtzman DM Three dimensions of the amyloid hypothesis: time, space and ‘wingmen’. *Nat. Neurosci* 18, 800–806 (2015). [PubMed: 26007213]
30. Liu P. et al. Quaternary structure defines a large class of amyloid- β oligomers neutralized by sequestration. *Cell Rep.* 11, 1760–1771 (2015). [PubMed: 26051935]
31. Ashe KH & Zahs KR Probing the biology of Alzheimer’s disease in mice. *Neuron* 66, 631–645 (2010). [PubMed: 20547123]

32. Morris M, Maeda S, Vossel K. & Mucke L. The many faces of tau. *Neuron* 70, 410–426 (2011). [PubMed: 21555069]
33. Huang Y. & Mucke L. Alzheimer mechanisms and therapeutic strategies. *Cell* 148, 1204–1222 (2012). [PubMed: 22424230]
34. Heppner FL, Ransohoff RM & Becher B. Immune attack: the role of inflammation in Alzheimer disease. *Nat. Rev. Neurosci* 16, 358–372 (2015). [PubMed: 25991443]
35. Maeda S. et al. Expression of A152T human tau causes age-dependent neuronal dysfunction and loss in transgenic mice. *EMBO Rep.* 17, 530–551 (2016). [PubMed: 26931567]
36. SantaCruz K. et al. Tau suppression in a neurodegenerative mouse model improves memory function. *Science* 309, 476–481 (2005). [PubMed: 16020737]
37. Mandelkow EM & Mandelkow E. Biochemistry and cell biology of tau protein in neurofibrillary degeneration. *Cold Spring Harb. Perspect. Med* 2, a006247 (2012).
38. Min SW et al. Critical role of acetylation in tau-mediated neurodegeneration and cognitive deficits. *Nat. Med* 21, 1154–1162 (2015). [PubMed: 26390242]
39. Mahley RW & Huang Y. Apolipoprotein E sets the stage: response to injury triggers neuropathology. *Neuron* 76, 871–885 (2012). [PubMed: 23217737]
40. Pike KE et al. β -Amyloid imaging and memory in non-demented individuals: evidence for preclinical Alzheimer's disease. *Brain* 130, 2837–2844 (2007). [PubMed: 17928318]
41. Savva GM et al. Age, neuropathology, and dementia. *N. Engl. J. Med* 360, 2302–2309 (2009). [PubMed: 19474427]
42. Ingelsson M. et al. Early A β accumulation and progressive synaptic loss, gliosis, and tangle formation in AD brain. *Neurology* 62, 925–931 (2004). [PubMed: 15037694]
43. Lesné SE et al. Brain amyloid- β oligomers in ageing and Alzheimer's disease. *Brain* 136, 1383–1398 (2013). [PubMed: 23576130]
44. Selkoe DJ Soluble oligomers of the amyloid β -protein impair synaptic plasticity and behavior. *Behav. Brain Res* 192, 106–113 (2008). [PubMed: 18359102]
45. Klein WL, Krafft GA & Finch CE Targeting small A β oligomers: The solution to an Alzheimer's disease conundrum. *Trends Neurosci.* 24, 219–224 (2001). [PubMed: 11250006]
46. Walsh DM & Selkoe DJ A β oligomers - a decade of discovery. *J. Neurochem* 101, 1172–1184 (2007). [PubMed: 17286590]
47. Shankar GM et al. Natural oligomers of the Alzheimer amyloid- β protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-dependent signaling pathway. *J. Neurosci* 27, 2866–2875 (2007). [PubMed: 17360908]
48. Cheng IH et al. Accelerating amyloid- β fibrillization reduces oligomer levels and functional deficits in Alzheimer disease mouse models. *J. Biol. Chem* 282, 23818–23828 (2007). [PubMed: 17548355]
49. Tomiyama T. et al. A mouse model of amyloid β oligomers: their contribution to synaptic alteration, abnormal tau phosphorylation, glial activation, and neuronal loss in vivo. *J. Neurosci* 30, 4845–4856 (2010). [PubMed: 20371804]
50. Lesné S. et al. A specific amyloid- β protein assembly in the brain impairs memory. *Nature* 440, 352–357 (2006). [PubMed: 16541076]
51. Meilandt WJ et al. Neprilysin overexpression inhibits plaque formation but fails to reduce pathogenic A β oligomers and associated cognitive deficits in human amyloid precursor protein transgenic mice. *J. Neurosci* 29, 1977–1986 (2009). [PubMed: 19228952]
52. Mably AJ et al. Anti- A β antibodies incapable of reducing cerebral A β oligomers fail to attenuate spatial reference memory deficits in J20 mice. *Neurobiol. Dis* 82, 372–384 (2015). [PubMed: 26215784]
53. De Strooper B. Loss-of-function presenilin mutations in Alzheimer disease. *EMBO Rep.* 8, 141–146 (2007). [PubMed: 17268505]
54. Pimplikar SW Neuroinflammation in Alzheimer's disease: from pathogenesis to a therapeutic target. *J. Clin. Immunol* 34, S64–S69 (2014). [PubMed: 24711006]
55. Xia D. et al. Presenilin-1 knockin mice reveal loss-of-function mechanism for familial Alzheimer's disease. *Neuron* 85, 967–981 (2015). [PubMed: 25741723]

56. Bredezen DE Neurodegeneration in Alzheimer's disease: caspases and synaptic element interdependence. *Mol. Neurodegener* 4, 27 (2009). [PubMed: 19558683]
57. Bentley WJ, Li JM, Snyder AZ, Raichle ME & Snyder LH Oxygen level and LFP in task-positive and task-negative areas: bridging BOLD fMRI and electrophysiology. *Cereb. Cortex*, 26, 346–357 (2014). [PubMed: 25385710],
58. Boyatzis RE, Rochford K. & Jack AI Antagonistic neural networks underlying differentiated leadership roles. *Front. Hum. Neurosci* 8, 114 (2014). [PubMed: 24624074]
59. Raichle ME et al. A default mode of brain function. *Proc. Natl. Acad. Sci. USA* 98, 676–682 (2001). [PubMed: 11209064]
60. Greicius MD, Krasnow B, Reiss AL & Menon V. Functional connectivity in the resting brain: a network analysis of the default mode hypothesis. *Proc. Natl. Acad. Sci. USA* 100, 253–258 (2003). [PubMed: 12506194]
61. Sperling RA et al. Functional alterations in memory networks in early Alzheimer's disease. *Neuromolecular Med.* 12, 27–43 (2010). [PubMed: 20069392]
62. Bookheimer SY et al. Patterns of brain activation in people at risk for Alzheimer's disease. *N. Engl. J. Med* 343, 450–456 (2000). [PubMed: 10944562] This study shows hippocampal hyperactivity during memory tasks in asymptomatic APOE ϵ 4 carriers.
63. Trivedi MA et al. fMRI activation during episodic encoding and metacognitive appraisal across the lifespan: risk factors for Alzheimer's disease. *Neuropsychologia* 46, 1667–1678 (2008). [PubMed: 18241895]
64. Kunz L. et al. Reduced grid-cell-like representations in adults at genetic risk for Alzheimer's disease. *Science* 350, 430–433 (2015). [PubMed: 26494756]
65. Quiroz YT et al. Hippocampal hyperactivation in presymptomatic familial Alzheimer's disease. *Ann. Neurol* 68, 865–875 (2010). [PubMed: 21194156]
66. Sepulveda-Falla D, Glatzel M. & Lopera F. Phenotypic profile of early-onset familial Alzheimer's disease caused by presenilin-1 E280A mutation. *J. Alzheimers Dis* 32, 1–12 (2012). [PubMed: 22766738]
67. Bakker A. et al. Reduction of hippocampal hyperactivity improves cognition in amnesic mild cognitive impairment. *Neuron* 74, 467–474 (2012). [PubMed: 22578498] This study shows that treatment with low doses of the antiepileptic drug levetiracetam reduces hippocampal hyperactivity and improves cognition in patients with MCI.
68. Dickerson BC et al. Increased hippocampal activation in mild cognitive impairment compared to normal aging and AD. *Neurology* 65, 404–411 (2005). [PubMed: 16087905]
69. Celone KA et al. Alterations in memory networks in mild cognitive impairment and Alzheimer's disease: an independent component analysis. *J. Neurosci* 26, 10222–10231 (2006). [PubMed: 17021177]
70. Persson J. et al. Altered deactivation in individuals with genetic risk for Alzheimer's disease. *Neuropsychologia* 46, 1679–1687 (2008). [PubMed: 18346764]
71. Dickerson BC et al. Medial temporal lobe function and structure in mild cognitive impairment. *Ann. Neurol* 56, 27–35 (2004). [PubMed: 15236399]
72. Putcha D. et al. Hippocampal hyperactivation associated with cortical thinning in Alzheimer's disease signature regions in non-demented elderly adults. *J. Neurosci* 31, 17680–17688 (2011). [PubMed: 22131428]
73. Bakker A, Albert MS, Krauss G, Speck CL & Gallagher M. Response of the medial temporal lobe network in amnesic mild cognitive impairment to therapeutic intervention assessed by fMRI and memory task performance. *NeuroImage Clin.* 7, 688–698 (2015). [PubMed: 25844322]
74. Sanchez PE et al. Levetiracetam suppresses neuronal network dysfunction and reverses synaptic and cognitive deficits in an Alzheimer's disease model. *Proc. Natl. Acad. Sci. USA* 109, E2895–E2903 (2012). [PubMed: 22869752] This study shows that levetiracetam reduces network hypersynchrony, synaptic deficits and behavioral abnormalities in hAPP-J20 mice.
75. Nygaard HB et al. Brivaracetam, but not ethosuximide, reverses memory impairments in an Alzheimer's disease mouse model. *Alzheimer Res. Ther* 7, 25 (2015).
76. Suberbielle E. et al. Physiologic brain activity causes DNA double-strand breaks in neurons, with exacerbation by amyloid- β . *Nat. Neurosci* 16, 613–621 (2013). [PubMed: 23525040]

77. Suberbielle E. et al. DNA repair factor BRCA1 depletion occurs in Alzheimer brains and impairs cognitive function in mice. *Nat. Commun* 6, 8897 (2015). [PubMed: 26615780]
78. Madabhushi R, Pan L. & Tsai LH DNA damage and its links to neurodegeneration. *Neuron* 83, 266–282 (2014). [PubMed: 25033177]
79. Kim J, Basak JM & Holtzman DM The role of apolipoprotein E in Alzheimer's disease. *Neuron* 63, 287–303 (2009). [PubMed: 19679070]
80. Saito T. et al. Single App knock-in mouse models of Alzheimer's disease. *Nat. Neurosci* 17, 661–663 (2014). [PubMed: 24728269]
81. Roh JH et al. Disruption of the sleep-wake cycle and diurnal fluctuation of β -amyloid in mice with Alzheimer's disease pathology. *Sci. Transl. Med* 4, 150ra122 (2012).
82. Kang JE et al. Amyloid- β dynamics are regulated by orexin and the sleep-wake cycle. *Science* 326, 1005–1007 (2009). [PubMed: 19779148]
83. Xie L. et al. Sleep drives metabolite clearance from the adult brain. *Science* 342, 373–377 (2013). [PubMed: 24136970]
84. Bero AW et al. Neuronal activity regulates the regional vulnerability to amyloid- β deposition. *Nat. Neurosci* 14, 750–756 (2011). [PubMed: 21532579]
85. Yamamoto K. et al. Chronic optogenetic activation augments A β pathology in a mouse model of Alzheimer disease. *Cell Rep.* 11, 859–865 (2015). [PubMed: 25937280]
86. Dolev I. et al. Spike bursts increase amyloid- β 40/42 ratio by inducing a presenilin-1 conformational change. *Nat. Neurosci* 16, 587–595 (2013). [PubMed: 23563578] This study shows that amyloid- β production and the amyloid- β 1–42/amyloid- β 1–40 ratio are regulated by neuronal action potential firing through calcium-dependent conformational changes in PSEN1 at presynaptic terminals.
87. Nitsch RM, Farber SA, Growdon HJ & Wurtman RJ Release of amyloid β -protein precursor derivatives by electrical depolarization of rat hippocampal slices. *Proc. Natl. Acad. Sci. USA* 90, 5191–5193 (1993). [PubMed: 8506366]
88. Kamenetz F. et al. APP processing and synaptic function. *Neuron* 37, 925–937 (2003). [PubMed: 12670422] This study shows that neuronal activity regulates amyloid- β production.
89. Abbott LF & Regehr WG Synaptic computation. *Nature* 431, 796–803 (2004). [PubMed: 15483601]
90. Laviv T. et al. Basal GABA regulates GABA β R conformation and release probability at single hippocampal synapses. *Neuron* 67, 253–267 (2010). [PubMed: 20670833]
91. Palop JJ & Mucke L. Amyloid- β -induced neuronal dysfunction in Alzheimer's disease: from synapses toward neural networks. *Nat. Neurosci* 13, 812–818 (2010). [PubMed: 20581818]
92. Cirrito JR et al. Synaptic activity regulates interstitial fluid amyloid- β levels in vivo. *Neuron* 48, 913–922 (2005). [PubMed: 16364896]
93. Abramov E. et al. Amyloid- β as a positive endogenous regulator of release probability at hippocampal synapses. *Nat. Neurosci* 12, 1567–1576 (2009). [PubMed: 19935655]
94. Puzzo D. et al. Picomolar amyloid- β positively modulates synaptic plasticity and memory in hippocampus. *J. Neurosci* 28, 14537–14545 (2008). [PubMed: 19118188]
95. Willem M. et al. η -Secretase processing of APP inhibits neuronal activity in the hippocampus. *Nature*, 526, 443–447 (2015). [PubMed: 26322584]
96. Palop JJ et al. Aberrant excitatory neuronal activity and compensatory remodeling of inhibitory hippocampal circuits in mouse models of Alzheimer's disease. *Neuron* 55, 697–711 (2007). [PubMed: 17785178] This study shows that hAPP-J20 mice have network hyperexcitability that is associated with extensive anatomical and physiological alterations in learning and memory centers.
97. Yang XF, Weisenfeld A. & Rothman SM Prolonged exposure to levetiracetam reveals a presynaptic effect on neurotransmission. *Epilepsia* 48, 1861–1869 (2007). [PubMed: 17521346]
98. Meehan AL, Yang X, McAdams BD, Yuan L. & Rothman SM A new mechanism for antiepileptic drug action: vesicular entry may mediate the effects of levetiracetam. *J. Neurophysiol* 106, 1227–1239 (2011). [PubMed: 21653714]
99. Kurudenkandy FR et al. Amyloid- β -induced action potential desynchronization and degradation of hippocampal gamma oscillations is prevented by interference with peptide conformation change

and aggregation. *J. Neurosci* 34, 11416–11425 (2014). [PubMed: 25143621] This study shows that amyloid- β reduces gamma oscillations by desynchronizing the action potentials of hippocampal pyramidal cells.

100. Sanchez-Mejia RO et al. Phospholipase A₂ reduction ameliorates cognitive deficits in mouse model of Alzheimer's disease. *Nat. Neurosci* 11, 1311–1318 (2008). [PubMed: 18931664]
101. Busche MA et al. Critical role of soluble amyloid- β for early hippocampal hyperactivity in a mouse model of Alzheimer's disease. *Proc. Natl. Acad. Sci. USA* 109, 8740–8745 (2012). [PubMed: 22592800] This study shows that amyloid- β increases hippocampal neuronal activity in vivo by calcium imaging.
102. Minkeviciene R. et al. Amyloid β -induced neuronal hyperexcitability triggers progressive epilepsy. *J. Neurosci* 29, 3453–3462 (2009). [PubMed: 19295151]
103. Um JW et al. Alzheimer amyloid- β oligomer bound to postsynaptic prion protein activates Fyn to impair neurons. *Nat. Neurosci* 15, 1227–1235 (2012). [PubMed: 22820466]
104. Bezzina C. et al. Early onset of hypersynchronous network activity and expression of a marker of chronic seizures in the Tg2576 mouse model of Alzheimer's disease. *PLoS ONE* 10, e0119910 (2015).
105. Harris JA et al. Transsynaptic progression of amyloid- β -induced neuronal dysfunction within the entorhinal-hippocampal network. *Neuron* 68, 428–441 (2010). [PubMed: 21040845]
106. Born HA et al. Genetic suppression of transgenic APP rescues hypersynchronous network activity in a mouse model of Alzheimer's disease. *J. Neurosci* 34, 3826–3840 (2014). [PubMed: 24623762]
107. Ittner AA, Gladbach A, Bertz J, Suh LS & Ittner LM p38 MAP kinase-mediated NMDA receptor-dependent suppression of hippocampal hypersynchronicity in a mouse model of Alzheimer inverted question marks disease. *Acta Neuropathol. Commun* 2, 149 (2014). [PubMed: 25331068]
108. Roberson ED et al. Amyloid- β /Fyn-induced synaptic, network, and cognitive impairments depend on tau levels in multiple mouse models of Alzheimer's disease. *J. Neurosci* 31, 700–711 (2011). [PubMed: 21228179]
109. Busche MA et al. Clusters of hyperactive neurons near amyloid plaques in a mouse model of Alzheimer's disease. *Science* 321, 1686–1689 (2008). [PubMed: 18802001] This study shows that APP23xPS45 mice have hyperactive neurons and synchronized neuronal activity by calcium imaging.
110. Roberson ED et al. Reducing endogenous tau ameliorates amyloid β -induced deficits in an Alzheimer's disease mouse model. *Science* 316, 750–754 (2007). [PubMed: 17478722] This study shows that endogenous tau promotes APP-dependent network hypersynchrony and cognitive decline.
111. Ittner LM et al. Dendritic function of tau mediates amyloid- β toxicity in Alzheimer's disease mouse models. *Cell* 142, 387–397 (2010). [PubMed: 20655099]
112. Palop JJ et al. Vulnerability of dentate granule cells to disruption of Arc expression in human amyloid precursor protein transgenic mice. *J. Neurosci* 25, 9686–9693 (2005). [PubMed: 16237173]
113. Chin J. et al. Fyn kinase induces synaptic and cognitive impairments in a transgenic mouse model of Alzheimer's disease. *J. Neurosci* 25, 9694–9703 (2005). [PubMed: 16237174]
114. Grienberger C. et al. Staged decline of neuronal function in vivo in an animal model of Alzheimer's disease. *Nat. Commun* 3, 774 (2012). [PubMed: 22491322]
115. Rudinskiy N. et al. Orchestrated experience-driven Arc responses are disrupted in a mouse model of Alzheimer's disease. *Nat. Neurosci* 15, 1422–1429 (2012). [PubMed: 22922786]
116. Spires TL & Hyman BT Neuronal structure is altered by amyloid plaques. *Rev. Neurosci* 15, 267–278 (2004). [PubMed: 15526551]
117. Koffie RM et al. Oligomeric amyloid β associates with postsynaptic densities and correlates with excitatory synapse loss near senile plaques. *Proc. Natl. Acad. Sci. USA* 106, 4012–4017 (2009). [PubMed: 19228947]

118. Siskova Z. et al. Dendritic structural degeneration is functionally linked to cellular hyperexcitability in a mouse model of Alzheimer's disease. *Neuron* 84, 1023–1033 (2014). [PubMed: 25456500]
119. Garcia-Cabrero AM et al. Hyperexcitability and epileptic seizures in a model of frontotemporal dementia. *Neurobiol. Dis* 58, 200–208 (2013). [PubMed: 23774255]
120. Hunter JM et al. Emergence of a seizure phenotype in aged apolipoprotein epsilon 4 targeted replacement mice. *Brain Res.* 1467, 120–132 (2012). [PubMed: 22682924]
121. Morris M. et al. Network dysfunction in α -synuclein transgenic mice and human Lewy body dementia. *Ann. Clin. Transl. Neurol* 2, 1012–1028 (2015). [PubMed: 26732627]
122. Siwek ME et al. Altered theta oscillations and aberrant cortical excitatory activity in the 5XFAD model of Alzheimer's disease. *Neural Plast.* 2015, 781731 (2015). [PubMed: 25922768]
123. Lalonde R, Dumont M, Staufenbiel M. & Strazielle C. Neurobehavioral characterization of APP23 transgenic mice with the SHIRPA primary screen. *Behav. Brain Res* 157, 91–98 (2005). [PubMed: 15617775]
124. Hsiao KK et al. Age-related CNS disorder and early death in transgenic FVB/N mice overexpressing Alzheimer amyloid precursor proteins. *Neuron* 15, 1203–1218 (1995). [PubMed: 7576662]
125. Vossel KA et al. Tau reduction prevents A β -induced impairments in axonal transport. *Science* 330, 198 (2010). [PubMed: 20829454]
126. Devos SL et al. Antisense reduction of tau in adult mice protects against seizures. *J. Neurosci* 33, 12887–12897 (2013). [PubMed: 23904623]
127. Romanelli MF, Morris JC, Ashkin K. & Coben LA Advanced Alzheimer's disease is a risk factor for late-onset seizures. *Arch. Neurol* 47, 847–850 (1990). [PubMed: 2375689]
128. Hauser WA, Morris ML, Heston LL & Anderson VE Seizures and myoclonus in patients with Alzheimer's disease. *Neurology* 36, 1226–1230 (1986). [PubMed: 3092131]
129. Irizarry MC et al. Incidence of new-onset seizures in mild to moderate Alzheimer disease. *Arch. Neurol* 69, 368–372 (2012). [PubMed: 22410444]
130. Risse SC et al. Myoclonus, seizures, and paratonia in Alzheimer disease. *Alzheimer Dis. Assoc. Disord* 4, 217–225 (1990). [PubMed: 2264979]
131. Horvath A, Szucs A, Barcs G, Noebels JL & Kamondi A. Epileptic seizures in Alzheimer disease: a review. *Alzheimer Dis. Assoc. Disord* 30, 186–192 (2016). [PubMed: 26756385]
132. Imfeld P, Bodmer M, Schuerch M, Jick SS & Meier CR Seizures in patients with Alzheimer's disease or vascular dementia: a population-based nested case-control analysis. *Epilepsia* 54, 700–707 (2013). [PubMed: 23215680]
133. Amatniek JC et al. Incidence and predictors of seizures in patients with Alzheimer's disease. *Epilepsia* 47, 867–872 (2006). [PubMed: 16686651] This study shows that early onset of AD is a risk factor for seizures.
134. Sherzai D, Losey T, Vega S. & Sherzai A. Seizures and dementia in the elderly: Nationwide Inpatient Sample 1999–2008. *Epilepsy Behav.* 36, 53–56 (2014). [PubMed: 24857809]
135. Scarmeas N. et al. Seizures in Alzheimer disease: who, when, and how common? *Arch. Neurol* 66, 992–997 (2009). [PubMed: 19667221]
136. Vossel KA et al. Seizures and epileptiform activity in the early stages of Alzheimer disease. *JAMA Neurol.* 70, 1158–1166 (2013). [PubMed: 23835471] This study shows that epileptic activity in patients with MCI or early AD can occur early in the disease course and is frequently non-convulsive in nature.
137. Hesdorffer DC, Hauser WA, Annegers JF, Kokmen E. & Rocca WA Dementia and adult-onset unprovoked seizures. *Neurology* 46, 727–730 (1996). [PubMed: 8618673]
138. Cretin B. et al. Epileptic prodromal Alzheimer's disease, a retrospective study of 13 new cases: expanding the spectrum of Alzheimer's disease to an epileptic variant? *J. Alzheimers Dis* 52, 1125–1133 (2016). [PubMed: 27104892]
139. Palop JJ & Mucke L. Epilepsy and cognitive impairments in Alzheimer disease. *Arch. Neurol* 66, 435–440 (2009). [PubMed: 19204149]

140. Noebels J. A perfect storm: converging paths of epilepsy and Alzheimer's dementia intersect in the hippocampal formation. *Epilepsia* 52 (Suppl. 1), 39–46 (2011).
141. Larner AJ Presenilin-1 mutation Alzheimer's disease: a genetic epilepsy syndrome? *Epilepsy Behav.* 21, 20–22 (2011). [PubMed: 21501974]
142. Larner AJ & Doran M. Clinical phenotypic heterogeneity of Alzheimer's disease associated with mutations of the presenilin-1 gene. *J. Neurol* 253, 139–158 (2006). [PubMed: 16267640] This study shows that seizures are a clinical feature of many pedigrees of FAD patients carrying PSEN1 mutations.
143. Jayadev S. et al. Alzheimer's disease phenotypes and genotypes associated with mutations in presenilin 2. *Brain* 133, 1143–1154 (2010). [PubMed: 20375137]
144. Cabrejo L. et al. Phenotype associated with APP duplication in five families. *Brain* 129, 2966–2976 (2006). [PubMed: 16959815] This study shows that seizures are a clinical feature of many FAD patients carrying wild-type APP duplications.
145. McNaughton D. et al. Duplication of amyloid precursor protein (APP), but not prion protein (PRNP) gene is a significant cause of early onset dementia in a large UK series. *Neurobiol. Aging* 33, e13–21 (2012).
146. Zarea A. et al. Seizures in dominantly inherited Alzheimer disease. *Neurology* 87, 912–919 (2016). [PubMed: 27466472]
147. Lai F. & Williams RS A prospective study of Alzheimer disease in Down syndrome. *Arch. Neurol* 46, 849–853 (1989). [PubMed: 2527024]
148. Snider BJ et al. Novel presenilin 1 mutation (S170F) causing Alzheimer disease with Lewy bodies in the third decade of life. *Arch. Neurol* 62, 1821–1830 (2005). [PubMed: 16344340]
149. Moehlmann T. et al. Presenilin-1 mutations of leucine 166 equally affect the generation of the Notch and APP intracellular domains independent of their effect on A β ₄₂ production. *Proc. Natl Acad. Sci. USA* 99, 8025–8030 (2002). [PubMed: 12048239]
150. Jensen O. & Colgin LL Cross-frequency coupling between neuronal oscillations. *Trends Cognit. Sci* 11, 267–269 (2007). [PubMed: 17548233]
151. Viney TJ et al. Network state-dependent inhibition of identified hippocampal CA3 axo-axonic cells in vivo. *Nat. Neurosci* 16, 1802–1811 (2013). [PubMed: 24141313]
152. Tukker JJ, Fuentealba P, Hartwich K, Somogyi P. & Klausberger T. Cell type-specific tuning of hippocampal interneuron firing during gamma oscillations in vivo. *J. Neurosci* 27, 8184–8189 (2007). [PubMed: 17670965]
153. Uhlhaas PJ et al. Neural synchrony in cortical networks: history, concept and current status. *Front. Integr. Neurosci* 3, 17 (2009). [PubMed: 19668703]
154. Matsumoto JY et al. Network oscillations modulate interictal epileptiform spike rate during human memory. *Brain* 136, 2444–2456 (2013). [PubMed: 23803305] This study shows that memory-induced oscillatory changes, including increased gamma power, reduce epileptiform discharges in humans with epilepsy.
155. Sederberg PB et al. Gamma oscillations distinguish true from false memories. *Psychol. Sci* 18, 927–932 (2007). [PubMed: 17958703]
156. Sederberg PB et al. Hippocampal and neocortical gamma oscillations predict memory formation in humans. *Cereb. Cortex* 17, 1190–1196 (2007). [PubMed: 16831858] This study shows that increases in cortical and hippocampal gamma oscillations during memory tasks predict successful encoding of new memories in humans.
157. Jensen O, Kaiser J. & Lachaux JP Human gamma-frequency oscillations associated with attention and memory. *Trends Neurosci.* 30, 317–324 (2007). [PubMed: 17499860]
158. Yamamoto J, Suh J, Takeuchi D. & Tonegawa S. Successful execution of working memory linked to synchronized high-frequency gamma oscillations. *Cell* 157, 845–857 (2014). [PubMed: 24768692]
159. Tallon-Baudry C, Bertrand O, Delpuech C. & Pernier J. Stimulus specificity of phase-locked and non-phase-locked 40 Hz visual responses in human. *J. Neurosci* 16, 4240–4249 (1996). [PubMed: 8753885]
160. Lachaux JP et al. The many faces of the gamma band response to complex visual stimuli. *Neuroimage* 25, 491–501 (2005). [PubMed: 15784428]

161. Harris KD & Thiele A. Cortical state and attention. *Nat. Rev. Neurosci* 12, 509–523 (2011). [PubMed: 21829219]
162. Crone NE, Miglioretti DL, Gordon B. & Lesser RP Functional mapping of human sensorimotor cortex with electrocorticographic spectral analysis. II. Event-related synchronization in the gamma band. *Brain*. 121, 2301–2315 (1998). [PubMed: 9874481]
163. Burke JF et al. Synchronous and asynchronous theta and gamma activity during episodic memory formation. *J. Neurosci* 33, 292–304 (2013). [PubMed: 23283342]
164. Sun L. et al. Evidence for dysregulated high-frequency oscillations during sensory processing in medication-naive, first episode schizophrenia. *Schizophr. Res* 150, 519–525 (2013). [PubMed: 24016727]
165. Herrmann CS & Demiralp T. Human EEG gamma oscillations in neuropsychiatric disorders. *Clin. Neurophysiol* 116, 2719–2733 (2005). [PubMed: 16253555]
166. Niessing J. et al. Hemodynamic signals correlate tightly with synchronized gamma oscillations. *Science* 309, 948–951 (2005). [PubMed: 16081740]
167. Logothetis NK, Pauls J, Augath M, Trinath T. & Oeltermann A. Neurophysiological investigation of the basis of the fMRI signal. *Nature* 412, 150–157 (2001). [PubMed: 11449264]
168. Goldman RI, Stern JM, Engel J Jr. & Cohen MS Simultaneous EEG and fMRI of the alpha rhythm. *Neuroreport* 13, 2487–2492 (2002). [PubMed: 12499854]
169. Laufs H. et al. EEG-correlated fMRI of human alpha activity. *Neuroimage* 19, 1463–1476 (2003). [PubMed: 12948703]
170. Scheeringa R. et al. Trial-by-trial coupling between EEG and BOLD identifies networks related to alpha and theta EEG power increases during working memory maintenance. *Neuroimage* 44, 1224–1238 (2009). [PubMed: 18840533]
171. Ossandon T. et al. Transient suppression of broadband gamma power in the default-mode network is correlated with task complexity and subject performance. *J. Neurosci* 31, 14521–14530 (2011). [PubMed: 21994368]
172. Verret L. et al. Inhibitory interneuron deficit links altered network activity and cognitive dysfunction in Alzheimer model. *Cell* 149, 708–721 (2012). [PubMed: 22541439] This study shows that impairments of inhibitory interneurons contribute to network hypersynchrony, altered oscillatory rhythms, and behavioural deficits in hAPP-J20 mice.
173. Maheshwari A, Marks RL, Yu KM & Noebels JL Shift in interictal relative gamma power as a novel biomarker for drug response in two mouse models of absence epilepsy. *Epilepsia* 57, 79–88 (2016). [PubMed: 26663261]
174. Crooks VC, Lubben J, Petitti DB, Little D. & Chiu V. Social network, cognitive function, and dementia incidence among elderly women. *Am. J. Public Health* 98, 1221–1227 (2008). [PubMed: 18511731]
175. Buzsaki G. & Draguhn A. Neuronal oscillations in cortical networks. *Science* 304, 1926–1929 (2004). [PubMed: 15218136]
176. Hestrin S. & Galarreta M. Electrical synapses define networks of neocortical GABAergic neurons. *Trends Neurosci.* 28, 304–309 (2005). [PubMed: 15927686]
177. Lapray D. et al. Behavior-dependent specialization of identified hippocampal interneurons. *Nat. Neurosci* 15, 1265–1271 (2012). [PubMed: 22864613]
178. Fu Y. et al. A cortical circuit for gain control by behavioral state. *Cell* 156, 1139–1152 (2014). [PubMed: 24630718]
179. Sohal VS, Zhang F, Yizhar O. & Deisseroth K. Parvalbumin neurons and gamma rhythms enhance cortical circuit performance. *Nature* 459, 698–702 (2009). [PubMed: 19396159] This study shows that parvalbumin cell activity is mechanistically linked to gamma oscillations.
180. Fazzari P. et al. Control of cortical GABA circuitry development by Nrg1 and ErbB4 signalling. *Nature* 464, 1376–1380 (2010). [PubMed: 20393464]
181. Kepecs A. & Fishell G. Interneuron cell types are fit to function. *Nature* 505, 318–326 (2014). [PubMed: 24429630]
182. Southwell DG et al. Interneurons from embryonic development to cell-based therapy. *Science* 344, 1240622 (2014).

183. Han S. et al. Autistic-like behaviour in *Scn1a*^{+/-} mice and rescue by enhanced GABA-mediated neurotransmission. *Nature* 489, 385–390 (2012). [PubMed: 22914087]
184. Chao HT et al. Dysfunction in GABA signalling mediates autism-like stereotypies and Rett syndrome phenotypes. *Nature* 468, 263–269 (2010). [PubMed: 21068835]
185. Hunt RF, Girsakis KM, Rubenstein JL, Alvarez-Buylla A. & Baraban SC GABA progenitors grafted into the adult epileptic brain control seizures and abnormal behavior. *Nat. Neurosci* 16, 692–697 (2013). [PubMed: 23644485]
186. Tong LM et al. Inhibitory interneuron progenitor transplantation restores normal learning and memory in ApoE4 knock-in mice without or with A β accumulation. *J. Neurosci* 34, 9506–9515 (2014). [PubMed: 25031394] This study shows that transplants of inhibitory interneurons restore learning and memory in APOE4-KI mice.
187. Freund TF & Katona I. Perisomatic inhibition. *Neuron* 56, 33–42 (2007). [PubMed: 17920013]
188. Schmidt MJ et al. Modulation of behavioral networks by selective interneuronal inactivation. *Mol. Psychiatry* 19, 580–587 (2014). [PubMed: 24322205]
189. Kvitsiani D. et al. Distinct behavioural and network correlates of two interneuron types in prefrontal cortex. *Nature* 498, 363–366 (2013). [PubMed: 23708967]
190. Cardin JA et al. Driving fast-spiking cells induces gamma rhythm and controls sensory responses. *Nature* 459, 663–667 (2009). [PubMed: 19396156]
191. Cramer PE et al. ApoE-directed therapeutics rapidly clear β -amyloid and reverse deficits in AD mouse models. *Science* 335, 1503–1506 (2012). [PubMed: 22323736]
192. Rubio SE et al. Accelerated aging of the GABAergic septohippocampal pathway and decreased hippocampal rhythms in a mouse model of Alzheimer’s disease. *FASEB J.* 26, 4458–4467 (2012). [PubMed: 22835830]
193. Gurevicius K, Lipponen A. & Tanila H. Increased cortical and thalamic excitability in freely moving APP^{swE}/PS1^{dE9} mice modeling epileptic activity associated with Alzheimer’s disease. *Cereb. Cortex* 23, 1148–1158 (2013). [PubMed: 22581851]
194. Kemere C, Carr MF, Karlsson MP & Frank LM Rapid and continuous modulation of hippocampal network state during exploration of new places. *PLoS ONE* 8, e73114 (2013).
195. Busche MA et al. Rescue of long-range circuit dysfunction in Alzheimer’s disease models. *Nat. Neurosci* 18, 1623–1630 (2015). [PubMed: 26457554]
196. Mander BA et al. β -Amyloid disrupts human NREM slow waves and related hippocampus-dependent memory consolidation. *Nat. Neurosci* 18, 1051–1057 (2015). [PubMed: 26030850]
197. van Deursen JA, Vuurman EF, van Kranen-Mastenbroek VH, Verhey FR & Riedel WJ 40-Hz steady state response in Alzheimer’s disease and mild cognitive impairment. *Neurobiol Aging* 32, 24–30 (2011). [PubMed: 19237225]
198. van Deursen JA, Vuurman EF, Verhey FR, van Kranen-Mastenbroek VH & Riedel WJ Increased EEG gamma band activity in Alzheimer’s disease and mild cognitive impairment. *J. Neural Transm. (Vienna)* 115, 1301–1311 (2008). [PubMed: 18607528]
199. Nimmrich V, Draguhn A. & Axmacher N. Neuronal network oscillations in neurodegenerative diseases. *Neuromolecular Med.* 17, 270–284 (2015). [PubMed: 25920466]
200. Gillespie AK et al. Apolipoprotein E4 causes age-dependent disruption of slow gamma oscillations during hippocampal sharp-wave ripples. *Neuron* 90, 740–751 (2016). [PubMed: 27161522] This study shows reduced gamma power during hippocampal sharp-wave ripples in APOE4-KI mice.
201. Pena-Ortega F, Solis-Cisneros A, Ordaz B, Balleza-Tapia H. & Javier Lopez-Guerrero J. Amyloid beta 1–42 inhibits entorhinal cortex activity in the beta-gamma range: role of GSK-3. *Curr. Alzheimer Res* 9, 857–863 (2012). [PubMed: 22631612]
202. Driver JE et al. Impairment of hippocampal gamma-frequency oscillations in vitro in mice overexpressing human amyloid precursor protein (APP). *Eur. J. Neurosci* 26, 1280–1288 (2007). [PubMed: 17767505]
203. Villette V. et al. Decreased rhythmic GABAergic septal activity and memory-associated theta oscillations after hippocampal amyloid- β pathology in the rat. *J. Neurosci* 30, 10991–11003 (2010). [PubMed: 20720106]

204. Royer S. et al. Control of timing, rate and bursts of hippocampal place cells by dendritic and somatic inhibition. *Nat. Neurosci* 15, 769–775 (2012). [PubMed: 22446878]
205. Tort AB, Komorowski RW, Manns JR, Kopell NJ & Eichenbaum H. Theta-gamma coupling increases during the learning of item-context associations. *Proc. Natl Acad. Sci. USA* 106, 20942–20947 (2009). [PubMed: 19934062]
206. Lakatos P, Karmos G, Mehta AD, Ulbert I. & Schroeder CE Entrainment of neuronal oscillations as a mechanism of attentional selection. *Science* 320, 110–113 (2008). [PubMed: 18388295]
207. Canolty RT et al. High gamma power is phase-locked to theta oscillations in human neocortex. *Science* 313, 1626–1628 (2006). [PubMed: 16973878]
208. Bragin A. et al. Gamma (40–100 Hz) oscillation in the hippocampus of the behaving rat. *J. Neurosci* 15, 47–60 (1995). [PubMed: 7823151]
209. Sirota A. et al. Entrainment of neocortical neurons and gamma oscillations by the hippocampal theta rhythm. *Neuron* 60, 683–697 (2008). [PubMed: 19038224]
210. Freund TF Interneuron diversity series: Rhythm and mood in perisomatic inhibition. *Trends Neurosci.* 26, 489–495 (2003). [PubMed: 12948660]
211. Belluscio MA, Mizuseki K, Schmidt R, Kempter R. & Buzsaki G. Cross-frequency phase-phase coupling between theta and gamma oscillations in the hippocampus. *J. Neurosci* 32, 423–435 (2012). [PubMed: 22238079]
212. Goutagny R. et al. Alterations in hippocampal network oscillations and theta-gamma coupling arise before A β overproduction in a mouse model of Alzheimer’s disease. *Eur. J. Neurosci* 37, 1896–1902 (2013). [PubMed: 23773058]
213. Klausberger T. et al. Brain-state- and cell-type-specific firing of hippocampal interneurons in vivo. *Nature* 421, 844–848 (2003). [PubMed: 12594513] This study shows that different types of interneurons contribute to specific oscillatory frequencies.
214. Andrews-Zwilling Y. et al. Hilar GABAergic interneuron activity controls spatial learning and memory retrieval. *PLoS ONE* 7, e40555 (2012).
215. Wang WZ et al. The developmental changes of Na_v1.1 and Na_v1.2 expression in the human hippocampus and temporal lobe. *Brain Res.* 1389, 61–70 (2011). [PubMed: 21377452]
216. Corbett BF et al. Sodium channel cleavage is associated with aberrant neuronal activity and cognitive deficits in a mouse model of Alzheimer’s disease. *J. Neurosci* 33, 7020–7026 (2013). [PubMed: 23595759]
217. Kim DY et al. BACE1 regulates voltage-gated sodium channels and neuronal activity. *Nat. Cell Biol* 9, 755–764 (2007). [PubMed: 17576410]
218. Yang LB et al. Elevated β -secretase expression and enzymatic activity detected in sporadic Alzheimer disease. *Nat. Med* 9, 3–4 (2003). [PubMed: 12514700]
219. Kim DY et al. The E280A presenilin mutation reduces voltage-gated sodium channel levels in neuronal cells. *Neurodegener. Dis* 13, 64–68 (2014). [PubMed: 24217025]
220. Ponomareva NV, Korovaitseva GI & Rogaev EI EEG alterations in non-demented individuals related to apolipoprotein E genotype and to risk of Alzheimer disease. *Neurobiol. Aging* 29, 819–827 (2008). [PubMed: 17293007]
221. Andrews-Zwilling Y. et al. Apolipoprotein E4 causes age- and Tau-dependent impairment of GABAergic interneurons, leading to learning and memory deficits in mice. *J. Neurosci* 30, 13707–13717 (2010). [PubMed: 20943911]
222. Terry RD et al. Physical basis of cognitive alterations in Alzheimer’s disease: synapse loss is the major correlate of cognitive impairment. *Ann. Neurol* 30, 572–580 (1991). [PubMed: 1789684]
223. DeKosky ST & Scheff SW Synapse loss in frontal cortex biopsies in Alzheimer’s disease: correlation with cognitive severity. *Ann. Neurol* 27, 457–464 (1990). [PubMed: 2360787]
224. Walsh DM et al. Naturally secreted oligomers of amyloid β protein potently inhibit hippocampal long-term potentiation in vivo. *Nature* 416, 535–539 (2002). [PubMed: 11932745]
225. Hsia AY et al. Plaque-independent disruption of neural circuits in Alzheimer’s disease mouse models. *Proc. Natl. Acad. Sci. USA* 96, 3228–3233 (1999). [PubMed: 10077666]

226. Mucke L. et al. High-level neuronal expression of A β _{1–42} in wild-type human amyloid protein precursor transgenic mice: synaptotoxicity without plaque formation. *J. Neurosci* 20, 4050–4058 (2000). [PubMed: 10818140]
227. Kim JH, Anwyl R, Suh YH, Djamgoz MB & Rowan MJ Use-dependent effects of amyloidogenic fragments of β -amyloid precursor protein on synaptic plasticity in rat hippocampus in vivo. *J. Neurosci* 21, 1327–1333 (2001). [PubMed: 11160403]
228. Hsieh H. et al. AMPAR removal underlies A β -induced synaptic depression and dendritic spine loss. *Neuron* 52, 831–843 (2006). [PubMed: 17145504]
229. Palop JJ & Mucke L. Synaptic depression and aberrant excitatory network activity in Alzheimer's disease: two faces of the same coin? *Neuromolecular Med.* 12, 48–55 (2010). [PubMed: 19838821]
230. Liu L. et al. Role of NMDA receptor subtypes in governing the direction of hippocampal synaptic plasticity. *Science* 304, 1021–1024 (2004). [PubMed: 15143284]
231. Campbell SL, Hablitz JJ & Olsen ML Functional changes in glutamate transporters and astrocyte biophysical properties in a rodent model of focal cortical dysplasia. *Front. Cell. Neurosci* 8, 425 (2014). [PubMed: 25565960]
232. Palop JJ et al. Neuronal depletion of calcium-dependent proteins in the dentate gyrus is tightly linked to Alzheimer's disease-related cognitive deficits. *Proc. Natl Acad. Sci. USA* 100, 9572–9577 (2003). [PubMed: 12881482]
233. Cheng JS et al. Collagen VI protects neurons against A β toxicity. *Nat. Neurosci* 12, 119–121 (2009). [PubMed: 19122666]
234. Chin J. et al. Reelin depletion in the entorhinal cortex of human amyloid precursor protein transgenic mice and humans with Alzheimer's disease. *J. Neurosci* 27, 2727–2733 (2007). [PubMed: 17360894]
235. Meilandt WJ et al. Enkephalin elevations contribute to neuronal and behavioral impairments in a transgenic mouse model of Alzheimer's disease. *J. Neurosci* 28, 5007–5017 (2008). [PubMed: 18463254]
236. Orr AG et al. Astrocytic adenosine receptor A_{2A} and G_s-coupled signaling regulate memory. *Nat. Neurosci* 18, 423–434 (2015). [PubMed: 25622143]
237. Albert MS et al. The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement.* 7, 270–279 (2011). [PubMed: 21514249]
238. Gainotti G, Quaranta D, Vita MG & Marra C. Neuropsychological predictors of conversion from mild cognitive impairment to Alzheimer's disease. *J. Alzheimers Dis* 38, 481–495 (2014). [PubMed: 24002185]
239. Budson AE & Solomon PR New criteria for Alzheimer disease and mild cognitive impairment: implications for the practicing clinician. *Neurologist* 18, 356–363 (2012). [PubMed: 23114667]
240. Jensen HS, Grunnet M. & Bastlund JF Therapeutic potential of Na_v1.1 activators. *Trends. Pharmacol. Sci* 35, 113–118 (2014). [PubMed: 24439681]
241. Tuszynski MH et al. A phase 1 clinical trial of nerve growth factor gene therapy for Alzheimer disease. *Nat. Med* 11, 551–555 (2005). [PubMed: 15852017]
242. Knoferle J. et al. Apolipoprotein E4 produced in GABAergic interneurons causes learning and memory deficits in mice. *J. Neurosci* 34, 14069–14078 (2014). [PubMed: 25319703]
243. Buckner RL et al. Molecular, structural, and functional characterization of Alzheimer's disease: evidence for a relationship between default activity, amyloid, and memory. *J. Neurosci* 25, 7709–7717 (2005). [PubMed: 16120771] This study shows that PiB+ amyloid preferentially deposits in brain regions of the DMN.
244. Axelman K, Basun H, Winblad B. & Lannfelt L. A large Swedish family with Alzheimer's disease with a codon 670/671 amyloid precursor protein mutation: a clinical and genealogical investigation. *Arch. Neurol* 51, 1193–1197 (1994). [PubMed: 7986173]
245. Godbolt AK et al. A second family with familial AD and the V717L APP mutation has a later age at onset. *Neurology* 66, 611–612 (2006). [PubMed: 16505331]

246. Abe M. et al. Phenotypical difference of amyloid precursor protein (APP) V717L mutation in Japanese family. *BMC Neurol.* 12, 38 (2012). [PubMed: 22702962]
247. Kennedy AM et al. Familial Alzheimer's disease. A pedigree with a mis-sense mutation in the amyloid precursor protein gene (amyloid precursor protein 717 valine→glycine). *Brain* 116, 309–324 (1993). [PubMed: 8461968]
248. Mullan M. et al. Clinical comparison of Alzheimer's disease in pedigrees with the codon 717 Val→Ile mutation in the amyloid precursor protein gene. *Neurobiol. Aging* 14, 407–419 (1993). [PubMed: 8247223]
249. Lindquist SG et al. Atypical early-onset Alzheimer's disease caused by the Iranian APP mutation. *J. Neurol. Sci* 268, 124–130 (2008). [PubMed: 18187157]
250. Edwards-Lee T. et al. An African American family with early-onset Alzheimer disease and an APP (T714I) mutation. *Neurology* 64, 377–379 (2005). [PubMed: 15668448]
251. Sleegers K. et al. APP duplication is sufficient to cause early onset Alzheimer's dementia with cerebral amyloid angiopathy. *Brain* 129, 2977–2983 (2006). [PubMed: 16921174]
252. Remes AM et al. Hereditary dementia with intracerebral hemorrhages and cerebral amyloid angiopathy. *Neurology* 63, 234–240 (2004). [PubMed: 15277614]
253. Rovelet-Lecrux A. et al. APP locus duplication in a Finnish family with dementia and intracerebral haemorrhage. *J. Neurol. Neurosurg. Psychiatry* 78, 1158–1159 (2007). [PubMed: 17442758]
254. Axelman K, Basun H. & Lannfelt L. Wide range of disease onset in a family with Alzheimer disease and a His163Tyr mutation in the presenilin-1 gene. *Arch. Neurol* 55, 698–702 (1998). [PubMed: 9605727]
255. Mangone CA et al. Early onset Alzheimer's disease in a South American pedigree from Argentina. *Acta Neurol. Scand* 91, 6–13 (1995). [PubMed: 7732777]
256. Mann DMA, et al. Amyloid angiopathy and variability in amyloid β deposition is determined by mutation position in presenilin-1-linked Alzheimer's disease. *Am. J. Pathol* 158, 2165–2175 (2001). [PubMed: 11395394]
257. Frommelt P, Schnabel R, Kuhne W, Nee LE & Polinsky RJ Familial Alzheimer disease: a large, multigeneration German kindred. *Alzheimer Dis. Assoc. Disord* 5, 36–43 (1991). [PubMed: 2025423]
258. Vossel KA et al. Incidence and impact of subclinical epileptiform activity in alzheimer's disease. *Ann. Neurol* 10.1002/ana.24794 (2016).




Figure 1 | Synchrony and functional states of networks.

The degree of correlated neuronal activity (synchrony) reflects the functional state of networks and circuits. **a** | To illustrate this relationship at the network level, the top panels show local field potentials (LFPs) recorded from four electrodes (1–4) inserted 1 mm apart into the cat parietal cortex (suprasylvian gyrus) during sleep (left) and wakefulness (right). Network activity during resting and non-active states is predominantly characterized by synchronized slow-frequency and high-amplitude fluctuations (red shading). By contrast, network activity during active states is characterized by desynchronized fast-frequency and low-amplitude fluctuations (blue shading). The bottom panels show hypothetical representations of the amplitude (red line) and frequency (blue line) of LFP fluctuations and of the associated network synchrony (pink line). **b** | Membrane potential recordings of two layer 2/3 (L2/3) pyramidal neurons from mouse parietal cortex (whisker barrel cortex) during resting and active (whisker use) periods illustrate that neurons desynchronize during active periods. Part **a** is republished with permission of Society for Neuroscience, from

Spatiotemporal analysis of local field potentials and unit discharges in cat cerebral cortex during natural wake and sleep states, Destexhe, A., Contreras, D. & Steriade, M., 19, 11, 1999; permission conveyed through Copyright Clearance Center Inc. Part **b** is from REF.12, Nature Publishing Group.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 2 | Encoding reveals network dysfunction in people with mild cognitive impairment. Memory encoding causes specific changes in brain activity-related functional MRI (fMRI) signals across different networks. People at increased risk for Alzheimer disease (AD) show early changes in the activation and deactivation of specific networks. **a** | Lateral (left panel) and medial (right panel) views of a healthy left hemisphere. During encoding in healthy people, network activity increases in task-related networks (blue) and decreases in non-task-related networks (yellow/orange). Anti-correlated activity (blue versus yellow/orange) in these two widely distributed and non-overlapping networks is also evident when spontaneous fluctuations of fMRI signals during resting states are examined (not shown). The default mode network (yellow and orange regions) includes brain regions that show decreased fMRI activity during attention-demanding tasks but become active during inwardly oriented mental activity. **b** | In healthy individuals whose brain activity was monitored by fMRI during a cognitive task, a relatively smaller extent of hippocampal activation and a relatively greater extent of precuneal deactivation were associated with better cognitive performance. **c** | Pattern separation and completion are cognitive functions that heavily rely on the hippocampal formation and allow us to discriminate (pattern separation) or merge (pattern completion) similar representations or episodes. In a pattern-

separation task, in which individuals were asked to discriminate between slightly different trowels (left panel), patients with amnesic mild cognitive impairment (MCI) made more pattern-completion errors (that is, they failed to discriminate slightly different trowels) and had aberrant hyperactivation of the dentate gyrus (DG) and CA3 regions of the hippocampus on fMRI (right panel). Treatment with the antiepileptic drug levetiracetam reversed the hippocampal hyperactivation and improved the patients' ability to discriminate between images that were similar but not identical. **d** | During a face–name association task, patients with MCI and amyloid deposits in the brain (as revealed by a Pittsburgh compound B-positive (PiB⁺) signal on positron emission tomography) and patients with AD showed deactivation deficits in the precuneus. Part **a** is adapted from REF. 58. Part **b** is adapted with permission from REF. 61, Springer. Part **c** is adapted with permission from REF. 67, Cell Press/Elsevier. Part **d** is adapted with permission from REF. 4, Cell Press/Elsevier.




Figure 3 | Neuronal activity regulates amyloid- β production and deposition.

a | Most patients with Alzheimer disease (AD) and some people without dementia have increased Pittsburgh compound B-positive amyloid deposits in the brain²⁴³ (red). Amyloid deposits predominate in brain regions of the default mode network (blue), which shows deactivation deficits in AD (FIG. 2), suggesting a potential link between aberrant neuronal activity and amyloid deposition. **b** | In APP-A7 mice, chronic, optogenetic stimulation of pyramidal cells in the entorhinal cortex triggered epileptiform activity and increased amyloid deposition in the molecular layer of the dentate gyrus, directly supporting the notion that aberrant neuronal activity can promote amyloid deposition *in vivo*. **c** | At the synaptic level, an increase in the frequency of action potentials (APs) proportionally enhances amyloid- β_{1-42} (A β_{1-42}) and amyloid- β_{1-40} (A β_{1-40}) production. Compared with regular firing, burst firing reduces the A β_{1-42} /A β_{1-40} ratio. **d** | Basal neurotransmitter release determines the ‘filter’ mode of synapses and regulates synaptic plasticity. The top panel indicates that excitatory synapses with lower release probability (high-pass-filter synapses) have greater presynaptic Ca $^{2+}$ build-up, produce lower A β_{1-42} /A β_{1-40} ratios, exhibit synaptic facilitation and primarily transfer potentiated responses. The bottom panel depicts synapses with higher release probability (low-pass-filter synapses), which have less presynaptic Ca $^{2+}$ build-up, produce higher A β_{1-42} /A β_{1-40} ratios and show synaptic depression as well as enhanced

spike transfer. In AD, synapses may shift from high- to low-pass synaptic filtering. **e** | Electroencephalography recordings capture the combined electrical output of neuronal ensembles. In such recordings, most familial AD (FAD) mice, including hAPP-J20 (REF. 96), APP/PSEN1dE9 (REFS 102,103), Tg2576 (REF. 104), 5xFAD¹²², 3xTg-AD⁷⁵, APP/TTA-EC¹⁰⁵, APP/TTA-CaMKII α ¹⁰⁶, and APP23 (REF. 107) mice (Supplementary information S1 (table)), show intermittent large-amplitude epileptiform discharges (denoted by the asterisk in hAPP-J20 mice; bottom panel), which provide evidence of network hypersynchrony. fMRI, functional MRI; PET, positron emission tomography. Part **a** is republished with permission of Society for Neuroscience, from Molecular, structural, and functional characterization of Alzheimer's disease: evidence for a relationship between default activity, amyloid, and memory, Buckner, R. L. *et al.*, 25, 34, 2005; permission conveyed through Copyright Clearance Center, Inc. Part **e** is adapted with permission from REF. 172, Cell Press/Elsevier.




Figure 4 | Neuronal ensembles generate oscillatory activity patterns.

a | Oscillatory rhythms (grey boxes) emerge from, and control, the activity of neuronal ensembles that are formed by excitatory neurons (blue) and inhibitory interneurons (red). The amplitude, or power, of low-frequency oscillations increases during rest, whereas the amplitude of high-frequency oscillations increases during activity. Action potentials (APs) of excitatory neurons are phase-locked to high-power, lower-frequency oscillations (top grey box), whereas APs of inhibitory cells are phase-locked to high-power, high-frequency oscillations (bottom grey box). **b** | The left panel shows local field potential (LFP) gamma oscillations (blue) and APs (red) for a parvalbumin-positive (PV⁺) cell in area CA3, revealing a close association between APs and the phase of gamma oscillations. In the right panel, the AP firing rate (colour coded) is shown as a function of gamma phase for the same PV⁺ cell, which prominently fires during the ascending phase of gamma oscillations. **c** | Optogenetic stimulation of PV⁺ and pyramidal cells at different frequencies resulted in a cell type- and frequency-specific generation of oscillatory activity. 40-Hz stimulation of PV⁺, but not pyramidal, cells increased gamma power, whereas 8-Hz stimulation of pyramidal, but not PV⁺, cells increased theta power. **d** | In macaques, visual stimulation (15s of dim 1 Hz illumination) increased the power of LFP gamma (30–150 Hz) oscillations in a task-related network (visual area V3; left panel) and decreased LFP power across multiple oscillatory frequencies in a default mode network region (posterior cingulate cortex; right panel). SEM, standard error of the mean. Part **b** is from REF.151, Nature Publishing Group. Part **c** is from

REF. 190, Nature Publishing Group. Part **d** is adapted from Bentley, W. J., Li, J.M., Snyder, A. Z., Raichle, M.E. & Snyder, L.H., Oxygen level and LFP in task-positive and task-negative areas: bridging BOLD fMRI and electrophysiology, *Cerebral Cortex*, 2014, 26, 1, 346–357, by permission of Oxford Journals.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 5 | Close association between behavioural state, gamma oscillations and epileptiform activity in mice and humans.

a,b | Behavioural activity, full-frequency range spectrograms, gamma oscillatory power, and distribution of epileptic discharges in cortical networks of an hAPP-J20 mouse (part **a**) and a human with epilepsy (part **b**). In mice, exploration (active) of a novel environment robustly increased gamma oscillatory power and reduced epileptiform discharges, suggesting that the brain state modulates brain rhythms and network hypersynchrony. In humans, successful, but not unsuccessful, memory encoding also increased gamma oscillatory power and reduced epileptiform discharges. **c** | In our hypothetical model, memory encoding requires frequency-specific modulation of oscillatory frequencies, which reduces network hypersynchrony. Part **a** is adapted with permission from REF. 172, Cell Press/Elsevier. Part **b** is adapted from Matsumoto, J.Y. *et al.*, Network oscillations modulate interictal epileptiform spike rate during human memory, *Brain*, 2013, **136**, 8, 2444–2456, by permission of Oxford Journals.




Figure 6 | Targeting interneurons to improve Alzheimer disease-related network dysfunction.

a | In hAPP-J20 mice (middle panel), reduced levels of the voltage-gated sodium channel subunit Nav1.1 in parvalbumin-positive (PV⁺) cells were associated with reduced gamma power, epileptiform activity and cognitive impairment (the level of cognitive performance is denoted by the plus symbols). Restoring Nav1.1 levels in PV⁺ cells with an Nav1.1-BAC (bacterial artificial chromosome) transgene (right panel) reduced all of these deficits. **b** | *APOE4*-KI mice have increased neuronal levels of phosphorylated tau, age-dependent loss of somatostatin-positive interneurons in the hilus of the dentate gyrus, and seizures. Memory deficits in these mice were reduced by pentobarbital treatment²²¹, tau removal²²¹ and transplantation of interneuron precursor cells¹⁶⁰. Part **a** is adapted with permission from REF. 172, Cell Press/Elsevier.

Table 1 |

Network hypersynchrony in humans with FAD caused by mutations used in mouse models.

Affected genes	FAD mutations	Number of patients with seizures (study population percentage)	Age at onset of cognitive deficits, years (mean)	Refs
<i>APP</i>	KM670/671NL	10 (50%)	44–61 (53)	244
<i>APP</i>	V717L, V717G or V717I	11 (40%)	40–67 (51)	245–248
<i>APP</i>	T714I or 714A	6 (85%)	33–55 (NA)	249,250
<i>APP</i>	Duplication	27 (45%)	39–62 (NA)	144,145,251–253
<i>APP; PSEN1</i>	KM670/671NL (<i>APP</i>); H163Y (<i>PSEN1</i>)	8 (89%)	44–65 (54)	254
<i>PSEN1</i>	M146L	7 (70%)	33–46 (39)	255
<i>PSEN1</i>	M146V	1 (100%)	39 (NA)	256
<i>PSEN1</i>	L286V	7 (64%)	39–56 (48)	257

APP, amyloid precursor protein; FAD, familial Alzheimer's disease; NA, not available; *PSEN1*, presenilin-1.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript