

Cigarette Smoking and Minority Stress Across Age Cohorts in a National Sample of Sexual Minorities: Results From the Generations Study

Allegra R. Gordon, ScD, MPH^{1,2,*} · Jessica N. Fish, PhD³ · Wouter J. Kiekens, MSc⁴ · Marguerita Lightfoot, PhD⁵ · David M. Frost, PhD⁶ · Stephen T. Russell, PhD⁷

Published online: 24 September 2020

© Society of Behavioral Medicine 2020. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com.

Abstract

Background Sexual minority populations in the United States have persistently higher rates of cigarette use than heterosexuals, partially driven by exposure to minority stressors (e.g., discrimination and victimization). Little is known about cigarette use across cohorts of sexual minority adults who came of age in distinctly different sociopolitical environments.

Purpose To examine cigarette use and minority stressors across three age cohorts of U.S. sexual minority adults.

Methods We used data from the Generations Study, a nationally representative sample ($N = 1,500$) of White, Black, and Latino/a sexual minority adults in three age cohorts (younger: 18–25 years; middle: 34–41 years; and older: 52–59 years). Survey data were collected from March 2016 to March 2017. We used sex-stratified

logistic regression models to estimate adjusted odds ratios (aORs) and 95% confidence intervals (CIs) for associations between age cohort, minority stressors (discrimination and victimization), and two indicators of cigarette smoking (lifetime use and current use).

Results Prevalence of current cigarette use in each age cohort was high (younger: 20%; middle: 33%; and older: 29%). Relative to the younger cohort, men and women in the middle- and older-age cohorts had significantly higher odds of lifetime and current smoking (e.g., men, current, aOR [95% CI]: middle = 2.47 [1.34, 4.52], older = 2.85 [1.66, 4.93]). Minority stressors were independently associated with higher odds of current smoking; when victimization was included, the magnitude of the association between age cohort and current smoking was diminished but remained significant.

Conclusions Smoking cessation interventions must consider the role of minority stress and the unique needs of sexual minority people across the life course.

✉ Allegra R. Gordon
allegrag@bu.edu

¹ Department of Community Health Sciences, Boston University School of Public Health, Boston, MA 02118, USA

² Division of Adolescent/Young Adult Medicine, Boston Children's Hospital, Boston, MA, USA

³ Department of Family Science, School of Public Health, University of Maryland, College Park, MD, USA

⁴ Department of Sociology/Interuniversity Center for Social Science Theory and Methodology (ICS), University of Groningen, Groningen, The Netherlands

⁵ Department of Medicine, University of California San Francisco, San Francisco, CA, USA

⁶ Department of Social Science, University College London, London, UK

⁷ Department of Human Development and Family Sciences, University of Texas at Austin, Austin, TX, USA

Keywords: Sexual minority · LGB · Cigarette use · Tobacco · Minority stress

Introduction

Tobacco smoking remains the leading cause of preventable death in the USA and a public health priority for addressing and eliminating health inequities [1, 2]. Community and probability samples have documented an elevated risk of cigarette smoking among sexual minority individuals (e.g., lesbian, gay, bisexual, and queer): sexual minorities in the USA have up to 2.5 times higher rates of cigarette smoking than heterosexuals [3, 4]. Notably, public health surveillance data reveal that

sexual orientation disparities in cigarette use persist [5, 6] despite (a) a decades-long decline in the prevalence of cigarette smoking in the USA [1] and (b) recent U.S. social and policy shifts toward greater social acceptance of sexual minorities (e.g., legalization of same-sex marriage) [7]. To address persistent disparities, there is a need to better understand differences in cigarette smoking within sexual minority populations that may drive these disparities.

Despite the recent exponential growth in research on sexual minority health disparities in general, and tobacco use in particular [8], population-based research on sexual minority tobacco use has often been hampered in its ability to explore heterogeneity within sexual minority populations, including differences by gender, sexual orientation identity, and age cohort. Several studies indicate variation by gender, but findings have been mixed. Although some studies have found sexual orientation disparities in cigarette smoking present among both men and women [5], others have found that sexual orientation disparities vary by gender, with greater sexual orientation disparities among girls and women compared to the disparity among boys and men [9–12]. In addition, some studies identify higher rates of cigarette smoking among bisexual compared to gay and lesbian adults [3, 13, 14], with bisexual women particularly at risk [15], although findings have not always been consistent [16, 17].

Minority stress theory posits that minority stress processes are fundamental drivers of sexual orientation disparities [18, 19]. Minority stress processes include exposure to *distal stressors*, such as exposure to discrimination and victimization, which, in turn, influence *proximal stressors*, such as hypervigilance about potential victimization. There is robust evidence that such stressors can trigger a cascade of stress responses, including maladaptive coping behaviors, such as cigarette use [20–22]. These minority stress processes must be considered against a backdrop of decades of targeted marketing to sexual minority communities by the tobacco industry [23–25], which also may take advantage of minority stress-related susceptibility to such targeted marketing [26] and contribute to sexual orientation disparities [27].

Furthermore, different cohorts of sexual minority adults came of age in distinct social and political climates with regards to social acceptance of sexual minority people, accompanying exposures to minority stressors and tobacco-related norms and regulation. Today's sexual minority young adults came of age in the 2000s, an era of increasing cultural inclusion of sexual minority people [7] and regulatory limitations on advertising tobacco to youth [28]. By comparison, sexual minority adults who came of age in the 1970s may have initiated cigarette use in an era when sexual minority visibility was just emerging (e.g., the first gay pride parades)

and tobacco industry regulation remained incipient [29]. Only recently has research begun to examine how tobacco use varies by age, with one recent study finding sexual orientation disparities in tobacco use disorder declining with increasing age [30].

Existing studies have been limited in their ability to examine whether there is variation across different age cohorts of sexual minority adults in their rates of cigarette use and the degree to which cigarette use is associated with minority stressors. Moreover, while several probability samples to date have been able to document cigarette use disparities between sexual minorities and heterosexuals, with one exception [31], most have not been able to examine within-group differences and how these may be related to distal minority stressors. In one notable exception, McCabe et al. found evidence in a nationally representative survey that discrimination attributed to sexual orientation was associated with past-year cigarette smoking among sexual minorities [31] but called for future research to further elucidate within-group differences. A better understanding of such within-group differences, such as those by gender and age cohort, is crucial to advancing efforts to identify and eliminate tobacco-related health disparities [32].

This paper presents data from the first national probability survey designed specifically for sampling U.S. sexual minority adults across three age cohorts. Drawing on minority stress theory [18], we examine current cigarette smoking in this sample and hypothesize that current smoking will vary by gender and sexual orientation with women and bisexual people having elevated odds of cigarette smoking. Furthermore, we hypothesize that lifetime and current cigarette smoking will be more common in the older cohort compared to the younger and middle cohorts and that these associations will be reduced in magnitude when adjusting for exposure to minority stressors.

Methods

Sample and Procedures

The current study includes data from White, Black, and Latino/a men and women in the Generations Study ($N = 1,518$). The Generations Study is a national probability sample of U.S. sexual minorities in three age cohorts, representing distinct sociopolitical environments in which sexual minorities came of age (younger: ages 18–25 years; middle: 34–41 years; and older: 52–59 years). Age cohorts were defined based on when respondents would have experienced key LGBT-related U.S. historical events (e.g., the Stonewall uprising, the emergence of the AIDS epidemic, and the legalization of same-sex marriage) in the course of their adolescent and emerging

adulthood development. For example, those in the oldest cohort were emerging adults during the post-Stonewall era of collective organizing around sexual identities and heightened visibility of sexual minority populations. The middle cohort, in contrast, experienced emerging adulthood during an era when the Internet was changing the landscape of social interactions and access to information, while the younger cohort experienced emerging adulthood in an era when same-sex marriage was newly becoming institutionalized in states across the country. More details about the sample design can be found in Meyer et al. [33] and Frost et al. [34].

Participants were recruited using a two-step process in which Gallup, Inc. collected a national probability sample using a dual-frame random-digit-dial sampling procedure to sample both landlines and mobile phones. Gallup screened all participants in the national probability sample; those who identified as lesbian, gay, bisexual, or transgender received follow-up screening questions. Participants were eligible for the Generations Study if they (a) identified as lesbian, gay, or bisexual, (b) were not transgender, (c) identified their race/ethnicity as Black, Latino/a, or White, (d) were aged 18–25, 34–41, or 52–59 years, (e) completed a sixth-grade education, and (f) answered the phone in English. Respondents who identified during screening as transgender were invited to participate in a separate study on transgender health, *TransPop* (www.transpop.org). Notably, some LGB respondents who identified as cisgender on the initial screener later identified with a gender identity other than male or female (e.g., genderqueer) on the survey itself and were included in analyses as described below. Eligible participants who consented received a self-administered online or mailed survey questionnaire covering a broad range of health behaviors, outcomes, and risk and protective factors. Study procedures were approved by the institutional review board of the University of California, Los Angeles.

Gallup screened a total of 366,644 participants between March 2016 and March 2017. Of these, 3.5% identified as lesbian, gay, or bisexual; 27% of these met eligibility criteria. Of those eligible, 80% agreed to participate in the survey and, of those, 48% completed the survey, for a total conditional participation rate of 39%. To increase the number of racial/ethnic minority respondents, Black and Latino/a respondents were oversampled using the same procedures in an extended recruitment period (April 1, 2017 to March 30, 2018). The final sample of 1,518 included 1,331 participants from the original sample and 187 from the oversample. There was less than 1% missing for all study variables except the discrimination scale (2% missing). We found no significant associations between missingness on the discrimination scale and all other study variables. The present analyses excluded participants who were missing on

self-reported lifetime and current cigarette use ($n = 18$), resulting in an analytic sample of 1,500.

Measures

Cigarette smoking

Outcomes were assessed using two items from the U.S. Behavioral Risk Factor Surveillance System. Participants were instructed not to include the use of electronic cigarettes, other tobacco products, or marijuana in their responses. Any lifetime cigarette smoking was assessed with the question “Have you smoked at least 100 cigarettes in your lifetime?” and coded dichotomously (0 = no cigarette smoking; 1 = any cigarette smoking). Participants were then asked, “Do you now smoke every day, some days, or not at all?”; current cigarette smoking was coded dichotomously following the approach of national surveillance surveys (0 = not at all or never smoked; 1 = currently smoke every day or some days) [6, 35].

Age cohort

Age cohort was assigned based on the date of birth: 18–25 years (younger), 34–41 years (middle), and 52–59 years (older). Cohort parameters were determined based on when respondents would have experienced key LGBT-related U.S. historical events [33, 34] as described above.

Distal minority stressors

Discrimination and victimization: Experiences of *day-to-day discrimination* were captured with a nine-item scale adapted from Williams et al.’s Everyday Discrimination Scale [36], which asked the frequency over the past year of experiences of unfair treatment. Response options ranged from 1 = never to 4 = often (Cronbach’s $\alpha = .91$). *Victimization* experiences in adulthood were assessed using a six-item scale developed by Herek et al. [37] that asks participants to report how often they had experienced verbal or physical violence victimization since they were 18 years old. Response options ranged from 1 = never to 4 = three or more times (Cronbach’s $\alpha = .82$). Both of these measures ask broadly about any unfair treatment or victimization, respectively, without requiring participants to attribute the discrimination to a particular characteristic (e.g., race/ethnicity or sexual orientation); we elected to use these measures of minority stress without attribution based on previous research suggesting that using such attribution may lead to underestimates of discriminatory experiences [38, 39]. For both the discrimination and victimization scales, frequency scores were summed to create continuous variables, with higher scores indicating more frequent discrimination in the previous year or

victimization since age 18, respectively. The survey did not include lifetime measures of discrimination and victimization; thus, these variables were not included in analyses related to lifetime history of smoking.

Sociodemographic covariates

Sexual orientation identity was assessed with a widely used measure [40] with response options: straight/heterosexual, lesbian, gay, bisexual, queer, same-gender loving, and other. These were coded as three categories: lesbian/gay, bisexual, and another sexual orientation; by design, those who identified as straight/heterosexual were not included in the Generations Study. *Race/ethnicity*: Eligible participants for the Generations Study were those who identified as Black or African American, Latino/a or Hispanic, or White; these were the three categories used for analysis. *Assigned sex* (female/woman or male/man) was the primary stratification variable. As all participants screened into this sample by identifying as nontransgender (i.e., reported their gender aligned with their assigned sex), in this paper, we refer to differences between women and men. However, in the main survey, some participants also identified with a *nonbinary gender identity* (e.g., genderqueer); this was included in models as a dichotomous variable (nonbinary or binary identity). *Annual household income* was collected using 12 categorical responses ranging from “Under \$720” to “\$240,000 and over.” Average values were calculated for each household income range (\$720 and \$240,000 representing the lowest and highest values, respectively). These household income estimates were adjusted for household size and scaled to represent three-person households based on the 2016 U.S. median household income (\$57,617) [41] following the Pew Research Center’s approach [42]. U.S. median household income from 2016 was selected to match the Generations Study data collection period. *Urbanicity* was measured based on participants’ residential zip codes and classified using the U.S. Department of Agriculture rural–urban commuting area (RUCA) coding system (coded 1–10, with each delineating a degree of urbanicity/rurality from major metropolitan areas to rural areas; higher numbers indicate greater rurality/lower commuting flow into the area) [43].

Statistical Analysis

Analyses were conducted from July 2018 to February 2019 with Stata 15.1 and weighted to produce nationally representative estimates of the target population. Base weights were first calculated for the Gallup sample frame for the timeframe included in this study (2016–2018) in multiple stages. The sample was initially weighted to represent the aged 18+ U.S. population; the weighting process then accounted for multiple stages of selection

and nonresponse (for more on the Generations Study methods, see www.generations-study.com/methods). This resulted in weights that allow estimates to be generalizable to the U.S. population of sexual minority adults aged 18–25, 34–41, and 52–59 during data collection.

Our analyses first examined cigarette smoking prevalence and bivariate associations among demographic characteristics, minority stressors, and cigarette smoking. We used stepwise logistic regression models to estimate adjusted odds ratios (aORs) and 95% confidence intervals (95% CIs) for associations between age cohort, minority stressors, and both cigarette smoking outcomes. Based on prior research indicating differences between women and men in sexual orientation disparities in cigarette use [9–11], we ran multivariable analyses separately by assigned sex. Models were adjusted for nonbinary gender, race/ethnicity, sexual orientation identity, scaled household income, and RUCA score. For current cigarette smoking only, Model 1 was adjusted for demographic characteristics and we then added the minority stressors to the model (Model 2) to examine the extent to which observed age cohort associations with cigarette smoking were affected by the inclusion of minority stressors. Minority stressors were not included in models for lifetime cigarette smoking due to the lack of temporally appropriate measures (i.e., participants were only asked to report exposure to minority stressors in the previous year; however, lifetime cigarette smokers may not have smoked in the previous year). Sensitivity analyses were conducted to examine the independent effects of each minority stressor in the model. Effect estimates were similar, so we present models with both variables included. Multiple imputation was used to account for missing data on covariates with 50 imputation draws using *mi estimate* in Stata.

Results

Weighted estimates indicated that 46.9% of the sample identified as gay/lesbian, 40.5% as bisexual, and 12.6% as another sexual orientation identity (Table 1). The weighted prevalence of lifetime smoking in the sample was 41.8%, while the prevalence of current cigarette smoking was 24.6%. As displayed in Table 1, both lifetime and current cigarette smoking prevalence varied across age cohorts at the bivariate level, with the majority of those in the older and middle cohorts (62.5% and 58.8%, respectively) reporting a lifetime history of smoking compared to less than one third of those in the younger cohort (30.4%; $p < .0001$). Prevalence of current smoking was highest among those in the middle cohort (33.2%), followed by the older (29.4%) and younger (20.3%) cohorts ($p < .0001$).

Table 1. Sample characteristics by age cohort^a in a probability sample of sexual minorities ($N = 1,500$)

	Total			Younger ($n = 665$)			Middle ($n = 367$)			Older ($n = 468$)			<i>p</i> -value*
	<i>n</i>	% _W	(95% CI)	<i>n</i>	% _W	(95% CI)	<i>n</i>	% _W	(95% CI)	<i>n</i>	% _W	(95% CI)	
Gender													
Women	804	59.9	(56.8, 62.9)	394	65.4	(61.2, 69.4)	203	58.8	(52.7, 64.7)	207	41.4	(36.4, 46.6)	<.0001
Men	696	40.1	(37.1, 43.2)	271	34.6	(30.6, 39.8)	164	41.1	(35.3, 47.3)	261	58.6	(53.4, 63.6)	<.0001
Nonbinary gender identity													
Binary gender identity	1,406	92.5	(90.5, 94.1)	604	90.1	(87.0, 92.5)	350	96.4	(94.2, 97.8)	452	96.4	(93.4, 97.9)	<.0001
Nonbinary/genderqueer	94	7.5	(5.9, 9.5)	61	9.9	(7.5, 13.0)	17	3.6	(2.2, 5.8)	16	3.6	(2.1, 6.2)	<.0001
Sexual orientation identity													
Gay/lesbian	821	46.9	(43.8, 50.1)	244	36.6	(32.5, 41.0)	201	50.3	(44.2, 56.6)	376	79.7	(75.1, 83.6)	
Bisexual	489	40.5	(37.3, 43.7)	300	47.7	(43.2, 52.1)	123	40.0	(34.0, 46.3)	66	15.2	(11.7, 19.4)	
Another sexual orientation	179	12.6	(10.7, 14.9)	115	15.7	(12.9, 19.0)	40	9.7	(6.9, 13.3)	24	5.1	(3.3, 8.0)	
Race/ethnicity													
White	970	62.2	(59.1, 65.2)	362	56.4	(51.2, 60.7)	231	64.9	(59.0, 70.5)	377	79.6	(74.9, 83.6)	<.0001
Black or African American	235	16.5	(14.3, 18.9)	125	17.9	(14.9, 21.4)	68	18.0	(13.8, 23.0)	42	9.6	(6.9, 13.4)	
Hispanic or Latino/a	295	21.3	(18.9, 24.0)	178	25.7	(22.1, 29.6)	68	17.1	(13.1, 22.0)	49	10.7	(7.8, 14.6)	
Household income^b													
Lower income	467	39.6	(36.5, 42.8)	281	45.9	(41.5, 50.4)	90	33.7	(27.7, 40.2)	96	24.2	(19.8, 29.4)	<.0001
Middle income	551	35.5	(32.6, 38.6)	273	37.5	(33.3, 41.8)	133	32.6	(27.3, 38.4)	146	31.9	(27.3, 36.9)	
Upper income	481	24.9	(22.4, 27.5)	111	16.6	(13.6, 20.2)	144	33.7	(28.4, 39.5)	226	43.8	(38.8, 49.0)	
Outcomes													
Lifetime smoking ^c	669	41.8	(38.8, 45.0)	190	30.4	(26.4, 34.7)	200	58.8	(52.7, 64.6)	279	62.5	(57.4, 67.2)	<.0001
Current smoking ^d	333	24.6	(21.9, 27.5)	120	20.3	(16.9, 24.3)	100	33.2	(27.4, 39.6)	113	29.4	(24.5, 34.7)	<.0001

Table 1. Continued

	Total			Younger (n = 665)			Middle (n = 367)			Older (n = 468)			p-value*
	n	%w	(95% CI)	n	%w	(95% CI)	n	%w	(95% CI)	n	%w	(95% CI)	
Urbanicity	M	(SD)		M	(SD)		M	(SD)		M	(SD)		
RUCA score ^e	1.72	(1.87)		1.75	(1.60)		1.58	(1.83)		1.76	(2.63)		.93
Minority stressors													
Day-to-day discrimination ^f	2.04	(0.73)		2.16	(0.61)		2.00	(0.78)		1.63	(0.78)		<.0001
Victimization ^g	1.97	(0.84)		1.85	(0.68)		2.22	(0.94)		2.11	(1.13)		<.0001

CI confidence interval.

^aAge cohorts defined as: younger = 18–25 years; middle = 34–41 years; older = 52–59 years.

^bIncome: Based on annual household (HH) income, adjusted for household size, and scaled in relation to median U.S. household size (per Pew Research Center 2015). Lower income = <2/3 U.S. median HH income; middle income = 2/3 – double U.S. median HH income; Upper income = >double U.S. median HH income.

^cRespondents were asked “Have you smoked at least 100 cigarettes in your entire life?” (Yes/No). Respondents were asked to *exclude* any use of e-cigarettes, cigars, or other tobacco products.

^dRespondents who endorsed smoking at least 100 cigarettes were asked “Do you now smoke cigarettes every day, some days, or not at all?” Responses were dichotomized: every day or some days = current smoker; not at all or never smoked = not current smoker.

^eRUCA = rural–urban commuting area (score range: 1–10). U.S. Department of Agriculture-created system to delineate degree of urbanicity/rurality based on level of commuting flow into an area; higher scores indicate greater rurality/lower commuting flow into area.

^fVictimization: mean score of six items about frequency of victimization experiences since age 18 (range: 1–4; higher scores = more victimization experiences).

^gDay-to-day discrimination: mean score of nine items about frequency of unfair treatment in a variety of settings in past year (range: 1–4; higher scores = more experiences of discrimination).

*Boldface indicates statistical significance ($p < .0001$). p-values are for differences across generations. p-values based on chi-square tests for demographic characteristics and categorical outcomes based on analysis of variance F-tests for continuous predictor variables.

Table 2. Prevalence of lifetime and current cigarette smoking by sociodemographic characteristics among U.S. sexual minorities ($N = 1,500$)

	Lifetime cigarette smoking ^a			Current cigarette smoking ^b		
	% (wtd)	95% CI	<i>p</i> -value	% (wtd)	95% CI	<i>p</i> -value
Gender			.87			.70
Women	42.0	(37.8, 46.3)		25.0	(21.4, 29.1)	
Men	41.5	(37.1, 46.0)		23.9	(20.2, 28.1)	
Nonbinary gender identity			.08			.29
Binary gender identity	42.6	(39.4, 45.9)		25.0	(22.2, 28.1)	
Genderqueer/nonbinary	32.0	(22.3, 43.6)		19.0	(11.3, 30.4)	
Sexual orientation identity			.70			.62
Gay/lesbian	42.7	(38.5, 47.0)		24.7	(21.1, 28.7)	
Bisexual	41.6	(36.4, 47.0)		24.8	(20.3, 29.9)	
Another sexual orientation	38.5	(30.5, 47.1)		20.6	(14.4, 28.7)	
Race/ethnicity			.02			.64
White	45.2	(41.2, 49.2)		25.4	(22.0, 29.2)	
Black or African American	35.2	(28.4, 42.7)		24.5	(18.6, 31.6)	
Hispanic or Latino/a	37.1	(30.9, 43.8)		22.2	(16.9, 28.5)	
Socioeconomic position ^c			<.0001			<.0001
Lower income	50.7	(45.3, 56.1)		36.1	(31.0, 41.6)	
Middle income	36.7	(32.0, 41.6)		19.5	(15.8, 23.8)	
Higher income	35.1	(30.1, 40.4)		13.5	(10.2, 17.6)	
Minority stressors	<i>M</i>	(<i>SD</i>)	<i>p</i> -value	<i>M</i>	(<i>SD</i>)	<i>p</i> -value
Everyday discrimination ^d			.18			.003
No to smoking	2.01	(0.03)		1.99	(0.72)	
Yes to smoking	2.07	(0.04)		2.17	(0.74)	
Victimization ^e			<.0001			<.0001
No to smoking	1.76	(0.72)		1.85	(0.79)	
Yes to smoking	2.26	(0.92)		2.36	(0.87)	

Boldface indicates statistical significance ($p < .05$). *p*-values are for differences within each demographic category. *p*-values based on chi-square tests for demographic characteristics and categorical outcomes and on analysis of variance *F*-tests for continuous variables. *CI* confidence interval.

^aRespondents were asked “Have you smoked at least 100 cigarettes in your entire life?” (Yes/No). Respondents were asked to *exclude* any use of e-cigarettes, cigars, or other tobacco products.

^bRespondents who endorsed smoking at least 100 cigarettes were asked “Do you now smoke cigarettes every day, some days, or not at all?” Responses were dichotomized: every day or some days = current smoker; not at all or never smoked = not current smoker.

^cSocioeconomic position: based on annual household (HH) income, adjusted for household size, and scaled in relation to median U.S. household size (per Pew Research Center 2015). Lower income = <2/3 U.S. median HH income; middle income = 2/3 – double U.S. median HH income; upper income = >double U.S. median HH income.

^dVictimization: mean score of six items about frequency of victimization experiences since age 18 (range: 1–4; higher scores = more victimization experiences).

^eDay-to-day discrimination: mean score of nine items about frequency of unfair treatment in a variety of settings in past year (range: 1–4; higher scores = more experiences of discrimination).

Lifetime and current cigarette smoking varied across demographic characteristics and exposure to minority stressors (see Table 2). For example, lifetime smoking prevalence was highest among White participants (45.2%) followed by Latino/a (37.1%) and Black (35.2%) participants ($p < .02$); however, no statistical differences were observed in current cigarette smoking.

Participants in the lower-income group had strikingly elevated prevalence of both lifetime and current cigarette smoking compared to those in the middle- and higher-income groups. Current smokers had higher mean levels of day-to-day discrimination ($M_s = 2.2$ vs. 2.0, $p = .003$) and victimization ($M_s = 2.4$ vs. 1.8, $p < .0001$).

Table 3. Multivariable models predicting any lifetime history of smoking^a in a sample of sexual minorities ($N = 1,500$)

	Women		Men	
	OR	(95% CI)	OR	(95% CI)
Age cohort				
Younger (Ref)	1.00		1.00	
Middle	5.56	(3.62, 8.54)	2.81	(1.63, 4.84)
Older	6.21	(3.70, 10.42)	4.96	(2.94, 8.34)
Sexual orientation identity				
Gay/lesbian (Ref)	1.00		1.00	
Bisexual	1.06	(0.68, 1.68)	1.20	(0.73, 1.97)
Another sexual orientation	1.40	(0.72, 2.72)	1.61	(0.65, 3.94)
Binary gender				
Binary woman/man (Ref)	1.00		1.00	
Nonbinary/genderqueer	0.56	(0.26, 1.20)	0.87	(0.34, 2.23)
Race/ethnicity				
White (Ref)	1.00		1.00	
Black or African American	0.50	(0.30, 0.84)	0.86	(0.46, 1.61)
Hispanic or Latino/a	0.67	(0.40, 1.12)	1.21	(0.73, 1.99)
Income				
Upper income (Ref)	1.00		1.00	
Middle income	1.68	(1.02, 2.79)	1.56	(0.99, 2.53)
Low income	4.29	(2.51, 7.33)	2.67	(1.51, 4.75)
Urbanicity				
RUCA Score	1.06	(0.97, 1.17)	1.15	(1.02, 1.30)

Boldface indicates statistical significance ($p < .05$). Models estimated with weights to be nationally representative and 50 imputations to account for missing data on covariates.

CI confidence interval; *OR* odds ratio; *RUCA* rural–urban commuting area.

^aLifetime smoking defined as report of ever having smoked at least 100 cigarettes (five packs).

In adjusted multivariable logistic regression models predicting any lifetime cigarette smoking (Table 3), women in the older cohort had over six times higher odds and those in the middle cohort had over five times higher odds of lifetime smoking relative to women in the younger cohort. Men in the older cohort had five times higher odds and those in the middle cohort had nearly three times higher odds of lifetime smoking relative to men in the younger cohort. No differences were observed in lifetime smoking history across sexual orientation, although variation was observed by race/ethnicity for women and socioeconomic position for both men and women (see Table 3).

In multivariable models predicting any current cigarette smoking, odds of current smoking were 2.4 times higher for women in the middle cohort relative to the younger cohort (Table 4; Model 1). Among men, patterns were similar, with significant elevated odds for men in both the middle and older cohorts relative to the younger cohort. When minority stressors were included in models, the observed associations between age cohort and smoking status were modestly lower (Table 4; Model

2). Among women, although day-to-day discrimination was associated with smoking status when added independently (not shown), once the victimization scale was added to the model, the effect estimate for day-to-day discrimination became nonsignificant and only victimization remained significant. Among men, when minority stressors were added to the model, there was a minor decrease in the magnitude of the association between age cohort and smoking status; the aORs for the minority stressors had CIs that crossed the null.

Discussion

Findings from this analysis of a national probability sample of sexual minority adults across three age cohorts offer further evidence of the persistence of longstanding sexual orientation disparities in cigarette use. Across the sample, nearly one in four (24.6%) reported current cigarette smoking compared to 15.5% of U.S. adults in the general population in the same year [6]. The prevalence was also higher in each age cohort compared to similar

Table 4. Multivariable models predicting current cigarette smoking^a in a sample of sexual minorities ($N = 1,500$)

	Women				Men			
	Model 1		Model 2		Model 1		Model 2	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Age cohort								
Younger (Ref)	1.00		1.00		1.00		1.00	
Middle	2.38	(1.45, 3.89)	1.64	(0.94, 2.84)	2.47	(1.34, 4.52)	2.23	(1.17, 4.28)
Older	1.59	(0.90, 2.82)	1.27	(0.66, 2.44)	2.85	(1.66, 4.93)	2.63	(1.44, 4.81)
Sexual orientation								
Gay/lesbian (Ref)	1.00		1.00		1.00		1.00	
Bisexual	0.77	(0.46, 1.27)	0.60	(0.36, 1.02)	1.09	(0.60, 1.96)	1.03	(0.56, 1.93)
Another sexual orientation	0.81	(0.40, 1.64)	0.73	(0.36, 1.50)	1.47	(0.56, 3.83)	1.41	(0.54, 3.74)
Binary gender								
Binary woman/man (Ref)	1.00		1.00		1.00		1.00	
Nonbinary/genderqueer	0.62	(0.25, 1.56)	0.47	(0.17, 1.23)	1.04	(0.35, 3.11)	1.02	(0.33, 3.13)
Race/ethnicity								
White (Ref)	1.00		1.00		1.00		1.00	
Black or African American	0.63	(0.36, 1.09)	0.64	(0.36, 1.16)	1.16	(0.60, 2.25)	1.12	(0.55, 2.30)
Hispanic or Latino/a	0.68	(0.37, 1.24)	0.74	(0.38, 1.43)	1.17	(0.64, 2.13)	1.19	(0.64, 2.21)
Income								
Upper income (Ref)	1.00		1.00		1.00		1.00	
Middle income	1.78	(0.90, 3.52)	1.55	(0.80, 3.00)	2.25	(1.33, 3.81)	2.07	(1.20, 3.56)
Low income	4.97	(2.52, 9.77)	3.81	(1.96, 7.43)	4.72	(2.56, 8.69)	3.96	(2.10, 7.46)
Urbanicity (<i>RUCA</i> score)	1.08	(0.98, 1.20)	1.10	(0.99, 1.22)	1.16	(1.02, 1.32)	1.16	(1.02, 1.30)
Day-to-day discrimination			1.08	(0.73, 1.62)			1.06	(0.68, 1.65)
Victimization			2.15	(1.59, 2.91)			1.31	(0.95, 1.80)

Boldface indicates statistical significance ($p < .05$). Models estimated with weights to be nationally representative and 50 imputations to account for missing data on covariates. Model 1 includes key demographic covariates. Model 2 includes the addition of distal minority stressors (day-to-day discrimination in the past year; victimization since age 18).

CI confidence interval; *OR* odds ratio; *RUCA* rural–urban commuting area.

^aCurrent smoking coded as: 0 = not at all or never smoked; 1 = smoke some days or every day.

(albeit not identical) age groups of U.S. adults: the 2016 National Health Interview Survey (NHIS) found that 14.7% of U.S. adults 18–24 years old, 20.6% of those 25–44 years old, and 19.3% of U.S. adults 45–64 years old were current smokers [6] (compared to our sample in which 20.3%, 33.2%, and 29.4% of sexual minorities in our younger-, middle-, and older-age cohorts, respectively, were current smokers). Our prevalence estimates are higher than NHIS prevalence estimates for sexual minority adults (20.5% [6]), possibly due to our restricted age range (NHIS lowest cigarette prevalence was among adults aged 65 years and older; the Generations Study only included those under 60 years).

Our study offers novel insights into differences in both lifetime and current cigarette smoking across three distinct age cohorts of sexual minority people (i.e., “age effects”). Although a cross-sectional study like this one is

not able to disentangle cohort effects or period effects from age effects, it is important to note that, by design, the three age cohorts in this study represent sexual minority people who came of age in distinctly different social and political environments with regard to sexual minority rights and societal acceptance. The majority of sexual minority adults in the middle and older cohorts reported ever having smoked cigarettes, as would be anticipated given that older cohorts have more life years in which to have tried smoking and were more likely to have initiated cigarette use during periods when smoking was normative in the USA [29]. However, even in the younger cohort (18–25 years in 2016), one in three reported a history of cigarette smoking. These prevalence estimates reflect well-documented [3, 11, 44] heightened risk of cigarette smoking in sexual minority compared to heterosexual populations. Notably, the gap between lifetime

and current cigarette use could indicate that cigarette smoking cessation has been successful for a substantial proportion of sexual minorities who were smokers. The gap between lifetime and current smoking was greatest in the older cohort but, even in the younger cohort, about 10% reported a lifetime history but no current smoking.

We also observed that age cohort differences in lifetime cigarette use were greater magnitude among sexual minority women, relative to sexual minority men, with older women having higher odds of smoking relative to younger women than the gap among men. This suggests a need to think not only about smoking cessation interventions targeted at sexual minority populations broadly [45] but also about targeting by gender. Given that cigarette use is a key risk factor for multiple cancers, practical implications of these gender differences underscore the need to ensure that cancer screening initiatives are effectively targeted to sexual minority women in general, and older sexual minority women in particular. Extensive research has demonstrated a range of stigma-related barriers to health care, including cancer screenings, for sexual minority women [46, 47]. Reducing stigma-related barriers to routine screening and health care could increase both cancer screening and access to health care provider-delivered smoking cessation interventions.

Notably, in this probability sample of sexual minorities, there were no differences in cigarette smoking among sexual minority identity subgroups in contrast to some previous research and national surveillance surveys that have found an elevated prevalence of current cigarette use among bisexual compared to lesbian/gay populations [13, 14], and particularly among adolescent girls and young women [9, 10, 17, 48]. Although several prior studies show sexual orientation disparities for bisexual relative to lesbian/gay adults, differences between these two groups are typically not statistically tested. These findings underscore the importance of continuing to explore heterogeneity within sexual minority populations to better understand variation in risk factors and underlying drivers of cigarette use in these vulnerable populations. For example, although not a focus of this analysis, we did observe notable differences by household income, with lower-income sexual minority people at the greatest risk of cigarette use. This finding aligns with a robust body of research on cigarette use and socioeconomic position [49] and extends this research to demonstrate the same patterns among sexual minorities. Future research should explore opportunities for tailored smoking cessation interventions for lower-income sexual minority populations.

Previous research on the role of distal minority stressors in cigarette smoking has indicated that discrimination and victimization are associated with cigarette smoking, particularly in adolescents; research in adult samples has been less conclusive [4, 50]. One recent study

with a nationally representative sample of adults observed associations between self-reported discrimination due to sexual orientation and past-year cigarette smoking [31]. Our findings support these findings and extend them by documenting the association between current smoking and overall self-reported discrimination, not just that which participants believe was linked to their sexual orientation. As has been discussed extensively by scholars of discrimination [39, 51], both approaches (i.e., discrimination with and without attribution to specific targets) are necessary for improving our understanding of the health consequences of discrimination. Our findings in concert with previous research affirm the importance of including distal minority stressors, such as discrimination and victimization, whether attributed to sexual orientation or not, in research on cigarette smoking disparities.

Additionally, in our analysis of the role that minority stressors may play in observed cohort differences in cigarette use within this sexual minority sample, we found that both day-to-day discrimination and victimization experiences were associated with current cigarette use at the bivariate level. In multivariable models, victimization partially contributed to the associations between age cohort and current cigarette smoking, particularly among women. This suggests that minority stressors may be one set of factors involved in the persistence of higher smoking prevalence among older sexual minorities, creating barriers to smoking cessation or exacerbating the need for cigarette use as a coping mechanism for recent or past experiences of discrimination.

Limitations

Findings must be interpreted in light of several limitations. Data are cross-sectional and based on self-report; we hypothesized that minority stressors may mediate the association between age cohort and cigarette smoking but, as with all survey research, we cannot establish temporal ordering and reverse causation remains a possibility (e.g., smokers may spend time in more public environments than nonsmokers and, thus, be more likely to be exposed to discrimination or be victimized). Furthermore, we did not have data on lifetime exposure to discrimination and victimization; thus, we were not able to include the role of minority stressors in the models looking at age cohort differences in lifetime cigarette smoking. Another key limitation is that our survey did not include questions on e-cigarette use/vaping; thus, we cannot know how much of the difference between the younger and older cohorts could be made up for by higher rates of e-cigarette use in the younger cohort. This form of tobacco use has been rising dramatically in recent years, especially among adolescents and young adults [52, 53]. Emerging evidence

suggests that e-cigarette prevalence may be disproportionately high among sexual minority adults relative to heterosexuals [53, 54], although findings are inconsistent [55] and more research on this topic is needed. These analyses also were not able to address and, thus, leave open some important questions for future research to address—particularly the entangled role of both minority stress experiences *and* the continued insidiousness of targeted tobacco industry marketing to sexual and gender minority youth and adults [56].

Conclusion

This study offers an important picture of cigarette smoking in a U.S. national probability sample of sexual minorities in three age cohorts. Our findings urge public health practitioners and health care providers to consider how the distinctive experiences of sexual minorities of different age cohorts may influence health-related behaviors, with important implications for smoking prevention and cessation interventions. There have been notable steps forward in recent years to produce smoking prevention campaigns tailored to sexual and gender minority youth (e.g., the U.S. Food and Drug Administration's This Free Life campaign [57]). However, given the elevated prevalence of cigarette use in the younger cohort in this study, compared to the general population prevalence for U.S. young adults, and given the rise in e-cigarette use and vaping in the USA, both tailored and universal prevention measures (e.g., regulations) will continue to be essential.

Smoking cessation interventions should consider the role of minority stress and the unique needs of sexual minorities across the life course. In particular, smoking cessation interventionists should consider ways to address unique generational experiences for sexual minority populations, including both current exposure to minority stressors and histories of exposure to anti-LGBT bias. Recognizing both shared experiences of minority stressors among sexual minorities, as well as variation in experiences and health-related behaviors linked to an array of intersecting identities and histories, will be essential in advancing health equity.

Acknowledgments:

Funding: A.R.G. acknowledges support by the National Institute on Drug Abuse (1F32DA042506). J.N.F. acknowledges support by the University of Maryland Prevention Research Center cooperative agreement U48DP006382 from the Centers for Disease Control and Prevention and by the Eunice Kennedy Shriver National Institute of Child Health and Human Development grant P2CHD041041 (Maryland Population Research Center). S.T.R. acknowledges support from grant P2CHD042849 awarded to the Population Research Center at The University of Texas at

Austin by the Eunice Kennedy Shriver National Institute of Child Health and Human Development and from the Priscilla Pond Flawn Endowment at the University of Texas at Austin. Generations Study is funded by a grant from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD grant 1R01HD078526) and through supplemental grants from the National Institutes of Health, Office of Behavioral and Social Sciences Research, and the Office of Research on Women's Health. The Generations investigators are Ilan H. Meyer, PhD (PI), D.M.F., PhD, Phillip L. Hammack, PhD, M.L., PhD, S.T.R., PhD, and Bianca D. M. Wilson, PhD (co-investigators, listed alphabetically). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or Centers for Disease Control and Prevention. Thank you to Rose Eiduson, Harvard SOGIE Collaborative, and all Generations Study participants.

Compliance with Ethical Standards

Authors' Statement of Conflict of Interest and Adherence to Ethical Standards The authors declare that they have no conflicts of interest.

Authors' Contributions A.R.G. conceptualized and designed the study, oversaw the analysis, and led the writing of the manuscript. J.N.F. assisted in study design, carried out quantitative analyses, drafted sections of the manuscript, and critically reviewed the manuscript. W.J.K. assisted in study design, carried out quantitative analyses, and critically reviewed the manuscript. M.L., D.M.F., and S.T.R. assisted in study design and data interpretation and critically reviewed the manuscript.

Ethical Approval The study had IRB approval.

Informed Consent Informed consent was obtained from all participants in the Generations Study.

References

1. U.S. Department of Health and Human Services. *The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General*. Atlanta, GA: Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2014. Available at <https://www.surgeongeneral.gov/library/reports/50-years-of-progress/index.html>. Accessibility verified March 6, 2019.
2. Simmons VN, Pineiro B, Hooper MW, Gray JE, Brandon TH. Tobacco-related health disparities across the cancer care continuum. *Cancer Control*. 2016;23:434–441.
3. Lee JG, Griffin GK, Melvin CL. Tobacco use among sexual minorities in the USA, 1987 to May 2007: A systematic review. *Tob Control*. 2009;18:275–282.
4. Blossnich J, Lee JG, Horn K. A systematic review of the aetiology of tobacco disparities for sexual minorities. *Tob Control*. 2013;22:66–73.
5. Johnson SE, Holder-Hayes E, Tessman GK, King BA, Alexander T, Zhao X. Tobacco product use among sexual minority adults: Findings from the 2012–2013 national adult tobacco survey. *Am J Prev Med*. 2016;50:e91–e100.
6. Jamal A, Phillips E, Gentzke AS, et al. Current cigarette smoking among adults—United States, 2016. *MMWR Morb Mortal Wkly Rep*. 2018;67:53–59.

7. McCarthy J. *Two in Three Americans Support Same-Sex Marriage*. Washington, DC: Gallup, Inc.; 2018. Available at <https://news.gallup.com/poll/234866/two-three-americans-support-sex-marriage.aspx>. Accessibility verified July 19, 2019.
8. Cochran SD, Mays VM. Advancing the LGBT health research agenda: Differential health trends within the lesbian, gay, and bisexual populations. *Am J Public Health*. 2017;107:497–498.
9. Dai H. Tobacco product use among lesbian, gay, and bisexual adolescents. *Pediatrics*. 2017;139(4):e20163276.
10. Li J, Haardörfer R, Vu M, Windle M, Berg CJ. Sex and sexual orientation in relation to tobacco use among young adult college students in the US: A cross-sectional study. *BMC Public Health*. 2018;18:1244.
11. Corliss HL, Wadler BM, Jun HJ, et al. Sexual-orientation disparities in cigarette smoking in a longitudinal cohort study of adolescents. *Nicotine Tob Res*. 2013;15:213–222.
12. Vu M, Li J, Haardörfer R, Windle M, Berg CJ. Mental health and substance use among women and men at the intersections of identities and experiences of discrimination: Insights from the intersectionality framework. *BMC Public Health*. 2019;19:108.
13. Matthews AK, Steffen A, Hughes T, Aranda F, Martin K. Demographic, healthcare, and contextual factors associated with smoking status among sexual minority women. *LGBT Health*. 2017;4:17–23.
14. Fallin A, Goodin A, Lee YO, Bennett K. Smoking characteristics among lesbian, gay, and bisexual adults. *Prev Med*. 2015;74:123–130.
15. Schuler MS, Collins RL. Sexual minority substance use disparities: Bisexual women at elevated risk relative to other sexual minority groups. *Drug Alcohol Depend*. 2020;206:107755.
16. Lunn MR, Cui W, Zack MM, Thompson WW, Blank MB, Yehia BR. Sociodemographic characteristics and health outcomes among lesbian, gay, and bisexual U.S. adults using healthy people 2020 leading health indicators. *LGBT Health*. 2017;4:283–294.
17. Delahanty J, Ganz O, Hoffman L, Guillory J, Crankshaw E, Farrelly M. Tobacco use among lesbian, gay, bisexual and transgender young adults varies by sexual and gender identity. *Drug Alcohol Depend*. 2019;201:161–170.
18. Meyer IH. Prejudice as stress: Conceptual and measurement problems. *Am J Public Health*. 2003;93:262–265.
19. Hatzenbuehler ML. How does sexual minority stigma “get under the skin”? A psychological mediation framework. *Psychol Bull*. 2009;135:707–730.
20. Gruskin EP, Byrne KM, Altschuler A, Dibble SL. Smoking it all away: Influences of stress, negative emotions, and stigma on lesbian tobacco use. *J LGBT Health Res*. 2008;4:167–179.
21. D’Avanzo PA, Halkitis PN, Yu K, Kapadia F. Demographic, mental health, behavioral, and psychosocial factors associated with cigarette smoking status among young men who have sex with men: The P18 cohort study. *LGBT Health*. 2016;3:379–386.
22. Blosnich JR, Horn K. Associations of discrimination and violence with smoking among emerging adults: Differences by gender and sexual orientation. *Nicotine Tob Res*. 2011;13:1284–1295.
23. Goebel K. Lesbian and gays face tobacco targeting. *Tob Control*. 1994;3(1):65.
24. Smith EA, Thomson K, Offen N, Malone RE. “If you know you exist, it’s just marketing poison”: Meanings of tobacco industry targeting in the lesbian, gay, bisexual, and transgender community. *Am J Public Health*. 2008;98:996–1003.
25. Emory K, Buchting FO, Trinidad DR, Vera L, Emery SL. Lesbian, gay, bisexual, and transgender (LGBT) view it differently than non-LGBT: Exposure to tobacco-related couponing, e-cigarette advertisements, and anti-tobacco messages on social and traditional media. *Nicotine Tob Res*. 2019;21:513–522.
26. Remafedi G. Lesbian, gay, bisexual, and transgender youths: Who smokes, and why? *Nicotine Tob Res*. 2007;9(suppl 1):S65–S71.
27. Amroussia N, Pearson JL, Gustafsson PE. What drives us apart? Decomposing intersectional inequalities in cigarette smoking by education and sexual orientation among U.S. adults. *Int J Equity Health*. 2019;18:109.
28. Husten CG, Deyton LR. Understanding the Tobacco Control Act: Efforts by the US Food and drug administration to make tobacco-related morbidity and mortality part of the USA’s past, not its future. *Lancet*. 2013;381:1570–1580.
29. Cummings KM, Proctor RN. The changing public image of smoking in the United States: 1964–2014. *Cancer Epidemiol Biomarkers Prev*. 2014;23(1):32–36.
30. Rice CE, Vasilenko SA, Fish JN, Lanza ST. Sexual minority health disparities: An examination of age-related trends across adulthood in a national cross-sectional sample. *Ann Epidemiol*. 2019;31:20–25.
31. McCabe SE, Hughes TL, Matthews AK, et al. Sexual orientation discrimination and tobacco use disparities in the United States. *Nicotine Tob Res*. 2019;21:523–531.
32. Schwartz S, Meyer IH. Mental health disparities research: The impact of within and between group analyses on tests of social stress hypotheses. *Soc Sci Med*. 2010;70:1111–1118.
33. Meyer IH, Marken S, Russell ST, Frost DM, Wilson BDM. An innovative approach to the design of a national probability sample of sexual minority adults. *LGBT Health*. 2020;7:101–108.
34. Frost DM, Hammack PL, Wilson BDM, Russell ST, Lightfoot M, Meyer IH. The qualitative interview in psychology and the study of social change: Sexual identity development, minority stress, and health in the generations study. *Qual Psychol*. 2019. doi:10.1037/qup0000148.
35. Fleary SA, Nigg CR. Trends in health behavior patterns among U.S. adults, 2003–2015. *Ann Behav Med*. 2019;53:1–15.
36. Williams DR, Yu Y, Jackson JS, Anderson NB. Racial differences in physical and mental health: Socio-economic status, stress and discrimination. *J Health Psychol*. 1997;2:335–351.
37. Herek GM. Hate crimes and stigma-related experiences among sexual minority adults in the United States: Prevalence estimates from a national probability sample. *J Interpers Violence*. 2009;24:54–74.
38. Williams DR, Mohammed SA. Discrimination and racial disparities in health: Evidence and needed research. *J Behav Med*. 2009;32:20–47.
39. Shariff-Marco S, Breen N, Landrine H, et al. Measuring everyday racial/ethnic discrimination in health surveys: How best to ask the questions, in one or two stages, across multiple racial/ethnic groups? *Du Bois Rev*. 2011;8:159–177.
40. Sexual Minority Assessment Research Team. *Best Practices for Asking Questions About Sexual Orientation on Surveys*. Los Angeles, CA: The Williams Institute; 2009. Available at <https://williamsinstitute.law.ucla.edu/wp-content/uploads/Best-Practices-SO-Surveys-Nov-2009.pdf>. Accessibility verified September 13, 2020.
41. Guzman GG. *Household Income: 2016 (American Community Survey Briefs)*. Washington, DC: U.S. Census Bureau; 2017. Available at <https://www.census.gov/content/dam/Census/library/publications/2017/acs/acsbr16-02.pdf>. Accessibility verified June 1, 2019.
42. Kochhar R, Fry R, Rohal M. America’s shrinking middle class: A close look at changes within metropolitan areas [Internet]. 2016. Available at <https://www.pewresearch.org/wp-content/uploads/sites/3/2016/05/Middle-Class-Metro-Areas-FINAL.pdf>. Accessibility verified March 15, 2019.

43. U.S. Department of Agriculture Economic Research Service. Rural–urban commuting area codes [Internet]. 2019. Available at <https://www.ers.usda.gov/data-products/rural-urban-commuting-area-codes.aspx>. Accessibility verified July 19, 2019.
44. McCabe SE, Hughes TL, Bostwick WB, West BT, Boyd CJ. Sexual orientation, substance use behaviors and substance dependence in the United States. *Addiction*. 2009;104:1333–1345.
45. Lee JG, Matthews AK, McCullen CA, Melvin CL. Promotion of tobacco use cessation for lesbian, gay, bisexual, and transgender people: A systematic review. *Am J Prev Med*. 2014;47:823–831.
46. Tracy JK, Schluterman NH, Greenberg DR. Understanding cervical cancer screening among lesbians: A national survey. *BMC Public Health*. 2013;13:442.
47. Matthews AK, Li C-C, Ross N, Ram J, Ramsey R, Aranda F. Breast and cervical cancer screening behaviors of African American sexual minority women. *J Gen Pract*. 2013;1:1–8.
48. Talley AE, Turner B, Foster AM, Phillips G, II. Sexual minority youth at risk of early and persistent alcohol, tobacco, and marijuana use. *Arch Sex Behav*. 2019;48:1073–1086.
49. Hiscock R, Bauld L, Amos A, Fidler JA, Munafò M. Socioeconomic status and smoking: A review. *Ann N Y Acad Sci*. 2012;1248:107–123.
50. Gamarel KE, Watson RJ, Mouzoon R, Wheldon CW, Fish JN, Fleischer NL. Family rejection and cigarette smoking among sexual and gender minority Adolescents in the USA. *Int J Behav Med*. 2020;27:179–187.
51. Harnois CE, Bastos JL, Campbell ME, Keith VM. Measuring perceived mistreatment across diverse social groups: An evaluation of the everyday discrimination scale. *Soc Sci Med*. 2019;232:298–306.
52. McMillen RC, Gottlieb MA, Shaefer RM, Winickoff JP, Klein JD. Trends in electronic cigarette use among U.S. adults: Use is increasing in both smokers and nonsmokers. *Nicotine Tob Res*. 2015;17:1195–1202.
53. Mirbolouk M, Charkhchi P, Kianoush S, et al. Prevalence and distribution of e-cigarette use among U.S. adults: Behavioral risk factor surveillance system, 2016. *Ann Intern Med*. 2018;169:429–438.
54. Coulter RWS, Bersamin M, Russell ST, Mair C. The effects of gender- and sexuality-based harassment on lesbian, gay, bisexual, and transgender substance use disparities. *J Adolesc Health*. 2018;62:688–700.
55. Azagba S, Latham K, Shan L. Cigarette smoking, e-cigarette use, and sexual identity among high school students in the USA. *Eur J Pediatr*. 2019;178:1343–1351.
56. Spivey JD. Tobacco policies and alcohol sponsorship at lesbian, gay, bisexual, and transgender pride festivals: Time for intervention. *Am J Public Health*. 2018;108(2):187–188.
57. U.S. Food and Drug Administration. This free life [Internet]. Available at <https://thisfreelife.betobaccofree.hhs.gov/>. Accessibility verified July 19, 2019.