

ARTICLE

Open Access

Plasma lipidome is dysregulated in Alzheimer's disease and is associated with disease risk genes

Yue Liu^{1,2}, Anbupalam Thalamuthu¹, Karen A. Mather^{1,3}, John Crawford¹, Marina Ulanova¹, Matthew Wai Kin Wong¹, Russell Pickford⁴, Perminder S. Sachdev^{1,5} and Nady Braidy^{1,6}

Abstract

Lipidomics research could provide insights of pathobiological mechanisms in Alzheimer's disease. This study explores a battery of plasma lipids that can differentiate Alzheimer's disease (AD) patients from healthy controls and determines whether lipid profiles correlate with genetic risk for AD. AD plasma samples were collected from the Sydney Memory and Ageing Study (MAS) Sydney, Australia (aged range 75–97 years; 51.2% male). Untargeted lipidomics analysis was performed by liquid chromatography coupled–mass spectrometry (LC–MS/MS). We found that several lipid species from nine lipid classes, particularly sphingomyelins (SMs), cholesterol esters (ChEs), phosphatidylcholines (PCs), phosphatidylethanolamines (PIs), phosphatidylinositols (PIs), and triglycerides (TGs) are dysregulated in AD patients and may help discriminate them from healthy controls. However, when the lipid species were grouped together into lipid subgroups, only the DG group was significantly higher in AD. ChEs, SMs, and TGs resulted in good classification accuracy using the Glmnet algorithm (elastic net penalization for the generalized linear model [glm]) with more than 80% AUC. In general, group lipids and the lipid subclasses LPC and PE had less classification accuracy compared to the other subclasses. We also found significant increases in SMs, PIs, and the LPE/PE ratio in human U251 astrogloma cell lines exposed to pathophysiological concentrations of oligomeric A β ₄₂. This suggests that oligomeric A β ₄₂ plays a contributory, if not causal role, in mediating changes in lipid profiles in AD that can be detected in the periphery. In addition, we evaluated the association of plasma lipid profiles with AD-related single nucleotide polymorphisms (SNPs) and polygenic risk scores (PRS) of AD. We found that *FERMT2* and *MS4A6A* showed a significantly differential association with lipids in all lipid classes across disease and control groups. *ABCA7* had a differential association with more than half of the DG lipids (52.63%) and PI lipids (57.14%), respectively. Additionally, 43.4% of lipids in the SM class were differentially associated with *CLU*. More than 30% of lipids in ChE, PE, and TG classes had differential associations with separate genes (ChE–*PICALM*, *SLC24A4*, and *SORL1*; PE–*CLU* and *CR1*; TG–*BIN1*) between AD and control group. These data may provide renewed insights into the pathobiology of AD and the feasibility of identifying individuals with greater AD risk.

Introduction

Alzheimer's disease (AD) is the most common cause of dementia, accounting for about 70% of total cases¹. This progressively neurodegenerative disease is characterized

by an insidious onset and is clinically defined by a progressive loss of memory and other cognitive deficits. AD has become one of the major challenges for the public health and economic system of the 21st century. There is still no international consensus on the etiology of this multifactorial disease, in which in addition to proteopathies, oxidative stress, inflammation, metabolic disorder, and other factors play a part^{2–4}.

The lack of effective treatments and the potential for prevention highlight the importance of identifying early

Correspondence: Nady Braidy (n.braidy@unsw.edu.au)

¹Centre for Healthy Brain Ageing (CHeBA), School of Psychiatry, University of New South Wales, Sydney, Australia

²Guangdong Mental Health Center, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences, Guangzhou, Guangdong, China
Full list of author information is available at the end of the article

© The Author(s) 2021



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

biomarkers for diagnosing AD. In addition, there is evidence that pathological processes associated with AD can also manifest in the peripheral system⁵, indicating the possibility of identifying non-invasive blood biomarkers. Lipids participate in important functions such as cell membrane formation, cellular transport, and energy storage, and act as essential signaling molecules. Beyond their structural roles, lipids have also been shown to act as modulators of transmembrane proteins, such as ion channels, whereby alteration of the composition or conformation of lipids surrounding ion channels can affect their function^{6,7}. Given the essential role of lipids in major biological processes, blood lipids have emerged as promising biomarkers for AD^{8–10}.

Although there have been many studies on the association between lipids and the pathobiology of AD, there are few studies on the plasma lipidome in AD. In contrast to classical biochemical approaches that focus on single metabolites or reactions, lipidomics approaches simultaneously identify and quantify hundreds of lipids. Measurement of large numbers of lipids enables network analysis approaches and provides means to identify critical metabolic drivers in disease pathophysiology. Lipidomics provides powerful tools for mapping global biochemical changes in disease and treatment.

In our current study, we examined differences in the plasma lipidome between AD and 'healthy' age-matched controls and compared the ability of different lipid profiles to discriminate between the two groups. We also showed that human astroglia cultures exposed to pathological levels of amyloid-beta ($A\beta$)₄₂ oligomers shared similar cellular lipidomic profiles to those observed in human plasma AD. We explored the effect of AD polygenic risk scores and AD-related SNPs on plasma lipid levels between AD and controls.

Method

Participants

Participants were a subsample from the population-based longitudinal Sydney Memory and Ageing Study (Sydney MAS)¹¹, an ongoing study that began in 2005 and focuses on cognitive decline in community-dwelling elderly. Participants were aged 70–90 years, initially without dementia, living in the community, and able to complete their assessments in English. There have been four Waves of data collection, two years apart. At each Wave, participants underwent an MRI scan, comprehensive neuropsychological assessment, medical examination, and blood collection for biochemistry analyses and DNA extraction. Written informed consent was obtained from all participants. In this study, AD (diagnosed by NINCDS-ADRDA criteria) and cognitively normal control samples

were collected at Waves 2 and 4. Ethics approval for this study was obtained from UNSW Sydney Australia and the South-Eastern Illawarra Area Health Service—Eastern sector¹¹. The investigators were blinded to sample allocation during the study and outcome assessment.

Plasma lipid extraction

Plasma lipids were extracted as previously described¹². Briefly, 10 μ L internal lipid standards (ISTDs) (Avanti lipids, <https://avantilipids.com/product/330707>) were added to 10 μ L aliquot of each plasma sample. 100 μ L of 1-butanol-Methanol (1:1 v/v) containing 5 mM ammonium formate were used to dissolve the mixture. Samples were vortexed for 10 s then sonicated for one hour. Afterward, samples were centrifuged at 13,000 $\times g$ for 10 min. The supernatant was transferred into a fresh Eppendorf tube. A further 100 μ L of 1-butanol/methanol (1:1 v/v) with 5 mM ammonium formate was added to the white pellet to re-extract any remaining lipids. The supernatant was dried in a speed vacuum centrifuge for 40–60 min. The lipids were reconstituted by adding 100 μ L of 1-butanol/methanol (1:1 v/v) containing 5 mM ammonium formate to each tube. The contents were transferred into a 300 μ L glass Chromacol vial with a glass insert prior to liquid chromatography/mass spectrometry (LC–MS).

Liquid chromatography/mass spectrometry

Lipid analysis was performed by LC ESI-MS/MS using a Thermo QExactive Plus Orbitrap mass spectrometer as previously described¹². Briefly, a Waters ACQUITY UPLC CSHTM C18 1.7 μ m, 2.1 \times 100 mm column was used for liquid chromatography at a flow rate of 260 μ L/min, using the following gradient condition: 32% solvent B to 100% over 25 min, a return to 32% B and finally 32% B for 5 min prior to the next injection. Solvents A and B consisted of acetonitrile: MilliQ water (6:4 v/v) and isopropanol:acetonitrile (9:1 v/v) respectively, both containing 10 mM ammonium formate and 0.1% formic acid. The first 3 min of eluent, containing the eluted salts, was diverted to waste. A product ion scan in positive and negative ion modes was performed to analyze the individual lipid species. The order of sampling was randomized prior to analysis. Ceramide (Cer), sphingomyelin (SM), phosphatidylcholines (PC), phosphatidylethanolamines (PE), phosphatidylinositol (PI), lyso-phosphatidylcholines (LPC), cholesterol esters (ChE), diacylglycerol (DG) and triacylglycerols (TG) were detected. The abundance of lipids was acquired using Lipidsearch software version 4.2 (Thermo Fisher Scientific, Sydney, NSW AU) according to accurate lipid mass and fragment matching¹². The LC–MS data were exported into Microsoft Excel and normalized by dividing the abundance

of internal standards to be used for further statistical analyses.

Cell lipidomics

Cell culture

U251 human astrogloma cell lines were purchased from the ATCC. These cells were cultivated in Roswell Park Memorial Institute (RPMI) 1640 Medium supplemented with 10% fetal bovine serum, 1% L-glutamax, and 1% antibacterial/antifungal. The cells were recently tested and found to be mycoplasma-free using the MycoAlert Mycoplasma Detection Kit (Lonza). The cells were grown at 37 °C in 95% humidified air and 5% CO₂. The culture medium was replaced every 2 days. U251 cells were seeded in a 12-well microtitre plate and given treatment when cells were nearly at confluency (0.5 × 10⁶ cells per well). Groups for evaluation are as follows: 6 wells containing naive U251 cells with no treatment (control) for 24 h, and 6 wells containing U251 cells treated for 24 h with 5 μM oligomeric Aβ₄₂ (see below for further details).

Preparation of recombinant Aβ₄₂ peptide

Aβ₄₂ peptide was purchased from Recombinant Peptide Technologies (Athens, GA, USA). The peptide was immediately stored in sealed glass vials at -80 °C in a lyophilized form. Consistent homogenous preparations of recombinant Aβ₄₂ oligomers or fibrils for use in cell culture stimulation experiments were obtained following a previously published protocol¹³. To avoid condensation upon opening, each vial was left at room temperature for 30 min prior to resuspension. Using 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP; Sigma, Castle Hill, Australia), the lyophilized peptide was initially dissolved to 1 mM and separated into 50 μL aliquots in sterile microcentrifuge tubes. Aliquots were left for 3 h in a fume hood, which allowed almost complete evaporation of HFIP. The resulting peptide films were further dried under vacuum using a Speed Vac (ThermoSavant, Patterson, CA, USA). This procedure is important as HFIP evaporated Aβ will form fibrils if exposed to moisture in the air, hence it is paramount to store HFIP evaporated Aβ films in desiccant. These preparations were then stored at -20 °C until required.

Production of recombinant Aβ₄₂ oligomers

Each HFIP aliquot of the peptide film was thoroughly resuspended to 5 mM in anhydrous dimethyl sulfoxide (DMSO; Sigma, Castle Hill, Australia) via pipette mixing, followed by 10-min bath sonication (Model: FX8, Unisonics, Sydney, Australia). Oligomers were formed by adding ice-cold Dulbecco's modified Eagle medium/Ham F-12 without phenol red (Sigma, Castle Hill, Australia) to a final concentration of 100 μM. Following a 30 s vortex,

the preparation was incubated at 4 °C for 6 weeks for the formation of Aβ₄₂ oligomers.

Cell lipidomics

Samples were collected using a cell scraper to 1 mL PBS solvent. After centrifuging 5 min at 13,000 × g, cell pellets were collected and reconstituted in 110 μL MilliQ water. 10 μL of cell solution was used for protein assay and the rest was used for lipid extraction. Lipids extraction and LC-MS methods were the same as plasma lipids, which were described above. 10 lipid classes including Cer, SM, PC, PE, PI, LPC, lyso-phosphatidylethanolamines (LPE), ChE, DG, and TG were detected.

Quantification of oligomeric Aβ

Oligomeric Aβ was quantified as previously described¹⁴. Briefly, plasma samples were thawed at 37 °C for 15 min. Afterwards, 10 μL of plasma, 4 μL of HAMA (human anti-murine antibody, HAMA) blocker (Scantibodies Laboratory, Santee, CA, USA), were mixed. 10 μL of PBR-1 (1% proprietary + 1.25% dimethyl sulfoxide (DMSO) + 96.75% phosphate-buffered saline contains Tween 20 (PBST) + 1% ultra-pure water) were further mixed into the plasma mixture. The mixtures were incubated for 48 and 1 h, respectively. The plasma sample mixture and serially diluted standards were added to separate wells of the plate in a total volume of 100 μL. The plates were incubated at RT for 1 h. The detection antibody was added to the wells, and the plate was incubated for 1 h at RT. Finally, 100 μL of 3,3',5,5'-tetramethylbenzidine (TMB) reagent was added as a substrate, and after 15 min, the reaction was stopped with 50 μL of 1 M H₂SO₄. Optical density (OD) values were measured using the BMG Fluostar Optima multimode plate reader (NY, USA), at a wavelength of 450 nm. Prior studies using this method detected the raw luminescence signal and used relative luminescence units (RLU) to present the oligomeric Aβ levels¹⁵.

Genotyping

DNA was extracted using standard procedures. Genotyping of the APOE ε2/3/4 polymorphism was performed as described¹⁶. Genome-wide genotyping was undertaken using the Affymetrix Genome-wide Human SNP Array 6.0 (California, USA) at the Ramaciotti Centre, UNSW Australia¹⁷. The CRLMM package (v1.10.0) in R (v2.12.1) was used to call genotypes. SNPs were excluded if the genotyping call rate was <95%, had a minor allele frequency <0.01 or if they failed a Hardy-Weinberg equilibrium threshold of <1 × 10⁻⁶. After further QC checks, there were 925 Sydney MAS participants with data for 734,550 SNPs. Imputation was undertaken to the 1000 Genome reference panel using the Michigan Imputation

Server. SNPs with poor imputation quality were omitted from any further analyses ($R^2 \leq 0.6$).

Polygenic Risk Scores (PRS) and AD implicated SNPs

PRS were generated using the PRSice program¹⁸ from summary statistics obtained from a previous Alzheimer's disease GWAS¹⁹. Linkage disequilibrium pruning was performed using the clumping option ($r^2 > 0.25$ and physical distance threshold of 250 kb KB). We present the association of lipids with the PRS calculated using the SNPs with the AD GWAS p -value threshold $\leq 5 \times 10^{-5}$. Associations between 33 individual AD-related SNPs²⁰ that passed QC checks (SNPs with MAF > 0.05 and imputation quality > 0.6) and lipids were also undertaken. Details regarding the SNPs and their associated genes utilized in the analyses are provided in Supplementary Table 1.

Statistical methods

Comparison of lipids between AD and controls

Inverse normal transformed residuals for the individual lipids and group lipids were obtained after regressing out the effect of possible confounders: age, sex, BMI, diabetes status, hypertension status, medication status for hypertension and hyperlipidemia, *APOE* e4 carrier status, education, and current smoking status. This transformed data was used in all subsequent analyses. T -tests were used to compare the mean value of lipids residuals between AD and controls. We used a significance threshold of 0.05 after False discovery rate (FDR) correction for all comparisons. Fold change (FC) was calculated as the ratio of the average lipids abundance in AD and controls. T -tests were also applied to compare lipid abundance between $A\beta$ treated and untreated cells. Between-group comparisons are done assuming unequal variance between groups with an approximation for degrees of freedom.

Classification of AD vs. Control using GLMnet

We used a machine-learning algorithm, glmnet (elastic net penalization for the generalized linear model [glm]) to classify AD versus control samples. A combination of two penalty functions with two tuning parameters was utilized to shrink the beta coefficients in the glm²¹. R (version 3.5.1)²² package caret²³ for fitting the elastic net glm model with default options was used to identify the optimum values for the tuning parameters.

For classification analysis, the data were randomly split into 70% training and 30% test samples maintaining the proportion of cases and controls in the training and test samples as in the full dataset. For the training data, the algorithms were run with three cross-validations with five repeats. To avoid bias due to a single random split of the

original data, we have repeated the analysis 10 times and the results were summarized over the 10 iterations.

The glmnet classification accuracy was examined based on several subgroups of the lipid species. The receiver operating curve (ROC) and area under ROC (AUC) were obtained using the R package pROC²⁴. Average sensitivity (proportion of AD cases predicted by the model in the test data), specificity (proportion of controls predicted by the model in the test data), and the AUC across 10 iterations are reported.

Genetic variation and lipid profiles among AD vs. controls

Linear regression was used to examine the association of AD risk variants and AD PRS with the lipids. The inverse normal transformed lipid residuals were used as the dependent variable and individual SNPs or PRS, case-controls (CC) status (AD vs. controls), and the relevant interaction term (CC \times SNP or CC \times PRS) were used as independent variables. Differential association of the AD risk variants and the PRS with lipids among cases and controls were examined based on the significance of the interaction term.

Results

Descriptive statistics

The sample comprised 82 plasma samples (40 AD patients and 42 cognitively normal 'healthy' controls) from the Sydney MAS cohort. The demographic information of participants is displayed in Table 1. The AD patients were significantly older than the controls with all aged more than 75 years old. There were no gender differences between the 2 groups ($\chi^2 = 1.233$, $p = 0.267$). As expected, cognitive scores on the Mini-Mental State Examination (MMSE) were significantly lower in AD patients. Education levels, which represent a protective factor for AD were measured in years and showed no statistical difference between the AD and control groups. AD and controls did not differ in vascular risk factors, including frequency of hypertension, diabetes, and current smoking status. Carriers of the *APOE**4 allele, a strong genetic risk factor for AD risk, were more common in AD patients than controls, although this did not reach statistical significance (AD 37.5% vs. control 19%, $\chi^2 = 3.457$, $p = 0.063$). The elevated levels of oligomeric $A\beta$ in the plasma distinguished the AD and control groups and were associated with increased MMSE, in patients with AD.

Comparison of lipids between AD and control patients

A total of 778 distinct lipid species from 9 lipid classes were analyzed, including neutral lipids such as 14 cholesteryl esters (ChE), 50 diglycerides (DG), 382 triglycerides (TG); sphingolipids including 43 ceramides (Cer), 90 sphingomyelins (SM); and phospholipid subclasses including 120 phosphatidylcholines (PC), 24 lyso-

Table 1 Characteristics of Alzheimer's disease and control participants.

	Healthy controls (n = 42)	Alzheimer's disease (n = 40)	Difference
Age(years)	81.27(2.48)	86.72(5.03)	$t = 6.270, p < 0.01$
Sex (males/females)	19/23	23/17	$\chi^2 = 1.233, p = 0.267$
Oligomeric A β (OD value)	0.25(0.05)	0.40(0.10)	$t = 6.391, p < 0.01$
Education (years)	11.30(3.59)	11.59(3.64)	$t = 0.363, p = 0.717$
Body mass index	26.69(3.19)	25.45(3.72)	$t = 1.631, p = 0.107$
Diabetes	7.10%	15%	$\chi^2 = 1.294, p = 0.255$
Hypertension	61.90%	77.50%	$\chi^2 = 2.351, p = 0.125$
Anti-hypertensive medication	60.00%	54.30%	$\chi^2 = 0.249, p = 0.618$
Anti-hyperlipidemia medication	64.30%	50.00%	$\chi^2 = 1.709, p = 0.791$
Current smoker	9.50%	5.00%	$\chi^2 = 0.618, p = 0.432$
APOE*4 carriers	19%	37.50%	$\chi^2 = 3.457, p = 0.063$
MMSE	28.21(1.58)	22.63(3.86)	$t = 8.557, p < 0.01$

Continuous variables are expressed as means and standard deviations. Categorical variables are expressed as %. APOE*4 apolipoprotein epsilon 4, MMSE Mini-mental state examination.

phosphatidylcholines(LPC), 38 phosphatidylethanolamines (PE), and 17 phosphatidylinositols (PI). The proportion of lipids significantly different between AD and controls according to each of the examined lipid classes are shown in Fig. 1. Volcano plots were drawn according to lipid fold changes of abundance comparing AD and controls and FDR-corrected p values in for all individual lipid species (see Fig. 2 and Supplementary Table 2).


Almost all ceramides were significantly different between control and AD groups after adjustment for age, sex, and vascular risk factors, even after multiple testing corrections. Molecular profiles of the ceramidome showed that dihydroxy Cer species comprising fatty acyls (18C) with 0 or 2 double bonds and all dihydroxy Cer with 19C were significantly higher in AD. On the other hand, dihydroxy Cer containing 16C or 18C with 1 double bond showed the opposite results. Most monohydroxy and trihydroxy Cer were significantly lower in AD subjects. Four AD positive ceramides including Cer(d18:0_16:0), Cer(d19:1_24:0), Cer(d18:0_23:0), and Cer(d18:2_25:0), and AD negatively correlated ceramides: Cer(d18:1_23:0), Cer(t16:1_14:0), Cer(m18:1_20:0), Cer(m18:0_22:0), and Cer(d19:1_22:0) showed the largest fold changes ($FC > 4$). Sphingomyelin is another type of sphingolipid. Most lipids in SM class showed significantly lower levels in AD plasma. The lipids with $FC > 4$ included SM(d35:4), SM(d34:1), SM(d31:1), SM(d18:1_21:0), SM(t36:2), and SM(d40:4). In contrast, SM(d41:1), SM(d41:4), and SM(d18:1_24:3) were largest downregulated in AD.

In the AD cases, there was a higher abundance of PC lipids compared to controls. They were comprised of very short and short-chain fatty acyls except for PC containing

C15 without double bonds and C16 with 2 double bonds. The largest fold changes were observed in PC(16:0_22:6), PC(18:2_18:2), PC(18:0_20:4), PC(36:2), PC(16:1_22:5) that showed upregulation in AD. PC (20:2_18:2) and PC (38:7) showed lower plasma concentrations in AD. LPCs with longer fatty acid chains tended to be lower in AD patients. No LPC lipids differed significantly between cognitively normal and AD groups. Most of PE lipids were lower in AD-affected patients. Major PE lipids were reduced in AD, including highly changed ($FC > 4$) PE (16:0p_22:6), PE(18:0p_20:4), PE(16:0p_18:1), and PE (18:0p_22:4). PE (18:0_18:1) was higher in plasma of Alzheimer's patients. PI showed an elevated abundance in AD and the PI lipids, with PI (18:0_18:3) and PI (18:1_20:4) the most significantly elevated in AD cases.

ChE (18:3), ChE (20:3), and ChE (22:3) showed the largest fold changes. Consistent with the result of group DGs, most DG lipids species were higher in AD patients, including DG(16:0_18:3), DG(22:4e), DG(17:1_18:1), DG (20:0_18:2), and DG(36:4e) with largest fold changes. Most downregulated DGs include DG(18:1_20:4), DG (16:0_18:1), and DG(18:0_18:1).

As well, around 30% of TG lipids were observed to be higher in AD patients. The most upregulated TG lipids include TG(18:1_17:1_18:3), TG(14:0_18:2_20:5), TG (16:1_20:1_22:4), TG(60:10), and TG(18:1_18:1_22:5). On the other hand, most of the long-chain polyunsaturated fatty acid-containing TGs were reduced in AD compared to age-matched controls. About 18% of plasma TG lipids are decreased in AD patients, and TG(18:1_12:0_14:0), TG (18:4_16:1_18:3), TG(16:0_20:4_22:6), TG(16:0_14:0_18:1), and TG(16:0_16:0_16:0) showed the largest fold changes. We



also identified suggestive group differences between almost all species containing more than 2 double bonds. Nine lipid groups calculated by summarizing all lipid species in the same lipid class were compared between AD and controls (Fig. 3 and Supplementary Table 3). Only the DG group was significantly higher in AD ($p = 0.009$, mean of residuals: Control: -0.276 ± 0.143 , AD: 0.290 ± 0.154). Other lipid group's lipids did not differ after adjusting for covariates.


Cell lipidomics

Results are plotted in Fig. 4. The concentration of oligomeric peptide used in this study is similar to that reported from human CSF and human cell culture conditioned medium^{25,26}. Additionally, it also represents a concentration closer to the levels found in brain^{25,26} offering a more physiologic impression of the effect of A β oligomers on astrocyte

glial lipid profiles. SMs were significantly elevated in astroglia cells treated with A β_{42} oligomers compared to non-treated cells ($p = 0.0015$). No significant differences were found in phospholipids except that the level of the PI group was higher in the A β_{42} group ($p = 0.041$), which is consistent with the result found in human AD plasma samples. Even though PE and LPE were not significantly different between treated and non-treated cells, the ratio of LPE/PE significantly increased in A β_{42} -treated astroglia cells (p value). TG was also increased in the A β_{42} group ($p = 0.005$).

Classification of AD vs. Control using lipid profiles


Several models were used for the classification of AD versus controls based on the plasma lipidome profile. The glmnet package was used in all the analyses. The analyses were repeated 10 times and results were summarized



across the 10 iterations. The algorithms were run in turn using the full list of individual lipids, all of the 9 subclasses of lipids, and group lipids. The average sensitivity, specificity, and AUC of the test data are summarized in Table 2. The ROC curves for the top 3 models, ChE, SM, and TG are presented in Fig. 5 and resulted in good classification accuracy with more than 80% AUC. In general, group lipids and the lipid subclasses LPC and PE had less classification accuracy compared to the other subclasses.

SNPs and lipids with AD versus control

The associations of AD PRS and AD-related SNPs with the lipidome were examined. The individual lipids that were significantly different between AD and controls ($n = 420$) were utilized to explore the association. No significant association of AD PRS with lipids was observed after FDR correction (Supplementary Table 4). However, the AD PRS \times CC interaction was nominally significant (p -value $< = 0.05$) for 57 lipid species (Table 3), which were mainly



PCs, SMs, and TGs. Out of the 57 significant interactions, only two PRS scores had marginal significant effects (p value < 0.05). The same analysis was also repeated using 33 AD implicated individual SNPs recently identified in another study²⁰. All of these individual SNPs were found to have different effects on a certain number of lipids between AD and controls before adjustment for multiple testing. Nominally significantly different associations of SNPs with lipids between AD and controls are listed in Supplementary Table 5.

We further explored the association of AD-associated genes ($n = 19$) derived from the list of 33 SNPs with lipids in each lipid class (Table 4). *FERMT2* and *MS4A6A* showed a significantly differential association with lipids in all lipid classes across disease and control groups. In addition, PC, SM, and TG lipids were differentially associated with the largest number of AD-related genes. The heatmap in Fig. 6 shows the proportion of lipids in each lipid class having differential associations with the AD-associated genes. *ABCA7* had a differential association with more than half of the DG

lipids (52.63%) and PI lipids (57.14%), respectively. Additionally, 43.4% of lipids in the SM class were differentially associated with *CLU*. More than 30% of lipids in ChE, PE, and TG classes had differential associations with separate genes (*ChE-PICALM*, *SLC24A4*, and *SORL1*; *PE-CLU* and *CRI*; *TG-BINI*) between AD and control group. Finally, we found 43 lipids that were significantly associated with both the AD-related genes and PRS (Table 5).

Discussion

In this study, significant differences in plasma lipids between AD cases and controls were observed. In vitro analyses demonstrated similar changes in astroglial cells when treated with $A\beta_{42}$. GLMnet algorithm was used to distinguish AD from controls using different plasma lipid-based models. SMs, ChEs, and TGs showed the greatest accuracy in discriminant analysis with AUC of more than 80%. Finally, associations between AD case-control status and genetic risk factors for AD were nominally significant when examining lipid profiles.





Fig. 4 Normalized lipid group abundance in amyloid beta-treated astrocytes versus controls. Cer ceramides, SM sphingomyelins, ChE Cholesteryl esters, DG diglycerides, TG triglycerides, PC phosphatidylcholines, LPC lyso-phosphatidylcholines, PE phosphatidylethanolamines, PI phosphatidylinositols. *T* tests were used to compare lipids between two groups. ns: no significance, * $p < 0.05$, ** $p < 0.01$.

Table 2 GLMnet results of lipid models in the classification of AD versus control.

Lipid models	Sensitivity	Specificity	AUC
All_lipids	0.66	0.73	0.78
Cer	0.7	0.79	0.79
ChE	0.78	0.75	0.82
DG	0.72	0.69	0.76
LPC	0.49	0.62	0.58
PC	0.69	0.76	0.78
PE	0.58	0.5	0.57
PI	0.7	0.76	0.79
SM	0.71	0.78	0.83
TG	0.65	0.82	0.83
SM_TG	0.69	0.75	0.81
ChE_SM	0.7	0.81	0.79
ChE_TG	0.71	0.78	0.82
ChE_SM_TG	0.67	0.77	0.79

Cer ceramides, SM sphingomyelins, ChE cholesteryl esters, DG diglycerides, TG triglycerides, PC phosphatidylcholines, LPC lyso-phosphatidylcholines, PE phosphatidylethanolamines, PI phosphatidylinositols, GLMnet elastic net penalization for the generalized linear model.



Identification of the roles of lipids associated with AD

Our study did not show a significant group ceramide difference between AD and controls. However, we found

that most individual ceramides differed significantly between these two groups. Molecular profiles of the ceramidome showed that dihydroxy Cer species, comprising fatty acyls (18C) with 0 or 2 double bonds and all dihydroxy Cers with 19C, were significantly increased in AD. On the other side, dihydroxy Cers containing 16C or 18C with 1 double bond showed an opposite result. Most monohydroxy and trihydroxy Cers decreased significantly in AD subjects, which might suggest the opposite pathobiology of different Cer subgroups. Ceramides are essential for the maintenance of membrane structure stabilization, cell-to-cell recognition, and secondary messenger signaling²⁷. Dysregulation of Cers has been linked to cognitive decline and brain atrophy²⁸. Higher levels of serum ceramide were associated with increased risk of AD compared to subjects with the lowest tertile of serum ceramide levels in another study²⁹. In addition, even though the sum of SM lipids was not significantly different between the two groups, we found that most individual SM lipids were lower in AD plasma. Altered sphingolipid metabolism has been associated with the pathogenesis of several neurodegenerative diseases including AD, and metabolic syndrome^{29–32}. The decrease in SM lipids has been shown to be strongly correlated with parameters of insulin resistance, and lipid metabolism, which are major risk factors for AD³³. A cross-sectional study ($n=26$ NC, 26 AD) observed reduced SM species in AD versus controls and long aliphatic chains (C22, C24) in particular³¹, which is consistent with our results.

We did not find any significant differences among group PC lipids. However, the AD population tended to present higher abundance in individual PC lipids comprising very short and short-chain fatty acyls except for PC containing C15 without double bonds and C16 with 2 double bonds. Recent studies have demonstrated altered PC metabolism in cognitively impaired elderly and AD patients^{9,34,35}. Similar to PC lipids, we did not observe significant changes in group PE. About 24% of individual PEs were all lower in AD patients, which is almost 2 fold of PEs that were higher in AD. PE has been previously reported to be significantly reduced in the brains of individuals with AD and HD³⁶. Since a multitude of neuropathological processes can lead to a decrease in PC and PE, it is likely that reduced levels of these phospholipids may be linked to the observed neuronal loss. Increased activities of the Kennedy pathway enzymes, phosphoethanolamine cytidyltransferase, phosphocholine cytidyltransferase, and PS synthase have been previously reported to be elevated in the diseased brain regions of patients with AD and PD³⁴. Contrary to the decreased trend of other phospholipids, including PC and PE, we observed an opposite change in PI in AD patients, with most of the individual PI lipids higher in AD subjects. PI is another phospholipid that is

Table 3 Nominally significantly differential association of the AD risk PRS with lipids between AD and controls control.

Lipid	Beta.PRS	Beta.PRS_CC	SE.PRS	SE.PRS_CC	t.PRS	t.PRS_CC	Pval.PRS	Pval.PRS_CC
Cer(d18:1_16:0)	0.197	-0.511	0.133	0.195	1.483	-2.624	0.142	0.011
Cer(t16:1_14:0)	0.268	-0.510	0.141	0.207	1.894	-2.461	0.062	0.016
ChE(22:5)	-0.080	0.539	0.137	0.200	-0.583	2.690	0.562	0.009
PC(16:0_20:4)	-0.107	0.495	0.144	0.211	-0.745	2.345	0.459	0.022
PC(16:0_20:5)	0.001	0.535	0.144	0.210	0.007	2.544	0.994	0.013
PC(16:0_22:6)	0.150	0.419	0.123	0.180	1.221	2.331	0.226	0.022
PC(16:2e_16:0)	0.089	-0.428	0.145	0.212	0.612	-2.018	0.543	0.047
PC(18:0_20:3)	0.131	0.433	0.135	0.198	0.973	2.187	0.334	0.032
PC(18:1_20:3)	0.025	0.441	0.143	0.209	0.174	2.108	0.863	0.038
PC(18:2_18:2)	-0.005	0.415	0.138	0.202	-0.039	2.053	0.969	0.044
PC(34:0)	0.057	-0.510	0.142	0.208	0.404	-2.447	0.688	0.017
PC(37:2)	0.285	-0.488	0.147	0.215	1.942	-2.267	0.056	0.026
PE(16:0_20:3)	-0.059	0.445	0.144	0.210	-0.413	2.117	0.681	0.038
PI(18:0_18:2)	0.026	-0.421	0.123	0.180	0.208	-2.346	0.836	0.022
PI(36:3)	-0.163	0.603	0.148	0.217	-1.105	2.784	0.273	0.007
SM(d17:1_13:0)	0.353	-0.668	0.142	0.209	2.476	-3.200	0.016	0.002
SM(d17:1_18:3)	0.041	-0.405	0.134	0.196	0.303	-2.064	0.763	0.043
SM(d18:2_24:3)	0.240	-0.470	0.145	0.213	1.653	-2.207	0.103	0.030
SM(d32:4)	0.166	-0.585	0.134	0.197	1.233	-2.971	0.222	0.004
SM(d33:1)	0.095	-0.521	0.139	0.204	0.683	-2.551	0.497	0.013
SM(d34:2)	0.132	-0.416	0.141	0.206	0.933	-2.014	0.354	0.048
SM(d35:1)	0.204	-0.462	0.148	0.216	1.384	-2.134	0.171	0.036
SM(d36:3)	0.365	-0.568	0.146	0.213	2.506	-2.662	0.014	0.010
SM(d37:2)	-0.012	-0.429	0.136	0.199	-0.090	-2.161	0.929	0.034
SM(d38:1)	0.006	-0.419	0.127	0.186	0.049	-2.250	0.961	0.027
SM(d38:2)	0.027	-0.458	0.137	0.200	0.197	-2.285	0.844	0.025
SM(d39:1)	-0.031	-0.413	0.131	0.192	-0.240	-2.153	0.811	0.035
SM(d39:2)	0.108	-0.516	0.139	0.203	0.776	-2.539	0.440	0.013
SM(d40:1)	0.038	-0.453	0.135	0.198	0.283	-2.294	0.778	0.025
SM(d40:3)	0.118	-0.458	0.149	0.218	0.789	-2.098	0.433	0.039
SM(d41:4)	-0.070	-0.397	0.135	0.198	-0.520	-2.003	0.605	0.049
SM(d42:1)	0.103	-0.435	0.138	0.202	0.746	-2.152	0.458	0.035
SM(d42:2)	0.028	-0.507	0.129	0.189	0.221	-2.688	0.826	0.009
SM(d42:3)	0.041	-0.494	0.140	0.205	0.296	-2.409	0.768	0.018
SM(d43:1)	0.106	-0.500	0.147	0.215	0.721	-2.328	0.473	0.023
SM(d43:2)	0.167	-0.606	0.140	0.204	1.200	-2.966	0.234	0.004
SM(d44:4)	0.019	-0.421	0.136	0.200	0.143	-2.107	0.887	0.038
SM(d44:5)	0.134	-0.575	0.131	0.192	1.028	-3.001	0.308	0.004
SM(d44:6)	0.054	-0.427	0.139	0.204	0.389	-2.092	0.699	0.040

Table 3 continued

Lipid	Beta.PRS	Beta.PRS_CC	SE.PRS	SE.PRS_CC	t.PRS	t.PRS_CC	Pval.PRS	Pval.PRS_CC
SM(t18:0_24:2)	0.163	-0.509	0.143	0.209	1.145	-2.436	0.256	0.017
SM(t42:1)	0.240	-0.578	0.139	0.204	1.724	-2.832	0.089	0.006
TG(12:0_17:1_18:2)	-0.122	0.473	0.144	0.211	-0.848	2.243	0.399	0.028
TG(12:0_18:2_18:2)	-0.159	0.555	0.134	0.196	-1.191	2.832	0.237	0.006
TG(14:0_14:3_18:2)	-0.136	0.470	0.143	0.209	-0.950	2.250	0.345	0.027
TG(14:0_18:3_18:3)	-0.046	0.518	0.135	0.197	-0.345	2.623	0.731	0.011
TG(15:0_14:1_16:1)	-0.048	0.463	0.129	0.189	-0.375	2.453	0.709	0.017
TG(15:0_16:0_20:5)	-0.063	0.480	0.133	0.195	-0.474	2.464	0.637	0.016
TG(16:0_14:0_18:3)	-0.142	0.468	0.138	0.202	-1.032	2.317	0.305	0.023
TG(16:0_14:1_16:1)	-0.198	0.579	0.146	0.213	-1.360	2.711	0.178	0.008
TG(16:1_12:0_18:2)	-0.046	0.421	0.142	0.208	-0.324	2.023	0.747	0.047
TG(16:1_20:5_20:5)	0.088	0.503	0.126	0.185	0.696	2.724	0.488	0.008
TG(18:3_14:1_18:2)	-0.045	0.412	0.141	0.206	-0.319	1.999	0.751	0.049
TG(18:4_14:0_16:1)	-0.155	0.616	0.143	0.209	-1.082	2.944	0.283	0.004
TG(18:4_16:0_20:4)	-0.081	0.525	0.141	0.207	-0.571	2.533	0.570	0.013
TG(20:1_20:4_20:4)	0.304	-0.531	0.144	0.210	2.117	-2.524	0.038	0.014
TG(22:5_18:2_18:2)	-0.078	0.454	0.147	0.215	-0.529	2.110	0.598	0.038
TG(33:4e)	-0.160	0.651	0.137	0.201	-1.168	3.241	0.247	0.002

Marginal effect: Beta.PRS/SE.PRS/t.PRS/Pval. PRS: Beta coefficient/standard error/t statistics/p value of association between PRS scores and individual lipids.

Interaction effect: Beta.PRS_CC/SE.PRS_CC/t.PRS_CC/Pval.PRS_CC: Beta coefficient/standard error/t statistics/p value of differential association of PRS with lipids between AD and controls control.

Cer ceramides, *SM* sphingomyelins, *ChE* cholesteryl esters, *DG* diglycerides, *TG* triglycerides, *PC* phosphatidylcholines, *LPC* lyso-phosphatidylcholines, *PE* phosphatidylethanolamines, *PI* phosphatidylinositols.


present in the membrane of almost all cell types and is involved in mediating Ca^{2+} mobilization in response to many hormones, neurotransmitters, and growth factors³⁷. Consistently, an animal study showed that lower levels of plasma PC and higher plasma PI were correlated with post-stroke cognitive impairment³⁸.

We also found significant differences in ChEs between AD and controls. ChE is produced in the plasma by the conversion of fatty acids to cholesterol from PC by the enzymatic activity of cholesterol acyl transferase (LCAT)^{9,39}. While free cholesterol can be taken up by APOE containing liposomes (e.g. HDL) and is bound to the outer particle surface, esterification enhances cholesterol uptake within the interior of the lipoproteins and enhances cholesterol transport through the bloodstream. LCAT has a preference for highly unsaturated fatty acid chain PC and can link the reduction in PC to dysregulation of specific steps in cholesterol metabolism in AD⁴⁰. Another enzyme, acyl-coenzyme A can also esterify cholesterol in other tissues. Inhibition of this enzyme has been reported to reduce amyloid plaque load in the brain of AD mice and improves cognitive outcomes⁴¹. Consistent with our results showing a significant increase of

DGs in Alzheimer's disease, previous studies have demonstrated augmented levels of DGs in the frontal cortex and plasma of AD patients⁴². DGs are important in maintaining structural integrity and signal transduction⁴². The conversion of DGs to phosphatidic acid by DG kinase has been reported to be decreased in the AD brain⁴³. We also found almost 2/3 plasma TG lipids in all significant TG lipids were higher in AD subjects, even though the abundance of group TGs were not different among disease groups. A longitudinal finding was reported in the Honolulu-Asia Aging Study in which a 1SD increase in TG levels during midlife significantly increased the risk of dementia a quarter of a decade later⁴⁴. The exploration of the exact pathobiology of individual TG lipids is needed in the future.

Relationship between plasma lipid profiles and the cell lipidome exposed to oligomeric $\text{A}\beta_{42}$ in vivo

Several studies have reported significant differences in lipid profiles between plasma and postmortem brain tissue⁴⁵. The separation of the brain from the periphery is a major challenge when examining the biological significance of plasma lipids in Alzheimer's disease since not



all lipids can be transported across the blood–brain barrier. This separation may likely be compromised by age-related changes in blood–brain barrier integrity, affecting lipid and Aβ kinetics. To determine whether our observed changes in the plasma lipidome were related to AD pathology we also quantified the cellular lipidome in human U251 astrogloma cell lines exposed to pathophysiological concentrations of oligomeric Aβ₄₂.

Astrocytes are important cells in the CNS and are involved in various physiological aspects⁴⁶. Neurons are thought to be the main source of Aβ in the adult brain. Moreover, the ability of neurons to generate cholesterol in the adult brain is impaired during development. However, astrocytes can produce cholesterol and transport it to neurons with apolipoprotein E (apoE)⁴⁷. ApoE is the major genetic risk factor for sporadic AD. This has led to the hypothesis that apoE regulates Aβ formation via modulation of lipid raft functions⁴⁸. Recently, astrocyte-derived cholesterol has been shown to regulate Aβ formation in vivo and influence processes involved in inflammation and neurodegeneration⁴⁹. In our study, we quantified the sum of lipid classes when the astrogloma cell line was exposed to subpathological amounts of oligomeric Aβ₄₂. We still found a similar trend of lipid changes between plasma and cells treated with oligomeric Aβ₄₂. For example, SM, PI, and TG were significantly higher oligomeric Aβ₄₂-treated cells, and some SM, PI, and TG species were upregulated in AD plasma. However,

Table 4 Differential interaction of genes with lipids between AD and control.

Genes	Cer	ChE	DG	PC	PE	PI	SM	TG
<i>ABCA7</i>			✓		✓	✓	✓	✓
<i>BIN1</i>			✓	✓	✓	✓	✓	✓
<i>CASS4</i>			✓	✓			✓	✓
<i>CD2AP</i>				✓	✓		✓	✓
<i>CD33</i>			✓	✓			✓	✓
<i>CELF1</i>	✓			✓		✓	✓	✓
<i>CLU</i>	✓			✓	✓		✓	✓
<i>CR1</i>	✓			✓	✓	✓	✓	✓
<i>EPHA1</i>			✓	✓				✓
<i>FERMT2</i>	✓	✓	✓	✓	✓	✓	✓	✓
<i>INPP5D</i>	✓					✓	✓	✓
<i>MEF2C</i>								✓
<i>MS4A6A</i>	✓	✓	✓	✓	✓	✓	✓	✓
<i>NME8</i>							✓	✓
<i>PICALM</i>	✓	✓		✓			✓	✓
<i>PTK2B</i>		✓	✓	✓		✓	✓	
<i>SLC24A4</i>	✓	✓		✓			✓	✓
<i>SORL1</i>		✓		✓	✓		✓	✓
<i>ZCWPW1</i>	✓	✓	✓	✓			✓	✓

“✓”: ≥1 lipid in each lipid class were differentially associated with the named gene between AD and controls.

Cer ceramides, SM sphingomyelins, ChE cholesteryl esters, DG diglycerides, TG triglycerides, PC phosphatidylcholines, LPC lyso-phosphatidylcholines, PE phosphatidylethanolamines, PI phosphatidylinositols.

no significant differences were observed between AD and control in plasma and cells for other lipid groups. The results in the cellular lipidome support findings that Aβ₄₂ plays a contributory if not causal role in mediating changes in plasma and cellular lipid profiles in AD.

The lipidome signature as a biomarker of AD

We also examined whether a battery of plasma lipids can be used to discriminate AD patients from controls. All individual lipids, 9 subgroup lipids, and group lipids were used in the GLMnet algorithm. We observed that three classification models, using either TG, ChE, and SM, had significant power in discriminating AD from controls, with >80% AUC. Additionally, individual lipids in TG, ChE, and SM showed even higher AUC than group lipids. Therefore, measurement of these group lipids using mass spectrometry will provide renewed insight in the underlying mechanisms of AD and provide additional drug targets. Taken together, our findings suggest that alterations in both cholesterol and sphingolipid metabolism may play an important role in the pathobiology of AD.

Table 5 Overlap of lipids with significant association with SNPs and PRS scores.

Lipid	Number of associated SNPs	Most significant Pval.SNPs	Gene (minimum Pvalue.SNPs)	Pval.PRS_CC
Cer(d18:1_16:0)	3	0.009	<i>CLU</i>	0.011
ChE(22:5)	5	0.014	<i>SORL1</i>	0.009
PC(16:0_20:4)	1	0.045	<i>CASS4</i>	0.022
PC(16:0_20:5)	2	0.033	<i>SORL1</i>	0.013
PC(18:0_20:3)	1	0.027	<i>MS4A6A</i>	0.032
PC(18:1_20:3)	1	0.042	<i>CELF1</i>	0.022
PC(18:2_18:2)	1	0.007	<i>FERMT2</i>	0.044
PE(16:0_20:3)	1	0.005	<i>CR1</i>	0.038
PI(36:3)	1	0.002	<i>INPP5D</i>	0.007
SM(d17:1_13:0)	2	0.014	<i>SORL1</i>	0.002
SM(d17:1_18:3)	3	0.040	<i>CLU</i>	0.043
SM(d32:4)	3	0.008	<i>MS4A6A</i>	0.004
SM(d33:1)	5	0.021	<i>CLU</i>	0.013
SM(d34:2)	2	0.045	<i>CLU</i>	0.048
SM(d35:1)	1	0.012	<i>MS4A6A</i>	0.036
SM(d37:2)	4	0.014	<i>CD2AP</i>	0.034
SM(d38:1)	3	0.041	<i>CLU</i>	0.027
SM(d38:2)	3	0.022	<i>CLU</i>	0.025
SM(d39:1)	3	0.027	<i>CLU</i>	0.035
SM(d39:2)	3	0.027	<i>CLU</i>	0.013
SM(d40:1)	5	0.021	<i>CLU</i>	0.025
SM(d41:4)	1	0.028	<i>MS4A6A</i>	0.049
SM(d42:1)	2	0.013	<i>MS4A6A</i>	0.035
SM(d42:2)	5	0.012	<i>MS4A6A</i>	0.009
SM(d42:3)	3	0.044	<i>ABCA7</i>	0.018
SM(d43:1)	2	0.030	<i>CD2AP</i>	0.023
SM(d43:2)	1	0.032	<i>CASS4</i>	0.004
SM(d44:4)	3	0.016	<i>CLU</i>	0.038
SM(d44:5)	4	0.026	<i>CLU</i>	0.004
SM(t18:0_24:2)	3	0.022	<i>MS4A6A</i>	0.017
SM(t42:1)	5	0.023	<i>MS4A6A</i>	0.006
TG(12:0_17:1_18:2)	1	0.013	<i>NME8</i>	0.028
TG(14:0_14:3_18:2)	1	0.037	<i>BIN1</i>	0.027
TG(14:0_18:3_18:3)	3	0.003	<i>NME8</i>	0.011
TG(15:0_14:1_16:1)	1	0.032	<i>NME8</i>	0.017
TG(15:0_16:0_20:5)	2	0.048	<i>BIN1</i>	0.016
TG(16:0_14:0_18:3)	5	0.011	<i>CLU</i>	0.023
TG(16:0_14:1_16:1)	1	0.018	<i>SORL1</i>	0.008
TG(18:3_14:1_18:2)	2	0.035	<i>ABCA7</i>	0.049

Table 5 continued

Lipid	Number of associated SNPs	Most significant Pval.SNPs	Gene (minimum Pvalue.SNPs)	Pval.PRS_CC
TG(18:4_14:0_16:1)	1	0.045	<i>ZCWPW1</i>	0.004
TG(18:4_16:0_20:4)	3	0.011	<i>CLU</i>	0.013
TG(22:5_18:2_18:2)	4	0.017	<i>NME8</i>	0.038
TG(33:4e)	4	0.011	<i>MS4A6A</i>	0.002

Only lipids with significant differential associations with both PRS and SNPs were included in the table. The most significant *p* value of lipids association with SNPs of the same gene was presented when the SNP had the most significant differential association with the lipid across AD and control.

AD genetic risk, lipids, and AD

We examined the relationships between AD PRS and the AD-associated lipids in our sample. Even though no significant association was found after FDR correction, we observed a number of lipid species that were nominally differentially affected between AD and controls, most of which were PC, SM, and TGs. Our data suggest that these lipids might mediate the effect of several SNPs linked to AD.

We also investigated individual AD-related SNPs and lipid levels. Significant associations between AD-related SNPs and individual lipids were only observed before FDR correction. However, we found all SNPs having nominally significant differential association with a certain number of lipid species between AD and control group.

FERMT2 and *MS4A6A* genes showed a significantly differential association with lipids in all lipid classes across disease groups. *FERMT2* is a member of the Fermitin family of proteins, which are involved in cell–matrix adhesion complexes. Shulman et al.⁵⁰ validated the association of *FERMT2* with AD risk after performing a gene screen and in vivo studies in *Drosophila melanogaster*. Their work showed altered expression of both *FERMT2* and *CELF1* homologs modulates tau neurotoxicity. It is also upregulated in atherosclerotic plaques, which is a risk factor for AD⁵¹. The correlation of *FERMT2* with post-stroke brain recovery⁵² and higher plasma PI in post-stroke mice³⁸ supports the relationship between *FERMT2* and PI in our study, i.e. the PI class had the highest proportion of lipids of differential association with *FERMT2* between AD and control. It has been hypothesized that *FERMT2* regulates APP internalization and degradation⁵³. For instance, *FERMT2* silencing induces increased amounts of full-length APP and by-products and *FERMT2* over-expression leads to a reduction of APP and its related metabolites. Recently, a direct interaction between *FERMT2* and APP—through the F3 domain of *FERMT2* and the NxTY motif within APP's intracellular domain was reported⁵⁴ and can influence axonal growth⁵⁵. Importantly, because amyloidogenic processing of APP is dependent on lipid rafts, we postulate that the effect of *FERMT2* might be—at least in part—due to

alterations in lipid profiles which alter the accumulation of full-length APP or its by-products within the growth cone, impairing vesicle trafficking and/or cell adhesion.

MS4A6A is membrane-spanning 4-domains, subfamily A, member 6A having been revealed to be associated with cortical and hippocampal atrophy independent of glucose metabolism and A β deposition⁵⁶. The important association between *MS4A6A* and PI (57.14% had differential association with *MS4A6A*) might indicate the involvement of PI mediating *MS4A6A*'s effect on brain atrophy. Another gene *ABCA7* had an important effect on more than 50% of lipids in the DG lipid class, but with little influence on other lipid classes. DG metabolism and cholesterol synthesis might be the key component linking *ABCA7* to Alzheimer's onset. Increasing evidence has demonstrated that *ABCA7* deficiency exacerbates A β pathology in vitro and in vivo models. In detail, *ABCA7* is involved in the microglial A β clearance pathway and accelerated A β production⁵⁷. *ABCA7* may also mediate the release of cellular cholesterol and phospholipids to generate HDL. Human *ABCA7* mediates the efflux of both cellular cholesterol and phospholipids to apoA-I, whereas mouse *ABCA7* mediates only phospholipids but not cholesterol efflux to apoA-I⁵⁸. Both *MS4A6A* and *ABCA7* have been previously linked to the immune and complement system⁵⁹, lipid metabolism, and immune system⁶⁰, immune response and inflammation⁶¹, cytokine signaling in immune system⁶², cholesterol/lipid metabolism, and immune and complement systems⁶³.

CLU-encoded apolipoprotein J associated with the clearance of cellular debris and apoptosis, which was suggested to be involved in A β -independent pathways as part of the cascade leading to Alzheimer pathology⁶⁴. In our study, SM and PE were revealed to be associated with *CLU* in the pathogenesis of AD. ChE lipids were differentially associated with *PICALM*, *SLC24A4*, and *SORL1* between AD and control. *PICALM* has been associated with reduced connectivity of the frontal gyrus with the hippocampus, as well as with the precuneus. The *SLC24A4* gene encodes the 24 solute carrier family member 4 protein, which is a member of the K⁺-dependent Na⁺/Ca²⁺ exchanger protein family. These exchanger proteins are

widely expressed in brain tissues, suggesting that SLC24A4 may play an important role in the nervous system. The association of this gene with blood pressure in African Americans suggests it may have relevance with AD through vascular disease⁶⁵. *SORL1* belongs to the low-density lipoprotein receptor family, binding ApoE and lipoprotein lipase on membranes⁶⁶. Decreased Sorl1 has been identified as related to AD⁶⁷. It is downregulated in lymphoblasts and cortical pyramidal neurons of AD patients. The mechanisms may include APP trafficking and recycling⁶⁸. Additionally, the association of *SORL1* with risk of amnesic mild cognitive impairment (aMCI) has been reported in the Han Chinese⁶⁹. TG lipids were correlated with *BIN1*, which functions in clathrin-mediated endocytosis and endocytic recycling, as does the AD risk gene *PICALM*. DNA methylation of the *BIN1* promoter has been suggested as a possible epigenetic mechanism influencing AD risk.

Several prior studies have been conducted to explore the potential pathological roles of AD-related SNPs. A review of GWAS-identified risk genes for AD found they were clustered into three main molecular pathways: lipid metabolism, immune function, and endosome vesicle cycling⁷⁰. The lipids that have shown marked differences between AD versus controls which been directly and indirectly linked to at least one of these mechanisms. The significant associations between lipid profiles and AD which were observed in our study encouraged us to further explore the possible mechanisms of lipids changes in AD patients.

Our results highlight that genetic pathway and SNPs related to AD influence lipids that are associated with AD. These results reinforce the importance of lipid metabolism and dysregulation in AD. It also suggests inflammation, ion channel modification, and A β pathways influence lipid levels that are dysregulated in AD. While studies examining the exact mechanisms linking these SNPs with reported lipid changes in AD patients are nascent in the current literature, our results provide renewed insights on potential lipid-related mechanisms in AD which may be examined further in preclinical and clinical studies.

Limitations

Firstly, the sample size was around 40 each in the AD and paired control group, which provides enough power for analyzing 9 lipid classes. However, the strength of the data is limited for examining a large number of individual lipids. FDR correction was used to reduce the bias, although a larger cohort is needed to replicate the findings.

Secondly, we did not subclassify AD to different stages. In addition, we did not look at overlapping cerebral vascular disease in AD patients because only 10 of 40 AD subjects had MRI scanning. Therefore, future studies will look at lipid differences between control, pure AD, and mixed AD in the same stage.

Thirdly, our study is useful for exploring clinical biomarkers of AD because plasma lipids are relatively easy to extract. However, plasma lipids may also be affected by damage to peripheral organs as well, which limits the accuracy of any association between changes in plasma lipid levels and brain pathology. The lipidomics of CSF or post-mortem brain samples from AD patients will be necessary to validate our findings in the future.

Conclusion

Our findings suggest that plasma lipids may differentiate between AD and cognitively normal controls and lipids models can be applied to discriminate the two groups. In addition, AD-related SNPs may have a different association with lipids between AD and controls and the underlying mechanisms need to be explored further in the future. In summary, our study provides evidence that certain lipids may be involved in AD pathogenesis and are associated with AD-related SNPs.

Author details

¹Centre for Healthy Brain Ageing (CHEBA), School of Psychiatry, University of New South Wales, Sydney, Australia. ²Guangdong Mental Health Center, Guangdong Provincial People's Hospital, Guangdong Academy of Medical Sciences, Guangzhou, Guangdong, China. ³Neuroscience Research Australia, Randwick, Australia. ⁴Mark Wainwright Analytical Centre, University of New South Wales, Sydney, Australia. ⁵Neuropsychiatric Institute, Euroa Centre, Prince of Wales Hospital, Sydney, Australia. ⁶School of Medicine, Huzhou University, Huzhou Central Hospital Huzhou, Huzhou, China

Conflict of interest

The authors declare no competing interests.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41398-021-01362-2>.

Received: 26 June 2020 Revised: 10 March 2021 Accepted: 6 April 2021
Published online: 07 June 2021

References

1. Reitz C, Brayne C, Mayeux R. Epidemiology of Alzheimer disease. *Nat. Rev. Neurol.* **7**, 137–152 (2011).
2. Smith M, A., Rottkamp C, A., Nunomura A, Raina A, K. & Perry G. Oxidative stress in Alzheimer's disease. *Biochim. Biophys. Acta* **1502**, 139–144 (2000).
3. Heppner F, L., Ransohoff R, M. & Becher B. Immune attack: the role of inflammation in Alzheimer disease. *Nat. Rev. Neurosci.* **16**, 358–372 (2015).
4. Xu J, et al. Graded perturbations of metabolism in multiple regions of human brain in Alzheimer's disease: Snapshot of a pervasive metabolic disorder. *Biochim. Biophys. Acta* **1862**, 1084–1092 (2016).
5. Morris J, K., Honea R, A., Vidoni E, D., Swerdlow R, H. & Burns J, M. Is Alzheimer's disease a systemic disease? *Biochim. Biophys. Acta* **1842**, 1340–1349 (2014).
6. Dart C. Lipid microdomains and the regulation of ion channel function. *J. Physiol.* **588**, 3169–3178 (2010).
7. Brugger B. Lipidomics: analysis of the lipid composition of cells and sub-cellular organelles by electrospray ionization mass spectrometry. *Annu. Rev. Biochem.* **83**, 79–98 (2014).

8. Walter, A. et al. Glycerophosphocholine is elevated in cerebrospinal fluid of Alzheimer patients. *Neurobiol. Aging* **25**, 1299–1303 (2004).
9. Mapstone, M. et al. Plasma phospholipids identify antecedent memory impairment in older adults. *Nat. Med.* **20**, 415–418 (2014).
10. Casanova, R. et al. Blood metabolite markers of preclinical Alzheimer's disease in two longitudinally followed cohorts of older individuals. *Alzheimers Dement.* **12**, 815–822 (2016).
11. Sachdev, P. S. et al. The Sydney Memory and Ageing Study (MAS): methodology and baseline medical and neuropsychiatric characteristics of an elderly epidemiological non-demented cohort of Australians aged 70–90 years. *Int. Psychogeriatr.* **22**, 1248–1264 (2010).
12. Wong, M. W. K. et al. Plasma lipidome variation during the second half of the human lifespan is associated with age and sex but minimally with BML. *PLoS ONE* **14**, e0214141 (2019).
13. Dahlgren, K. N. et al. Oligomeric and fibrillar species of amyloid-beta peptides differentially affect neuronal viability. *J. Biol. Chem.* **277**, 32046–32053 (2002).
14. Meng, X. et al. Association between increased levels of amyloid-beta oligomers in plasma and episodic memory loss in Alzheimer's disease. *Alzheimers Res Ther.* **11**, 89 (2019).
15. Wang, M. J. et al. Oligomeric forms of amyloid-beta protein in plasma as a potential blood-based biomarker for Alzheimer's disease. *Alzheimers Res. Ther.* **9**, 98 (2017).
16. Mather, K. A. et al. Investigating the genetics of hippocampal volume in older adults without dementia. *PLoS ONE* **10**, e0116920 (2015).
17. Mather, K. A. et al. Genome-wide significant results identified for plasma apolipoprotein H levels in middle-aged and older adults. *Sci. Rep.* **6**, 23675 (2016).
18. Euesden, J., Lewis, C. M. & O'Reilly, P. F. PRSice: Polygenic Risk Score software. *Bioinformatics* **31**, 1466–1468 (2015).
19. Lambert, J. C. et al. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nat. Genet.* **45**, 1452–1458 (2013).
20. Apostolova, L. G. et al. Associations of the Top 20 Alzheimer Disease Risk Variants With Brain Amyloidosis. *JAMA Neurol.* **75**, 328–341 (2018).
21. Friedman, J., Hastie, T. & Tibshirani, R. Regularization paths for generalized linear models via coordinate descent. *J. Stat. Softw.* **33**, 1–22 (2010).
22. Team, R. C. R: A language and environment for statistical computing <https://www.R-project.org/> (2018).
23. Kuhn, M. et al. Caret: Classification and Regression Training. R package version. (2018).
24. Robin, X. et al. pROC: an open-source package for R and S+ to analyze and compare ROC curves. *BMC Bioinforma.* **12**, 77 (2011).
25. Podlisny, M. B. et al. Aggregation of secreted amyloid beta-protein into sodium dodecyl sulfate-stable oligomers in cell culture. *J. Biol. Chem.* **270**, 9564–9570 (1995).
26. Seubert, P. et al. Isolation and quantification of soluble Alzheimer's beta-peptide from biological fluids. *Nature* **359**, 325–327 (1992).
27. Olsen, A. S. B. & Faergeman, N. J. Sphingolipids: membrane microdomains in brain development, function and neurological diseases. *Open Biol.* **7**, <https://doi.org/10.1098/rsob.170069> (2017).
28. Mielke, M. M. et al. Plasma ceramides are altered in mild cognitive impairment and predict cognitive decline and hippocampal volume loss. *Alzheimers Dement.* **6**, 378–385 (2010).
29. Mielke, M. M. & Lyketsos, C. G. Alterations of the sphingolipid pathway in Alzheimer's disease: new biomarkers and treatment targets? *Neuromol. Med.* **12**, 331–340 (2010).
30. Varma, V. R. et al. Brain and blood metabolite signatures of pathology and progression in Alzheimer disease: a targeted metabolomics study. *PLoS Med.* **15**, e1002482 (2018).
31. Han, X. et al. Metabolomics in early Alzheimer's disease: identification of altered plasma sphingolipidome using shotgun lipidomics. *PLoS ONE* **6**, e21643 (2011).
32. Hussain, G. et al. Role of cholesterol and sphingolipids in brain development and neurological diseases. *Lipids Health Dis.* **18**, 26 (2019).
33. Li, Z. et al. Reducing plasma membrane sphingomyelin increases insulin sensitivity. *Mol. Cell Biol.* **31**, 4205–4218 (2011).
34. Whitley, L. et al. Evidence of altered phosphatidylcholine metabolism in Alzheimer's disease. *Neurobiol. Aging* **35**, 271–278 (2014).
35. Atri, A. Imaging of neurodegenerative cognitive and behavioral disorders: practical considerations for dementia clinical practice. *Handb. Clin. Neurol.* **136**, 971–984 (2016).
36. Ojo, J. O. et al. Converging and differential brain phospholipid dysregulation in the pathogenesis of repetitive mild traumatic brain injury and Alzheimer's disease. *Front. Neurosci.* **13**, 103 (2019).
37. Arancio, O. PIP2: a new key player in Alzheimer's disease. *Cellscience* **5**, 44–47 (2008).
38. Sabogal-Guaqueta, A. M., Villamil-Ortiz, J. G., Arias-Londono, J. D. & Cardona-Gomez, G. P. Inverse phosphatidylcholine/phosphatidylinositol levels as peripheral biomarkers and phosphatidylcholine/lysophosphatidylethanolamine-phosphatidylserine as hippocampal indicator of posts ischemic cognitive impairment in rats. *Front. Neurosci.* **12**, 989 (2018).
39. Gerl, M. J. et al. Cholesterol is inefficiently converted to cholesteryl esters in the blood of cardiovascular disease patients. *Sci. Rep.* **8**, 14764 (2018).
40. Proitsi, P. et al. Plasma lipidomics analysis finds long chain cholesteryl esters to be associated with Alzheimer's disease. *Transl. Psychiatry* **5**, e494 (2015).
41. Hejazi, L. et al. Mass and relative elution time profiling: two-dimensional analysis of sphingolipids in Alzheimer's disease brains. *Biochem. J.* **438**, 165–175 (2011).
42. Wood, P. L. et al. Targeted lipidomics of frontal cortex and plasma diacylglycerols (DAG) in mild cognitive impairment and Alzheimer's disease: validation of DAG accumulation early in the pathophysiology of Alzheimer's disease. *J. Alzheimer's Dis.* **48**, 537–546 (2015).
43. Shin, J., Xie, D. & Zhong, X. P. MicroRNA-34a enhances T cell activation by targeting diacylglycerol kinase zeta. *PLoS ONE* **8**, e77983 (2013).
44. Kalmijn, S. et al. Metabolic cardiovascular syndrome and risk of dementia in Japanese-American elderly men. The Honolulu-Asia aging study. *Arterioscler. Thromb. Vasc. Biol.* **20**, 2255–2260 (2000).
45. Tajima, Y. et al. Lipidomic analysis of brain tissues and plasma in a mouse model expressing mutated human amyloid precursor protein/tau for Alzheimer's disease. *Lipids Health Dis.* **12**, 68 (2013).
46. Ben Haim, L. & Rowitch, D. H. Functional diversity of astrocytes in neural circuit regulation. *Nat. Rev. Neurosci.* **18**, 31–41 (2017).
47. Jeong, W., Lee, H., Cho, S. & Seo, J. ApoE4-induced cholesterol dysregulation and its brain cell type-specific implications in the pathogenesis of Alzheimer's disease. *Mol. Cells* **42**, 739–746 (2019).
48. Zhang, J. & Liu, Q. Cholesterol metabolism and homeostasis in the brain. *Protein Cell* **6**, 254–264 (2015).
49. Oksanen, M. et al. Astrocyte alterations in neurodegenerative pathologies and their modeling in human induced pluripotent stem cell platforms. *Cell. Mol. Life Sci.* **76**, 2739–2760 (2019).
50. Shulman, J. M. et al. Functional screening in *Drosophila* identifies Alzheimer's disease susceptibility genes and implicates Tau-mediated mechanisms. *Hum. Mol. Genet.* **23**, 870–877 (2014).
51. Oksala, N. et al. Kindlin 3 (FERMT3) is associated with unstable atherosclerotic plaques, anti-inflammatory type II macrophages and upregulation of beta-2 integrins in all major arterial beds. *Atherosclerosis* **242**, 145–154 (2015).
52. Ellison, J. A. et al. Osteopontin and its integrin receptor alpha(v)beta3 are upregulated during formation of the glial scar after focal stroke. *Stroke* **29**, 1698–1706 (1998). discussion 1707.
53. Chapuis, J. et al. Genome-wide, high-content siRNA screening identifies the Alzheimer's genetic risk factor FERMT2 as a major modulator of APP metabolism. *Acta Neuropathol.* **133**, 955–966 (2017).
54. Eysert, F. et al. Alzheimer's genetic risk factor FERMT2 (Kindlin-2) controls axonal growth and synaptic plasticity in an APP-dependent manner. *Mol. Psychiatry* <https://doi.org/10.1038/s41380-020-00926-w> (2020).
55. Sosa, L. J. et al. Amyloid precursor protein is an autonomous growth cone adhesion molecule engaged in contact guidance. *PLoS ONE* **8**, e64521 (2013).
56. Ramirez, L. M. et al. Common variants in ABCA7 and MS4A6A are associated with cortical and hippocampal atrophy. *Neurobiol. Aging* **39**, 82–89 (2016).
57. Aikawa, T., Holm, M. L. & Kanekiyo, T. ABCA7 and pathogenic pathways of Alzheimer's disease. *Brain Sci.* **8**, <https://doi.org/10.3390/brainsci8020027> (2018).
58. Wang, X. & Paigen, B. Genetics of variation in HDL cholesterol in humans and mice. *Circ. Res.* **96**, 27–42 (2005).
59. Lambert, J. C. et al. Genome-wide association study identifies variants at CLU and CR1 associated with Alzheimer's disease. *Nat. Genet.* **41**, 1094–1099 (2009).
60. Sims, R. et al. Rare coding variants in PLCG2, ABI3, and TREM2 implicate microglial-mediated innate immunity in Alzheimer's disease. *Nat. Genet.* **49**, 1373–1384 (2017).

61. MahmoudianDehkordi, S. et al. Altered bile acid profile associates with cognitive impairment in Alzheimer's disease—An emerging role for gut microbiome. *Alzheimers Dement.* **15**, 76–92 (2019).
62. Walker, D. G., Tang, T. M. & Lue, L. F. Studies on colony stimulating factor receptor-1 and ligands colony stimulating factor-1 and interleukin-34 in Alzheimer's disease brains and human microglia. *Front. Aging Neurosci.* **9**, 244 (2017).
63. Hollingworth, P. et al. Common variants at ABCA7, MS4A6A/MS4A4E, EPHA1, CD33 and CD2AP are associated with Alzheimer's disease. *Nat. Genet.* **43**, 429–435 (2011).
64. Slot, R. E. R. et al. ApoE and clusterin CSF levels influence associations between APOE genotype and changes in CSF tau, but not CSF Abeta42, levels in non-demented elderly. *Neurobiol. Aging* **79**, 101–109 (2019).
65. Adeyemo, A. et al. A genome-wide association study of hypertension and blood pressure in African Americans. *PLoS Genet.* **5**, e1000564 (2009).
66. Jacobsen, L. et al. Activation and functional characterization of the mosaic receptor SorLA/LR11. *J. Biol. Chem.* **276**, 22788–22796 (2001).
67. Scherzer, C. R. et al. Loss of apolipoprotein E receptor LR11 in Alzheimer disease. *Arch. Neurol.* **61**, 1200–1205 (2004).
68. Rogaeva, E. et al. The neuronal sortilin-related receptor SORL1 is genetically associated with Alzheimer disease. *Nat. Genet.* **39**, 168–177 (2007).
69. Gao, X. et al. SORL1 genetic variants modulate risk of amnesic mild cognitive impairment in northern Han Chinese. *Int. J. Neurosci.* **124**, 296–301 (2014).
70. Van Cauwenberghe, C., Van Broeckhoven, C. & Sleegers, K. The genetic landscape of Alzheimer disease: clinical implications and perspectives. *Genet. Med.* **18**, 421–430 (2016).