



Correction to: A novel method for alveolar bone grafting assessment in cleft lip and palate patients: cone-beam computed tomography evaluation

Marcin Stasiak¹ · Anna Wojtaszek-Słomińska¹ · Bogna Racka-Pilszak¹

Published online: 15 September 2020

© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: *Clinical Oral Investigations*

<https://doi.org/10.1007/s00784-020-03505-z>

In figure 2, the description of the 3rd score has been amended:

From: Thickness of the alveolar bone bridge is more than the labiolingual width of central incisor's root.

To: Thickness of the alveolar bone bridge amounts to at least the labiolingual width of central incisor's root.

Below is the corrected figure 2:

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Score	CBCT cross-sectional image (cleft side is on the left side)	Description
0		No alveolar bone bridge.
1		Thickness of the alveolar bone bridge < 1/2 of the labiolingual width of central incisor's root.
2		Thickness of the alveolar bone bridge \geq 1/2 of the labiolingual width of central incisor's root and less than the labiolingual width of central incisor's root.
3		Thickness of the alveolar bone bridge amounts to at least the labiolingual width of central incisor's root.

The online version of the original article can be found at <https://doi.org/10.1007/s00784-020-03505-z>

✉ Marcin Stasiak
marcinstasiak@gumed.edu.pl

¹ Department of Orthodontics, Faculty of Medicine, Medical University of Gdańsk, Aleja Zwycięstwa 42c, 80-210 Gdańsk, Poland