

Different Characteristics of Serum Alfa Fetoprotein and Serum Des-gamma-carboxy Prothrombin in Resected Hepatocellular Carcinoma

MASAMICHI HAYASHI¹, SUGURU YAMADA¹, NAO TAKANO¹, YUKIYASU OKAMURA²,
HIDEKI TAKAMI¹, YOSHIKUNI INOKAWA¹, FUMINORI SONOHARA¹, NOBUTAKE TANAKA¹,
DAI SHIMIZU¹, NORIFUMI HATTORI¹, MITSURO KANDA¹, CHIE TANAKA¹,
GORO NAKAYAMA¹, MASAHIKO KOIKE¹ and YASUHIRO KODERA¹

¹Department of Gastroenterological Surgery, Nagoya University Graduate School of Medicine, Nagoya, Japan;
²Division of Hepato-Biliary-Pancreatic Surgery, Shizuoka Cancer Center, Shizuoka, Japan

Abstract. *Background/Aim:* Hepatocellular carcinoma (HCC) mainly develops in the damaged liver from hepatitis C virus (HCV) or hepatitis B virus (HBV) infection in Japan. On the other hand, the occurrence of HCCs derived from the liver without viral infection has recently been increasing. Our aim was to identify characteristics specific to HCCs with virus-infected liver (HCC-BC) or those with non-B- and non-C-infected liver (HCC-NBNC). *Patients and Methods:* We collected preoperative serum α -fetoprotein (AFP) and Des-Gamma-Carboxy Prothrombin (DCP), also known as PIVKA-II values from surgically resected HCC cases during 1994-2017 in our department. *Results:* Preoperative serum AFP values of HCC-BC cases ($n=284$) were higher compared to HCC-NBNC cases ($n=88$) ($p=0.016$), whereas serum DCP values of HCC-NBNC cases were higher compared to HCC-BC cases ($p<0.001$). Multivariable analyses indicated that abnormal serum AFP [hazard ratio (HR)=1.46, 95% confidence interval (CI)=1.03-2.07, $p=0.035$] was one of the significant recurrence-free survival predictors of HCC-BC cases, while abnormal serum DCP (HR=4.99, 95%CI=1.91-13.01, $p=0.001$) was one of the significant recurrence-free survival predictors of HCC-NBNC cases. *Conclusion:* HCC-NBNC cases have a different tumor marker profile from HCC-BC

cases. Elevated DCP could be both a diagnostic and prognostic marker of HCC-NBNC patients.

Hepatocellular carcinoma (HCC) is the 6th most frequently occurring cancer globally and still has a high likelihood of recurrence and a poor prognosis (1). HCCs are mainly derived from the damaged liver caused by various etiological factors, including hepatitis C virus (HCV) or hepatitis B virus (HBV) infection, as well as chronic alcohol abuse (2, 3). Among them, HCV (65%) and HBV (15%) are the two major pathogenic factors in Japan (4). Recently, the occurrence of HCCs derived from non-B non-C livers (HCC-NBNC) have been relatively increasing because HBV or HCV treatments have dramatically improved. HCC-NBNC lesions typically arise from non-alcoholic steatohepatitis (NASH) or alcoholic liver disease.

To characterize the background liver status, whole-genome analyses have been widely performed (5, 6). Some mutational signatures and altered pathways have been associated with certain histological characteristics of background livers or tumor stages (7, 8). For instance, the mutation of catenin beta 1 (*CTNNB1*), one of the critical cluster of Wnt-signaling, has been related to alcohol-damaged liver. Telomerase reverse transcriptase (*TERT*), cyclin dependent kinase inhibitor 2A (*CDKN2A*), SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 2 (*SMARCA2*) and hepatocyte growth factor (*HGF*) alterations are also enriched in alcohol-related HCC patients. Tumor protein p53 (*TP53*) mutations are frequently associated with HBV infection. The integration of HBV into the host genome (9, 10) induces upregulation of cancer-related genes, such as *TERT*, lysine methyltransferase 2B (*MLL4*), and cyclin E1 (*CCNE1*) genes. This leads to alterations in the genes functioning downstream of all these genes or cause whole genome

This article is freely accessible online.

Correspondence to: Suguru Yamada, Department of Gastroenterological Surgery, Nagoya University Graduate School of Medicine, 65 Tsurumai-cho, Showa-ku, Nagoya, 466-8550, Japan. Tel: +41 0527442249, Fax: +41 0527442252, e-mail: suguru@med.nagoya-u.ac.jp

Key Words: AFP, DCP, hepatocellular carcinoma, survival predictor.

chromosomal instability (10, 11). Concerning the HCC-NBNC and background liver, Kutlu *et al.* have reported several molecular characteristics (12), including a patatin-like phospholipase domain containing 3 (*PNPLA3*) gene mutation, epigenetic changes of phosphodiesterase 1B (*PDE1B*) and chromodomain helicase DNA-binding protein 1 (*CHD1*), micro RNA deregulation including miR-122, metabolic pathway activating insulin receptor signaling and mitochondrial dysfunction caused by reactive oxygen species and endoplasmic reticulum stress.

We hypothesized that some molecular characteristics distinguishing HCC-NBNC from HCC with virus-infected liver (HCC-BC) may affect the positivity of well-known tumor markers of HCC, such as alpha-fetoprotein (AFP) and des-gamma-carboxy prothrombin (DCP) (13). In this study, we used the HCC resection cohort in our institution and retrospectively compared HCC-NBNC cases with HCC-BC cases from the viewpoint of these well-known HCC serum tumor markers.

Patients and Methods

Patient cohort. Among surgically resected HCC cases from 1994 to 2017 in the Department of Gastroenterological Surgery, at Nagoya University (Aichi, Japan), 372 cases with available preoperative AFP and DCP markers were included (Institute Review Board approval number: 2013-0295). Of these, 284 patients were categorized as HCC-BC and 88 patients as HCC-NBNC. The average follow-up period was 51.4 months. Clinical factors including age, gender, liver damage scores, tumor size and numbers, and pathological factors of tumor differentiation, growth pattern, capsule formation, serosal and vascular invasion were categorically compared between the two groups.

Serum marker collection. Each serum marker was checked by peripheral blood examination preoperatively. The standard institutional cut-off values were 10 ng/ml for AFP and 40 mAU/ml for DCP.

Statistical analysis. Patient clinicopathological characteristics were compared using Fisher's exact test for categorical variables and Mann-Whitney *U*-test for continuous variables. Overall survival (OS) was defined as the time from surgery to the date of HCC disease-related death. Recurrence-free survival (RFS) was defined as the time from surgery to the date of recurrence diagnosis. Those who remained alive were censored at the last date they were known to be alive. A log-rank test was applied to compare the survival outcomes of the two groups. The Cox proportional hazards model was used for univariate and multivariable analysis for survival outcomes. All tests were considered statistically significant and clinically promising at $p < 0.05$. Statistical analyses were carried out using the JMP 15 software (SAS Institute Japan, Tokyo, Japan).

Results

Patients characteristics. Clinicohistological characteristics of both HCC-BC cases (n=284) and HCC-NBNC cases (n=88) are shown in Table I. Due to the viral hepatic damage, liver damage score B/C cases were more frequently

Table I. Patient characteristics.

Factors	HCC-BC (n=284)	HCC-NBNC (n=88)	p-Value
Age			0.582
≥60	207	67	
<60	77	21	
Gender			0.348
Female	56	13	
Male	228	75	
Liver Damage			0.071
A	218	76	
B/C	66	12	
Tumor number			0.121
Single	207	72	
Multiple	77	16	
Tumor size			<0.001
≥2.0 cm	151	70	
<2.0 cm	133	18	
Differentiation			0.753
Well	50	17	
Moderate/Poor	230	71	
Unknown	4	0	
Growth pattern			0.197
Expansive	236	67	
Invasive	45	19	
Unknown	3	2	
Capsule formation			0.896
Positive	192	58	
Negative	92	29	
Unknown	0	1	
Infiltration to capsule			0.806
Positive	155	46	
Negative	128	41	
Unknown	1	1	
Septal formation			0.294
Positive	182	61	
Negative	97	24	
Unknown	5	3	
Serosal invasion			0.045
Positive	48	25	
Negative	202	57	
Unknown	34	6	
Portal vein invasion			1.000
Positive	56	17	
Negative	226	69	
Unknown	2	2	
Hepatic vein invasion			0.027
Positive	30	18	
Negative	248	68	
Unknown	6	2	
LCSGJ stage			1.000
I-II	178	55	
III-IV	105	33	
Unknown	1	0	
Liver cirrhosis			<0.001
Positive	126	19	
Negative	158	69	
AFP			0.061
≤10 ng/ml	107	43	
>10 ng/ml	171	43	
Unknown	6	2	
DCP			<0.001
≤40 mAU/ml	139	23	
>40 mAU/ml	115	62	
Unknown	30	3	

HCC-BC: Hepatocellular carcinoma with virus-infected liver; HCC-NBNC: hepatocellular carcinoma with no virus-infected liver; LCSGJ: Liver Cancer Study Group of Japan; AFP: α-fetoprotein; DCP: des-gamma-carboxy prothrombin. Significant *p*-Values are shown in bold.




Figure 1. Preoperative AFP and DCP values of HCC-BC and HCC-NBNC cases. HCC-BC cases ($n=284$) had significantly higher AFP values compared to the NBNC cohort, while HCC-NBNC cases ($n=88$) had significantly higher DCP values compared to the BC cohort. AFP: Alpha-fetoprotein; DCP: des-gamma-carboxy prothrombin; HCC-BC: hepatocellular carcinoma with virus-infected liver; HCC-NBNC: hepatocellular carcinoma with no virus-infected liver.

found in HCC-BC rather than in HCC-NBNC cases ($p=0.071$). Histologically advanced cases with large diameter ($p<0.001$), serosal invasion ($p=0.045$) and hepatic vein invasion ($p=0.027$) were frequently found in HCC-NBNC cases, while the cancer stage distributions of Liver Cancer Study Group of Japan (LCSGJ) between the two groups were comparable ($p=1.000$). The distribution of actual serum values for AFP and DCP were compared between HCC-BC and HCC-NBNC cases, as depicted in Figure 1. AFP values were inclined to exceed the cut-off value in HCC-BC cases ($p=0.061$), whereas DCP values were significantly higher in HCC-NBNC cases compared to HCC-BC cases ($p<0.001$).

Serum tumor marker and survival outcomes. We compared high and low tumor marker cases based on the cut-off values in each HCC-BC and HCC-NBNC cohort to ascertain the markers' impact on postoperative RFS and OS. With regards to RFS (Figure 2), cases with aberrantly high values of tumor markers showed significantly poor survival outcomes in both cohorts. Concerning OS (Figure 3), high AFP was associated with a significantly poor prognosis in the HCC-BC cohort. In contrast, patients with high DCP had significantly lower OS in both cohorts, with a vast difference in OS between high and low values in the HCC-NBNC cohort. Then, we compared AFP high with AFP low (Table II) as well as DCP high with DCP low (Table III) in the HCC-BC and HCC-NBNC cohorts to examine the

characteristics associates with these values in detail. High AFP cases were related to aged people, with i) moderate or poor differentiation, ii) portal vein invasion, iii) advanced tumor stage and iv) positive liver cirrhosis, while high AFP cases were also specific to the HCC-NBNC cohort with both i) portal vein invasion and ii) advanced tumor stage. On the contrary, high DCP cases were significantly correlated with HCC-BC cases with i) a large tumor size, ii) moderate or poor differentiation, iii) infiltration to a capsule, iv) serosal invasion, v) vascular invasion, vi) advanced tumor stage and vii) liver cirrhosis. Also, they were associated with HCC-NBNC with i) large tumor size and ii) moderate or poor differentiation.

Univariate and multivariable analyses of survival outcomes.

Univariate and multivariable analyses of survival outcomes were performed. All significant factors in the univariate analysis were put into the multivariable analysis. The backward stepwise method was performed until the p -Values of all remaining factors became significant. Tables IV and V summarize the results of RFS in the HCC-BC and HCC-NBNC cohorts. In HCC-BC cases, i) tumor size, ii) AFP elevation, iii) serosal invasion, iv) portal vein invasion and v) hepatic vein invasion were detected as significant prognostic factors of RFS in multivariable analysis. On the other hand, i) DCP elevation and ii) portal vein invasion were significant factors in HCC-NBNC cases.




Figure 2. Recurrence-free survival curves (RFS). RFS was compared between high AFP cases (AFP>10ng/ml) and low AFP cases, as well as high DCP cases (DCP>40 mAU/ml) and low DCP cases in both HCC-BC and HCC-NBNC cohorts. Both serum markers indicated significantly poor survival outcomes in both cohorts. AFP: Alpha-fetoprotein; DCP: des-gamma-carboxy prothrombin; HCC-BC: hepatocellular carcinoma with virus-infected liver; HCC-NBNC: hepatocellular carcinoma with no virus-infected liver.

Tables VI and VII demonstrate the results of OS in each cohort. In the multivariable analysis of HCC-BC cases i) tumor number, ii) serosal invasion, iii) portal vein invasion and iv) hepatic vein invasion were significant predictors. In contrast, i) DCP elevation was an extremely significant predictor of HCC-NBNC cases in addition to ii) serosal invasion and iii) portal vein invasion. None of the low DCP cases died from the disease in our cohort.

Clinical characteristics of AFP and DCP elevation. AFP values of HCC-BC cases increased depending on tumor T stage, while DCP values of HCC-NBNC cases increased

depending on the T stage (Figure 4). Besides, the association of both markers with background liver are shown in Figure 5. AFP does not decrease in the cirrhotic liver, while DCP decreases in them.

Discussion

Clinically, the measurement of both AFP and DCP has been strongly recommended in the Clinical Practice Guidelines for Hepatocellular Carcinoma (14); however, the mechanism of each tumor marker elevation is unknown and may differ between tumor types. HCCs derived from NBNC are




Figure 3. Overall survival (OS). OS curves were compared between high AFP cases (AFP>10ng/ml) and low AFP cases, and high DCP cases (DCP>40 mAU/ml) and low DCP cases in the HCC-BC cohort and HCC-NBNC cohort, respectively. High AFP indicated significantly poor survival outcomes in the HCC-BC cohort, while high DCP displayed significantly poor survival outcomes in both cohorts. AFP: Alpha-fetoprotein; DCP: des-gamma-carboxy prothrombin; HCC-BC: hepatocellular carcinoma with virus-infected liver; HCC-NBNC: hepatocellular carcinoma with no virus-infected liver.

reported to have relatively low serum AFP levels compared to hepatitis B-derived HCCs (15). Also, hepatitis C-infected livers usually have high serum AFP levels (16). These findings suggest that AFP elevation is commonly influenced by a viral infection of the background liver. AFP is a glycoprotein derived from the embryonic endoderm. It is closely related to the growth of malignant tumors (17). During embryonic development, AFP is initially produced in the fetal liver and yolk sac. The serum AFP concentration increases during the period between 12-16 weeks of gestation and then it gradually reduces to normal range till adulthood (18). AFP increases again during early stages of

hepatocytes' malignant transformation, and it is activated in the malignant cells. Zheng Y *et al.*, have summarized the AFP production mechanism in HBV-derived hepatitis-based HCCs (17), where the HBV X protein promotes the acceleration of AFP's accretion, which induces growth signal activation, metastases and bears an immunosuppressive role.

Instead, DCP is abnormal prothrombin and produced due to the defect of the post-translational carboxylation of prothrombin's precursor (19); however, the detailed mechanism of its production is unclear. Taniguchi T *et al.* have used mass spectrometry analysis of hepatoma cell lines to reveal that PARP-1 activates prothrombin gene

Table II. Clinico-histological features of AFP high cases.

Factors	HCC-BC		p-Value	HCC-NBNC		p-Value
	AFP high	AFP low		AFP high	AFP low	
Age			0.018			1.000
≥60	116	87		32	33	
<60	55	20		11	10	
Gender			0.759			0.549
Female	35	20		35	38	
Male	136	87		8	5	
Liver damage			0.773			0.351
A	132	81		39	35	
B/C	39	26		4	8	
Tumor number			0.332			1.000
Single	121	82		35	35	
Multiple	50	25		8	8	
Tumor size			0.462			0.792
≥2.0 cm	88	60		33	35	
<2.0 cm	83	47		10	8	
Differentiation			0.006			0.103
Well	21	28		5	12	
Moderate/Poor	147	78		38	31	
Growth pattern			0.739			0.186
Expansive	139	91		31	35	
Invasive	29	16		12	6	
Capsule formation			0.114			0.818
Positive	122	66		29	28	
Negative	49	41		13	15	
Infiltration to capsule			0.047			0.517
Positive	101	50		24	21	
Negative	69	57		18	22	
Septal formation			0.796			1.000
Positive	110	66		30	29	
Negative	59	38		12	12	
Serosal invasion			0.415			0.144
Positive	31	17		15	9	
Negative	113	84		24	32	
Portal vein invasion			0.019			0.003
Positive	41	13		14	3	
Negative	129	93		27	40	
Hepatic vein invasion			0.555			0.113
Positive	20	10		12	6	
Negative	145	97		29	37	
LCSGJ stage			0.040			0.026
I-II	99	76		21	32	
III-IV	71	31		22	11	
Liver cirrhosis			0.047			0.604
Positive	84	39		11	8	
Negative	87	68		32	35	
DCP			0.796			0.465
≤40 mAU/ml	80	56		13	9	
>40 mAU/ml	68	44		30	31	

HCC-BC: Hepatocellular carcinoma with virus-infected liver; HCC-NBNC: hepatocellular carcinoma with no virus-infected liver; LCSGJ: Liver Cancer Study Group of Japan; AFP: α-fetoprotein; DCP: des-gamma-carboxy prothrombin. Significant p-Values are shown in bold.

transcription and that this excessive transcription induces DCP production (20). PARP-1 inhibition is also reported as a candidate therapeutic strategy for hepatic triglyceride

accumulation, metabolic dysregulation, inflammation and fibrosis in mouse NASH models (21). DCP elevation reflects vascular invasion and tumor recurrences following

Table III. *Clinicohistological features of DCP high cases.*

Factors	HCC-BC		<i>p</i> -Value	HCC-NBNC		<i>p</i> -Value
	DCP high	DCP low		DCP high	DCP low	
Age			0.267			1.000
≥60	86	94		15	6	
<60	29	45		47	17	
Gender			0.204			0.742
Female	18	31		9	4	
Male	97	108		53	19	
Liver damage			0.174			0.727
A	94	103		54	19	
B/C	21	36		8	4	
Tumor number			0.162			0.750
Single	77	105		50	20	
Multiple	38	34		12	3	
Tumor size			<0.001			0.002
≥2.0 cm	85	60		55	13	
<2.0 cm	30	79		7	10	
Differentiation			<0.001			0.013
Well	9	35		8	9	
Moderate/Poor	104	103		54	14	
Growth pattern			0.736			0.771
Expansive	97	112		48	17	
Invasive	18	24		13	6	
Capsule formation			0.285			0.439
Positive	81	88		43	14	
Negative	34	51		18	9	
Infiltration to capsule			0.043			0.469
Positive	69	66		35	11	
Negative	45	73		26	12	
Septal formation			0.227			0.268
Positive	77	86		46	14	
Negative	33	53		14	8	
Serosal invasion			0.013			0.177
Positive	30	16		21	4	
Negative	79	102		37	18	
Portal vein invasion			0.001			1.000
Positive	33	17		12	4	
Negative	81	122		48	19	
Hepatic vein invasion			<0.001			0.134
Positive	22	7		16	2	
Negative	89	132		44	21	
LCSGJ stage			<0.001			0.459
I-II	55	101		37	16	
III-IV	60	38		25	7	
Liver cirrhosis			0.030			0.379
Positive	40	68		12	7	
Negative	75	71		50	16	

HCC-BC: Hepatocellular carcinoma with virus-infected liver; HCC-NBNC: hepatocellular carcinoma with no virus-infected liver; LCSGJ: liver cancer study group of Japan; AFP: α -fetoprotein; DCP: des-gamma-carboxy prothrombin. Significant *p*-Values are shown in bold.

hepatectomy (22). It has also been reported to increase during epithelial to mesenchymal transition in tumors (23). In other words, DCP goes up by tumor factors.

Interestingly, Suzuki H *et al.*, have reported that mild hypoxia induces HCC to produce DCP, while long-lasting hypoxia impaires DCP production in HCC cells (23), which

could partly explain why DCP is elevated in HCC-NBNCs rather than in HCC-BCs. In our study tumor sizes of HCC-NBNCs were significantly larger than HCC-BCs because no intensive follow-up examination was usually performed for NBNC patients. The relatively large HCC-NBNCs sometimes induce intratumoral hypoxia, which is easy to

Table IV. Univariate and multivariable analyses of RFS in HCC-BC cases.

Clinicopathological factors		Univariate analysis			Multivariable analysis		
		HR	95%CI	p-Value	HR	95%CI	p-Value
Age	≥65 years	1.00	0.71-1.41	0.983			
Gender	Male	1.34	0.88-2.02	0.169			
Tumor number	Multiple	1.66	1.20-2.31	0.003			
Tumor size	≥2.0 cm	1.89	1.37-2.61	<0.001	1.66	1.15-2.39	0.007
AFP	≥10 ng/ml	1.42	1.02-1.96	0.036	1.46	1.03-2.07	0.035
DCP	≥40 ng/ml	1.72	1.25-2.38	0.001			
Differentiation	Poor, Moderate	1.32	0.87-2.02	0.193			
Growth form	Infiltrative	1.70	1.15-2.52	0.008			
Serosal invasion	Positive	2.46	1.67-3.62	<0.001	1.94	1.30-2.89	0.001
Portal vein invasion	Positive	2.34	1.63-3.34	<0.001	1.88	1.26-2.81	0.002
Hepatic vein invasion	Positive	2.99	1.87-4.78	<0.001	2.67	1.65-4.32	<0.001
Liver cirrhosis	Present	1.09	0.80-1.48	0.598			

RFS: Recurrence-free survival time; HCC-BC: hepatocellular carcinoma with virus-infected liver; AFP: α-fetoprotein; DCP: des-gamma-carboxy prothrombin; HR: hazard ratio; CI: confidence interval.

Table V. Univariate and multivariable analyses of RFS in HCC-NBNC cases.

Clinicopathological factors		Univariate analysis			Multivariable analysis		
		HR	95%CI	p-Value	HR	95%CI	p-Value
Age	≥65 years	1.09	0.55-2.16	0.805			
Gender	Male	1.54	0.61-3.91	0.366			
Tumor number	Multiple	1.90	0.90-4.00	0.090			
Tumor size	≥2.0 cm	2.23	0.94-5.30	0.068			
AFP	≥10 ng/ml	2.04	1.11-3.74	0.022			
DCP	≥40 ng/ml	3.79	1.49-9.67	0.005	4.99	1.91-13.01	0.001
Differentiation	Poor, Moderate	1.50	0.67-3.36	0.330			
Growth form	Infiltrative	1.63	0.82-3.24	0.164			
Serosal invasion	Positive	2.00	1.06-3.77	0.033			
Portal vein invasion	Positive	3.22	1.67-6.19	<0.001	5.41	2.69-10.87	<0.001
Hepatic vein invasion	Positive	3.19	1.65-6.18	<0.001			
Liver cirrhosis	Present	0.99	0.48-2.07	0.989			

RFS: Recurrence-free survival time ; HCC-NBNC: hepatocellular carcinoma with no virus-infected liver; AFP: α-fetoprotein; DCP: des-gamma-carboxy prothrombin; HR: hazard ratio; CI: confidence interval.

produce DCP (24). Our clinical data clearly indicate that DCP values increased depending on the T stage of HCC-NBNCs. Besides HCC-BCs are derived from the damaged background liver, which is chronically exposed to long-lasting hypoxia (25). Actually, DCP values of the cirrhotic liver tumors were significantly decreased.

Exome sequences of hepatocellular carcinomas have identified new mutational signatures and potential therapeutic targets (7). Depending on the risk factors of hepatocarcinogenesis, responsible gene signatures vary. For instance, *CTNNB1*, *TERT*, *CDKN2A*, *SMRCA2* and *HGF* gene alterations can be frequently found in alcohol-based hepatitis. *TP53* mutation was dominant in hepatitis B cases.

In contrast, no distinct signature was identified in hepatitis C or NASH-based HCCs. Totoki *et al.*, have revealed 30 candidate driver genes and 11 core pathway modules from 503 liver cancer genomes (8). *TERT* or *ATRX* chromatin remodeler (*ATRX*) genes are widely mutated in all virus-induced HCCs. For NBNC HCCs, AT-rich interaction domain 1A (*ARID1A*) mutation is frequently found. Moore *et al.*, have demonstrated that *ARID1A*-deficient livers are more susceptible to high-fat diet-induced liver steatosis and fibrosis in mice models (26). As a detailed mechanism, Qu YL *et al.*, have revealed that *ARID1A* deficiency impairs fatty acid oxidation by epigenetically downregulating Peroxisome proliferator-activated receptor alpha (*PPARα*)

Table VI. Univariate and multivariable analyses of OS in HCC-BC cases.

Clinicopathological factors		Univariate analysis			Multivariable analysis		
		HR	95%CI	<i>p</i> -Value	HR	95%CI	<i>p</i> -Value
Age	≥65 years	1.06	0.67-1.67	0.813			
Gender	Male	1.01	0.59-1.71	0.983			
Tumor number	Multiple	2.23	1.46-3.41	<0.001	1.94	1.18-3.17	0.008
Tumor size	≥2.0 cm	1.21	0.78-1.86	0.391			
AFP	≥10 ng/ml	1.65	1.04-2.61	0.034			
DCP	≥40 ng/ml	1.63	1.04-2.55	0.034			
Differentiation	Poor, Moderate	1.50	0.84-2.65	0.169			
Growth form	Infiltrative	2.41	1.48-3.91	<0.001			
Serosal invasion	Positive	2.61	1.56-4.37	<0.001	1.82	1.06-3.13	0.031
Portal vein invasion	Positive	3.93	2.54-6.07	<0.001	2.63	1.58-4.38	<0.001
Hepatic vein invasion	Positive	5.17	2.97-9.00	<0.001	4.73	2.62-8.54	<0.001
Liver cirrhosis	Present	1.37	0.90-2.09	0.138			

OS: Overall survival time; HCC-BC: hepatocellular carcinoma with virus-infected liver; AFP: α -fetoprotein; DCP: des-gamma-carboxy prothrombin; HR: hazard ratio; CI: confidence interval.

Table VII. Univariate and multivariable analyses of OS in HCC-NBNC cases.

Clinicopathological factors		Univariate analysis			Multivariable analysis		
		HR	95%CI	<i>p</i> -Value	HR	95%CI	<i>p</i> -Value
Age	≥65 years	2.14	0.48-9.60	0.320			
Gender	Male	1.16	0.26-5.22	0.845			
Tumor number	Multiple	1.91	0.53-6.89	0.321			
Tumor size	≥2.0 cm	4.46	0.58-34.22	0.150			
AFP	≥10 ng/ml	2.91	0.91-9.27	0.071			
DCP	≥40 ng/ml	N/A	N/A	N/A	N/A	N/A	N/A
Differentiation	Poor, Moderate	1.64	0.36-7.38	0.522			
Growth form	Infiltrative	2.53	0.83-7.75	0.104			
Serosal invasion	Positive	7.42	1.96-28.05	0.003	4.74	1.13-19.87	0.033
Portal vein invasion	Positive	5.07	1.70-15.17	0.004	8.26	1.88-36.22	0.005
Hepatic vein invasion	Positive	5.89	1.87-18.52	0.002			
Liver cirrhosis	Present	1.60	0.50-5.12	0.424			

OS: Overall survival time; HCC-NBNC: Hepatocellular carcinoma with no virus-infected liver; AFP: α -fetoprotein; DCP: des-gamma-carboxy prothrombin; HR: hazard ratio; CI: confidence interval; N/A: not adequate.

and other metabolism-related genes, such as carnitine palmitoyltransferase 1A (*CPT1A*) and acyl-CoA oxidase 1 (*ACOX1*) (27).

This study has some limitations. First, this is a retrospective study from a single-institution with a modest sample size. Further confirmation with large multicenter data is required. Second, the mechanism of DCP elevation in HCC-NBNC should be explained by specific molecular characteristics, including *PNPLA3* mutation, *ARID1A* deficiency or lipid metabolism-related genes in non-hepatitis livers in future studies.

In conclusion, AFP elevation and DCP elevation were differentially observed depending on the background liver

status. Hepatocarcinogenesis in NASH liver was specific to DCP elevation, rather than AFP. DCP seems to be a significant predictive serum marker of survival outcomes, especially for HCC-NBNC cases.

Conflicts of Interest

The Authors declare no conflicts of interest.

Authors' Contributions

MH, SY and YO designed the project. MH, NT, YO, HT, YI, FS, NT and MK collected the clinical data. MH, NT and YO analyzed the data.




Figure 4. Distribution of preoperative AFP and DCP values according to histological T grades. AFP values of HCC-BC cases are gradually increased in parallel with T grades, whereas of HCC-NBNC cases did not. On the contrary, HCC-NBNC cases showed a steady increase in DCP values with T stage, while HCC-BC cases showed no increase. AFP: Alpha-fetoprotein; DCP: des-gamma-carboxy prothrombin; HCC-BC: hepatocellular carcinoma with virus-infected liver; HCC-NBNC: hepatocellular carcinoma with no virus-infected liver.




Figure 5. Association between each tumor marker and liver cirrhosis (LC). AFP values showed no decrease in LC cases, whereas DCP values decreased in LC cases. AFP: Alpha-fetoprotein, DCP: des-gamma-carboxy prothrombin.

MH and SY checked and approved all the statistical analyses. MH prepared the manuscript and DS, NH, CT, GN, MK and YK revised it.

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA and Jemal A: Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 68(6): 394-424, 2018. PMID: 30207593. DOI: 10.3322/caac.21492
- Forner A, Llovet JM and Bruix J: Hepatocellular carcinoma. *Lancet* 379(9822): 1245-1255, 2012. PMID: 22353262. DOI: 10.1016/S0140-6736(11)61347-0
- El-Serag HB: Hepatocellular carcinoma. *N Engl J Med* 365(12): 1118-1127, 2011. PMID: 21992124. DOI: 10.1056/NEJMr1001683
- Utsunomiya T, Shimada M, Kudo M, Ichida T, Matsui O, Izumi N, Matsuyama Y, Sakamoto M, Nakashima O, Ku Y, Takayama T, Kokudo N and Liver Cancer Study Group of Japan.: A comparison of the surgical outcomes among patients with HBV-positive, HCV-positive, and non-B non-C hepatocellular carcinoma: a nationwide study of 11,950 patients. *Ann Surg* 261(3): 513-520, 2015. PMID: 25072437. DOI: 10.1097/SLA.0000000000000821
- Brunner SF, Roberts ND, Wylie LA, Moore L, Aitken SJ, Davies SE, Sanders MA, Ellis P, Alder C, Hooks Y, Abascal F, Stratton MR, Martincorena I, Hoare M and Campbell PJ: Somatic mutations and clonal dynamics in healthy and cirrhotic human liver. *Nature* 574(7779): 538-542, 2019. PMID: 31645727. DOI: 10.1038/s41586-019-1670-9
- Letouzé E, Shinde J, Renault V, Couchy G, Blanc JF, Tubacher E, Bayard Q, Bacq D, Meyer V, Semhoun J, Bioulac-Sage P, Prévôt S, Azoulay D, Paradis V, Imbeaud S, Deleuze JF and Zucman-Rossi J: Mutational signatures reveal the dynamic interplay of risk factors and cellular processes during liver tumorigenesis. *Nat Commun* 8(1): 1315, 2017. PMID: 29101368. DOI: 10.1038/s41467-017-01358-x
- Schulze K, Imbeaud S, Letouzé E, Alexandrov LB, Calderaro J, Rebouissou S, Couchy G, Meiller C, Shinde J, Soysouvanh F, Calatayud AL, Pinyol R, Pelletier L, Balabaud C, Laurent A, Blanc JF, Mazzaferro V, Calvo F, Villanueva A, Nault JC, Bioulac-Sage P, Stratton MR, Llovet JM and Zucman-Rossi J: Exome sequencing of hepatocellular carcinomas identifies new mutational signatures and potential therapeutic targets. *Nat Genet* 47(5): 505-511, 2015. PMID: 25822088. DOI: 10.1038/ng.3252
- Totoki Y, Tatsuno K, Covington KR, Ueda H, Creighton CJ, Kato M, Tsuji S, Donehower LA, Slagle BL, Nakamura H, Yamamoto S, Shinbrot E, Hama N, Lehmkuhl M, Hosoda F, Arai Y, Walker K, Dahdouli M, Gotoh K, Nagae G, Gingras MC, Muzny DM, Ojima H, Shimada K, Midorikawa Y, Goss JA, Cotton R, Hayashi A, Shibahara J, Ishikawa S, Guiteau J, Tanaka M, Urushidate T, Ohashi S, Okada N, Doddapaneni H, Wang M, Zhu Y, Dinh H, Okusaka T, Kokudo N, Kosuge T, Takayama T, Fukayama M, Gibbs RA, Wheeler DA, Aburatani H and Shibata T: Trans-ancestry mutational landscape of hepatocellular carcinoma genomes. *Nat Genet* 46(12): 1267-1273, 2014. PMID: 25362482. DOI: 10.1038/ng.3126
- Edman JC, Gray P, Valenzuela P, Rall LB and Rutter WJ: Integration of hepatitis B virus sequences and their expression in a human hepatoma cell. *Nature* 286(5772): 535-538, 1980. PMID: 6250075. DOI: 10.1038/286535a0
- Jhunjhunwala S, Jiang Z, Stawiski EW, Gnad F, Liu J, Mayba O, Du P, Diao J, Johnson S, Wong KF, Gao Z, Li Y, Wu TD, Kapadia SB, Modrusan Z, French DM, Luk JM, Seshagiri S and Zhang Z: Diverse modes of genomic alteration in hepatocellular carcinoma. *Genome Biol* 15(8): 436, 2014. PMID: 25159915. DOI: 10.1186/s13059-014-0436-9
- Sung WK, Zheng H, Li S, Chen R, Liu X, Li Y, Lee NP, Lee WH, Ariyaratne PN, Tennakoon C, Mulawadi FH, Wong KF, Liu AM, Poon RT, Fan ST, Chan KL, Gong Z, Hu Y, Lin Z, Wang G, Zhang Q, Barber TD, Chou WC, Aggarwal A, Hao K, Zhou W, Zhang C, Hardwick J, Buser C, Xu J, Kan Z, Dai H, Mao M, Reinhard C, Wang J and Luk JM: Genome-wide survey of recurrent HBV integration in hepatocellular carcinoma. *Nat Genet* 44(7): 765-769, 2012. PMID: 22634754. DOI: 10.1038/ng.2295
- Kutlu O, Kaleli HN and Ozer E: Molecular Pathogenesis of Nonalcoholic Steatohepatitis- (NASH-) Related Hepatocellular Carcinoma. *Can J Gastroenterol Hepatol* 2018: 8543763, 2018. PMID: 30228976. DOI: 10.1155/2018/8543763
- Song T, Wang L, Xin R, Zhang L and Tian Y: Evaluation of serum AFP and DCP levels in the diagnosis of early-stage HBV-related HCC under different backgrounds. *J Int Med Res* 48(10): 300060520969087, 2020. PMID: 33135527. DOI: 10.1177/0300060520969087
- Makuuchi M and Kokudo N: Clinical practice guidelines for hepatocellular carcinoma: the first evidence based guidelines from Japan. *World J Gastroenterol* 12(5): 828-829, 2006. PMID: 16521207. DOI: 10.3748/wjg.v12.i5.828
- Wakiyama S, Matsumoto M, Haruki K, Gocho T, Sakamoto T, Shiba H, Futagawa Y, Ishida Y and Yanaga K: Clinical features and outcome of surgical patients with non-b non-c hepatocellular carcinoma. *Anticancer Res* 37(6): 3207-3213, 2017. PMID: 28551666. DOI: 10.21873/anticancer.11682
- El Raziky M, Attia D, El Akel W, Shaker O, Khatab H, Abdo S, Elsharkawy A and Esmat G: Hepatic fibrosis and serum alpha-fetoprotein (AFP) as predictors of response to HCV treatment and factors associated with serum AFP normalisation after treatment. *Arab J Gastroenterol* 14(3): 94-98, 2013. PMID: 24206736. DOI: 10.1016/j.ajg.2013.08.005
- Zheng Y, Zhu M and Li M: Effects of alpha-fetoprotein on the occurrence and progression of hepatocellular carcinoma. *J Cancer Res Clin Oncol* 146(10): 2439-2446, 2020. PMID: 32725355. DOI: 10.1007/s00432-020-03331-6
- Zhang H, Cao D, Zhou L, Zhang Y, Guo X, Li H, Chen Y, Spear BT, Wu JW, Xie Z and Zhang WJ: ZBTB20 is a sequence-specific transcriptional repressor of alpha-fetoprotein gene. *Sci Rep* 5: 11979, 2015. PMID: 26173901. DOI: 10.1038/srep11979
- Yue P, Gao ZH, Xue X, Cui SX, Zhao CR, Yuan Y, Yin Z, Inagaki Y, Kokudo N, Tang W and Qu XJ: Des-γ-carboxyl prothrombin induces matrix metalloproteinase activity in hepatocellular carcinoma cells by involving the ERK1/2 MAPK signalling pathway. *Eur J Cancer* 47(7): 1115-1124, 2011. PMID: 21349701. DOI: 10.1016/j.ejca.2011.01.017
- Taniguchi T, Kishi K, Nakagawa T, Tanaka H, Tanaka T, Tomonari T, Okamoto K, Sogabe M, Miyamoto H, Okahisa T, Muguruma N, Kajimoto M, Sagawa I and Takayama T: Poly-(ADP-Ribose) Polymerase-1 promotes prothrombin gene transcription and produces des-gamma-carboxy prothrombin in

- hepatocellular carcinoma. *Digestion* 95(3): 242-251, 2017. PMID: 28384634. DOI: 10.1159/000470837
- 21 Mukhopadhyay P, Horváth B, Rajesh M, Varga ZV, Gariani K, Ryu D, Cao Z, Holovac E, Park O, Zhou Z, Xu MJ, Wang W, Godlewski G, Paloczi J, Nemeth BT, Persidsky Y, Liaudet L, Haskó G, Bai P, Boulares AH, Auwerx J, Gao B and Pacher P: PARP inhibition protects against alcoholic and non-alcoholic steatohepatitis. *J Hepatol* 66(3): 589-600, 2017. PMID: 27984176. DOI: 10.1016/j.jhep.2016.10.023
- 22 Yamazaki S, Takayama T, Kurokawa T, Shimamoto N, Mitsuka Y, Yoshida N, Higaki T and Sugitani M: Next-generation des- γ -carboxy prothrombin for immunohistochemical assessment of vascular invasion by hepatocellular carcinoma. *BMC Surg* 20(1): 201, 2020. PMID: 32928172. DOI: 10.1186/s12893-020-00862-0
- 23 Suzuki H, Murata K, Gotoh T, Kusano M, Okano H, Oyamada T, Yasuda Y, Imamura M, Kudo M, Mizokami M and Sakamoto A: Phenotype-dependent production of des- γ -carboxy prothrombin in hepatocellular carcinoma. *J Gastroenterol* 46(10): 1219-1229, 2011. PMID: 21744129. DOI: 10.1007/s00535-011-0432-8
- 24 Höckel M and Vaupel P: Biological consequences of tumor hypoxia. *Semin Oncol* 28(2 Suppl 8): 36-41, 2001. PMID: 11395851.
- 25 Zhu C, Liu X, Wang S, Yan X, Tang Z, Wu K, Li Y and Liu F: Hepatitis C virus core protein induces hypoxia-inducible factor 1 α -mediated vascular endothelial growth factor expression in Huh7.5.1 cells. *Mol Med Rep* 9(5): 2010-2014, 2014. PMID: 24626461. DOI: 10.3892/mmr.2014.2039
- 26 Moore A, Wu L, Chuang JC, Sun X, Luo X, Gopal P, Li L, Celen C, Zimmer M and Zhu H: Arid1a loss drives nonalcoholic steatohepatitis in mice through epigenetic dysregulation of hepatic lipogenesis and fatty acid oxidation. *Hepatology* 69(5): 1931-1945, 2019. PMID: 30584660. DOI: 10.1002/hep.30487
- 27 Qu YL, Deng CH, Luo Q, Shang XY, Wu JX, Shi Y, Wang L and Han ZG: Arid1a regulates insulin sensitivity and lipid metabolism. *EBioMedicine* 42: 481-493, 2019. PMID: 30879920. DOI: 10.1016/j.ebiom.2019.03.021

Received January 16, 2021

Revised February 6, 2021

Accepted February 10, 2021