



The *Rhox* gene cluster suppresses germline *LINE1* transposition

Kun Tan^{a,1}, Matthew E. Kim^{a,1}, Hye-Won Song^a, David Skarbrevik^a, Eric Babajanian^a, Tracy A. Bedrosian^b, Fred H. Gage^b, and Miles F. Wilkinson^{a,c,2}

^aDepartment of Obstetrics, Gynecology, and Reproductive Sciences, School of Medicine, University of California San Diego, La Jolla, CA 92093; ^bLaboratory of Genetics, The Salk Institute for Biological Studies, La Jolla, CA 92037; and ^cInstitute of Genomic Medicine, University of California San Diego, La Jolla, CA 92093

Edited by Martin M. Matzuk, Baylor College of Medicine, Houston, TX, and approved May 3, 2021 (received for review December 2, 2020)

Transposable elements (TEs) are mobile sequences that engender widespread mutations and thus are a major hazard that must be silenced. The most abundant active class of TEs in mammalian genomes is long interspersed element class 1 (*LINE1*). Here, we report that *LINE1* transposition is suppressed in the male germline by transcription factors encoded by a rapidly evolving X-linked homeobox gene cluster. *LINE1* transposition is repressed by many members of this *RHOX* transcription factor family, including those with different patterns of expression during spermatogenesis. One family member—*RHOX10*—suppresses *LINE1* transposition during fetal development in vivo when the germline would otherwise be susceptible to *LINE1* activation because of epigenetic reprogramming. We provide evidence that *RHOX10* suppresses *LINE* transposition by inducing *Piwil2*, which encodes a key component in the Piwi-interacting RNA pathway that protects against TEs. The ability of *RHOX* transcription factors to suppress *LINE1* is conserved in humans but is lost in *RHOXF2* mutants from several infertile human patients, raising the possibility that loss of *RHOXF2* causes human infertility by allowing uncontrolled *LINE1* expression in the germline. Together, our results support a model in which the *Rhox* gene cluster is in an evolutionary arms race with TEs, resulting in expansion of the *Rhox* gene cluster to suppress TEs in different biological contexts.

LINE1 | piRNA | transposon | *RHOX10* | *Piwil2*

Mammalian genomes are dominated by transposable elements (TEs)—parasitic genetic units that can reach copy numbers in the hundreds of thousands (1). The most abundant active class of TEs in mammals is long interspersed element class 1 (*LINE1*), which comprises ~20% of mammalian genomes (2). *LINE1* elements are autonomous TEs that propagate in the genome by a copy-and-paste mechanism through retrotransposition. *LINE1* propagation depends on the *LINE1*-encoded proteins, ORF1p and ORF2p, which can also mobilize nonautonomous retrotransposons, other noncoding RNAs, and messenger RNAs, leading to the generation of processed pseudogenes (3, 4). All told, *LINE1*-mediated transposition has been estimated to have generated at least a third of the human genome (5).

LINE1 elements are detrimental to cells. One obvious negative consequence of *LINE1* elements is they engender mutations when they undergo transposition to a new genomic site. Indeed, insertional mutagenesis elicited by *LINE1* and other TEs has been shown to cause more than 65 human genetic diseases (6). In addition, the *LINE1* open reading frames (ORFs) are themselves known to encode deleterious endonuclease and reverse transcriptase activities (7–9). The ultimate effect of overly active *LINE1*s is accumulation of DNA damage, checkpoint activation, and cell death (8, 9). Thus, limiting *LINE1* expression and translocation is fundamental to genome integrity and health.

To protect against these negative effects, multiple epigenetic and RNA-mediated mechanisms have been postulated to have evolved (10). Such silencing mechanisms are particularly critical for the germline, as they reduce the transmission of TE-induced

mutations to subsequent generations. A major opportunity for TEs to undergo transposition is when the male germline undergoes genome-wide demethylation (between embryonic day [E]7.5 and E12.5) in primordial germ cells (PGCs) (11). This hypomethylated state is maintained when PGCs become nonproliferative cells called prospermatogonia (ProSG, also known as gonocytes) at ~E13.5 (12). While genome-wide DNA demethylation provides the benefit of reprogramming the genome for the next generation, it opens the door for a major hazard, as it allows an opportunity for TE activation and transposition. Thus, mechanisms have evolved to defend against TEs becoming active when PGCs and ProSG are in this hypomethylated state. For example, evidence suggests that PGCs and ProSG make use of the histone modification, H2A/H4R3me2, to suppress TE expression and protect genomic integrity (13). Another major mechanism that protects against TEs in PGCs and ProSG is the Piwi-interacting (pi) RNA pathway. This pathway is mediated by short (24 to 31 nt) noncoding RNAs called “piRNAs” that associate with members of the PIWI RNA-binding Argonaute protein family to suppress transposons and maintain germline genome integrity (14). In mice, there are three PIWI family members—PIWIL1, PIWIL2, and PIWIL4 (also known as MIWI, MILL, and MIWI2, respectively)—the latter two of which are expressed in PGCs and/or ProSG. *Piwil2* is first detected in gonadal tissue at E12.5, around the PGC-to-ProSG transition (15), whereas *Piwil4* is not detectable until E15.5 (16). Both PIWIL2 and PIWIL4

Significance

Long interspersed element class 1 (*LINE1*) elements are transposable elements that comprise ~20% of mammalian genomes. Their activity must be tightly controlled or genome integrity will be compromised, leading to DNA damage and cell death. Here, we report that several members of a rapidly evolving X-linked homeobox gene cluster suppress *LINE1* transposition. One family member of this *Rhox* gene cluster—*Rhox10*—silences *LINE1* expression and transposition in the male germline when it is hypomethylated and thus highly susceptible to *LINE1* activation. *Rhox10* acts by driving the expression of *Piwil2*, which encodes a key component in the Piwi-interacting RNA pathway. The ability of *Rhox* genes to suppress *LINE1* elements is evolutionarily conserved and perturbed by mutation in infertility patients.

Author contributions: K.T., M.E.K., H.-W.S., and M.F.W. designed research; K.T., M.E.K., H.-W.S., D.S., E.B., and T.A.B. performed research; F.H.G. contributed new reagents/analytic tools; K.T. analyzed data; and K.T. and M.F.W. wrote the paper.

The authors declare no competing interest.

This article is a PNAS Direct Submission.

Published under the PNAS license.

¹K.T. and M.E.K. contributed equally to this work.

²To whom correspondence may be addressed. Email: mfwilkinson@health.ucsd.edu.

This article contains supporting information online at <https://www.pnas.org/lookup/suppl/doi:10.1073/pnas.2024785118/-DCSupplemental>.

Published June 3, 2021.

are critical for suppression of *LINE1* elements and other TEs in ProSG, based on analysis of *Piwil2*- and *Piwil4*-knockout (KO) mice (16, 17). Both KO strains suffer from male sterility (16, 17).

In this communication, we report evidence for a new line of defense against *LINE1* elements: the RHOX homeobox transcription factor family. Our laboratory identified a member of this family—*Pem* (*Rhox5*)—as an oncofetal gene expressed during normal embryogenesis and in diverse tumors (18). Later, we and others discovered that *Rhox5* is part of a very large (33-member) homeobox gene cluster on the X chromosome in mice (19–23). All known members of this *Rhox* gene cluster are primarily expressed in the placenta and in the male and female reproductive tracts (24). By definition, all homeobox genes encode a DNA-binding homeo-domain, suggesting that the RHOX proteins are transcription factors devoted to reproductive functions. Indeed, this possibility has been supported by several studies, including those showing that loss of *Rhox5*, *Rhox8*, *Rhox10*, or *Rhox13* in mice causes spermatogenic defects (25–28). Relevant to this communication, we found that knock out of the entire *Rhox* cluster (either globally or specifically in germ cells) causes progressive spermatogenic decline, consistent with a defect in spermatogonial stem cells (SSCs) (26). We then demonstrated that one particular *Rhox* gene—*Rhox10*—is critical for this defect, as we found that *Rhox10*-null mice have the same progressive spermatogenic decline as *Rhox* cluster KO mice (26). Through a battery of assays, including germ cell transplantation and single-cell RNA sequencing (RNA-seq) analyses, the defect in *Rhox10*-null mice was pinpointed to be an inability of ProSG to efficiently undergo differentiation and thereby generate SSCs. Thus, *Rhox10* functions in the “gateway cells” that drive the initiation of spermatogenesis: ProSG.

In the present study, we demonstrate that RHOX transcription factors have an unexpected role in TE defense in the germline. We initially postulated this possibility for several reasons: First, all *Rhox* genes are expressed in the testis and ovary (19, 24). In humans, the *RHOX* genes are expressed exclusively in germ cells in the testes and ovary, suggesting they have specific roles in the germline (29). Second, the *Rhox* genes encode DNA-binding homeobox proteins (19, 24), suggesting obvious mechanisms by which they could act on TEs. Third, *Rhox* genes are dramatically transcriptionally derepressed in response to DNA hypomethylation, based on many lines of evidence both in vitro and in vivo (30–35). We reasoned that this regulatory trigger would generate RHOX transcription factors precisely when needed for TE defense—during genome-wide demethylation of the germline. Consistent with this reasoning, most *Rhox* genes are induced in PGCs precisely when they undergo genome-wide hypomethylation (36, 37).

In this communication, we report our experiments to directly test whether RHOX transcription factors have a role in TE defense. Our experiments revealed that, indeed, several members of the mouse RHOX homeobox family are capable of suppressing *LINE1* transposition. We demonstrate that one RHOX family member—RHOX10—strongly suppresses *LINE1* transposition during the critical period when the germline is largely hypomethylated and thus highly susceptible to *LINE1* aggression. We determined the mechanism by which this suppression occurs and then asked whether *LINE1* defense extends to the human *RHOX* cluster. Together, our results suggest that the *Rhox* gene cluster is engaging in an evolutionary arms race with *LINE1* elements to suppress the expression and expansion of these parasitic TEs.

Results

***Rhox10* Represses *LINE1* Transposition in the Male Germline.** To determine whether the *Rhox* cluster represses *LINE1* transposition, we crossed *SN1 LINE1* reporter mice (38) with *Rhox* cluster (*Rhox-c*)^{fl/fl};Vasa-Cre mice, which delete the entire *Rhox* cluster in germ cells (26). We chose to use the *SN1* single-copy *LINE1* transgene reporter mouse line for several reasons, including that

it contains codon-optimized mouse ORF1 and ORF2 for improved translation and an intron-containing split EGFP for assaying retrotransposition (38). Furthermore, unlike past *LINE1* reporters (39–45), *SN1* is under the control of an endogenous mouse *LINE1* promoter, allowing for more physiological evaluation of *LINE1* regulation (38). We used the digital droplet (dd) PCR assay to assay reporter genomic copy number. This assay showed that *SN1 LINE1* reporter copy number was high in the brains of wild-type (WT) mice (Fig. 1A), consistent with previous reports that *LINE1* elements undergo high rates of transposition in the brain and neural progenitor cells (42, 46, 47). ddPCR analysis of *Rhox-c* conditional KO (cKO) mice revealed *SN1 LINE1* copy number was significantly higher in the testes from these cKO mice than control (Vasa-Cre;*SN1*^{+/+}) mice, indicative of greatly increased *LINE1* transposition as a result of *Rhox* cluster loss (Fig. 1A). In contrast, *SN1* copy number was not significantly increased in other adult tissues we tested, including brain (Fig. 1A). Time-course analysis showed that these *Rhox-c* cKO mice exhibited a dramatic increase in *SN1* copy number in testes between E13.5 and E16.5, which was sustained at postnatal (P) day 0 and adult stages (Fig. 1B). This indicated that the *Rhox* cluster suppresses *LINE1* transposition in ProSG, as this is the only germ cell type present from ~E13.5 to birth (12).

The *Rhox* cluster contains 33 homeobox genes (19, 24), any of whose loss could potentially contribute to *LINE1* suppression. Since *Rhox10* is expressed in the germ cell stage in which the *Rhox* cluster acts to suppress *LINE1* transposition—ProSG (26)—we examined the role of *Rhox10*. To this end, we generated *Rhox10*-null;*SN1* transgenic mice. Like *Rhox-c* cKO mice, these *Rhox10*-null mice had greatly elevated *SN1* transgene copy number compared to control (*SN1*^{+/+}) mice. Furthermore, both *Rhox10*-null and *Rhox-c* cKO mice had elevated *SN1* transgene copy number in precisely the same contexts: E16.5, P0, and adult testes but not in E13.5 testes or nontesticular adult tissues (Fig. 1A and B). This finding suggested that *Rhox10* is largely responsible for the ability of the *Rhox* cluster to suppress *LINE1* transposition during fetal germ cell development in vivo.

The notion that *Rhox10* suppresses *LINE1* transposition was validated by several lines of evidence. First, using the *ORFeus* dual luciferase reporter (48), we confirmed the suppressive effect of *Rhox10* on *LINE1* transposition in the GC1 spermatogonial cell line (Fig. 1C). Second, qPCR analysis showed that *SN1 LINE1* transgene expression was elevated in *Rhox10*-null mice testes (Fig. 1D). Third, to examine whether regulation extends to endogenous *LINE1* elements, we used primer pairs specific for several different *LINE1* subfamilies. This analysis revealed that the *L1a* *LINE1* subfamily exhibited strongly elevated expression in *Rhox10*-null mouse testes (Fig. 1D). Specificity was demonstrated by the finding that *L1g* and *L1t* subfamilies did not exhibit statistically significant changes in expression (Fig. 1D). Fourth, to assess whether *Rhox10* might suppress *LINE1* transposition by repressing *LINE1* ORF1p protein, we examined ORF1p expression and found it was significantly up-regulated in *Rhox10*-null mouse testes, as detected by Western blot analysis of both fetal and adult testes (Fig. 1E). Fifth, we confirmed ORF1 up-regulation using immunofluorescence analysis: at E16.5, most germ cells (TRA98+ cells) in WT testes had a very dim anti-ORF1p signal, whereas most germ cells in *Rhox10*-null mice testes had a strong signal (Fig. 1F). In adult WT mice, all germ cells in all seminiferous tubules examined had undetectable or weak expression of ORF1p, while in adult *Rhox10*-null mice, 18% of tubules (11 of 61) had germ cells with a strong ORF1p signal (*SI Appendix*, Fig. S1).

Evidence for a *Rhox10*–*Piwil2*–*LINE1* Circuit. To identify candidate genes that act downstream of the RHOX10 transcription factor to suppress *LINE1* elements in the male germline, we performed RNA-seq analysis on fluorescence-activated cell sorting (FACS)-purified germ cells from fetal *Rhox10*-null;Oct4-eGFP^{+/+} (KO)




Fig. 1. *Rhox10* represses *LINE1* transposition in the male germline. (A) ddPCR analysis of SN1 *LINE1* copy number in adult tissues from SN1 reporter mice mated with *Rhox* cluster (c) cKO (*Rhox-c^{fl/y};Vasa-Cre;SN1^{+/+}*), *Rhox10*-null (*Rhox10^{-Y};SN1^{+/+}*) (KO), or WT (*Vasa-Cre;SN1^{+/+}*) mice ($n = 6$). $*P < 0.05$. (B) SN1 *LINE1* copy number in testes from mice of the ages indicated (adult testis is the same as in A). The genotypes and analysis are described in A ($n = 3$). $*P < 0.05$. (C) Luciferase analysis of *LINE1* transposition activity in the GC1 spermatogonia cell line cotransfected with the mouse *ORF1p* reporter and a *Rhox10*-expression vector or an empty expression vector ($n = 3$). $*P < 0.05$. (D) qPCR analysis of testes from E16.5 *Rhox10*-null (KO) and littermate WT mice. The primers used are specific for the indicated *LINE1* subfamilies ($n = 3$). $*P < 0.05$. (E) Western blot analysis of *LINE1* ORF1p protein expression in testes lysates from *Rhox10*-null (KO) and littermate WT mice of the indicated ages. (Right) ORF1p quantification when normalized to GAPDH ($n = 3$). $*P < 0.05$. (F) Immunofluorescence analysis of ORF1p (green) in *Rhox10*-null (KO) and littermate WT testes from E16.5 mice. The sections were costained with an antibody against TRA98 to detect all germ cells (red) and DAPI to detect all cells (blue). $n = 3$.

and littermate Oct4-eGFP^{+/+} (control) mice. Principal component analysis of quadruplicate replicates of *Rhox10^{-Y};Oct4-eGFP* and *Rhox10^{+/Y};Oct4-eGFP* germ cells showed that their transcriptomes segregated separately, indicative of a significant

change in gene expression elicited by loss of *Rhox10* (Fig. 2A). In total, 370 genes exhibited significantly altered expression in *Rhox10*-null germ cells compared to control germ cells ($q < 0.05$, Basemean > 10, $|\text{Log}_2\text{FC}| > 0.25$). Of these 370 differentially

expressed genes (DEGs), 213 were up-regulated and 157 were down-regulated (Dataset S1). Fig. 2B shows signaling pathways associated with these DEGs.

Among the functional categories associated with genes regulated by *Rhox10* loss (Fig. 2C and D) was “piRNA biogenesis.” The piRNA pathway is known to silence *LINE1* elements (49, 50), which led us to postulate that *Rhox10* acts through the piRNA pathway to suppress *LINE1* elements. Among piRNA biogenesis genes, *Piwil2* and *Tdrd1* were significantly down-regulated in *Rhox10*-null germ cells (Fig. 3A), which we validated by qPCR analysis (Fig. 3B) and Western blot analysis for PIWIL2 (Fig. 3C). To determine whether RHOX10 might directly regulate their expression, we first employed reporter analysis. The *Piwil2* and *Tdrd1* promoter regions were separately cloned into a firefly luciferase reporter vector and cotransfected with a *Rhox10*-expression vector into HEK-293T cells. *Rhox10* dramatically increased expression from the *Piwil2* promoter (by ~seven-fold), demonstrating that RHOX10 acts through the *Piwil2* promoter to increase *Piwil2* expression (Fig. 3D). In contrast, RHOX10 likely acts by a different mechanism to up-regulate *Tdrd1*, as forced *Rhox10* expression did not significantly increase reporter expression from the *Tdrd1* promoter.

The finding that the RHOX10 transcription factor strongly up-regulates *Piwil2* expression in the male germline in vivo raised the possibility that RHOX10 acts in a circuit with PIWIL2 to suppress *LINE1* transposition. In support of this model, we found that RHOX10 occupies the *Piwil2* promoter, as shown by chromatin immunoprecipitation (ChIP) analysis (Fig. 3E). This finding, coupled with 1) the coexpression of *Rhox10* and *Piwil2* in fetal germ cells (16, 51, 36), 2) the ability of RHOX10 to drive expression of the *Piwil2* promoter in vitro (Fig. 3D), and 3) the ability of RHOX10 to positively regulate *Piwil2* in germ cells in vivo (Fig. 3B and C), strongly suggests that RHOX10 directly drives transcription of the *Piwil2* gene in fetal germ cells. To assess whether RHOX10 suppresses *LINE1* expression via *Piwil2*, we performed a rescue experiment. In this experiment, we asked whether reduced *LINE1* defense caused by RHOX10 knockdown could be rescued by adding back the depleted PIWIL2. Indeed, we found that forced expression of PIWIL2 in germline stem (GS) cells reversed the defect in *LINE1* transposition caused by RHOX10 knockdown (Fig. 3F), as judged by a cotransfected dual luciferase *LINE1* reporter plasmid (*ORFeus* reporter) (48) (Fig. 3G). Together, these data support a model in which RHOX10 directly




Fig. 2. RHOX10-regulated genes in fetal germ cells. (A) Principal component analysis plot of the RNA-seq datasets from quadruplicate testicular germ cell samples obtained from E16.5 *Rhox10*-null; Oct4-eGFP^{+/+} (KO) and Oct4-eGFP^{+/+} (WT) mice. (B) Heatmap of DEGs identified from the analysis of the samples in A (the data from each biological replicate are shown). Selected signaling pathways enriched in DEGs down- and up-regulated in *Rhox10*-null germ cells are shown in *Upper* and *Lower*, respectively. (C and D) STRING analysis of DEGs defined in B.




Fig. 3. RHOX10 suppresses *LINE1* expression via regulating *Piwil2* expression. (A) Heatmap of piRNA pathway gene expression determined by the RNA-seq analysis shown in Fig. 2. *, $q < 0.05$, Basemean > 10 , $|\text{Log}_2\text{FC}| > 0.25$. (B) qPCR validation of *Rhox10*-mediated regulation of *Piwil2* and *Tdrd1* expression in germ cells from E16.5 *Rhox10*-null; Oct4-eGFP^{+/+} (KO) and Oct4-eGFP^{+/+} (WT) mice ($n = 3$). * $P < 0.05$. (C, Top) Western blot analysis of PIWIL2 protein expression in testes lysates from E16.5 *Rhox10*-null (KO) and littermate WT mice. (Bottom) PIWIL2 quantification when normalized to GAPDH ($n = 3$). * $P < 0.05$. (D) Luciferase analysis of *Piwil2* or *Tdrd1* promoter activity in HEK-293T cells cotransfected with a luciferase reporter vector harboring the *Piwil2* or *Tdrd1* promoter and a *Rhox10*-expression vector or an empty expression vector as a control (Ctrl) ($n = 4$). * $P < 0.05$. (E) ChIP-qPCR analysis of GS cells transfected with an expression vector encoding HA-tagged RHOX10 and analyzed using an HA antibody. Endogenous RHOX10 was knocked down using a *Rhox10*-shRNA lentivirus ($>80\%$ efficiency). IgG antibody was used as a negative control. The values shown are the DNA signal in the immunoprecipitation samples relative to the input. ($n = 3$). * $P < 0.05$. (F) qPCR analysis of *Rhox10* expression in GS cells transfected with a *Rhox10*-shRNA lentivirus or a scramble-shRNA lentivirus as a control ($n = 2$). * $P < 0.05$. (G) Luciferase analysis of *LINE1* transposition activity in GS cells cotransfected with the ORF5 mouse *LINE1* reporter, a *Piwil2* expression vector, and a *Rhox10*-shRNA vector, as shown ($n = 3$). * $P < 0.05$. (H) TaqMan-qPCR analysis of the indicated piRNAs in FACS-purified germ cells from E16.5 *Rhox10*-null; Oct4-eGFP^{+/+} (*Rhox10* KO) and littermate Oct4-eGFP^{+/+} (WT) mice ($n = 3$). * $P < 0.05$. U6 small nuclear RNA levels were used for normalization. (I) Model: the RHOX10 transcription factor activates *Piwil2* transcription to suppress *LINE1* elements.

transcriptionally activates the *Piwil2* gene to provide defense against *LINE1* TEs in the developing male germline (Fig. 3I).

PIWIL2 is an endonuclease that produces piRNA intermediates, leading to the formation of mature piRNAs (52). This raises the possibility that by stimulating PIWIL2 expression, RHOX10 also up-regulates the expression of piRNAs. To test this, we selected three piRNAs known to be bound to PIWIL2 in ProSG, each in a different subfamily (L1Md type A, D, and F2) (53). TaqMan analysis showed that all of these piRNAs are significantly down-regulated in FACS-purified germ cells from *Rhox10*-null E16.5 Oct4-eGFP^{+/+} mice compared to germ cells from littermate Oct4-eGFP^{+/+} controls (Fig. 3H). These data indicate that *Rhox10* promotes piRNA expression in fetal germ cells, which may contribute to *Rhox10*'s role in *LINE1* defense.

***Rhox10* Drives *LINE1* Promoter DNA Methylation.** Our evidence that *Rhox10* acts, at least in part, through *Piwil2* to suppress *LINE1* transposition predicts that *Rhox10*-null mice testes will have hypomethylated *LINE1* promoters. This prediction follows from

the considerable evidence that the piRNA pathway suppresses *LINE1* expression by methylating *LINE1* promoters (16, 17). To test this possibility, we performed bisulfite analysis of the *SN1* transgene promoter in testes from fetal (E16.5) *Rhox10*-null and control mice. Among 12 CpG sites, three showed significantly decreased methylation levels in *Rhox10*-null testes when compared with control testes (Fig. 4A). The total DNA methylation level of all CpG sites was also significantly decreased in *Rhox10*-null compared to control mice (Fig. 4A). To test whether the methylation of the *LINE1* promoter is regulated by *Rhox10* specifically in germ cells, we purified germ cells from E16.5 testes by magnetic-activated cell sorting using epithelial cell adhesion molecule, an early germ cell marker (54, 55). Bisulfite analysis showed that 5 of 12 CpG sites in the *SN1* transgene promoter exhibited significantly decreased methylation in *Rhox10*-null germ cells compared to control germ cells (Fig. 4B). Total methylation level was also significantly decreased (Fig. 4B). Together, these results indicated that *Rhox10* drives the methylation of *LINE1* promoters in germ cells.




Fig. 4. *Rhox10* drives *LINE1* promoter methylation. (A, Top) The degree of methylation at each CpG site in the *SN1* transgene promoter in testes from E16.5 *Rhox10*-null; *SN1*^{+/-} (KO) and *SN1*^{+/-} (WT) mice. (Bottom) Bisulfite analysis data of individual samples. **P* < 0.05. (B) Bisulfite analysis, performed as in A, of purified germ cells from E16.5 *Rhox10*-null; *SN1*^{+/-} (KO) and *SN1*^{+/-} (WT) mice testes. **P* < 0.05.

Many Members of the Mouse *Rhox* Gene Cluster Repress *LINE1* Transposition. The evidence described above indicated that mouse *Rhox10* suppresses *LINE1* transposition in fetal germ cells, including ProSG. Other members of the mouse *Rhox* cluster are expressed in other contexts, including at later stages of male germ cell development, in female germ cells, and in somatic cells in the testes and ovary (24, 56). To determine whether other *Rhox* cluster genes suppress *LINE1* transposition, we used the *ORFeus* dual luciferase reporter assay (48). We cotransfected this *LINE1* reporter with expression vectors encoding individual *Rhox* gene in NIH 3T3 cells, revealing that several of the mouse *Rhox* genes repress *LINE1* transposition (Fig. 5A). As described in Discussion, this finding raises the possibility that individual members of the *Rhox* cluster coordinate *LINE1* suppression across male germline development, as well as acting to suppress *LINE1* elements in different cell types in the reproductive tract.

Human *RHOXF2* Represses *LINE1* Transposition. *Rhox* homeobox genes are also present in other mammals, including humans (19, 57), raising the possibility that they encode transcription factors that suppress *LINE1* elements in the male germline in humans. Humans harbor only three *RHOXF* genes: *RHOXF1*, *RHOXF2A*, and *RHOXF2B* (also called *hPEPP1* and *hPEPP2*) (58). The latter two are likely to be functionally equivalent, as they are 99.8% identical in sequence in their exons and introns, and only differ by two amino acids in their deduced protein sequence. Both the *RHOXF1* and *RHOXF2A/B* genes are most highly expressed in testes; their encoded proteins are specifically expressed in germ, not somatic, cells in the human testes and ovary (29). To assess whether human *RHOXF1* and/or *RHOXF2* have *LINE1* silencing activity, we cotransfected a human *LINE1* reporter plasmid (*L1_{RP}* reporter) (48) with individual expression vectors encoding *RHOXF1* and *RHOXF2*. This revealed that *RHOXF2*, but not *RHOXF1*, has the ability to suppress *LINE1* transposition in human HEK-293T cells (Fig. 5B) and the human germ cell line Tcam-2 (Fig. 5C). To supplement these gain-of-function studies, we took a loss-of-function approach in K562 cells, which highly express *RHOXF2* (58). We generated a *RHOXF2*-small hairpin (sh)RNA lentivirus to knockdown *RHOXF2* in a K562 L1-GFP reporter line that harbors an EGFP transposition cassette inserted into the 3' untranslated region of a *LINE1* element to assay *LINE1* transposition (59). Transient knockdown of *RHOXF2* with this *RHOXF2*-shRNA lentivirus increased *LINE1* transposition, as

judged using this reporter (Fig. 5D). We conclude that human *RHOXF2* suppresses *LINE1* transposition.

***RHOXF* Mutants from Infertility Patients Lack the Ability to Repress *LINE1* Transposition.** Mutations in *RHOXF* genes have been implicated in causing male human infertility (60). Coupled with our findings described above, this raised the possibility that *RHOXF* mutations disrupt the ability of *RHOXF* proteins to suppress *LINE1* elements. To test this possibility, we first examined two missense *RHOXF2* mutants found in patients with severe oligozoospermia (60). We cotransfected these c.202G > A and c.679G > A mutants with the *L1_{RP}* reporter (48) into HEK-293T cells and found that neither repressed *LINE1* transposition (Fig. 5E). These mutations did not destabilize *RHOXF2* (Fig. 5F); thus, we conclude that instead they disrupt the ability of this protein to suppress *LINE1* elements. We also tested a frameshift mutant that encodes a truncated *RHOXF2* protein—c.381dupG (61)—and found it also lacked the ability to repress *LINE1* element transposition (Fig. 5G). Because this mutant encodes a truncated protein completely lacking the homeodomain, we considered the possibility that it is a dominant-negative mutant. In support, cotransfection of a constant amount of WT *RHOXF2* with increasing amounts of the c.381dupG *RHOXF2* mutant lead to a progressive loss in *LINE1*-reporter transposition inhibition (Fig. 5G).

Discussion

The *Rhox* homeobox genes and their encoded proteins have been widely used as cell type-specific and stage-specific germ cell markers in both humans and mice (27, 51, 29, 62, 63). *Rhox* genes also are widely used to measure androgen signaling (25). Despite their usefulness as markers, little is known about the function of most *Rhox* genes. With only one exception (*Rhox10*), KO or knockdown of *Rhox* genes leads to either relatively subtle defects in spermatogenesis or no detectable defects at all (19, 27, 28, 64). Our discovery that many *Rhox* genes function in *LINE1* defense provides an explanation for such weak phenotypes. Reduced *LINE1* germline defense would not be expected to necessarily cause overt defects in the first generation. Instead, the selection pressure to evolve defense against TEs would likely come from the progressive accumulation of mutations in the germline over subsequent generations.

We previously reported that loss of the *Rhox10* homeobox gene causes greatly impaired differentiation and migration of ProSG, leading to the formation of very few SSCs (26). Our findings




Fig. 5. Many *Rhox* family members represses *LINE1* transposition. (A) Luciferase analysis of *LINE1* transposition activity in NIH/3T3 cells cotransfected with the mouse *ORFeus LINE1* reporter and vectors expressing the indicated *Rhox* genes or an empty expression vector as a control ($n = 3$). $*P < 0.05$. (B) Luciferase analysis of *LINE1* transposition activity in HEK-293T cells cotransfected with the human $L1_{RP}$ *LINE1* reporter and vectors expressing the indicated human *RHOX* genes or an empty expression vector as a control ($n = 4$). $*P < 0.05$. (C) Luciferase analysis of *LINE1* transposition activity in TCam-2 cells cotransfected with the human $L1_{RP}$ *LINE1* reporter and vectors expressing the indicated human *RHOX* genes or an empty expression vector ($n = 3$). $*P < 0.05$. (D) Luciferase analysis of *LINE1* transposition activity in K562 cells cotransfected with the human $L1_{RP}$ *LINE1* reporter and the *RHOXF1* or *RHOXF2* shRNA vector or a randomly scrambled shRNA vector ($n = 3$). $*P < 0.05$. (E) Luciferase analysis of *LINE1* transposition activity in HEK-293T cells cotransfected with the human $L1_{RP}$ *LINE1* reporter and vectors expressing WT or mutated *RHOXF2* or an empty expression vector ($n = 3$). $*P < 0.05$. (F) Western blot analysis of HEK-293T cells transiently transfected with vectors expressing WT *RHOXF2* or the *RHOXF2* mutants. β -actin was used for the internal control ($n = 2$). (G, Left) Luciferase analysis of *LINE1* transposition activity in HEK-293T cells cotransfected with the human $L1_{RP}$ *LINE1* reporter and vectors expressing WT or truncated non-functional *RHOXF2* (c.381dupG). (Right) Luciferase analysis of *LINE1* transposition activity in HEK-293T cells cotransfected with the human *LINE1* ($L1_{RP}$) reporter, a constant amount of WT *RHOXF2*, and different amount of c.381dupG *RHOXF2*. Cells cotransfected with the human *LINE1* ($L1_{RP}$) reporter and an empty expression vector serve as the control ($n = 3$). $*P < 0.05$. (H) Western blot analysis of HEK-293T cells transiently transfected with a vector expressing WT *RHOXF2* or the *RHOXF2* mutants shown. Cells that were transiently transfected with an empty expression vector is the control ($n = 2$).

reported herein indicate that *Rhox10* also has a second function—*LINE1* defense—which raises the possibility that two distinct *Rhox10*-associated regulatory circuits evolved: one to support germ cell progression and another to defend against TEs. In support of the notion that *Rhox10* evolved to regulate ProSG differentiation and migration, we previously showed that *Rhox10* regulates numerous differentiation and migration genes (26). However, we cannot rule out that the ProSG differentiation and migration defects observed in *Rhox10*-null mice are a secondary consequence of elevated *LINE1* expression in ProSG. This is an interesting possibility, but if true, it is unlikely to be due to *LINE1*-induced mutations, as random mutations would presumably not consistently cause the stage-specific ProSG progression defect found in *Rhox10*-null mice (26). A more likely scenario is that the elevated *LINE1* ORF1 expression we observed in *Rhox10*-null germ cells is responsible, as this and elevated ORF2p expression have both been found to lead to cell toxicity in other contexts (7–9). Another possibility is that *LINE1* activation in *Rhox10*-null ProSG perturbs the epigenetic chromatin landscape in these cells, leading to impaired differentiation. This

putative mechanism is supported by studies showing that aberrant TE activation perturbs the chromatin landscape (65).

Our finding that RHOX10 positively regulates *Piwi2* expression is intriguing given that *Piwi2* encodes a key endonuclease that silences TEs and greatly curbs *LINE1* mobilization and thus is essential for germline integrity and normal spermatogenesis (14). Our findings support a model in which RHOX10 transcriptionally activates the *Piwi2* gene during the PGC–ProSG transition, leading to production of PIWIL2 protein, which then acts to drive piRNA pathway-mediated suppression of *LINE1* promoters and consequent reduced transposition (Fig. 3I). Several findings support this model: 1) *Piwi2* is down-regulated in *Rhox10*-null fetal germ cells (as shown by RNA-seq analysis and verified by qPCR and Western analysis); 2) *Rhox10* dramatically increases expression from the *Piwi2* promoter (as demonstrated by reporter analysis); 3) RHOX10 occupies the *Piwi2* promoter (as shown by ChIP analysis); 4) forced expression of PIWIL2 in a germ cell line reverses the defect in *LINE1* transposition caused by RHOX10 knockdown; 5) *Piwi2* exhibits enriched expression

in T1-ProSG (66); 6) *Rhox10* and *Piwil2* have both been reported to be first detectably expressed in male PGCs on precisely the same day of fetal development: E12.5 (16, 36); and 7) piRNAs known to be bound to PIWIL2 (53) exhibit reduced expression in *Rhox10*-null ProSG.

Rhox10 is expressed not only in ProSG but also in spermatogonia and early spermatocytes (51), raising the possibility that it functions in *LINE1* defense at these germ cell stages as well. Our results suggest that other *Rhox* genes might also function in *LINE1* defense. For example, our in vivo *LINE1* reporter analysis of *Rhox10*-null versus whole *Rhox* cluster-null mice suggested that *Rhox10* is only responsible for ~1/2 of *LINE1* suppression mediated by the entire *Rhox* cluster in fetal germ cells. Candidates to also play a role are *Rhox1*, *Rhox2*, *Rhox4*, *Rhox5*, and *Rhox7*, all of which are expressed in male fetal germ cells between E12.5 and 15.5 (36, 37). We also demonstrated that mouse *Rhox* family members expressed in other biological contexts suppress *LINE1* transposition. Interestingly, these *Rhox* genes—*Rhox3*, *Rhox6*, *Rhox8*, and *Rhox11*—each have a different pattern of expression in the reproductive tract (24). *Rhox3* messenger (m)RNA and RHOX3 protein have a postnatal expression pattern consistent with selective expression in spermatids (67), as does *Rhox11* mRNA and RHOX11 protein. This raises the possibility that these two RHOX transcription factors function in *LINE1* defense in postmeiotic germ cells. *Rhox6* is primarily expressed in female PGCs (36), suggesting that it might suppress *LINE1* elements in the female germline. Indeed, many *Rhox* genes are expressed in the female germline (19). *Rhox8* is expressed in Sertoli cells and is necessary for normal spermatogenesis (68), raising the possibility that *Rhox8* suppresses *LINE1* elements in these testicular nurse cells. Finally, we suggest the placenta is another site where *Rhox* genes may provide *LINE1* defense. The placenta has an intimate relationship with *LINE1* elements, as this organ has widely co-opted these parasitic elements for its own purposes (69, 70) and has a hypomethylated genome potentially permissive to *LINE1* element mobilization (71). Given that all known *Rhox* genes are expressed in the placenta (19), we suggest they may function to reduce *LINE1* mobilization to a manageable level to avoid overt toxicity. Consistent with this, loss of the entire *Rhox* cluster causes embryonic lethality (26) and mice lacking the orphan *Rhox* gene, *Esx1* (24), exhibit an embryonic growth defect (72). Surprisingly, we obtained in vitro evidence that unlike other *Rhox* family members, *Rhox4* promotes, rather than inhibits, *LINE1* transposition. This raises the possibility that individual RHOX transcription factors have heterogeneous roles in *LINE1* defense, which requires future testing in vivo.

Interestingly, humans have only three RHOX genes—*RHOXF1* and two almost identical *RHOXF2* genes—all of which are expressed only in the male and female germline (29). Our finding that *RHOXF2* represses *LINE1* transposition in several different cell lines raises the possibility it also has this activity in humans in vivo at the sites where it is normally expressed: prespermatogonia in human fetal testes, spermatogonia and leptotene spermatocytes in adult male human testes, and oocytes in adult human female ovaries (29).

While several lines of evidence indicated that a PIWIL2-dependent mechanism is responsible, at least in part, for RHOX10-mediated suppression of *LINE1* transposition in vivo, RHOX transcription factors may also use other mechanisms to suppress *LINE1* transposition. Our in vitro reporter experiments demonstrated that several RHOX transcription factors suppress *LINE1* transposition in NIH 3T3, HEK-293T, and K562 cells, despite the fact that these somatic cell lines have trace or undetectable *Piwil2*/*PIWIL2* mRNA expression (SI Appendix, Fig. S2). This suggests that RHOX transcription factors also suppress *LINE1* transposition by a PIWIL2-independent mechanism, which may work in conjunction with the PIWIL2-dependent mechanism we uncovered.

The ability of RHOX transcription factors to suppress *LINE1* elements provides an explanation for the paradoxical finding that *Rhox* gene copy number varies greatly (by more than 10-fold) in different mammalian species, including expansion in mice compared to rats, and differential copy number in different primate species (19, 24, 57). In particular, we postulate that the *Rhox* gene cluster is in an evolutionary arms race with *LINE1* elements, leading to *Rhox* cluster expansion as a means to better repress the deleterious effects of *LINE1* elements, including mutagenetic effects to the germline. We suggest that the differential copy number of *Rhox* genes and their different expression patterns in different species reflect unique selective forces acting in different species to suppress *LINE1* elements.

In conclusion, we report that a large set of DNA-binding factors encoded by the X chromosome have the potential to provide defense against TEs in both the germline and soma of mammals. It remains for future studies to determine the full extent of their repertoire, both in terms of the types of TEs targeted and the cellular contexts in which they act.

Methods

Mice. This study was carried out in strict accordance with the guidelines of the Institutional Animal Care and Use Committee at the University of California San Diego (protocol S09160). All mice were housed under a 12-h light:12-h dark cycle and provided with food and water ad libitum. All mouse strains used for analysis were backcrossed to C57BL/6J for at least eight passages.

Mammalian Cell Culture, Plasmids, Transfections, and Luciferase Analysis. NIH/3T3, GC1, HEK-293T, and GS cells were used for transfection. For transient transfection experiments, the coding sequences of mouse *Rhox* and human RHOX genes were cloned into the plasmid cloning (pc)DNA 5/FRT expression vector (Invitrogen). The *Piwil2* coding sequence was cloned into pcDNA 3.1 (–) expression vector. *RHOXF1* and *RHOXF2* shRNA plasmids were generated by inserting small interfering RNA sequences into the pLLU2G backbone. Mouse (*ORFeus*) and human (L1_{RP}) *LINE1* reporter constructs were generated as previously described (48). The mutant *RHOXF2* constructs were obtained from two previous studies (60, 61) and cloned into the pRES-hrGFP expression vector. The *Piwil2* and *Tdrd1* promoters were separately cloned into the pGL3 vector (Promega). For a detailed description, reference SI Appendix, Materials and Methods.

ChIP Analysis. GS cells (5 million) were cross-linked, lysed, and homogenized for chromatin preparation. Detailed protocols are described in SI Appendix, Materials and Methods.

qRT-PCR and TaqMan Assay. Total cellular RNA was isolated as previously described (73). Reverse transcription and qPCR analysis were performed following the manufacturers' protocol. The primers used are listed in SI Appendix, Table S1. Results were from at least three independent replicates. Statistical significance was determined using the paired Student's *t* test. For detailed description, reference SI Appendix, Materials and Methods.

Western Blotting Analysis. Western blot analysis was performed as previously described (74). Quantification of the blots was performed using NIH ImageJ (1.8.0). Statistical significance was determined using the paired Student's *t* test. For detailed description, reference SI Appendix, Materials and Methods.

RNA-Seq Analysis. Single testicular cells were isolated from fetal testes using a two-step enzymatic digestion protocol described previously (75). For each sample analyzed, testes from three to four fetuses were pooled. RNA-seq was performed as described previously (76). The average number of reads per sample ranged from ~38 to 46 million. Detailed RNA-seq analyses are described in SI Appendix, Materials and Methods.

ddPCR. ddPCR reactions were performed as previously described (38). For detailed description, reference SI Appendix, Materials and Methods.

Bisulfite Sequencing Analysis. Cells were processed for methylation analysis using the EpiTect Plus LyseAll Bisulfite Kit (Qiagen). Bisulfite treatment of fetal DNA was conducted with the EpiTect Plus DNA Bisulfite Kit (Qiagen). PCR amplification, gel extraction, thymine and adenine cloning, and sequence

analysis were performed as previously described (38). For detailed description, reference *SI Appendix, Materials and Methods*.

Immunofluorescence Analysis. Immunofluorescence analysis was performed as previously described (66, 77). For a detailed description, reference *SI Appendix, Materials and Methods*.

Data Availability. RNA-seq data have been deposited in the National Center for Biotechnology Information's Gene Expression Omnibus database ([GSE160600](https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE160600)) (78). All other study data are included in the article and/or supporting information.

1. R. N. Platt II, M. W. Vandeweghe, D. A. Ray, Mammalian transposable elements and their impacts on genome evolution. *Chromosome Res.* **26**, 25–43 (2018).
2. E. S. Lander *et al.*; International Human Genome Sequencing Consortium, Initial sequencing and analysis of the human genome. *Nature* **409**, 860–921 (2001). Corrected in: *Nature* **412**, 565 (2001).
3. C. Ensnaut, J. Maestre, T. Heidmann, Human LINE retrotransposons generate processed pseudogenes. *Nat. Genet.* **24**, 363–367 (2000).
4. J. L. Garcia-Perez, A. J. Doucet, A. Bucheton, J. V. Moran, N. Gilbert, Distinct mechanisms for trans-mediated mobilization of cellular RNAs by the LINE-1 reverse transcriptase. *Genome Res.* **17**, 602–611 (2007).
5. P. Tristan-Ramos *et al.*, sRNA/L1 retrotransposition: Using siRNAs and miRNAs to expand the applications of the cell culture-based LINE-1 retrotransposition assay. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **375**, 20190346 (2020).
6. J. L. Goodier, H. H. Kazazian Jr, Retrotransposons revisited: The restraint and rehabilitation of parasites. *Cell* **135**, 23–35 (2008).
7. B. Tiwari *et al.*, p53 directly represses human LINE1 transposons. *Genes Dev.* **34**, 1439–1451 (2020).
8. N. A. Wallace, V. P. Belancio, P. L. Deininger, L1 mobile element expression causes multiple types of toxicity. *Gene* **419**, 75–81 (2008).
9. S. L. Gasior, T. P. Wakeman, B. Xu, P. L. Deininger, The human LINE-1 retrotransposon creates DNA double-strand breaks. *J. Mol. Biol.* **357**, 1383–1393 (2006).
10. Ö. Deniz, J. M. Frost, M. R. Branco, Regulation of transposable elements by DNA modifications. *Nat. Rev. Genet.* **20**, 417–431 (2019).
11. S. A. Smallwood, G. Kelsey, De novo DNA methylation: A germ cell perspective. *Trends Genet.* **28**, 33–42 (2012).
12. J. R. McCarrey, Toward a more precise and informative nomenclature describing fetal and neonatal male germ cells in rodents. *Biol. Reprod.* **89**, 47 (2013).
13. S. Kim *et al.*, PRMT5 protects genomic integrity during global DNA demethylation in primordial germ cells and preimplantation embryos. *Mol. Cell* **56**, 564–579 (2014).
14. Y. W. Iwasaki, M. C. Siomi, H. Siomi, PIWI-interacting RNA: Its biogenesis and functions. *Annu. Rev. Biochem.* **84**, 405–433 (2015).
15. S. Kuramochi-Miyagawa *et al.*, Two mouse Piwi-related genes: Miwi and Mili. *Mech. Dev.* **108**, 121–133 (2001).
16. S. Kuramochi-Miyagawa *et al.*, DNA methylation of retrotransposon genes is regulated by Piwi family members MILI and MIWI2 in murine fetal testes. *Genes Dev.* **22**, 908–917 (2008).
17. K. Kojima-Kita *et al.*, MIWI2 as an effector of DNA methylation and gene silencing in embryonic male germ cells. *Cell Rep.* **16**, 2819–2828 (2016).
18. M. F. Wilkinson, J. Kleeman, J. Richards, C. L. MacLeod, A novel oncofetal gene is expressed in a stage-specific manner in murine embryonic development. *Dev. Biol.* **141**, 451–455 (1990).
19. J. A. Maclean II *et al.*, RhoX: A new homeobox gene cluster. *Cell* **120**, 369–382 (2005).
20. M. Jackson *et al.*, A murine specific expansion of the RhoX cluster involved in embryonic stem cell biology is under natural selection. *BMC Genomics* **7**, 212 (2006).
21. L. Morris, J. Gordon, C. C. Blackburn, Identification of a tandem duplicated array in the RhoX alpha locus on mouse chromosome X. *Mamm. Genome* **17**, 178–187 (2006).
22. X. Wang, J. Zhang, Remarkable expansions of an X-linked reproductive homeobox gene cluster in rodent evolution. *Genomics* **88**, 34–43 (2006).
23. C. B. Geyer, E. M. Eddy, Identification and characterization of RhoX13, a novel X-linked mouse homeobox gene. *Gene* **423**, 194–200 (2008).
24. J. A. Maclean II, M. F. Wilkinson, The RhoX genes. *Reproduction* **140**, 195–213 (2010).
25. Z. Hu *et al.*, Androgen-induced RhoX homeobox genes modulate the expression of AR-regulated genes. *Mol. Endocrinol.* **24**, 60–75 (2010).
26. H. W. Song *et al.*, The homeobox transcription factor RHOX10 drives mouse spermatogonial stem cell establishment. *Cell Rep.* **17**, 149–164 (2016).
27. J. P. Welborn *et al.*, RhoX8 ablation in the Sertoli cells using a tissue-specific RNAi approach results in impaired male fertility in mice. *Biol. Reprod.* **93**, 8 (2015).
28. J. T. Busada *et al.*, RhoX13 is required for a quantitatively normal first wave of spermatogenesis in mice. *Reproduction* **152**, 379–388 (2016).
29. H. W. Song *et al.*, The RHOX homeobox gene cluster is selectively expressed in human oocytes and male germ cells. *Hum. Reprod.* **28**, 1635–1646 (2013).
30. A. W. Sasaki *et al.*, The oncofetal gene Pem encodes a homeodomain and is regulated in primordial and pre-muscle stem cells. *Mech. Dev.* **34**, 155–164 (1991).
31. M. Oda *et al.*, DNA methylation regulates long-range gene silencing of an X-linked homeobox gene cluster in a lineage-specific manner. *Genes Dev.* **20**, 3382–3394 (2006).
32. Q. Li, D. L. Bartlett, M. C. Gorro, M. E. O'Malley, Z. S. Guo, Three epigenetic drugs upregulate homeobox gene RhoX5 in cancer cells through overlapping and distinct molecular mechanisms. *Mol. Pharmacol.* **76**, 1072–1081 (2009).
33. J. A. Maclean *et al.*, The rhoX homeobox gene cluster is imprinted and selectively targeted for regulation by histone h1 and DNA methylation. *Mol. Cell. Biol.* **31**, 1275–1287 (2011).
34. M. E. Richardson, A. Bleiziffer, F. Tüttelmann, J. Gromoll, M. F. Wilkinson, Epigenetic regulation of the RHOX homeobox gene cluster and its association with human male infertility. *Hum. Mol. Genet.* **23**, 12–23 (2014).
35. M. L. Maeder *et al.*, Targeted DNA demethylation and activation of endogenous genes using programmable TALE-TET1 fusion proteins. *Nat. Biotechnol.* **31**, 1137–1142 (2013).
36. H. Daggag *et al.*, The rhoX homeobox gene family shows sexually dimorphic and dynamic expression during mouse embryonic gonad development. *Biol. Reprod.* **79**, 468–474 (2008).
37. J. L. Pitman, T. P. Lin, J. E. Kleeman, G. F. Erickson, C. L. MacLeod, Normal reproductive and macrophage function in Pem homeobox gene-deficient mice. *Dev. Biol.* **202**, 196–214 (1998).
38. S. J. Newkirk *et al.*, Intact piRNA pathway prevents L1 mobilization in male meiosis. *Proc. Natl. Acad. Sci. U.S.A.* **114**, E5635–E5644 (2017).
39. E. M. Ostertag *et al.*, A mouse model of human L1 retrotransposition. *Nat. Genet.* **32**, 655–660 (2002).
40. W. An *et al.*, Conditional activation of a single-copy L1 transgene in mice by Cre. *Genesis* **46**, 373–383 (2008).
41. W. An *et al.*, Active retrotransposition by a synthetic L1 element in mice. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 18662–18667 (2006).
42. A. R. Muotri *et al.*, Somatic mosaicism in neuronal precursor cells mediated by L1 retrotransposition. *Nature* **435**, 903–910 (2005).
43. D. V. Babushok, E. M. Ostertag, C. E. Courtney, J. M. Choi, H. H. Kazazian Jr, L1 integration in a transgenic mouse model. *Genome Res.* **16**, 240–250 (2006).
44. H. Kano *et al.*, L1 retrotransposition occurs mainly in embryogenesis and creates somatic mosaicism. *Genes Dev.* **23**, 1303–1312 (2009).
45. F. C. Grandi, J. M. Rosser, W. An, LINE-1-derived poly(A) microsatellites undergo rapid shortening and create somatic and germline mosaicism in mice. *Mol. Biol. Evol.* **30**, 503–512 (2013).
46. N. G. Coufal *et al.*, L1 retrotransposition in human neural progenitor cells. *Nature* **460**, 1127–1131 (2009).
47. J. K. Baillie *et al.*, Somatic retrotransposition alters the genetic landscape of the human brain. *Nature* **479**, 534–537 (2011).
48. Y. Xie, J. M. Rosser, T. L. Thompson, J. D. Boeke, W. An, Characterization of L1 retrotransposition with high-throughput dual-luciferase assays. *Nucleic Acids Res.* **39**, e16 (2011).
49. S. De Fazio *et al.*, The endonuclease activity of Mili fuels piRNA amplification that silences LINE1 elements. *Nature* **480**, 259–263 (2011).
50. M. Reuter *et al.*, Miwi catalysis is required for piRNA amplification-independent LINE1 transposon silencing. *Nature* **480**, 264–267 (2011).
51. H. W. Song *et al.*, Dynamic expression pattern and subcellular localization of the RhoX10 homeobox transcription factor during early germ cell development. *Reproduction* **143**, 611–624 (2012).
52. J. M. Wenda *et al.*, Distinct roles of RNA helicases MVH and TDRD9 in PIWI slicing-triggered mammalian piRNA biogenesis and function. *Dev. Cell* **41**, 623–637.e9 (2017).
53. A. A. Aravin *et al.*, A piRNA pathway primed by individual transposons is linked to de novo DNA methylation in mice. *Mol. Cell* **31**, 785–799 (2008).
54. K. Inoue, K. Ichyanagi, K. Fukuda, M. Glinka, H. Sasaki, Switching of dominant retrotransposon silencing strategies from posttranscriptional to transcriptional mechanisms during male germ-cell development in mice. *PLoS Genet.* **13**, e1006926 (2017).
55. T. Watanabe, X. Cui, Z. Yuan, H. Qi, H. Lin, MIWI2 targets RNAs transcribed from piRNA-dependent regions to drive DNA methylation in mouse prospermatogonia. *EMBO J.* **37**, e95329 (2018).
56. S. E. Lee, S. Y. Lee, K. A. Lee, RhoX in mammalian reproduction and development. *Clin. Exp. Reprod. Med.* **40**, 107–114 (2013).
57. A. L. Niu *et al.*, Rapid evolution and copy number variation of primate RHOXF2, an X-linked homeobox gene involved in male reproduction and possibly brain function. *BMC Evol. Biol.* **11**, 298 (2011).
58. C. M. Wayne, J. A. MacLean, G. Cornwall, M. F. Wilkinson, Two novel human X-linked homeobox genes, hPEPP1 and hPEPP2, selectively expressed in the testis. *Gene* **301**, 1–11 (2002).
59. E. M. Ostertag, E. T. Prak, R. J. DeBerardinis, J. V. Moran, H. H. Kazazian Jr, Determination of L1 retrotransposition kinetics in cultured cells. *Nucleic Acids Res.* **28**, 1418–1423 (2000).

60. J. Borgmann *et al.*, The human RHOX gene cluster: Target genes and functional analysis of gene variants in infertile men. *Hum. Mol. Genet.* **25**, 4898–4910 (2016).
61. C. Frainais *et al.*, RHOXF2 gene, a new candidate gene for spermatogenesis failure. *Basic Clin. Androl.* **24**, 3 (2014).
62. Y. S. Hwang *et al.*, Reconstitution of prospermatogonial specification in vitro from human induced pluripotent stem cells. *Nat. Commun.* **11**, 5656 (2020).
63. M. K. Rao, C. M. Wayne, M. L. Meistrich, M. F. Wilkinson, Pem homeobox gene promoter sequences that direct transcription in a Sertoli cell-specific, stage-specific, and androgen-dependent manner in the testis in vivo. *Mol. Endocrinol.* **17**, 223–233 (2003).
64. N. Takasaki, T. Rankin, J. Dean, Normal gonadal development in mice lacking GPBOX, a homeobox protein expressed in germ cells at the onset of sexual dimorphism. *Mol. Cell. Biol.* **21**, 8197–8202 (2001).
65. N. Zamudio *et al.*, DNA methylation restrains transposons from adopting a chromatin signature permissive for meiotic recombination. *Genes Dev.* **29**, 1256–1270 (2015).
66. K. Tan, H. W. Song, M. F. Wilkinson, Single-cell RNAseq analysis of testicular germ and somatic cell development during the perinatal period. *Development* **147**, dev183251 (2020).
67. H. W. Song *et al.*, shRNA off-target effects in vivo: Impaired endogenous siRNA expression and spermatogenic defects. *PLoS One* **10**, e0118549 (2015).
68. R. M. Brown, M. G. Davis, K. Hayashi, J. A. MacLean, Regulated expression of Rhox8 in the mouse ovary: Evidence for the role of progesterone and RHOX5 in granulosa cells. *Biol. Reprod.* **88**, 126 (2013).
69. D. Haig, Going retro: Transposable elements, embryonic stem cells, and the mammalian placenta (retrospective on DOI 10.1002/bies.201300059). *BioEssays* **37**, 1154 (2015).
70. D. Jangam, C. Feschotte, E. Betrán, Transposable element domestication as an adaptation to evolutionary conflicts. *Trends Genet.* **33**, 817–831 (2017).
71. D. I. Schroeder *et al.*, Early developmental and evolutionary origins of gene body DNA methylation patterns in mammalian placentas. *PLoS Genet.* **11**, e1005442 (2015).
72. Y. Li, R. R. Behringer, Esx1 is an X-chromosome-imprinted regulator of placental development and fetal growth. *Nat. Genet.* **20**, 309–311 (1998).
73. M. Ramaiah *et al.*, A microRNA cluster in the Fragile-X region expressed during spermatogenesis targets FMR1. *EMBO Rep.* **20**, e46566 (2019).
74. K. Tan *et al.*, Impaired imprinted X chromosome inactivation is responsible for the skewed sex ratio following in vitro fertilization. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 3197–3202 (2016).
75. A. Sohni *et al.*, The neonatal and adult human testis defined at the single-cell level. *Cell Rep.* **26**, 1501–1517.e4 (2019).
76. K. Tan *et al.*, Transcriptome profiling reveals signaling conditions dictating human spermatogonia fate in vitro. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 17832–17841 (2020).
77. K. Tan *et al.*, The role of the NMD factor UPF3B in olfactory sensory neurons. *eLife* **9**, e57525 (2020).
78. K. Tan, M. F. Wilkinson, The *Rhox* gene cluster suppresses germline *LINE1* transposition. *Gene Expression Omnibus*. <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE160600>. Deposited 1 November 2020.