

Commentary

Local Contrasts in Concentration of Ambient Particulate Air Pollution (PM_{2.5}) and Incidence of Alzheimer's Disease and Dementia: Results from the Betula Cohort in Northern Sweden

Daniel Oudin Åström^a, Rolf Adolfsson^b, David Segersson^c, Bertil Forsberg^a and Anna Oudin^{a,d,*}

^a*Sustainable health, Department for Public Health and Clinical Medicine, Umeå University, Umeå, Sweden*

^b*Division of Psychiatry, Department of Clinical Sciences, Umeå University, Umeå, Sweden*

^c*Swedish Meteorological and Hydrological Institute, Norrköping, Sweden*

^d*Environment, Society and Health, Occupational and Environmental Medicine, Department for Laboratory Medicine, Lund University, Lund, Sweden*

Accepted 19 February 2021

Pre-press 18 March 2021

Abstract. Exposure to fine particulate air pollution (PM_{2.5}) is emerging as a risk factor for Alzheimer's disease (AD), but existing studies are still limited and heterogeneous. We have previously studied the association between dementia (AD and vascular dementia) and PM_{2.5} stemming from vehicle exhaust and wood-smoke in the Betula cohort in Northern Sweden. The aim of this commentary is to estimate the association between total PM_{2.5} and dementia in the Betula cohort, which is more relevant to include in future meta-estimates than the source-specific estimates. The hazard ratio for incident dementia associated with a 1 µg/m³ increase in local PM_{2.5} was 1.38 (95% confidence interval: 0.99 – 1.92). The interpretation of our results is that they indicate an association between local contrasts in concentration of PM_{2.5} at the residential address and incidence of dementia in a low-level setting.

Keywords: Air pollution, Alzheimer's disease, dementia, particulate air pollution, PM_{2.5}

Worldwide, Alzheimer's disease (AD) is one of the leading causes of impaired health and mortality and as populations continue to age the share of individuals with AD is projected to increase in the future. Recently, the Lancet Commission of 2020 added air pollution as one of three new modifiable risk factors

to the 2017 Lancet Commission on dementia prevention, intervention, and care life-course model, now consisting of twelve factors [1].

We have previously reported results from exposure to air pollutants and dementia from the prospective Betula cohort study from Northern Sweden, in detail described elsewhere [2]. In the Betula cohort, we observed clear associations between traffic-related air pollution (nitrogen oxides) and incidence of AD [3] as well as associations between fine particulate

*Correspondence to: Anna Oudin, Sustainable Health, Department of Public Health and Clinical Medicine, Umeå University, 901 87, Umeå, Sweden. Tel.: +46 90 785 08 73; E-mail: anna.oudin@umu.se.

air pollution (PM_{2.5}) stemming from both vehicle exhaust and domestic wood burning [4]. We also observed that residential noise did not seem to be a risk factor for AD, or confounding the association between air pollution and AD [5], and that adverse effects of air pollution on AD incidence were not limited to, or stronger in, *APOE* ϵ 4 carriers than in the total population [6]. We have not yet presented an estimate for the total effect of PM_{2.5} on dementia incidence, however. Such an estimate is important to communicate now, as air pollution is beginning to be an established risk factor for AD and meta-analysis of effect estimates are sought for since substantial heterogeneity in published studies still remain [7, 8].

The aim of this commentary is thus to describe the association between total PM_{2.5} and dementia in the Betula cohort, using the same data and methods as when investigating source-specific exposure [4]. In short, a Cox regression model assuming proportional hazards is used, adjusting for the same individual level factors as previously (physical activity, smoking, sex, body mass index, waist-hip ratio, alcohol, and age) in the Betula study of approximate 1,806 individuals over a 15-year period, where the mean age at start of follow-up was 68 years, 57% were women, and the proportion of *APOE* ϵ 4 carriers were 28%. The hazard ratio for incident dementia associated with a 1 $\mu\text{g}/\text{m}^3$ increase in total PM_{2.5}, was 1.38 (95% confidence interval: 0.99 – 1.92). The interpretation of our results is that they indicate an association between total concentration of PM_{2.5} at residential address and incidence of dementia in the Betula cohort. Supported by other epidemiological studies, as well as laboratory studies [1, 9, 10], the results add to already existing evidence for air pollution to increase dementia risk. The estimate is quite high compared to the meta-estimate of a relative risk of 1.08 per 5 $\mu\text{g}/\text{m}^3$ increase in PM_{2.5} calculated by Yu and colleagues [7]. This is most likely due to the fine spatial resolution in our air pollution model, meaning that we model local contrasts in air pollution concentrations. The study area, Umeå municipality, is furthermore rather small (33.5 km²), and we have not access to any data on possible contrasts in long-range-transported PM_{2.5}. Our results should therefore be interpreted as risks associated with differences in local exposures, and may therefore be in line with what has been observed by others in terms of local air pollution having a substantially higher impact on mortality per unit compared to regional air pollution [11]. This could have implications on health impact assessments attributing dementia cases to air

pollution exposure, when local models are available, and populations are similar, it might be relevant to consider if a higher dose-response-function should be used than, for example, applied by Rittner and colleagues [12].

Based on strong evidence, the Lancet Commission of 2020 added air pollution as a modifiable risk factor and adding our result in future meta-analyses will help to gain a better understanding of the magnitude and precision on evidence suggesting total air pollution levels to be associated with incident dementia. Our findings further suggest the need to consider ambient air pollution associated with dementia pathology in public health policy decisions.

ACKNOWLEDGMENTS

This work was supported by Swedish Research Council Formas under grant agreement number 2017-00898 (AO).

Authors' disclosures available online (<https://www.j-alz.com/manuscript-disclosures/20-1538r1>).

REFERENCES

- [1] Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S, Brayne C, Burns A, Cohen-Mansfield J, Cooper C (2020) Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet* **396**, 413-446.
- [2] Nilsson L-G, Bäckman L, Erngrund K, Nyberg L, Adolfsson R, Bucht G, Karlsson S, Widing M, Winblad B (1997) The Betula prospective cohort study: Memory, health, and aging. *Aging Neuropsychol Cogn* **4**, 1-32.
- [3] Oudin A, Forsberg B, Adolfsson AN, Lind N, Modig L, Nordin M, Nordin S, Adolfsson R, Nilsson L-G (2015) Traffic-related air pollution and dementia incidence in Northern Sweden: A longitudinal study. *Environ Health Perspect* **124**, 306-312.
- [4] Oudin A, Segerström D, Adolfsson R, Forsberg B (2018) Association between air pollution from residential wood burning and dementia incidence in a longitudinal study in Northern Sweden. *PLoS One* **13**, e0198283.
- [5] Andersson J, Oudin A, Sundström A, Forsberg B, Adolfsson R, Nordin M (2018) Road traffic noise, air pollution, and risk of dementia—results from the Betula project. *Environ Res* **166**, 334-339.
- [6] Oudin A, Andersson J, Sundström A, Nordin Adolfsson A, Oudin -ström D, Adolfsson R, Forsberg B, Nordin M (2019) Traffic-related air pollution as a risk factor for dementia: No clear modifying effects of *APOE* ϵ 4 in the Betula cohort. *J Alzheimers Dis* **71**, 733-740.
- [7] Yu X, Zheng L, Jiang W, Zhang D (2020) Exposure to air pollution and cognitive impairment risk: A meta-analysis of longitudinal cohort studies with dose-response analysis. *J Glob Health* **10**, 010417.
- [8] Peters R, Ee N, Peters J, Booth A, Mudway I, Anstey KJ (2019) Air pollution and dementia: A systematic review. *J Alzheimers Dis* **70**(s1), S145-S163.

- [9] Oudin A (2017) Air pollution and dementia. *J Public Health Emerg* **1**, doi: 10.21037/jphe.2017.04.05
- [10] Oudin A (2020) Short review: Air pollution, noise and lack of greenness as risk factors for Alzheimer's disease-epidemiologic and experimental evidence. *Neurochem Int* **134**, 104646.
- [11] Turner MC, Jerrett M, Pope III CA, Krewski D, Gapstur SM, Diver WR, Beckerman BS, Marshall JD, Su J, Crouse DL (2016) Long-term ozone exposure and mortality in a large prospective study. *Am J Respir Crit Care Med* **193**, 1134-1142.
- [12] Rittner R, Flanagan E, Oudin A, Malmqvist E (2020) Health impacts from ambient particle exposure in southern Sweden. *Int J Environ Res Public Health* **17**, 5064.