



“Lung volume reduction in emphysema: a pragmatic prospective cohort study”.  
 Christophe Doms, Astrid Blondeel, Laurens J. Ceulemans, Johan Coolen,  
 Stephanie Everaerts, Heleen Demeyer, Thierry Troosters Geert Verleden, Dirk Van  
 Raemdonck and Wim Janssens. *ERJ Open Res* 2021; 7: 00877-2020

This article was originally published with an error in the key to the symbols in figure 3. The corrected figure is shown below and has been corrected in the article itself.




FIGURE 3 Absolute changes from baseline in key outcome measures at 3 and 6 months post-allocation to endoscopic lung volume reduction (ELVR) group, no-ELVR group and lung volume reduction surgery (LVRS) group. Data presented are mean±SEM for a) forced expiratory volume in 1 s (FEV<sub>1</sub>), b) residual volume (RV), c) 6-min walk distance (6MWD) and d) St George’s Respiratory Questionnaire (SGRQ) total score.

ERJ Open Res 2021; In press | DOI: 10.1183/23120541.50877-2020 | Copyright ©The authors 2021. This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact [permissions@ersnet.org](mailto:permissions@ersnet.org)

