



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Original Contribution

Effect of prone versus supine position in COVID-19 patients: A systematic review and meta-analysis



Ee Xin Chua, MBBS^a, Syed Mohd Ikhmal Syed Mohd Zahir, MBBS^a, Ka Ting Ng, MBChB^{a,*}, Wan Yi Teoh, MBChB^b, Mohd Shahnaz Hasan, MANaes^a, Shairil Rahayu Binti Ruslan, MANaes^a, Mohammed F. Abosamak, MD^c

^a Department of Anaesthesiology, Faculty of Medicine, University of Malaya, Jalan Universiti, Kuala Lumpur, Malaysia

^b Faculty of Medicine, University of Liverpool, Liverpool L69 3BX, United Kingdom

^c Department of Anaesthesia and Intensive care medicine, Faculty of medicine, Tanta University, Egypt

ARTICLE INFO

Keywords:

Prone
Supine
Coronavirus Disease 2019
Oxygenation
Meta-analysis

ABSTRACT

Study objective: To review the effects of prone position and supine position on oxygenation parameters in patients with Coronavirus Disease 2019 (COVID-19).

Design: Systematic review and meta-analysis of non-randomized trials.

Patients: Databases of EMBASE, MEDLINE and CENTRAL were systematically searched from its inception until March 2021.

Interventions: COVID-19 patients being positioned in the prone position either whilst awake or mechanically ventilated.

Measurements: Primary outcomes were oxygenation parameters (PaO₂/FiO₂ ratio, PaCO₂, SpO₂). Secondary outcomes included the rate of intubation and mortality rate.

Results: Thirty-five studies ($n = 1712$ patients) were included in this review. In comparison to the supine group, prone position significantly improved the PaO₂/FiO₂ ratio (study = 13, patients = 1002, Mean difference, MD 52.15, 95% CI 37.08 to 67.22; $p < 0.00001$) and SpO₂ (study = 11, patients = 998, MD 4.17, 95% CI 2.53 to 5.81; $p \leq 0.00001$). Patients received prone position were associated with lower incidence of mortality (study = 5, patients = 688, Odd ratio, OR 0.44, 95% CI 0.24 to 0.80; $p = 0.007$). No significant difference was noted in the incidence of intubation rate (study = 5, patients = 626, OR 1.20, 95% CI 0.77 to 1.86; $p = 0.42$) between the supine and prone groups.

Conclusion: Our meta-analysis demonstrated that prone position improved PaO₂/FiO₂ ratio with better SpO₂ than supine position in COVID-19 patients. Given the limited number of studies with small sample size and substantial heterogeneity of measured outcomes, further studies are warranted to standardize the regime of prone position to improve the certainty of evidence.

PROSPERO Registration: CRD42021234050

1. Introduction

The surge of Coronavirus Disease 2019 (COVID-19) cases has overwhelmed the healthcare services in some countries [1]. Persistent hypoxemia is a common presentation in patients with severe COVID-19. Large numbers of hospitalized COVID-19 patients fulfilled the criteria of acute respiratory distress syndrome (ARDS), which require invasive mechanical ventilation and high level of patient care [2]. The need for invasive mechanical ventilation has exceeded the capacity of healthcare

systems, especially for those heavily affected countries.

Prior to this pandemic of COVID-19, the application of prone position has been studied in awake patients with acute respiratory failure to avoid endotracheal intubation and reduce the need for intensive care units (ICU) admissions [3–5]. In view of a huge demand for mechanical ventilation among COVID-19 patients, the method of prone position has been extensively utilized to improve oxygenation, minimize incidence of intubation rate and ICU admission. The method of awake self-pronation can be applied in spontaneously breathing patients

* Corresponding author at: Department of Anaesthesiology, Faculty of Medicine, University of Malaya, Jalan Universiti, 50603 Kuala Lumpur, Malaysia.

E-mail address: katingng1@gmail.com (K.T. Ng).

<https://doi.org/10.1016/j.jclinane.2021.110406>

Received 6 March 2021; Received in revised form 24 May 2021; Accepted 29 May 2021

Available online 22 June 2021

0952-8180/© 2021 Elsevier Inc. All rights reserved.

receiving non-invasive ventilation or high-flow nasal cannula (HFNC) in the setting of outside ICU. Some studies have reported an improved oxygenation with a decrease in respiratory effort when applied early in COVID-19 patients with acute respiratory failure [6]. Prone position improves the ventilation/perfusion ratio and recruitment of the dorsal lung segments, resulting in the opening of collapsed dorsal alveoli with better gas exchange and oxygenation [7,8]. In mechanically ventilated non COVID-19 patients with severe ARDS, those who received prone ventilation had a lower mortality rate [9]. However, the clinical outcomes of prone position in COVID-19 (intubated/non-intubated) patients remain unclear from the literature. Therefore, a systematic review and meta-analysis is warranted to examine the effectiveness and safety of prone position in COVID-19 patients before any recommendation can be made.

We hypothesized that prone position improved oxygenation parameters in COVID-19 patients. Our primary aim was to investigate the effect of prone position on partial pressure of arterial oxygen/fraction of inspired oxygen (PaO₂/FiO₂) ratio. Secondary aims were to examine the effect of prone position on peripheral oxygen saturation (SpO₂), arterial partial pressure of carbon dioxide (PaCO₂), incidence of intubation, mortality rate and number of patients discharged alive.

2. Method

2.1. Data sources and searches

The review was conducted and reported in accordance to the Cochrane Handbook Systematic Reviews of Interventions [10] and the Preferred Reporting Items for Systematic Review and Meta-Analyses (PRISMA statement) 2015 [11], respectively. Our review protocol was published in a public database (PROSPERO CRD42021234050) prior to the literature search. The Population (COVID-19 patients), Intervention (prone position), Comparison (supine position) and Outcomes approach was used to formulate our review questions (Online Supplement eTable 1). Databases of EMBASE (OvidSP), MEDLINE (OvidSP) and Cochrane Central Register of Controlled Trials (CENTRAL) were systematically searched from their inception until March 2021. The [ClinicalTrials.gov](https://www.clinicaltrials.gov) and WHO International Clinical Trials Registry Platform Search Portal were searched for any ongoing or unpublished trials. The search terms and strategy used are showed in Online Supplement eTable 2. Inclusion criteria were:

1. Cohort or case-control observational studies (prospective or retrospective)
2. Randomized controlled trials
3. Examining prone position versus supine position
4. Adults (≥ 18 years old) diagnosed with COVID-19 infection

Case series, case reports and editorials were excluded. There was no restriction applied to the publication language and publication date. The reference lists of all the included studies were manually searched for any new or additional papers. The authors of relevant studies were contacted at least twice for any missing or unclear data.

2.2. Study selection, data items, data collection, and assessment of validity

Titles and abstracts were independently screened by two authors (EC and SMI) based on the pre-defined inclusion and exclusion criteria. Studies coded with “no” were excluded. Any studies coded with “yes” were retrieved for full text screening and screened independently by two authors (EC and SMI). Studies coded with “maybe” at any stage of study selection were resolved by third author (KN) via a discussion to achieve consensus. The selection of final included studies was discussed and agreed by all the authors.

Data of all included studies were extracted independently by two

authors (EC and SMI) using a standardized online data extraction form, which was piloted prior to the data extraction. Any disagreements were resolved by a third author (KN). In addition to the measured outcome, the following data, namely author, year of publication, country, study design, total sample size, mean age, gender, body mass index, citation, severity of APACHE II, duration of prone position and types of oxygen delivered were extracted.

2.3. Outcomes

Primary outcome of this review was PaO₂/FiO₂ ratio after prone and supine position. Secondary outcomes included SpO₂, PaCO₂, mortality rate, incidence of intubation rate and number of patients discharged alive.

2.4. Risk of bias in individual studies, summary of findings and GRADE assessment

The risk of bias for all included observational studies was independently assessed by two authors (EC and SMI) using the Newcastle-Ottawa Scale [12]. Study which scored ≥ 7 was considered as low risk of bias. Any disagreements were discussed and resolved by a third author (KN). The certainty of evidence (GRADE) was conducted by two authors (EC and SMI) independently. Several domains, namely the risk of bias, inconsistency, imprecision, indirectness and publication bias were assessed for all the measured outcomes. Any conflicts were discussed with a third author (KN).

2.5. Summary measures and synthesis of results

Review Manager version 5.3 (The Cochrane Collaboration, Copenhagen, Denmark) was used for all statistical analyses. A two-sided *p*-value of <0.05 was denoted as statistically significant. Findings were reported as odds ratio (OR) for dichotomous outcomes and mean difference (MD) for continuous outcomes, along with 95% confidence interval (CI). When the data were presented as median and range/interquartile range, these data were converted to mean and standard deviation [13]. I-square (I²) test was used to assess heterogeneity of studies. I² of less than 40%, 40–60% and more than 60% were categorized as low, moderate and substantial heterogeneity, respectively [10]. In view of the limited number of studies with high heterogeneity, the random-effect model (DerSimonian-Laird method) was used to pool data for all the measured outcomes. Subgroup analysis based on intubated and non-intubated COVID-19 studies was performed on the measured outcomes to explore the substantial degree of heterogeneity. Funnel plots were used to assess for publication bias.

3. Results

3.1. Study selection

The study selection process is summarized in the PRISMA flow diagram (Fig. 1). The search strategy yielded 1012 articles for titles and abstracts screening, of which 47 articles were retrieved for full text screening. After applying inclusion and exclusion criteria, thirty-five studies with a total of 1712 patients were included in this review. The list of excluded studies is showed in the Online Supplement eTable 3. Searching of trial registries found 9 ongoing studies (Online Supplement eTable 4).

3.2. Clinical characteristics of eligible studies

The clinical characteristics of included studies are demonstrated in Table 1. All the included studies were cohort studies. There were no randomized clinical studies investigating the effect of supine and prone position in COVID-19 patients. Of all, seven studies [14–20] had two




Fig. 1. PRISMA flow diagram.

separate cohorts for intervention and a control group whilst the rest of them [21–48] only had one cohort with pre-intervention (supine position) and post-intervention (prone position). The majority of included studies were of single centered, except two which were multi-centered [14,29]. The total sample size varied across all the included studies, ranging from 9 patients to 261 patients. Most of the recruited COVID-19 patients were from general wards, emergency departments and ICUs. The mean age of all the included studies ranged between 53 and 68 years old. Out of the 35 studies, twenty-one studies [14–16,18–20,23–26,29–36,38–40] utilized non-invasive ventilation (nasal cannula, simple face mask, venturi mask, non-rebreather mask, continuous positive airway pressure, bi-level positive airway pressure or high flow nasal cannula) and 14 studies [17,21,22,27,28,37,41–48] included mechanically ventilated COVID-19 patients. The duration of prone position sessions varied across all the studies from 30 min, 4 h and 16 h. The risk of bias for 27 studies [14–23,27,29–34,38–40,42–48] were assessed as low risk bias (≥ 7) based on the three domains of study selection, comparability and exposure (eTable 5). Five [24,26,36,37,41] and three [25,28,35] studies were rated as 6 and 5, respectively due to high risk of bias in the domain of comparability. The PRISMA checklist is outlined in the Online Supplementary eTable 6. The summary of findings and certainty of evidence assessment are showed in the Tables 2 and

3, respectively.

3.3. Primary outcome: PaO_2/FiO_2 ratio

In the combined data of thirteen studies [14,21,22,30,32,34,37,40,41,43,45–47] involving 1002 patients, our review demonstrated that COVID-19 patients with prone position were associated with higher PaO_2/FiO_2 ratio as compared to the supine group (MD 52.15, 95% CI 37.08 to 67.22; $p < 0.00001$, Fig. 2). However, statistical heterogeneity was observed as substantial ($I^2 = 87\%$). The magnitude and effect size of subgroup analysis for both intubated (studies = 8, $n = 579$ patients, MD 46.74, 95% CI 33.34 to 60.15; $p < 0.00001$) and non-intubated (studies = 5, $n = 423$ patients, MD 68.81, 95% CI 15.94 to 121.69; $p = 0.01$) studies remained significant, which was consistent with the overall finding of PaO_2/FiO_2 ratio. Funnel plot was asymmetry, suggestive of publication bias.

3.4. Secondary outcomes: SpO_2 , $PaCO_2$, incidence of mortality, incidence of intubation, number of patients discharged alive

Eleven studies [16,18,21,24,25,28,32,34,38,39,43] ($n = 998$ patients) examined the SpO_2 in COVID-19 patients who received prone and

Table 1
Baseline characteristics of all the included studies.

Author	Year	Design	N	Country	Setting	Age (mean + -SD)	BMI (mean + -SD)	Severity		Types of Oxygen Delivery Interface	Intervention	Successful duration of proning (standard)	Average/ Total duration of proning
								APACHE II	SOFA				
Ferrando et al	2020	Multicenter Cohort study (Non-intubated)	199	Spain & Andorra	NR	62.2 (12.1)	Int: 27.6 (4.9) Con: 27.2 (3.2)	Int: 11 (4.5) Con: 9.2 (5.4)	Int: 4.3 (0.8) Con: 4.0 (0)	Int & Con: High flow nasal oxygen therapy	Awake PP	>16 h/ day	NR
Jagan et al	2020	Single Center Cohort study (Non-intubated)	105	USA	NR	60.9 (15.4)	Int: 60.0% obese Con: 40.0% obese	Int: 6.7 (3.8) Con: 11 (6.8)	Int: 2.3 (0.8) Con: 3.7 (2.3)	Int & Con: Non-invasive ventilation	Awake PP	≥ 1 continuous hr on ≥ 5 occasions per day & for ≥ 1 continuous hr overnight at least 4 h twice daily up to 15 days	NR
Padrao et al	2020	Single Center Cohort study (Non-intubated)	166	Brazil	ED	58.1 (14.1)	Int: 58% BMI > 30 Con: 51% BMI > 30	NR	NR	Int: 34% nasal cannula 5% venturi mask 61% non-rebreathable mask Con: 49% nasal cannula 13% venturi mask 39% non-rebreathable mask	Awake PP	at least 4 h twice daily up to 15 days	NR
Shelhamer et al	2020	Single Center Cohort study (Intubated)	261	USA	Across all hospital services (medicine, surgery, ICU)	64 (13.4)	Int: 31.7 (5.8) Con: 31.6 (7.8)	Int: 17.9 (8.9) Con: 19 (11.9)	Int: QuickSOFA 61.3% Con: QuickSOFA 61.3%	Int & Con: Mechanical ventilation	PP	at least 16 h	NR
Weiss et al	2020	Single Center Cohort study (Intubated)	42	USA	ICU	59.9 (13.4)	34.2 (7.5)	NR	6.8 (2.5)	Mechanical ventilation	PP	at least 16 h	NR
Berrill et al	2020	Single Center Cohort study (Intubated)	34	UK	ICU	58.5 (11.1)	31 (5.1)	14 (4.7)	NR	Mechanical ventilation	PP	NR	Average: 16.5 (2.7) hrs Total duration: 63.5 (38.2), 2 days Median duration: 3 h (IQR 2)/CYCLE Cycles per patient: 5(6.25) NR
Burton et al	2020	Single Center Cohort study (Non-intubated)	20	UK	ICU	53.4 (8.3)	55% BMI >30	11.5 (IQR 5)	3 (IQR 0)	NIV: 20% CPAP, 8% CPAP+ BiPAP	Self-prone	NR	NR
Caputo et al	2020	Single Center Cohort study (Non-intubated)	50	USA	ED	59 (13.7)	NR	NR	NR	76% non-rebreathable mask 24% nasal cannula at 5 L/min	PP	NR	NR
Ishak et al	2020	Single Center Cohort study (Non-intubated)	21	Turkey	Ambulance	69.2 (13.1)	NR	NR	NR	2 L/min nasal cannula	PP	NR	<15mins: 7 >15 mins: 14
Kelly et al	2020	Single Center Cohort study (Non-intubated)	17	UK	Ward & critical care	62.3 (11.3)	26.2 (3)	NR	NR	Non-invasive ventilation	Awake PP	4 h twice a day	Total: 10.7 (10.0) hrs/ cycle 2.3 (1.6) days 15.4 ± 2.5 h/ session Total: 6.2 (3.4) days NR
Mittermaier et al	2020	Single Center Cohort study (Intubated)	9	Germany	ICU	62 (14.2)	30.4 (6.5)	26.2 (6.5)	7.4 (4.9)	Mechanical ventilation	PP	NR	NR
Perrier et al	2020	Single Center Cohort study (Intubated)	9	France	NR	54 0.3 (8.7)	32.8 (5.1)	NR	NR	Mechanical ventilation	PP	NR	NR
Solverson et al	2020	Multicenter Cohort study (Non-intubated)	17	Canada	ICU & ward	56 (38)	18% BMI > 35	NR	NR	Non-invasive ventilation	Awake PP	NR	Total: 3 (4.9) days
Winearls et al	2020		24	UK		62 (13)	NR	11 (5)	NR	Non-invasive ventilation including CPAP	PP	NR	Mean duration in 1st 24 h: 8 (5)

(continued on next page)

Table 1 (continued)

Author	Year	Design	N	Country	Setting	Age (mean + -SD)	BMI (mean + -SD)	Severity		Types of Oxygen Delivery Interface	Intervention	Successful duration of proning (standard)	Average/ Total duration of proning
								APACHE II	SOFA				
Thompson et al	2020	Single Center Cohort study (Non-intubated)	25	USA	COVID-19 Respiratory high care unit	Intu: 68.7 (26.7) Non-intu: 61 (20.4)	Intu: 27.5 (8.6) Non-intu: 32.3 (20.4)	NR	NR	1) 6 L/min nasal cannula 2) 15 L/min non-rebreather face mask	Awake PP	NR	hrs Total: 10(5) days Total duration: Intubated: 2.0 (1.7) days Non-intubated: 2.7 (3.3) days
Coppo et al	2020	Single Center Cohort study (Non-intubated)	56	Italy	Medical wards, ED & respiratory HDU	57.4 (7.4)	27.5 (3.7)	NR	NR	79% Helmet CPAP 16% Reservoir mask 5% venturi mask	PP	at least 3 h	Total duration: 1 day
Elharrar et al	2020	Single Center Cohort study (Non-intubated)	24	France	Non-ICU	66.1 (10.2)	23% BMI > 30	NR	2.8 (0.9)	67% <4 L/min 33% ≥4 L/min or HFNC	PP	NR	<1 h (n = 4) to <3 h (n = 5) ≥ 3 h (n = 15)
Taboada et al	2020	Single Center Cohort study (Non-intubated)	29	Spain	Non-ICU	64 (12)	29.2 (3.6)	NR	NR	Nasal cannula or face mask	PP	At least 30 min 3 times a day	NR
Golestani- ragh et al	2020	Cohort study (Non-intubated)	10	NR	NR	NR	NR	NR	NR	Non-invasive ventilation	PP	NR	Mean duration: 9 h
Sartini et al	2020	Single Center Cohort study (Non-intubated)	15	Italy	General wards	59 (6.5)	24 (3.4)	NR	NR	Non-invasive ventilation	PP	NR	Per session: 3.3 (4.1) hrs
Astua et al	2020	Single Center Cohort study (Intubated)	31	USA	NR	58.3 (1.7)	27.9 (3.8)	NR	NR	Mechanical ventilation	PP	16 h each day	NR
Zang et al	2020	Single Center Cohort study (Non-intubated)	60	China	ICU	Int: 62.65 (10.83) Con: 66.14 (9.19)	NR	NR	NR	High Flow Nasal Cannula	PP	10 or 30 min	30 min
Wormser et al	2021	Single Center Cohort study (Non-intubated)	27	France	Wards	70.67 (14.87)	28.8 (5.8)	NR	NR	74% > 6 L/min	PP	61 sessions	NR
Wendt et al	2020	Single Center Cohort study (Non-intubated)	31	USA	ED	62 (12)	31 (5)	NR	NR	2 to 21 L/min by nasal cannula and/or nonrebreather mask	PP	At least 30 min	NR
Prud'homme et al	2021	Single Center Bicentric study (Non-intubated)	96	France	Outside ICU	Int: 62 (11) Con: 61 (18)	Int: 27 (5) Con: 28 (5)	NR	NR	Conventional Oxygen therapy or High Flow Nasal Cannula	Awake PP	At least 3 h a day	3 consecutive days
Taylor et al	2020	Single Center Pilot study (Non-intubated)	40	USA	NR	Int: 55.7 (16.4) Con: 59 (7.5)	Int: 31.3 (10.2) Con: 32.3 (7.8)	NR	NR	Nasal cannula, BiPAP	PP	At least 48 h	Lie prone between 10 and 120 minutes per day
Dubosh et al	2020	Single Center Cohort study (Non-intubated)	22	USA	ED	58.7 (11.9)	NR	NR	NR	Nasal cannula or nonrebreather	Awake PP	At least 30 min	Mean duration: 111 (74.5)
Khullar et al.	2020	Single Center Cohort Study (Intubated)	23	USA	NR	53.5 (13.0)	32.3 (6.0)	NR	4.78	NR	PP	≥16 consecutive hrs of PP for ≥1 d	
Douglas et al	2020		61	USA	ICU	56.7 (13.5)	33.39 (8.9)	NR	14.2 (3.4)	Mechanical ventilation	PP	NR	Mean duration: 5.1 (4.6)

(continued on next page)

Table 1 (continued)

Author	Year	Design	N	Country	Setting	Age (mean + -SD)	BMI (mean + -SD)	Severity		Types of Oxygen Delivery Interface	Intervention	Successful duration of proning (standard)	Average/ Total duration of proning
								APACHE II	SOFA				
Clarke et al	2020	Single Center Cohort study (Intubated)	20	Ireland	ICU	52.8 (11.6)	36.5 (10.7)	NR	8.2 (3.4)	Mechanical ventilation	PP	At least 16 h	NR
Osama et al	2020	Single Center Cohort study (Intubated)	25	France	ICU	61.0 (5.5)	30.0 (3.1)	NR	NR	Mechanical ventilation	PP	At least 16 h	NR
Doussot et al	2020	Single Center Cohort study (Intubated)	67	France	ICU	67.5 (8.3)	30.0 (6.1)	NR	NR	Mechanical ventilation	PP	NR	Mean duration: 17 min (6.9)
Gleissman et al	2020	Single Center Cohort study (Intubated)	44	Sweden	ICU	61.0 (13.0)	50% BMI > 30	NR	NR	Mechanical ventilation	PP	1 h each session	Median for 5 sessions: 14 (12–17)
Sang et al	2020	Single Center Cohort Study (Intubated)	20	China	ICU	69.5 (9.5)	NR	24.1 (3.4)	NR	Mechanical ventilation	PP	18–20 h	NR
Sharp et al	2020	Retrospective Observational Study (Intubated)	12	UK	ICU	56.5 (14.1)	NR	NR	NR	Mechanical ventilation	PP	NR	Successful 2 or more full proning cycles

USA: United States of America; UK: United Kingdom; ED: Emergency Department; ICU: Intensive Care Unit; HDU: High Dependency Unit; Intu: Intubated; Non-intu: Non-intubated; SD: Standard Deviation; NR: Not reported; Int; Intervention; Con: Control; BMI: Body Mass Index; APACHE II: Acute Physiology and Chronic Health Evaluation II; SOFA: Sequential Organ Failure Assessment; CPAP: Continuous positive airway pressure; BiPAP: Bilevel Positive Airway Pressure; PP: Prone Positioning; min: minute; hr: hour.

Table 2
Summary of findings for primary and secondary outcomes.

No	Outcomes	Trials	N	I ² (%)	MD/OR (95% CI)	p-value
1.	PaO ₂ /FiO ₂	13	1002	87	52.15 (37.08, 67.22)	<0.00001
	- Intubated group	8	579	78	46.74 (33.34, 60.15)	<0.00001
	- Non-intubated group	5	423	91	68.81 (15.94, 121.69)	0.01
Heterogeneity: Tau ² = 541.13; Chi ² = 95.88, df = 12 (P < 0.00001); I ² = 87% Test for overall effect: Z = 6.78 (P < 0.00001) Test for subgroup differences: Chi ² = 0.63, df = 1 (P = 0.43), I ² = 0%						
2.	PaCO ₂ (mmHg)	10	793	92	0.57 (-2.74, 3.88)	0.74
	- Intubated group	6	436	93	1.07 (-4.12, 6.26)	0.69
	- Non-intubated group	4	357	0	-0.08 (-1.41, 1.26)	0.91
Heterogeneity: Tau ² = 24.70; Chi ² = 107.86, df = 9 (P < 0.00001); I ² = 92% Test for overall effect: Z = 0.34 (P = 0.74) Test for subgroup differences: Chi ² = 0.17, df = 1 (P = 0.68), I ² = 0%						
3.	SpO ₂ (%)	11	998	94	4.17 (2.53, 5.81)	<0.00001
	- Intubated group	3	432	0	1.67 (1.08, 2.26)	<0.00001
	- Non-intubated group	8	566	95	5.51 (3.17, 7.85)	<0.00001
Heterogeneity: Tau ² = 6.57; Chi ² = 158.60, df = 10 (P < 0.00001); I ² = 94% Test for overall effect: Z = 4.98 (P < 0.00001) Test for subgroup differences: Chi ² = 9.72, df = 1 (P = 0.002), I ² = 89.7%						
4.	Incidence of intubation	5	626	25	1.20 (0.77, 1.86)	0.42
	- Intubated group	-	-	-	-	-
	- Non-intubated group	5	626	25	1.20 (0.77, 1.86)	0.42
Heterogeneity: Tau ² = 0.06; Chi ² = 5.37, df = 4 (P = 0.25); I ² = 25% Test for overall effect: Z = 0.81 (P = 0.42) Test for subgroup differences: Not applicable						

Table 2 (continued)

No	Outcomes	Trials	N	I ² (%)	MD/OR (95% CI)	p-value
5.	Mortality rate	5	688	31	0.44 (0.24, 0.80)	0.007
	- Intubated group	1	261	-	0.66 (0.32, 1.33)	0.24
	- Non-intubated group	4	427	28	0.35 (0.16, 0.75)	0.007
Heterogeneity: Tau ² = 0.14; Chi ² = 5.78, df = 4 (P = 0.22); I ² = 31% Test for overall effect: Z = 2.69 (P = 0.007) Test for subgroup differences: Chi ² = 1.43, df = 1 (P = 0.23), I ² = 30.2%						
6.	Number of patients discharged alive	3	532	63	1.72 (0.85, 3.48)	0.13
	- Intubated group	1	261	-	1.49 (0.72, 3.08)	0.28
	- Non-intubated group	2	271	82	1.90 (0.55, 6.63)	0.31
Heterogeneity: Tau ² = 0.25; Chi ² = 5.47, df = 2 (P = 0.07); I ² = 63% Test for overall effect: Z = 1.50 (P = 0.13) Test for subgroup differences: Chi ² = 0.11, df = 1 (P = 0.74), I ² = 0%						

PaO₂/ FiO₂: Ratio of partial pressure of arterial oxygen/fractional concentration of inspired oxygen; PaCO₂: Partial Pressure of Carbon Dioxide; SpO₂: Peripheral capillary oxygen saturation; I²: heterogeneity; MD: Mean difference; OR: Odds ratio; REM: Random effect model; FEM: Fixed effect model.

supine position. In the pooled data, our review showed that the prone group had a higher SpO₂ reading than the supine group (MD 4.17, 95% CI 2.53 to 5.81; $p \leq 0.0001$, I² = 94%). In term of PaCO₂, the combined data of ten studies [14,21,25,32,33,37,42,43,46,47], (n = 793 patients) found no significant difference in both the prone and supine group (MD 0.57, 95% CI -2.74 to 3.88; $p = 0.74$, I² = 92%). Of note, both outcomes were noted to have substantial degree of heterogeneity.

Five studies [15–19] (n = 688 patients) assessed the mortality rate in COVID-19 patients who received prone and supine position. The pooled data demonstrated that the prone group had a lower mortality rate than the supine group (OR 0.44, 95% CI 0.24 to 0.80; $p = 0.007$, I² = 31%), which was statistically significant. In term of incidence of intubation rate (studies = 5 [14–16,18,19], n = 626 patients, OR 1.20, 95% CI 0.77 to 1.86; $p = 0.42$) and number of patients discharged alive (studies = 3 [15–17], n = 532, OR 1.72, 95% CI 0.85 to 3.48; $p = 0.13$), no significant differences were noted between the prone and supine groups.

In the subgroup analysis of secondary outcomes, the effect size and direction of findings remained unchanged in both intubated and non-intubated groups for the outcomes of SpO₂, PaCO₂, incidence of intubation and number of patients discharged alive, except the incidence of mortality. The incidence of mortality became non-significant in the subgroup analysis of intubated group (studies = 1, n = 215, OR 0.66, 95% CI 0.32 to 1.33; $p = 0.24$, Fig. 3). Funnel plots were suggestive of publication bias for all the secondary outcomes, except for the number of patients discharged alive.

Table 3
Certainty of evidence assessment.

Certainty assessment							N ^o of patients		Effect		Certainty
N ^o of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Prone	supine	Relative (95% CI)	Absolute (95% CI)	
PaO ₂ /FiO ₂											
13	observational studies	not serious	very serious ^a	not serious	not serious	publication bias strongly suspected ^b	496	506	–	MD 52.15 higher (37.08 higher to 67.22 higher)	⊕xxx VERY LOW
PaCO ₂ (mmHg)											
10	observational studies	not serious	very serious ^a	not serious	not serious	publication bias strongly suspected ^b	355	438	–	MD 0.57 higher (2.74 lower to 3.88 higher)	⊕xxx VERY LOW
SpO ₂ (%)											
11	observational studies	not serious	very serious ^a	not serious	not serious	publication bias strongly suspected ^b	489	509	–	MD 4.17 higher (2.53 higher to 5.81 higher)	⊕xxx VERY LOW
Incidence of intubation											
5	observational studies	not serious	not serious	serious ^c	serious ^d	publication bias strongly suspected ^b	80/223 (35.9%)	143/403 (35.5%)	OR 1.20 (0.77 to 1.86)	43 more per 1000 (from 57 fewer to 151 more)	⊕xxx VERY LOW
Incidence of death											
5	observational studies	not serious	not serious	not serious	not serious	publication bias strongly suspected ^b	68/230 (29.6%)	239/458 (52.2%)	OR 0.44 (0.24 to 0.80)	197 fewer per 1000 (from 314 fewer to 56 fewer)	⊕xxx VERY LOW
Number of patients discharged alive											
3	observational studies	not serious	serious ^a	not serious	serious ^d	none	65/159 (40.9%)	100/373 (26.8%)	OR 1.72 (0.85 to 3.48)	118 more per 1000 (from 31 fewer to 292 more)	⊕xxx VERY LOW

CI: Confidence interval; MD: Mean difference; OR: Odds ratio.

Explanations.

^a Substantial heterogeneity I² > 60%.

^b Funnel plot suggested of publication bias.

^c Different threshold of intubation criteria varied across all included studies.

^d Total number of events is less than 300.

4. Discussion

Our meta-analysis demonstrated that COVID-19 patients with prone position were associated with higher PaO₂/FiO₂ ratio and SpO₂ than those with supine position. However, no significant differences were observed in the PaCO₂ level, incidence of intubation and number of patients discharged alive. The certainty of evidence for our measured outcomes was very low due to observational studies in nature, inconsistency, indirectness and suggestive of publication bias. It is believed that prone position expands the collapsed dorsal lung region, resulting in a better ventilation/perfusion ratio and more homogenous distribution of lung ventilation [49–51]. The changes of regional ventilation in the prone and supine position can be observed in both normal and ARDS lung, indicating an even distribution of distending forces throughout the lung tissues [52]. The distribution of pulmonary blood flow in the normal or diseased lung is mainly directed dorsally whether one is in supine or prone position. With this relatively constant regional perfusion in the prone position along with a significant improvement in lung homogeneity, the effect of shunt fraction is expected to reduce and lead to a marked improvement in oxygenation. This has been demonstrated in animal and human studies that the relative shunt fraction of prone

position was reduced by 30% than the supine group with injured lungs [53,54]. However, Gattinoni and colleagues reported that the improvement in oxygenation during prone position did not persist after returning to supine position and the PaO₂/FiO₂ ratio returned to baseline at 6 h following re-supination [55]. This may be explained by the re-collapsed of the previously opened dorsal lung units during prone position, resulting in ventilation/perfusion mismatch and rebound hypoxemia.

Our finding did not show an improvement in PaCO₂ after the application of prone position in COVID-19 patients. This finding has to be interpreted with caveat as our review included both spontaneously breathing and mechanically ventilated patients. The outcome of PaCO₂ may not be accurate in awake spontaneously breathing patients due to the variation in respiratory rate, tidal volume and minute ventilation. However, in the mechanically ventilated patients, controlled ventilation in prone position decreased the shunt fraction and promoted elimination of carbon dioxide [56]. Prone position is believed to reduce areas of over-distension in lung, resulting in a reduction in physiological dead-space and possibly PaCO₂. Gattinoni et al. stated that the effect of PaCO₂ in the prone and supine position is also affected by the alveolar ventilation and its relationship with the total ventilated lung volume [56].




Fig. 2. PaO₂/FiO₂ ratio.




Fig. 3. Incidence of mortality.

In this review, we demonstrated no significant differences in the rate of intubation and number of patients discharged alive between the prone and supine groups. Ferrando et al. emphasized that it may be due to non-standardized or non-protocolized intubation criteria, which may limit the generalizability of the results [14]. Among all the three included studies, there was only one paper reporting their intubation criteria [16]. The duration of prone position varied at the discretion of treating physicians across all the included studies, which introduced significant bias to our findings. Pavlov and colleagues reported that COVID-19

patients with longer duration of awake prone position were associated with lower incidence of intubation [57]. The absence of standardized intubation criteria can limit the interpretation of our findings [58].

Our meta-analysis reported that the mortality rate was lower in those who received prone ventilation. The level of evidence for mortality rate was low due to limited number of studies with small sample size. Among the five included studies, four studies included awake spontaneously breathing patients and only one study examined mechanically ventilated patients. The benefit of prone position was not significant in the

subgroup analysis of intubated group. However, a cohort study by Mathews and team showed that prone ventilation reduced mortality rate in mechanically ventilated COVID-19 patients with ARDS [59]. Prone position is a cornerstone treatment for intubated classical ARDS patients. The evidence of prone position on mortality rate in intubated or non-intubated COVID-19 patients remains uncertain as the pathophysiology of classical ARDS and COVID-induced ARDS were different.

In view of the absence of randomized controlled trials in the literature, our review included observational studies only to summarize the evidence of prone and supine position on oxygenation parameters in COVID-19 patients. In this meta-analysis, a set of strict criteria were applied to assess the included observational studies, which utilized complete full prone position as an intervention group with a control group of supine ventilation. Studies with prone position sessions that were not conducted in full prone position, namely lateral position were excluded [60,61]. There are currently two randomized controlled trials (NCT04350723–350 patients; NCT04395144–346 patients) assessing the effect of awake prone position in COVID-19 patients. The safety on prone position remains uncertain, especially in the hospital setting where back-up for invasive ventilation is not readily available. Most of our included studies did not report adverse events of both prone and supine position. In a recently published article evaluating the safety of prolonged prone position in mechanically ventilated COVID-19 patients, there were nearly 72% (38/61) developed ventral pressure wounds and 95% (58/61) had limb weakness with 8% (5/61) suffered from brachial plexus injury [47]. In another case-control study by Ibarra and team, patient's nutritional status and duration of more than 24 h in prone position were found to be the risk factors for the development of pressure sores in mechanically ventilated COVID-19 patients [62]. It is believed that awake prone position may be associated with lesser adverse events as the duration of prone position would be determined by patients as tolerated. Hyman et al. showed that COVID-19 patients who were intubated and mechanically ventilated earlier in the course of hospital admission were associated with an improved survival rate [63]. Hence, one should consider early invasive ventilation in a select group of awake prone position patients, especially when there is an excessive respiratory drive which may lead to further lung damage through patient self-inflicted lung injury [64].

The recommendations on prone position in mechanically ventilated ARDS patients are clear, but it remains uncertain for awake prone position in COVID-19 patients. The updated ‘Surviving Sepsis Campaign Guidelines on the Management of Adults with Coronavirus Disease 2019 (COVID-19) in the ICU’ stated that there is insufficient evidence to issue a recommendation on the use of awake prone position in non-intubated adults with severe COVID-19 [65]. There were several confounding factors, namely indication, timing, duration, frequency of prone position and type of oxygen supplementation techniques were not standardized across all the included studies. However, the World Health Organisation COVID-19 Clinical Management Living Guidance (25 Jan 2021) suggested a conditional recommendation for awake prone position of severely ill Covid-19 patients requiring supplemental oxygen (including high-flow nasal oxygen) [66]. The recently updated ‘British Thoracic Society/ Intensive Care Society Guidance: Respiratory care in patients with Acute Hypoxaemic Respiratory Failure associated with COVID-19’ also suggested prone position or side repositioning for COVID-19 patients in the respiratory support pathway [67]. A recent expert consensus for the management of COVID-19 related acute respiratory failure (16 Mar 2021) concluded that awake self-prone position may be considered to improve oxygenation and it should be used when supplemental oxygenation is required to maintain $SpO_2 > 90\%$ [68]. At present, awake prone position may be used to delay the respiratory deterioration in selected patients who require oxygen supplementation. This will decrease the demand for invasive mechanical ventilation and further offload the pressure placed on intensive care services worldwide, especially in the resource-limited countries. In the meantime, we await the findings of ongoing high-quality clinical trials to address the

uncertainties surrounding this intervention.

Our review had several limitations, which included observational studies in nature, high degree of heterogeneity, high risk of publication bias, low level of evidence, non-standardized regime of prone position and different mode of ventilation across all the included studies. Previous studies have shown that awake prone position may be most effective when started early during the exudative phase, as opposed to the intermediate phase of ARDS [49,69]. An optimum duration of prone ventilation needs to be studied further to ensure no delay in intubation if hypoxemia condition deteriorates. The clinical criteria for endotracheal intubation should be defined to minimize heterogeneity of our measured outcomes. Different adjuvant of respiratory assist devices, namely high flow nasal cannula, continuous positive airway pressure or bi-level positive airway pressure were used across included studies, which may introduce variances to our findings.

5. Conclusion

In this meta-analysis, prone position improved PaO_2/FiO_2 ratio with better SpO_2 than supine position in COVID-19 patients. Given the limited number of studies with small sample size and substantial heterogeneity of measured outcomes, further studies are warranted to standardize the regime of prone position to improve the certainty of evidence.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Contributions

EC: Data curation; Methodology; Project administration; Writing - original draft; Writing - review & editing. SMI: Data curation; Methodology; Project administration; Writing - original draft. KN: Conceptualization; Formal analysis; Investigation; Methodology; Project administration; Software; Supervision; Validation; Visualization; Writing - original draft; Writing - review & editing. WT: Conceptualization; Formal analysis; Investigation; Methodology; Project administration; Software; Supervision; Validation; Visualization; Writing - original draft; Writing - review & editing. MS: Supervision; Validation; Visualization; Writing - original draft; Writing - review & editing. SR: Supervision; Validation; Visualization; Writing - original draft; Writing - review & editing. MA: Validation; Visualization; Writing - original draft; Writing - review & editing.

Source of funding

No funding was received in support of this project.

Declaration of Competing Interest

All authors have declared that they do not have any conflicts of interest in this review.

Acknowledgements

We would like thank Mr. Bryan Allan for his effort in proof-reading our manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jclinane.2021.110406>.

References

- [1] Hussein NR, Saleem ZS, Ibrahim N, Musa DH, Naqid IA. The impact of COVID-19 pandemic on the care of patients with kidney diseases in Duhok City, Kurdistan region of Iraq. *Diabetes Metab Syndr Clin Res Rev* 2020;14:1551–3.
- [2] Ziehr DR, Alladina J, Petri CR, et al. Respiratory pathophysiology of mechanically ventilated patients with COVID-19: a cohort study. *Am J Respir Crit Care Med* 2020;201:1560–4.
- [3] Ding L, Wang L, Ma W, He H. Efficacy and safety of early prone positioning combined with HFNC or NIV in moderate to severe ARDS: a multi-center prospective cohort study. *Crit Care* 2020;24:8.
- [4] Scaravilli V, Grasselli G, Castagna L, et al. Prone positioning improves oxygenation in spontaneously breathing nonintubated patients with hypoxicemic acute respiratory failure: a retrospective study. *J Crit Care* 2015;30:1390–4.
- [5] Bellone A, Basile A. Prone positioning in severe acute hypoxicemic respiratory failure in the emergency ward. *Emerg Care J* 2018;14.
- [6] Ng ZQ, Tay WC, Ho CHB. Awake prone position for non-intubated oxygen dependent COVID 19 pneumonia patients. *Eur Respir J* 2020;56:2001198.
- [7] Musch G, Layfield JDH, Harris RS, et al. Topographical distribution of pulmonary perfusion and ventilation, assessed by PET in supine and prone humans. *J Appl Physiol* 2002;93:1841–51.
- [8] Lamm WJ, Graham MM, Albert RK. Mechanism by which the prone position improves oxygenation in acute lung injury. *Am J Respir Crit Care Med* 1994;150:184–93.
- [9] Guérin C, Reignier J, Richard JC, et al. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med* 2013;368:2159–68.
- [10] Higgins JPT, Thomas J, Chandler J, et al. *Cochrane handbook for systematic reviews of interventions version 6.1*. Cochrane Train 2020.
- [11] Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ Br Med J* 2015;349:g7647.
- [12] Wells GA, Shea B, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. The Ottawa Hospital, n.d., http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp (accessed on 20 March 2021).
- [13] Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014;14:135.
- [14] Ferrando C, Mellado-Artigas R, Gea A, et al. Awake prone positioning does not reduce the risk of intubation in COVID-19 treated with high-flow nasal oxygen therapy: a multicenter, adjusted cohort study. *Crit Care* 2020;24:597.
- [15] Jagan N, Morrow LE, Walters RW, et al. The positioned study: prone positioning in nonventilated coronavirus disease 2019 Patients—a retrospective analysis. *Crit Care Explor* 2020;2(10):e0229.
- [16] Padrão EMH, Valente FS, Besen BAMP, et al. Awake prone positioning in COVID-19 hypoxicemic respiratory failure: exploratory findings in a single-center retrospective cohort study. *Acad Emerg Med* 2020;27:1249–59.
- [17] Shelhamer MC, Wesson PD, Solari IL, et al. Prone positioning in moderate to severe acute respiratory distress syndrome due to COVID-19: a cohort study and analysis of physiology. *J Intensive Care Med* 2021;36:241–52.
- [18] Zang XF, Wang Q, Zhou H, Liu SH, Xue XY. Efficacy of early prone position for COVID-19 patients with severe hypoxia: a single-center prospective cohort study. *Intensive Care Med* 2020. <https://doi.org/10.1007/s00134-020-06182-4>.
- [19] Prud'homme E, Trigui Y, Elharrar X, et al. Effect of prone positioning on the respiratory support of nonintubated patients with coronavirus disease 2019 and acute hypoxicemic respiratory failure: a retrospective matching cohort study. *Chest* 2021. <https://doi.org/10.1016/j.chest.2021.01.048>. S0012-3692(21)00125-2.
- [20] Taylor SP, Bundy H, Smith WM, Skavronck S, Taylor B, Kowalkowski MA. Awake-prone positioning strategy for non-intubated hypoxic patients with COVID-19: a pilot trial with embedded implementation evaluation. *Ann Am Thorac Soc* 2020. <https://doi.org/10.1513/AnnalsATS.202009-1164OC>.
- [21] Weiss TT, Cerda F, Scott JB, et al. Prone positioning for patients intubated for severe acute respiratory distress syndrome (ARDS) secondary to COVID-19: a retrospective observational cohort study. *Br J Anaesth* 2021;126:48–55.
- [22] Berrill M. Evaluation of oxygenation in 129 proning sessions in 34 mechanically ventilated COVID-19 patients. *J Intensive Care Med* 2020;885066620955137.
- [23] Burton-Papp HC, Jackson AIR, Beecham R, et al. Conscious prone positioning during non-invasive ventilation in COVID-19 patients: experience from a single centre. *F1000Research* 2020;9:859.
- [24] Caputo ND, Strayer RJ, Levitan R. Early self-proning in awake, non-intubated patients in the Emergency Department: a single ED's experience during the COVID-19 pandemic. *Acad Emerg Med* 2020;27:375–8.
- [25] Şan İ, Yıldırım Ç, Bekgöz B, Gemcioglu E. Transport of awake hypoxic probable COVID 19 patients in the prone position. *Am J Emerg Med* 2021. <https://doi.org/10.1016/j.ajem.2020.10.036>.
- [26] Kelly NL, Curtis A, Douthwaite S, et al. Effect of awake prone positioning in hypoxaemic adult patients with COVID-19. *J Intensive Care Soc* 2020. <https://doi.org/10.1177/1751143720961244>.
- [27] Mittermaier M, Pickeroth P, Kurth F, et al. Evaluation of PEEP and prone positioning in early COVID-19 ARDS. *E Clin Med* 2020;100579.
- [28] Perier F, Tuffet S, Maraffi T, et al. Effect of positive end-expiratory pressure and proning on ventilation and perfusion in COVID-19 acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2020;202:1713–7.
- [29] Solverson K, Weatherald J, Parhar KKS. Tolerability and safety of awake prone positioning COVID-19 patients with severe hypoxicemic respiratory failure TT - Tolérabilité et sécurité de la position ventrale éveillée chez des patients atteints de la COVID-19 et d'insuffisance respiratoire hypoxém. *Can J Anaesth* 2021;68:64–70.
- [30] Winearls S, Swingwood EL, Hardaker CL, et al. Early conscious prone positioning in patients with COVID-19 receiving continuous positive airway pressure: a retrospective analysis. *BMJ Open Respir Res* 2020;7:e000711.
- [31] Thompson AE, Ranard BL, Wei Y, et al. Prone positioning in awake, nonintubated patients with COVID-19 hypoxicemic respiratory failure. *JAMA Intern Med* 2020. <https://doi.org/10.1001/jamainternmed.2020.3030>.
- [32] Coppo A, Bellani G, Winterton D, et al. Feasibility and physiological effects of prone positioning in non-intubated patients with acute respiratory failure due to COVID-19 (PRON-COVID): a prospective cohort study. *Lancet Respir Med* 2020;8:765–74.
- [33] Elharrar X, Trigui Y, Dols A-M, et al. Use of prone positioning in nonintubated patients with COVID-19 and hypoxicemic acute respiratory failure. *JAMA* 2020;323:2336–8.
- [34] Taboada M, González M, Álvarez A, et al. Effectiveness of prone positioning in nonintubated intensive care unit patients with moderate to severe acute respiratory distress syndrome by coronavirus disease 2019. *Anesth Analg* 2021;132:25–30.
- [35] Golestani-Eraghi M, Golestani-Eraghi M, Mahmoodpoor A. Early application of prone position for management of Covid-19 patients. *J Clin Anesth* 2020;66:109917.
- [36] Sartini C, Tresoldi M, Scarpellini P, et al. Respiratory parameters in patients with COVID-19 after using noninvasive ventilation in the prone position outside the intensive care unit. *JAMA* 2020;323:2338–40.
- [37] Astua A, Michaels E, Michaels A. Proning During Pandemic: The Rapid Institution of a Safe, Transferable, and Effective Prone Positioning Program at Nychhc/elmhurst Hospital, A Situationally Resource Limited Facility, During the Peak of the Covid 19 Surge. 2020. p. 1–16.
- [38] Dubosh NM, Wong ML, Grossestreuer AV, et al. Early, awake proning in emergency department patients with COVID-19. *Am J Emerg Med* 2020. <https://doi.org/10.1016/j.ajem.2020.11.074>. S0735-6757(20)31105-0.
- [39] Wendt C, Mobus K, Weiner D, Eskin B, Allegra JR. Prone positioning of patients with coronavirus disease 2019 who are nonintubated in hypoxic respiratory distress: single-site retrospective health records review. *J Emerg Nurs* 2021;47:279–87. e1.
- [40] Wormser J, Romanet C, Philippart F. Prone position in wards for spontaneous breathing COVID-19 patients: a retrospective study. *Irish J Med Sci* 2021. <https://doi.org/10.1007/s11845-020-02479-x>.
- [41] Sharp T, Al-Faham Z, Brown M, Martin-Lazaro J, Cevallos Morales J. Prone positioning in covid-19: can we tackle rising dead space? *J Intensive Care Soc* 2020. <https://doi.org/10.1177/1751143720975317>.
- [42] Sang L, Zheng X, Zhao Z, et al. Lung recruitment, individualized PEEP, and prone position ventilation for COVID-19-associated severe ARDS: a single center observational study. *Front Med* 2021;7:1098.
- [43] Gleissman H, Forsgren A, Andersson E, et al. Prone positioning in mechanically ventilated patients with severe acute respiratory distress syndrome and coronavirus disease 2019. *Acta Anaesthesiol Scand* 2020. <https://doi.org/10.3389/fmed.2020.603943>.
- [44] Doussot A, Ciceron F, Cerutti E, et al. Prone positioning for severe acute respiratory distress syndrome in COVID-19 patients by a dedicated team: a safe and pragmatic reallocation of medical and surgical work force in response to the outbreak. *Ann Surg* 2020;272:e311–5.
- [45] Abou-Arab O, Haye G, Beyls C, et al. Hypoxemia and prone position in mechanically ventilated Covid-19 patients: a prospective cohort study. *Can J Anesth* 2020. <https://doi.org/10.1007/s12630-020-01844-9>.
- [46] Clarke J, Geoghegan P, McEvoy N, et al. Prone positioning improves oxygenation and lung recruitment in patients with SARS-CoV-2 acute respiratory distress syndrome; a single centre cohort study of 20 consecutive patients. *BMC Res Notes* 2021;14:20. <https://doi.org/10.1186/s13104-020-05426-2>.
- [47] Douglas IS, Rosenthal CA, Swanson DD, et al. Safety and outcomes of prolonged usual care prone position mechanical ventilation to treat acute coronavirus disease 2019 hypoxicemic respiratory failure. *Crit Care Med* 2021;49:490–502.
- [48] Khullar R, Shah S, Singh G, et al. Effects of prone ventilation on oxygenation, inflammation, and lung infiltrates in COVID-19 related acute respiratory distress syndrome: a retrospective cohort study. *J Clin Med* 2020;9:4129.
- [49] Pelosi P, Brazzi L, Gattinoni L. Prone position in acute respiratory distress syndrome. *Eur Respir J* 2002;20:1017–28.
- [50] Nyrén S, Radell P, Lindahl SGE, et al. Lung ventilation and perfusion in prone and supine postures with reference to anesthetized and mechanically ventilated healthy volunteers. *Anesthesiology* 2010;112:682–7.
- [51] Zaranonello F, Andreatta G, Navalesi P. Prone position and lung ventilation and perfusion matching in acute respiratory failure due to COVID-19. *Images Pulmonary, Crit Care, Sleep Med Sci* 2020;202:278–9.
- [52] Tobin MJ. Mechanical ventilation. *N Engl J Med* 1994;330:1056–61.
- [53] Richter T, Bellani G, Scott Harris R, et al. Effect of prone position on regional shunt, aeration, and perfusion in experimental acute lung injury. *Am J Respir Crit Care Med* 2005;172:480–7.
- [54] Pelosi P, Tubiolo D, Mascheroni D, et al. Effects of the prone position on respiratory mechanics and gas exchange during acute lung injury. *Am J Respir Crit Care Med* 1998;157:387–93.
- [55] Gattinoni L, Tognoni G, Pesenti A, et al. Effect of prone positioning on the survival of patients with acute respiratory failure. *N Engl J Med* 2001;345:568–73.
- [56] Gattinoni L, Vagginelli F, Carlesso E, et al. Decrease in PaCO₂ with prone position is predictive of improved outcome in acute respiratory distress syndrome. *Crit Care Med* 2003;31:2727–33.

- [57] Pavlov I, He H, Menicholas B, et al. Awake Prone Positioning in Non-intubated Patients With Acute Hypoxemic Respiratory Failure Due to COVID-19: A Systematic Review and Meta-analysis. *Research Square COVID-19 Preprints*. 2020. <https://doi.org/10.21203/rs.3.rs-125426/v1>.
- [58] Ponnappa Reddy M, Subramaniam A, Lim ZJ, et al. Prone positioning of non-intubated patients with COVID-19 - a systematic review and meta-analysis. *MedRxiv* 2020. <https://doi.org/10.1101/2020.10.12.20211748>.
- [59] Mathews KS, Soh H, Shaefi S, et al. Prone positioning and survival in mechanically ventilated patients with coronavirus disease 2019-related respiratory failure. *Crit Care Med* 2021. <https://doi.org/10.1097/CCM.0000000000004938>.
- [60] Schifano G, de Grauw AJ, Daniele F, Comellini V, Fasano L, Pisani L. Effects of prone and lateral position in non-intubated patients with 2019 novel coronavirus (COVID-19) pneumonia. *Pulmonology* 2021;27:167–71.
- [61] Retucci M, Aliberti S, Ceruti C, et al. Prone and lateral positioning in spontaneously breathing patients with COVID-19 pneumonia undergoing noninvasive helmet CPAP treatment. *Chest* 2020;158:p2431–5.
- [62] Ibarra G, Rivera A, Fernandez-Ibarburu B, Lorca-García C, Garcia-Ruano A. Prone position pressure sores in the COVID-19 pandemic: the Madrid experience. *J Plast Reconstr Aesthet Surg* 2021. <https://doi.org/10.1016/j.bjps.2020.12.057>.
- [63] Hyman JB, Leibner ES, Tandon P, et al. Timing of intubation and in-hospital mortality in patients with coronavirus disease 2019. *Crit Care Explor* 2020;2:e0254.
- [64] Grieco DL, Menga LS, Eleuteri D, Antonelli M. Patient self-inflicted lung injury: implications for acute hypoxemic respiratory failure and ARDS patients on non-invasive support. *Minerva Anestesiol* 2019;85:1014–23.
- [65] Alhazzani W, Evans L, Alshamsi F, et al. Surviving Sepsis Campaign Guidelines on the Management of Adults with Coronavirus Disease 2019 (COVID-19) in the ICU: first update. *Crit Care Med* 2021;49.
- [66] World Health Organisation. Clinical Management Clinical Management : Living Guidance COVID-19. World Health Organisation; 2021. <https://www.who.int/publications/i/item/WHO-2019-nCoV-clinical-2021-1> (accessed on 27th February 2021).
- [67] Messer B, Antoine-Pitterson P, Blundell A, et al. BTS/ICS Guidance: Respiratory Care in Patients with Acute Hypoxaemic Respiratory Failure Associated with COVID-19. 2021. [file:///C:/Users/User/Downloads/BTS ICS%20Guidance%20respiratory%20care%20in%20patients%20with%20acute%20hypoxaemic%20respiratory%20failure%20associated%20with%20COVID-19.pdf](file:///C:/Users/User/Downloads/BTS%20ICS%20Guidance%20respiratory%20care%20in%20patients%20with%20acute%20hypoxaemic%20respiratory%20failure%20associated%20with%20COVID-19.pdf) (Accessed on 27th February 2021).
- [68] Nasa P, Azoulay E, Khanna AK, et al. Expert consensus statements for the management of COVID-19-related acute respiratory failure using a Delphi method. *Crit Care* 2021;25:106.
- [69] Kallet RH. A comprehensive review of prone position in ARDS. *Respir Care* 2015; 60:1660–87.