

Significance of albumin to globulin ratio as a predictor of febrile urinary tract infection after ureteroscopic lithotripsy

Seung Yun Yi¹, Dong Jin Park², Kyungchan Min¹, Jae-Wook Chung³, Yun-Sok Ha³, Bum Soo Kim⁴, Hyun Tae Kim⁴, Tae-Hwan Kim³, Eun Sang Yoo⁴

¹Department of Urology, Kyungpook National University Hospital, Daegu, Korea

²Department of Urology, Dongguk University Gyeongju Hospital, Gyeongju, Korea

³Department of Urology, School of Medicine, Kyungpook National University, Kyungpook National University Chilgok Hospital, Daegu, Korea

⁴Department of Urology, School of Medicine, Kyungpook National University, Kyungpook National University Hospital, Daegu, Korea

Received: January 31, 2021

Revised: March 18, 2021

Accepted: March 19, 2021

Corresponding author:

Eun Sang Yoo, MD

Department of Urology, School of Medicine, Kyungpook National University, Kyungpook National University Hospital, 680 Gukchaebosang-ro, Jung-gu, Daegu 41944, Korea

Tel: +82-53-420-5843

Fax: +82-53-421-9618

E-mail: uroyoo@knu.ac.kr

Background: We aimed to analyze the effectiveness of albumin to globulin ratio (AGR) in predicting postoperative febrile urinary tract infection (fUTI) after ureteroscopic lithotripsy (URS) and retrograde intrarenal surgery (RIRS).

Methods: From January 2013 to May 2018, 332 patients underwent URS and RIRS. The rate of postoperative fUTI and risk factors for postoperative fUTI were analyzed using logistic regression. Patients were divided into postoperative fUTI and non-postoperative fUTI (non-fUTI) groups. AGR with other demographic and perioperative data were compared between the two groups to predict the development of fUTI after URS.

Results: Of the 332 patients, postoperative fUTI occurred in 41 (12.3%). Preoperative pyuria, microscopic hematuria, diabetes mellitus, hypoalbuminemia, and hyperglobulinemia were more prevalent in the fUTI group. Patients in the fUTI group had larger stone size, lower preoperative AGR, longer operation time, and longer preoperative antibiotic coverage period. In a multivariable logistic analysis, preoperative pyuria, AGR, and stone size were independently correlated with postoperative fUTI ($p < 0.001$, $p = 0.008$, and $p = 0.041$, respectively). Receiver operating curve analysis showed that the cutoff value of AGR that could predict a high risk of fUTI after URS was 1.437 (sensitivity, 77.3%; specificity, 76.9%), while the cutoff value of stone size was 8.5 mm (sensitivity, 55.3%; specificity, 44.7%).

Conclusion: This study demonstrated that preoperative pyuria, AGR, and stone size can serve as prognostic factors for predicting fUTI after URS.

Keywords: Albumin to globulin ratio; Biomarkers; Nephrolithiasis; Ureteroscopy; Urinary tract infections

Introduction

Advancements in endourology have introduced the era of retrograde intrarenal surgery (RIRS) and ureteroscopic lithotripsy

(URS), with a higher stone-free rate and lower morbidity [1]. However, postoperative urinary tract infection (UTI) and subsequent sepsis remain a burden for postoperative care, despite proper administration of prophylactic intravenous antibiotics. UTI is one

of the most common postoperative complications after RIRS, and it has been reported that the rate varies widely between studies [2]. Postoperative UTI can be life-threatening because it can lead to sepsis and septic shock. If it advances to septic shock, higher postoperative mortality and longer hospital stays would be expected [3]. Attempts have been made to determine the risk factors to minimize postoperative febrile UTI. Some studies have suggested that the neutrophil-to-lymphocyte ratio (NLR) and platelet-to-lymphocyte ratio (PLR) can be effective biomarkers for predicting postoperative UTI after percutaneous nephrolithotomy [4,5].

Albumin and globulin are the main components of serum proteins, and these proteins play a pivotal role in acute inflammatory reactions and chronic inflammation. In this study, we validated the effectiveness of the albumin to globulin ratio (AGR) in predicting postoperative febrile UTI (fUTI). However, to date, few studies have evaluated AGR as a predictor of postoperative infection after URS and RIRS. AGR has already been studied in different types of cancer, and it has been found to be a possible prognostic marker for cancer [6]. However, so far, there have been limited studies investigating whether AGR can be used as a prognostic marker to predict postoperative fUTI in URS and RIRS.

Materials and methods

Patients who underwent URS and RIRS from January 2013 to May 2018 in a tertiary general hospital in South Korea were included in this study. Patients with bilateral stones and those without preoperative serum albumin and globulin data were excluded from this study. A total of 332 patients were included in the study. Patients were divided into the postoperative fUTI group (fUTI group) and non-postoperative fUTI group (non-fUTI group). Postoperative fUTI was diagnosed if the patient's body temperature was above 38°C with pyuria and other fever foci that should have been excluded within a week after the surgery. Preoperative clinical data including age, sex, and underlying diseases such as hypertension, diabetes mellitus (DM), chronic renal failure (CRF) defined as an estimated glomerular filtration rate of less than 60 mL/min per 1.73 m², body mass index (BMI), presence of pyuria, presence of microscopic hematuria, presence of preoperative ureteral stent and percutaneous nephrostomy (PCN), and history of previous URS and RIRS were collected. Stone size prior to surgery was measured using noncontrast computed tomography, and the largest diameter was recorded. Preoperative pyuria was defined as more than five to 10 white blood cells (WBC) per high-power field (HPF), while preoperative microscopic hematuria was defined as more than three red blood cells per HPF on laboratory urinalysis. Blood samples were collected within 1 month prior to surgery, and

AGR was calculated using the equation of 'AG ratio = albumin/total protein without albumin.' WBC counts were collected to demonstrate their effectiveness as an inflammatory indicator. Sterile preoperative urine culture was performed before surgery.

The patients were admitted to the hospital a day before surgery. None of the patients had preoperative fever. Fluoroquinolone was administered from the day of admission to the day of surgery. Under spinal or general anesthesia, we used a semi-rigid ureteroscope and/or flexible ureteroscope for lithotripsy. A hydrophilic guidewire was used to engage the semi-rigid ureteroscope into the ureter. For renal stones, we used Amplatz Super Stiff guidewire (Cook Medical Inc., Bloomington, IN, USA) and Flexor Ureteral Access Sheath (Cook Medical Inc.) prior to the engagement of the flexible ureteroscope. A 200- or 365-micro holmium laser lithotripter was used to fragment the target stone into pieces, with a power of 20 joules and 0.5 frequency. We used a stone basket to remove the fragments larger than 2 mm., and six French ureteral stents were kept for 2 weeks after the surgery. Patients were discharged on postoperative day one, and antibiotics were prescribed for 5 days if there was no evidence of postoperative UTI.

Chi-square test for categorical variables and Student t-test for continuous variables were used to analyze the baseline characteristics. Multivariate logistic regression analysis was performed using the forward-likelihood ratio method, and *p*-values, odds ratios (ORs), and 95% confidence intervals (CIs) were collected. A receiver operating characteristic (ROC) curve was used to determine the optimal cutoff value and its sensitivity and specificity. Statistical significance was set at *p* < 0.05. IBM SPSS version 18.0 for Windows (IBM Corp., Armonk, NY, USA) was used for the analysis.

Results

Table 1 shows the baseline characteristics of the patients and the chi-square test between the two groups. Of the 332 patients, 41 were included in the fUTI group and 291 were included in the non-fUTI group. Of all the patients, 135 had hypertension, 70 had DM, and nine had CRF. The mean age was 56.9 ± 13.3 years, mean stone size was 10.1 ± 4.2 mm, and mean preoperative AGR was 1.64 ± 0.43. Preoperative pyuria, microscopic hematuria, hypoalbuminemia, hyperglobulinemia, and DM were more prevalent in the fUTI group (*p* = 0.001, *p* = 0.004, *p* = 0.001, *p* = 0.041, and *p* = 0.037, respectively). Patients in the fUTI group had larger stone size, lower preoperative AGR, longer operation time, and longer preoperative antibiotic coverage period (*p* = 0.004, *p* = 0.003, *p* = 0.028, and *p* = 0.003, respectively). There were no statistically significant differences in hypertension, CRF, age, BMI, operation histo-

Table 1. Comparison of demographic, clinical, and postoperative data between postoperative fUTI group and non-fUTI group

Variable	Total	Postoperative fUTI		p-value
		Yes	No	
Patient	332 (100)	41 (12.3)	291 (87.7)	
Age (yr)	56.9 ± 13.3	57.2 ± 12.0	56.9 ± 13.3	0.869
Sex				0.128
Male	227 (68.4)	32 (78.0)	195 (67.0)	
Female	105 (31.6)	9 (22.0)	96 (33.0)	
Hypertension	135 (40.7)	17 (41.5)	118 (40.5)	0.979
Diabetes mellitus	70 (21.1)	14 (34.1)	56 (19.2)	0.037
Chronic renal failure	9 (2.7)	1 (2.4)	8 (2.7)	0.888
History of stone operation	59 (17.8)	9 (22.0)	50 (17.2)	0.507
Body mass index (kg/m ²)	25.3 ± 4.0	25.6 ± 5.1	25.3 ± 3.9	0.688
Procedure				0.445
URS	131 (39.5)	18 (43.9)	113 (38.8)	
RIRS	180 (54.2)	22 (53.7)	158 (54.3)	
URS+RIRS	21 (6.3)	1 (2.4)	20 (6.9)	
Preoperative pyuria	54 (16.3)	9 (22.0)	45 (15.5)	< 0.001
Preoperative ureteral stent	6 (1.8)	2 (4.9)	4 (1.4)	0.124
Preoperative PCN	51 (15.4)	8 (19.5)	43 (14.8)	0.478
Stone size (mm)	10.1 ± 4.2	11.7 ± 3.9	9.9 ± 4.1	0.004
Operation time (min)	75.1 ± 31.6	83.3 ± 32.2	74.0 ± 31.3	0.028
Preoperative AGR	1.64 ± 0.43	1.46 ± 0.35	1.66 ± 0.44	0.003
Preoperative WBC count (10 ³ /μL)	6,850 ± 2,240	6,874 ± 2,353	6,847 ± 2,228	0.493
Preoperative serum albumin (g/dL)	4.5 ± 0.4	4.2 ± 0.6	4.5 ± 0.4	0.001
Preoperative serum globulin (g/dL)	2.9 ± 0.5	3.0 ± 0.6	2.8 ± 0.4	0.041
Preoperative microscopic hematuria	142 (42.8)	26 (63.4)	116 (39.9)	0.004
Preoperative hydronephrosis	111 (33.4)	19 (46.3)	92 (31.6)	0.061
Stone site				0.144
Renal	198 (59.6)	23 (56.1)	175 (60.1)	
Upper	70 (21.1)	10 (24.4)	60 (20.3)	
Mid	16 (4.8)	3 (7.3)	13 (4.5)	
Low	48 (14.5)	5 (12.2)	43 (14.8)	
Preoperative antibiotics cover (day)	1.5 ± 2.3	2.5 ± 4.3	1.4 ± 1.8	0.003

Values are presented as number (%) or mean ± standard deviation.

AGR, albumin to globulin ratio; fUTI, febrile urinary tract infection; low, low ureter stone; mid, mid ureter stone; PCN, percutaneous nephrostomy; renal, renal stone; RIRS, retrograde intrarenal surgery; upper, upper ureter stone; URS, ureteroscopy lithotripsy; WBC, white blood cell.

ry, sex, preoperative ureteral stenting, preoperative PCN, preoperative WBC counts, and operation sites.

Table 2 shows the results of the multivariate logistic regression analysis for predicting postoperative fUTI. In multivariate logistic regression analysis for the factors that were statistically significant on univariate logistic regression analysis, stone size (OR, 1.1; 95% CI, 1.003–1.186; $p = 0.041$), preoperative AGR (OR, 0.4; 95% CI, 0.170–0.771; $p = 0.008$), and preoperative pyuria (OR, 5.7; 95% CI, 2.727–12.342; $p < 0.001$) were found to be significant factors.

According to ROC curves for predicting postoperative fUTI, the optimal cutoff value of AGR was 1.437 (area under the curve [AUC], 0.646; sensitivity, 77.3%; specificity, 76.9%) (Table 3), and the optimal

cutoff value of stone size was 8.5 mm (AUC, 0.640; sensitivity, 55.3%; specificity, 44.7%) (Table 3).

Discussion

Advancements in laser technology and ureteroscopy have made lithotripsy less invasive and have yielded a higher stone-free rate with fewer complications. One of the most common postoperative complications is UTI, even when prophylactic antibiotics are adequately administered. Fan et al. [7] found that preoperative pyuria was the most important parameter for UTI, and similarly, our study showed that the presence of pyuria prior to surgery signifi-

Table 2. Multivariate logistic regression analysis for predicting postoperative febrile urinary tract infection after ureteroscopic lithotripsy and/or retrograde intrarenal surgery

Variable	Univariate analysis, <i>p</i> -value	Multivariable analysis	
		OR (95% CI)	<i>p</i> -value
Preoperative pyuria	<0.001	5.8 (2.727–12.342)	<0.001
Stone size	0.004	1.1 (1.003–1.186)	0.041
Preoperative AGR	0.003	0.4 (0.170–0.771)	0.008
Diabetes mellitus	0.001	0.5 (0.215–1.064)	0.071
Operation time	0.041	1.0 (0.993–1.018)	0.389
Preoperative albumin	0.037	0.5 (0.198–1.434)	0.212
Preoperative globulin	0.028	0.7 (0.245–2.276)	0.608
Preoperative microscopic hematuria	0.004	2.1 (0.951–4.434)	0.064
Preoperative antibiotics cover	0.003	0.9 (0.890–1.134)	0.941

AGR, albumin to globulin ratio; CI, confidence interval; OR, odds ratio.

Table 3. Receiver operating characteristics curve analysis

Variable	AUC	SD	<i>p</i> -value	95% CI	Cutoff value	Sensitivity (%)	Specificity (%)
AGR	0.646	0.048	0.002	0.552–0.741	1.437	77.3	76.9
Stone size	0.640	0.039	0.004	0.563–0.717	8.5	55.3	44.7

AGR, albumin to globulin ratio; AUC, area under the curve; CI, confidence interval; SD, standard deviation.

cantly contributed to the rate of UTI. Lai and Assimos [8] found that preoperative hydronephrosis was significantly associated with postoperative infection; however, our study showed that hydronephrosis was not a contributing factor to fUTI. In our study, some patients with preoperative hydronephrosis underwent PCN prior to the operation due to ongoing infection or acute renal failure, which may lower the risk of fUTI in patients with preoperative hydronephrosis.

In our study, the stone size was found to be a risk factor for postoperative fUTI because larger stones may require a longer operation time. Moreover, larger stones might require excessive intrarenal and intraureteral irrigation, which can lead to increased renal pelvic pressure, and the whole sequence would contribute to a higher risk of UTI, which is more likely to increase the absorption of bacteriotoxins and true pathogens [9].

Our results also showed that preoperative AGR is a significant factor that is highly associated with the incidence of postoperative fUTI. Patients with a lower AGR (< 1.437) were more susceptible to UTI after URS and RIRS. Previous studies have shown the effectiveness of AGR as a postoperative prognostic factor in patients with different cancers [10–12]. Moreover, Jian et al. [13] and Xun et al. [14] showed the validity of low AGR to predict postoperative fUTI in patients who underwent stone removal operation. Albumin and globulins are major serum proteins that reflect nutritional status and systemic inflammation. Low albumin levels usually reflect malnutrition, which makes one more vulnerable to systemic

infections, including UTI. In addition, hypoalbuminemia has been found to lower target organ exposure to antibiotics, which decreases the effectiveness of antibiotics [15]. Non-albumin serum proteins are involved in various inflammatory reactions and consist of proteins such as globulin, C-reactive protein (CRP), and complements [16,17]. High globulin components are highly associated with worse prognosis in chronic inflammatory conditions, especially in patients with cancer [18]. High globulin levels reflect the inflammatory and active immune response of a host due to highly accumulated immunoglobulins and CRPs [19]. Hypoalbuminemia and hyperglobulinemia were more prevalent in the fUTI group, but each variable was not found to be a useful predictor of fUTI. Albumin can be influenced by various confounding factors such as stress, illness, hepatic insufficiency, and changes in the volume of body fluids, and this could affect the efficiency and accuracy of albumin as a prognostic marker [20]. Previous studies have proposed that AGR provides an assessment of albumin and globulin together, which could reflect the body's nutritional status and inflammatory states comprehensively, and this increases its value as a novel biochemical index [20,21]. Thus, patients with a lower AGR could be more susceptible to acute inflammatory conditions (especially fUTI) after surgery.

Our study had some limitations. First, our study had the potential for selection bias due to the retrospective nature of the study design. Second, more inflammatory markers such as NLR, PLR, CRP, and erythrocyte sedimentation rate were not estimated in

our study due to the retrospective nature of our study design. Finally, because our study population included patients from only one institution, ethnic and geographic variations in stone disease were not reflected in the study population.

Predicting postoperative fUTI after URS and RIRS is important because this complication can be life-threatening and can increase the cost of healthcare, mortality of patients, and hospital stay. Evaluation of preoperative AGR is cost-effective and reliable for predicting postoperative fUTI. According to our study, patients with preoperative pyuria, AGR below 1.437, or stone size greater than 8.5 mm should be monitored closely after surgery. Monitoring vital signs and giving careful attention to clinical symptoms would be necessary to manage postoperative fUTI. Further large-scale prospective studies are required to validate the efficacy of AGR in predicting postoperative fUTI.

Notes

Ethical statements

This was a retrospective study using personal medical records, and informed consent was waived. The study was approved by the Institutional Review Board (IRB) of Kyungpook National University Hospital (IRB No: 2020-10-009-001).

Conflicts of interest

No potential conflict of interest relevant to this article was reported.

Funding

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) & funded by the Korean government (MSIT) (2019R1F1A1044473) (2019R1H1A1079839) (2020R111A3071568).

Author contributions

Conceptualization: SY, YSH, ESY; Data curation: DJP, KM; Investigation: SY, THK; Formal analysis: DJP, KM; Funding acquisition: YSH, BSK; Validation: ESY; Writing-original draft: SY; Writing-review & editing: JWC, HTK, ESY.

ORCID

Seung Yun Yi, <https://orcid.org/0000-0003-4736-8198>
 Dong Jin Park, <https://orcid.org/0000-0003-1667-2146>
 Kyungchan Min, <https://orcid.org/0000-0001-5238-6265>
 Jae-Wook Chung, <https://orcid.org/0000-0002-1055-2357>
 Yun-Sok Ha, <https://orcid.org/0000-0003-3732-9814>
 Bum Soo Kim, <https://orcid.org/0000-0002-4873-3049>

Hyun Tae Kim, <https://orcid.org/0000-0002-4730-3776>

Tae-Hwan Kim, <https://orcid.org/0000-0003-4924-4826>

Eun Sang Yoo, <https://orcid.org/0000-0002-7442-6886>

References

1. Baseskioglu B. The prevalence of urinary tract infection following flexible ureteroscopy and the associated risk factors. *Urol J* 2019;16:439–42.
2. Senocak C, Ozcan C, Sahin T, Yilmaz G, Ozyuvali E, Sarikaya S, et al. Risk factors of infectious complications after flexible uretero-renaloscopy with laser lithotripsy. *Urol J* 2018;15:158–63.
3. Knopf HJ, Graff HJ, Schulze H. Perioperative antibiotic prophylaxis in ureteroscopic stone removal. *Eur Urol* 2003;44:115–8.
4. Cetinkaya M, Buldu I, Kurt O, Inan R. Platelet-to-lymphocyte ratio: a new factor for predicting systemic inflammatory response syndrome after percutaneous nephrolithotomy. *Urol J* 2017;14:4089–93.
5. Sen V, Bozkurt IH, Aydogdu O, Yonguc T, Yarimoglu S, Sen P, et al. Significance of preoperative neutrophil-lymphocyte count ratio on predicting postoperative sepsis after percutaneous nephrolithotomy. *Kaohsiung J Med Sci* 2016;32:507–13.
6. He J, Pan H, Liang W, Xiao D, Chen X, Guo M, et al. Prognostic effect of albumin-to-globulin ratio in patients with solid tumors: a systematic review and meta-analysis. *J Cancer* 2017;8:4002–10.
7. Fan S, Gong B, Hao Z, Zhang L, Zhou J, Zhang Y, et al. Risk factors of infectious complications following flexible ureteroscopy with a holmium laser: a retrospective study. *Int J Clin Exp Med* 2015;8:11252–9.
8. Lai WS, Assimos D. Factors associated with postoperative infection after percutaneous nephrolithotomy. *Rev Urol* 2018;20:7–11.
9. Li T, Sun XZ, Lai DH, Li X, He YZ. Fever and systemic inflammatory response syndrome after retrograde intrarenal surgery: Risk factors and predictive model. *Kaohsiung J Med Sci* 2018; 34:400–8.
10. Azab BN, Bhatt VR, Vonfrolio S, Bachir R, Rubinshteyn V, Alkaied H, et al. Value of the pretreatment albumin to globulin ratio in predicting long-term mortality in breast cancer patients. *Am J Surg* 2013;206:764–70.
11. Du XJ, Tang LL, Mao YP, Sun Y, Zeng MS, Kang TB, et al. The pretreatment albumin to globulin ratio has predictive value for long-term mortality in nasopharyngeal carcinoma. *PLoS One* 2014;9:e94473.
12. Duran AO, Inanc M, Karaca H, Dogan I, Berk V, Bozkurt O, et al. Albumin-globulin ratio for prediction of long-term mortality

- in lung adenocarcinoma patients. *Asian Pac J Cancer Prev* 2014;15:6449–53.
13. Jian ZY, Ma YC, Liu R, Li H, Wang K. Preoperative positive urine nitrite and albumin-globulin ratio are independent risk factors for predicting postoperative fever after retrograde Intra-renal surgery based on a retrospective cohort. *BMC Urol* 2020;20:50.
 14. Xun Y, Yang Y, Yu X, Li C, Lu J, Wang S. A preoperative nomogram for sepsis in percutaneous nephrolithotomy treating solitary, unilateral and proximal ureteral stones. *PeerJ* 2020;8:e9435.
 15. Ulldemolins M, Roberts JA, Rello J, Paterson DL, Lipman J. The effects of hypoalbuminaemia on optimizing antibacterial dosing in critically ill patients. *Clin Pharmacokinet* 2011;50:99–110.
 16. Codina Cazador A, Salvá Lacombe JA, Fernández-Llamazares Rodríguez J, Ruiz Feliu B, Codina Barreras A, Moreno Aguado V. Immunoglobulins and the complement system in colorectal cancer. *Rev Esp Enferm Apar Dig* 1989;75:143–8.
 17. Cohen MH, Makuch R, Johnston-Early A, Ihde DC, Bunn PA Jr, Fossieck BE Jr, et al. Laboratory parameters as an alternative to performance status in prognostic stratification of patients with small cell lung cancer. *Cancer Treat Rep* 1981;65:187–95.
 18. Zhang W, Zhangyuan G, Wang F, Zhang H, Yu D, Wang J, et al. High preoperative serum globulin in hepatocellular carcinoma is a risk factor for poor survival. *J Cancer* 2019;10:3494–500.
 19. Du XJ, Tang LL, Mao YP, Guo R, Sun Y, Lin AH, et al. Circulating EBV DNA, globulin and nodal size predict distant metastasis after intensity-modulated radiotherapy in stage II nasopharyngeal carcinoma. *J Cancer* 2016;7:664–70.
 20. Zhou T, He X, Fang W, Zhan J, Hong S, Qin T, et al. Pretreatment albumin/globulin ratio predicts the prognosis for small-cell lung cancer. *Medicine (Baltimore)* 2016;95:e3097.
 21. Zhang Y, Wang L, Lin S, Wang R. Preoperative albumin-to-globulin ratio as a significant prognostic indicator in urologic cancers: a meta-analysis. *Cancer Manag Res* 2018;10:4695–708.