



Research Paper

Global changes in maternity care provision during the COVID-19 pandemic: A systematic review and meta-analysis

Rosemary Townsend^{a,b}, Barbara Chmielewska^c, Imogen Barratt^{c,e}, Erkan Kalafat^{d,e},
Jan van der Meulen^f, Ipek Gurol-Urganci^f, Pat O'Brien^{f,g,h}, Edward Morris^{g,i}, Tim Draycott^{g,j},
Shakila Thangaratinam^k, Kirsty Le Doare^l, Shamez Ladhani^{l,m,n}, Peter von Dadelszen^o,
Laura A Magee^o, Asma Khalil^{b,c,*}

^a Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Edinburgh, United Kingdom

^b Vascular Biology Research Centre, Molecular and Clinical Sciences Research Institute, St George's University of London, United Kingdom

^c Fetal Medicine Unit, St George's Hospital, St George's University of London, St. George's University Hospitals NHS Foundation Trust, Blackshaw Road, London SW17 0QT, United Kingdom

^d Middle East Technical University, Faculty of Arts and Sciences, Department of Statistics, Ankara, Turkey

^e Koc University, School of Medicine, Department of Obstetrics and Gynaecology, Istanbul, Turkey

^f Department of Health Service Research and Policy, London School of Hygiene and Tropical Medicine, United Kingdom

^g The Royal College of Obstetricians and Gynaecologists, London, United Kingdom

^h University College London Hospitals NHS Foundation Trust, London, United Kingdom

ⁱ Norfolk and Norwich University Hospitals NHS Foundation Trust, Norwich, Norfolk, United Kingdom

^j North Bristol NHS Trust Department of Women's Health, Westbury on Trym, United Kingdom

^k Institute of Metabolism and Systems Research, WHO Collaborating Centre for Women's Health, University of Birmingham, Birmingham, United Kingdom

^l Paediatric Infectious Diseases Research Group and Vaccine Institute, Institute of Infection and Immunity, St George's University of London, London, United Kingdom

^m Immunisation and Countermeasures Division, Public Health England, United Kingdom

ⁿ British Paediatric Surveillance Unit, Royal College of Paediatrics and Child Health, London, United Kingdom

^o Department of Women and Children's Health, School of Life Course Sciences, King's College London, London, United Kingdom

ARTICLE INFO

Article History:

Received 6 April 2021

Revised 17 May 2021

Accepted 18 May 2021

Available online xxx

ABSTRACT

Background: The COVID-19 pandemic has had a profound impact on healthcare systems globally, with a worrying increase in adverse maternal and foetal outcomes. We aimed to assess the changes in maternity healthcare provision and healthcare-seeking by pregnant women during the COVID-19 pandemic.

Methods: We performed a systematic review and meta-analysis of studies of the effects of the pandemic on provision of, access to and attendance at maternity services (CRD42020211753). We searched MEDLINE and Embase in accordance with PRISMA guidelines from January 1st, 2020 to April 17th 2021 for controlled observational studies and research letters reporting primary data comparing maternity healthcare-seeking and healthcare delivery during compared to before the COVID-19 pandemic. Case reports and series, systematic literature reviews, and pre-print studies were excluded. Meta-analysis was performed on comparable outcomes that were reported in two or more studies. Data were combined using random-effects meta-analysis, using risk ratios (RR) or incidence rate ratios (IRR) with 95% confidence intervals (CI).

Findings: Of 4743 citations identified, 56 were included in the systematic review, and 21 in the meta-analysis. We identified a significant decrease in the number of antenatal clinic visits (IRR 0.614, 95% CI 0.486–0.776, $P < 0.00001$, $I^2 = 54.6\%$) and unscheduled care visits (IRR 0.741, 95% CI 0.602–0.911, $P = 0.0046$, $I^2 = 0\%$) per week, and an increase in virtual or remote antenatal care (IRR 1.4656, 95% CI 1.118–1.911, $P < 0.00001$, $I^2 = 90.6\%$) and hospitalisation of unscheduled attendees (IRR 1.214, 95% CI 1.118–1.319, $P < 0.00001$, $I^2 = 0\%$). There was a decrease in the use of GA for category 1 Caesarean sections (CS) (RR 0.529, 95% CI 0.407–0.690, $P < 0.00001$, $I^2 = 0\%$). There was no significant change in intrapartum epidural use ($P = 0.0896$) or the use of GA for elective CS ($P = 0.079$).

* Corresponding author.

E-mail addresses: akhilil@sgul.ac.uk, asma@hscfw.co.uk (A. Khalil).

Interpretation: Reduced maternity healthcare-seeking and healthcare provision during the COVID-19 pandemic has been global, and must be considered as potentially contributing to worsening of pregnancy outcomes observed during the pandemic.

© 2021 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Research in context

Evidence before this study

During the period of the COVID-19 pandemic significant increases in maternal mortality, stillbirth and maternal depression have been identified. At the same time there has been a reduction in preterm birth in high income settings. The mechanisms underlying the observed changes are unclear, but proposed drivers include the widespread behavioural change associated with national 'lockdowns' and other pandemic mitigation measures as well as the restructuring of clinical services that might have led to a reduction of pregnancy care contacts and increased barriers to accessing care.

Added value of this study

This study brings together reports of changes in healthcare usage and provision associated with the COVID-19 pandemic. We show that antenatal care contacts, both planned and emergent, have reduced during the pandemic. The included studies also suggest a reduction in companionship in labour. We also highlight some potentially positive changes – an increase in remote or virtual care provision and reduced postpartum length of stay in hospital. Reassuringly, despite widespread concern at the start of the pandemic response about access to intrapartum analgesia, we have found no evidence of a change in the rate of epidural analgesia associated with the COVID-19 pandemic. A reduction in the use of general anaesthesia for emergency Caesareans was observed.

Implications of all the available evidence

There is a clear change in perinatal outcomes that is contemporaneous with a fall in pregnancy care attendance during the COVID-19 pandemic, especially in low resource settings. While this does not establish a cause, this does support the hypothesis that reduced access to care may have worsened pregnancy outcomes during the pandemic. This finding highlights the need to further investigate the observed changes in perinatal outcomes during the pandemic response while developing robust and equitable maternity care pathways that centre the needs of vulnerable women.

1. Introduction

Over the past year maternity services worldwide have faced an unprecedented challenge from the precipitous global spread of the SARS-CoV-2 virus and the attendant societal and healthcare disruption. Initially, the potential effects of this novel virus on pregnancy outcomes, mothers and newborns were a significant concern. Early reports suggested an increase in iatrogenic preterm birth and caesarean birth in infected mothers [1], and there is evidence of an increased risk of maternal intensive care unit (ICU) admission and maternal mortality due to COVID-19 in some settings [2]. Furthermore, multiple reports have raised concerns about the indirect effects of the pandemic on pregnant women and babies, over and above the direct effects of viral infection. An MBRRACE-UK rapid response

highlighted an increased number of maternal deaths due to mental health illness, including suicide [3]. Other reports have suggested an increase in the population risk of stillbirth [4,5] but a reduction in overall preterm birth of undetermined mechanism [6–9]. Our recent meta-analysis has demonstrated an increase in maternal mortality, stillbirth, ruptured ectopic pregnancy and maternal depression during the pandemic [9], and suggested disproportionate adverse effects in lower resource settings.

In response to the pandemic national governments and healthcare providers implemented sweeping changes. In maternity care, face-to-face consultations were widely curtailed with rapid implementation of home blood pressure and blood glucose monitoring programs and telephone antenatal clinics where possible, mostly in high income countries [10]. In many contexts, partners and visitors were restricted from attending outpatient appointments, ultrasound scans or even providing support during intrapartum care [11,12]. Women's healthcare-seeking behaviour has changed; women have reported being less willing to attend hospital due to fear of contracting COVID-19 [13,14]. In addition to misinterpretations of local and national 'stay at home guidance', these factors may have impacted on the maternity care provided to mothers during pregnancy and the postpartum period [9]. A similar effect was seen during the Ebola epidemic in West Africa [15].

We undertook a systematic review to evaluate reported changes in maternity care provision and uptake during the global COVID-19 pandemic.

2. Methods

2.1. Overview

A prospective protocol for this systematic review and meta-analysis was developed in accordance with PRISMA guidelines [16] and registered with PROSPERO (CRD42020211753). MEDLINE, Embase and the COVID-19 database were searched electronically, without language restrictions, from 1st January 2020 to 17th April 2021, using combinations of the relevant medical subject heading (MeSH) terms, key words and word variants for pregnancy, antenatal and intrapartum care, and COVID-19 (Supplementary Table 1).

2.2. Search strategy, selection criteria, and data extraction

We included observational studies or research letters reporting primary data on the change in maternity service use (e.g. routine antenatal care attendance or unscheduled attendance) by pregnant women and/or maternity healthcare provision (e.g. virtual antenatal care or postpartum hospital length of stay) during the COVID-19 pandemic, compared to periods before. We excluded case reports and series, guidelines and papers describing mitigation strategies and service adaptation that did not include data on resource use. Two authors reviewed all abstracts and full texts independently (any two of IB, BC and RT), with any conflicts resolved by reference to a third reviewer (AK or EK). Data were then extracted from full texts by two reviewers independently using Covidence systematic review software (Veritas Health Innovation, Melbourne, Australia. Available at www.covidence.org). Pandemic mitigation response measures were extracted from The Oxford COVID-19 Government Response Tracker [17]. We recorded the maximum restrictions implemented during the study time frame. Quantitative assessment of severity of mitigation measures was recorded according to the

Government Response Stringency Index (GRSI) developed by The Blavatnik School of Government, University of Oxford [17].

2.3. Quality assessment

Quality assessment was performed by two reviewers independently (any two of IB, BC and RT) using the Newcastle-Ottawa Scale (NOS), with any conflicts resolved by a third reviewer (AK or EK). According to the scale, each study is judged on three broad perspectives: the selection of the study groups, the comparability of the groups and the ascertainment of outcome of interest [18].

2.4. Statistical analysis

Extracted data were combined in a two stage meta-analysis approach. In the first step, incidence rate ratios (IRR) with their 95% confidence intervals (CI) were estimated from individual studies reporting count data such as number of visits per given time period. Likewise, risk ratios (RR) with 95% CI were estimated from individual studies reporting binary outcome data, such as epidural use. In the second stage, a restricted Maximum Likelihood (REML) random-effects meta-analysis was employed to combine RRs and IRRs from individual studies. Statistical heterogeneity was quantified using the I^2 statistic for both analyses; $I^2 < 40\%$ may not be important, $30\text{--}60\%$ may represent moderate heterogeneity, $50\text{--}90\%$ may represent substantial heterogeneity, and $\geq 75\%$ represents considerable heterogeneity [19]. Summary statistics were reported as RR for binary outcomes and IRR for count data. Funnel plots displaying the outcome rate from individual studies were created for the exploration of publication bias. Tests for funnel plot asymmetry were not used when the total number of publications included for each outcome was less than ten. In this case, the power of the tests is too low to distinguish chance from real asymmetry [20–22]. All analyses were conducted using R for Windows software (version 4.0.1) metaphor package.

2.5. Role of the funding source

There was no funding source for this study. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

3. Results

The literature search identified 4743 potentially relevant citations, of which 204 were retrieved for full text review; 56 were included in the qualitative review and 21 in the meta-analysis (Fig. 1, Table 1). Excluded studies and reasons for their exclusion are provided in Supplementary Table 2. The methodological quality of included studies was generally robust, with a median NOS score of 7 out of 9, with only 10 studies scoring less than 7 (Table 2). Importantly, the Comparability domain of the exposed and un-exposed cohorts was good in just over half of cases and the ascertainment of outcomes of interest was of good quality in all studies.

The majority of studies reported findings from a single hospital site or group of facilities relating to specific and highly variable changes in protocols made during the pandemic [13,23–39]. This ranged from the institution of new telehealth services [29,33,35,40–47], altered hospital admission and discharge protocols [27,30,48–50], variance in anaesthetic management [27,36,51,52], and harmonisation of regional antenatal screening services [32,53,54]. Only 14 of the 56 papers reported data from low- or middle-income (LMIC) countries according to World Bank classification [13,28,37,54–64]. Lockdown measures in countries included in the studies varied from a GRSI [17] of 6667 to 1000. Comparison periods were most commonly from a similar timeframe in the preceding

year/s, with fewer studies reporting data on the months immediately prior to the pandemic response (Table 3). Where two or more studies reported comparable outcomes, meta-analysis was undertaken. Publication bias could not be formally assessed for any of the outcomes due to low number of studies for each outcome.

3.1. Maternity service use

3.1.1. Antenatal clinic attendance

Twenty-five studies [13,26,28,29,33,35–37,40,42,44–47,50,56–58,60,65–70] reported on antenatal clinic attendance during the pandemic using a variety of metrics. In several settings, no alteration was made to the standard antenatal care protocol, but decreased antenatal clinic attendance was reported in the majority of studies; in high income settings a decline in face to face contacts was offset by an increase in remote or virtual clinic appointments and the nature but not the number of the appointments varied [26,29,35,40,42–45,47,66,70]. Reports from low resource settings noted a particularly profound reduction in antenatal care contacts. One hospital in Ethiopia noted a fall in antenatal clinic attendance of over 29% [37] (from an average to 86 pregnant women per week to 61) even while delivery rates were maintained, whilst another found that only 293% of (114 out of 389) women giving birth had accessed all recommended antenatal visits [61]. A multicentre study identified reduced antenatal clinic attendance in Bangladesh, Nigeria and South Africa during the pandemic response [56] with similar findings in two additional reports from India [13,58]. Women cited both difficulties in travel and fear of contracting COVID-19 in healthcare settings as their reasons for not attending. A study in Ghana revealed over one third (25 of 71; 362%) missed an antenatal clinic appointment; [60] public transport was seriously restricted during lockdown and virtual appointments were not possible. In contrast, a report from a New York program serving primarily women of low socioeconomic status reported no change in clinic attendance during the pandemic response [67].

Seven studies reported on the number of scheduled antenatal visits in person per week [29,37,42,47,58,66,70]. Quantitative synthesis showed that overall there was a 38.6% drop in care appointments during the pandemic period (pooled IRR: 0.614, 95% CI 0.486 – 0.776; $P < 0.0001$) (Table 3, Fig. 2a) with evidence of moderate heterogeneity amongst the included studies ($I^2 = 54.6\%$).

3.1.2. Antenatal screening

Of particular concern is the impact of the pandemic on routine antenatal screening for infection, anaemia and foetal anomaly, none of which can be offered virtually. In one Italian study, attendance for a variety of routine outpatient encounters was reduced in comparison with the equivalent period in 2019 [25]. The reduction was smallest for antenatal toxoplasmosis screening (740 vs. 1005 visits, 26% reduction), and greatest for non-obstetric outpatient encounters (799 vs. 4253 visits, 81% reduction). One unit in Israel and two from the USA reported reductions in antenatal ultrasound visits during the pandemic period (absolute numbers not given) [33,34,50]. This finding raises concern that women may have been less able to access foetal anomaly screening where desired. In three studies reporting on invasive prenatal genetic testing all noted changes in the timing of presentation and type of procedures performed [32,53,54]. In one study from Turkey, fewer women took up the offer of chorionic villus sampling and amniocentesis - during ($n = 56$, 434%) compared with before ($n = 88$, 638%) the pandemic [54]. Nevertheless, more invasive testing in later pregnancy was offered via cordocentesis ($n = 6$, 11%), raising the possibility that screening tests had been delayed during early pregnancy. One foetal medicine centre in Sardinia, Italy, observed an increase in first trimester screening attendance (70% of population during vs. 50% pre-pandemic) and invasive procedure rates ($n = 150$ during vs $n = 146$ pre-pandemic), attributed to




Fig. 1. PRISMA flow chart.

increased referrals from other centres that were unable to offer COVID-secure testing [53]. Second and third trimester visits were unaffected. A third study from the US reported a significant decrease in CVS although amniocentesis rates were unchanged, attributed in this case to later referral or access to prenatal screening [32]. Meta-analysis showed a non-significant change in prenatal diagnostic procedures performed per year (pooled IRR: 0.880, 95% CI: 0.645–1.199; $P = 0.419$).

3.1.3. *Unscheduled care attendance*

In seven studies, quantitative synthesis showed a decrease in unscheduled care attendance at maternity triage, urgent care or obstetric emergency departments, in Italy, Israel, the USA and the UK (pooled IRR 0.741, 95% CI: 0.602–0.911, $I^2=0%$, $P = 0.0046$) (Table 4, Fig. 2b).^{23,24,31,36,39,68,71} In the two studies that reported the outcome, from Israel and Italy, the associated risk of hospital admission amongst 1042 pregnant women who presented at the emergency department significantly increased, by 214% (pooled RR 1.214, 95% CI:

1.118–1.319, $I^2=0%$, $P<0.0001$) (Table 4, Fig. 2c) [68,71]. Variable results were found in three studies that reported on measures of delay in care-seeking during labour. In one report from Brazil of 81 patients in spontaneous labour, the proportion who delivered within three hours of hospital presentation increased from 268% in 2019 to 40% in the equivalent period in 2020 [55]. While one Californian hospital reported no change in mean admission to delivery time, both spontaneous and induced labours were included in the comparison [27]. In a large Irish study no change in births before arrival (BBA) was noted [38].

3.2. *Maternity healthcare provision*

3.2.1. *Virtual antenatal care protocols*

Many reports described new provision of virtual services that enabled clinical contact to continue while reducing in-person clinic visits. A wide variety of protocols were described encompassing both routine care and specialist clinics providing care for the hypertensive

Table 1
Characteristics and summary of the findings of the included studies.

Author (year)	Country	Study Population	GRSI	Country Development Index	Outcomes Reported	Findings in exposed group	NOS Score
Abdela (2020) [37]	Ethiopia	Single Centre	80•56	LIC	Family planning visits Antenatal clinic attendance Deliveries per week	Decreased Decreased No change	7*
Abel (2021) [23]	USA	Single Centre	72.69	HIC	Obstetric emergency department visits	Decreased	6*
Ahmed (2021) [56]	Bangladesh, Nigeria, South Africa	Multi-centre	Bangladesh: 93.52 Nigeria: 85.65 South Africa: 87.96	LMIC	Antenatal care (ANC) clinic attendance Family planning clinic attendance Total deliveries	Decreased Decreased Decreased Mix Mix	7*
Albert (2020) [72]	Spain	Single Centre	85•19	HIC	Report on virtual review of gestational diabetes patients	NA	5*
Anderson (2020) [77]	USA	Single Centre	72•69	HIC	Obstetric hospitalisations	Decreased	7*
Baptiste (2021) [32]	USA	Single Centre	72.69	HIC	Number of prenatal genetic diagnostic procedures performed Chorionic villus sampling Amniocentesis	Decreased No change	9*
Bernstein (2021) [33]	USA	Single centre	72.69	HIC	Number of in-person visits GA of first prenatal visit GA of dating scan GA of anatomy scan Number of triage visits Total number of ultrasound scans Total number of visits (telehealth and in-person) Total number of no-shows Rates of standard prenatal care metrics (pap smear, genetic screening, GDM screening, GBS screening) Post-partum readmission	Decreased No change No change Later GA age No change No change Increased No change No change No change	7*
Bertozzi-Villa (2021) [34]	USA	Single Centre	72•69	HIC	Intake obstetric ultrasound scans	Decreased	7*
Bhatia (2020) [48]	UK	Multi-Centre	79•63	HIC	General anaesthetic for Caesarean section Regional anaesthesia to general anaesthesia conversion during caesarean section	Decreased Decreased	7*
Biviá-Rovig (2020) [69]	Spain	Regional (Valencia)	79•17	HIC	Cancellation of Antenatal Classes Attendance of Online Classes Cancellation of Appointments due to fear of contagion	52•2%* 24•4%* 22•5%*	4*
Bornstein (2020) [52]	USA	Multi-Centre	72•69	HIC	Postpartum length of stay (hours) CS Postpartum length of stay (hours) VD	Decreased Decreased	7*
Chen (2020) [59]	China	Regional	75•46	UMIC	Online consultations according to severity of pandemic		
Online consultations according to trimester Reason for online consultation	High rate in severely affected areas for obstetric care-seeking Majority in third trimester Majority for routine care	7*					
Dell'Utri (2020) [24]	Italy	Single Centre	75•46	HIC	Overall pregnancy related admissions		

(continued)

Table 1 (Continued)

Author (year)	Country	Study Population	GRSI	Country Development Index	Outcomes Reported	Findings in exposed group	NOS Score
Admissions for delivery Postpartum admissions Duryea (2021) [35]	Decreased Increased Decreased USA	7* Single Centre	72.69	HIC	GA at first prenatal visit Total number of prenatal encounters (in-person and virtual) Attendance to prenatal care visits Number of audio-only virtual prenatal visits attended	Decreased Increased No change Increased	9*
Facco (2021) [36]	USA	Single Centre	72.69	HIC	Number of prenatal visits (PNV) Number of postpartum visits (PPV) Length of hospital stay (hours from delivery to discharge) All deliveries VD Maternal post-partum readmission Maternal post-partum emergency department (ED) visits Infant readmission Infant ED visits	No change No change Decreased Decreased No change Decreased (when stratified for insurance type – only for those with Medicaid insurance) No change Decreased (when stratified for insurance type – only for those with commercial insurance)	8*
Filice (2020) [25]	Italy	Single Centre	93•52	HIC	Uptake of toxoplasmosis screening	No change	7*
Futtermann (2020) [26]	USA	Single Centre	72•69	HIC	Satisfaction with in-person antenatal appointments Satisfaction with virtual antenatal appointments	SAPS ₁ score 24 SAPS ₁ score 20	9*
Gildner (2020) [73]	USA	National Survey	72•69	HIC	Changes in birth plan Changes in labour companionship Changes in location Changes in birth plan in hospital	45•3% reported a change in plan* Qualitative Qualitative	5*
Goyal (2020) [13]	India	Single Centre	100•0	LMIC	Pregnancy related admissions	Decreased Decreased	6*
Greco (2021) [44]	USA	Single Centre	72.69	HIC	Antenatal clinic attendance Total in-person prenatal hypertensive disorders of pregnancy (HDP) visits Total virtual HDP visits Total number of in-person postpartum HDP visits Total number of no postpartum HDP visits Total number of virtual postpartum HDP visits Diagnosis site of HDP HDP diagnosis timing Post-partum readmission rate	Decreased No change Decrease Increase Increase (not statistically significant due to 0 in control group) No change No change Increase	9*
Greene (2020) [27]	USA	Single Centre	72•69	HIC	Admission to delivery time (hours) Postpartum length of stay (nights) Epidural use	No change Decreased No change	8*
Gu (2020) [28]	China	Single Centre	81•02	UMIC	Outpatient visits per week Hospitalisations per week Emergency Department attendance	Decreased Decreased Decreased	9*
	USA	Single Centre	72•69	HIC			7*

(continued)

Table 1 (Continued)

Author (year)	Country	Study Population	GRSI	Country Development Index	Outcomes Reported	Findings in exposed group	NOS Score
Holcomb (2020) [29]					Percentage of appointments conducted virtually Clinic waiting time for in-person visits Clinic attendance for virtual appointments compared to in-person appointments Satisfaction with telemedicine (WHC) Satisfaction with telemedicine (MFM)	Increased Decreased Higher 95% good or very good* 87% good or very good*	
Hui (2020) [30]	Hong Kong	Single Centre	66•67	HIC	Rate of hospital births Proportion of women with labour companionship Epidural use Administration of pethidine injections	Decreased Decreased No change Increased	5*
Hussain (2021) [45]	USA	Single centre	72.69	HIC	GA at diagnosis of GDM Total GDM antenatal visits Visits with self-reported blood glucose data (compared to downloaded) Total antenatal GDM ultrasounds In-person and telemedicine attendance	No change Decreased Increased Decreased Increased	9*
Jeganathan (2020) [70]	USA	Multi-Centre	72•69	HIC	Antenatal clinic attendance Antenatal clinic 'no show' Antenatal clinic cancellation Antenatal clinic cancellation by patient Patient satisfaction with telemedicine Provider satisfaction with telemedicine	Decreased Decreased Increased Decreased 86•9% satisfied* 87•8% satisfied*	8*
Justman (2020) [50]	Israel	Single Centre	94•44	HIC	Hospital admissions Triage attendance High risk clinic visits Ultrasound visits Total number of births Epidural use	Decreased Decreased Decreased Decreased Decreased Increased	9*
KC (2020) [64]	Nepal	Multi-Centre	96•3	LMIC	Births per week Attendance at childbirth services by disadvantaged groups Companionship during labour Hand hygiene practices by clinicians during childbirth Clinician use of gloves and gowns during childbirth Intrapartum foetal heart rate monitoring Preparation of equipment used during childbirth Skin to skin after birth	Decreased Decreased Decreased Increased Decreased Decreased Decreased No change Increased	9*
Khalil (2020) [31]	UK	Single Centre	79•63	HIC	Antenatal bookings per week Obstetric triage attendance per week Births per week	Decreased Decreased Decreased	7*
Krishnamurti (2021) [81]	USA	Single Centre	72.69	HIC	Women completing the onboarding process for a prenatal care app Use of in-app intimate partner violence (IPV) risk assessment Rates of IPV	Decreased Increased Increased (non-significant)	7*
Kugelman (2020) [71]	Israel	Single Centre	94•44	HIC	Hospitalisation from obstetric triage Admission to Delivery Suite	Increased Increased Increased	9*

(continued)

Table 1 (Continued)

Author (year)	Country	Study Population	GRSI	Country Development Index	Outcomes Reported	Findings in exposed group	NOS Score
					from triage Presentation with reduced foetal movements Presentation with premature rupture of membranes Home birth Admission in second stage of labour Obstetric Emergency Department Visits	Increased No change No change Decreased	
Kumari (2020) [63]	India	Multi-Centre	100•0	LMIC	Hospitalisation Referred obstetric emergencies	Decreased Decreased	9*
Limaye (2020) [43]	USA	Single Centre	72•69	HIC	Percentage of telehealth visits	Higher proportion in those with private health insurance	8*
Liu (2020) [62]	China	Multi-Centre	77•31	UMIC	Patient request for online consultation Change of planned mode of birth from vaginal to elective caesarean section due to the pandemic Change of planned mode of birth from caesarean section to vaginal delivery due to the pandemic	75•4% Wuhan, 69•5% Chongqing* 12•7% Wuhan, 6•0% Chongqing* 5•6% Wuhan, 3•1% Chongqing*	6*
Madden (2020) [66]	USA	Multi-centre	72•69	HIC	Proportion of antenatal clinic visits conducted virtually Proportion of booked visits that were 'no shows'	Increased Decreased	7*
McDonnell (2020) [38]	Ireland	Single Centre	90•74	HIC	Unbooked mothers presenting in labour	No change No change	8*
Meyer (2020) [39]	Israel	Single Centre	94•44	HIC	Babies born before arrival Referral indications from Emergency Department Duration of treatment until decision Emergency Department referrals	No change No change Decreased Increased	
Monni (2020) [53]	Italy	Single Centre	93•52	HIC	Admission in active labour First trimester prenatal screening Second trimester prenatal screening Third trimester prenatal screening	Increased No change No change Increased	7*
Moyer (2020) [60]	Ghana	National Survey	52•78	LMIC	Performance of invasive foetal testing Missed antenatal visit Plan to deliver in hospital/health centre	36•2%* Decreased	6*
Ozalp (2020) [54]	Turkey	Single Centre	77•78	UMIC	Rate of women accepting offered invasive testing Number of procedures performed Chorionic villus sampling Amniocentesis Cordocentesis	Decreased Decreased Decreased Increased	9*
Patkar-Kattimani (2021) [49]	UK	Single Centre	79.63	HIC	Epidural use Epidural response time <30 min Emergency general anaesthesia rate General anaesthesia for elective CS	No change No change Reduction Increase	7*
Peahl (2020) [42]	USA	Single Centre	72•69	HIC	Average total antenatal clinic visit volume Proportion of antenatal clinic visits conducted	Decreased Increased 77•5%* 83•1%*	8*

(continued)

Table 1 (Continued)

Author (year)	Country	Study Population	GRSI	Country Development Index	Outcomes Reported	Findings in exposed group	NOS Score
Racine (2021) [51]	USA	Single Centre	72.69	HIC	virtually Patient satisfaction with telemedicine Provider satisfaction with telemedicine Likelihood of attending in spontaneous labour Likelihood of need for induction Maternal length of stay Neonatal length of stay Delivery >41 weeks	Increased Decreased Decreased Decreased Increased	7*
Sarkar (2021) [58]	India	Single Centre	100	LMIC	Total antenatal attendance New patients Old patients Total gynaecology outpatient attendance (including infertility, postpartum and termination of pregnancy)	Decreased Decreased Decreased Decreased	7*
Sakowicz (2021) [74]	USA	Single centre	72.69	HIC	Post-partum visit attendance (virtually and in-person) Likelihood of having post-partum depression screening	Decreased Decreased	7*
Sakowicz (2) (2021) [75]	USA	Single centre	72.69	HIC	Long acting reversible contraceptive use postpartum	Decreased	8*
Salsi (2020) [68]	Italy	Single Centre	91•67	HIC	Self-referrals to the Emergency Department Number of admissions Proportion of admissions	Decreased Decreased Increased	8*
Selinger (2021) [46]	UK	Multi-centre	79.63	HIC	Face to face IBD clinic during pregnancy Telephone IBD clinic during pregnancy	Decreased Increased	8*
Shields (2020) [41]	USA	Single Centre	72•69	HIC	Visits per day after conversion to telehealth 'No shows' after implementation of telehealth Invasive prenatal testing	Decreased Decreased No change	7*
Silverman (2020) [67]	USA	Single Centre	72•69	HIC	Antenatal clinic attendance	No change	6*
Soffer (2021) [47]	USA	Single Centre	72.69	HIC	In-person prenatal care visits Third trimester ultrasound scans Detection of foetal growth restriction (FGR) Telehealth visits GA at diagnosis of FGR	Decreased Decreased Increased Increased No change	7*
Sun (2020) [55]	Brazil	Single Centre	81•02	UMIC	Delivery within 3 h of admission	Increased	6*
Tadesse (2020) [61]	Ethiopia	Single Centre	80•56	LIC	Missed/delayed access to antenatal services Full utilisation of antenatal services Age of patients Education level of patients Urban residency	55•5%* 29•3%* Positively associated with utilisation Positively associated with utilisation Positively associated with utilisation	8*
Wanyana (2021) [57]	Rwanda	Multi-centre	90.74	LIC	ANC first standard visit utilisation rate Deliveries at health facility Mothers in labour referred to higher level for delivery 1ST PNC visit (maternal and infant) utilisation rate 4TH PNC visit (maternal and	Decrease Decrease No change No change Increase Decrease	7*

(continued)

Table 1 (Continued)

Author (year)	Country	Study Population	GRSI	Country Development Index	Outcomes Reported	Findings in exposed group	NOS Score
Weingarten (2021) [40]	USA	Single centre	72.69	HIC	infant) utilisation rate Vaccination uptake Virtual prenatal diabetic visits In person prenatal diabetic visits	Increased Decreased	9*
Zarasvand (2020) [65]	UK	Single Centre	79•63	HIC	Number of face-to-face appointments Number of telephone appointments Use of regional anaesthetic for cerclage placement New referrals to preterm birth services Inappropriate referral percentage		
Total number of clinic appointments	Decreased Increased Increased No change Increased	7*					

*Results from survey, no comparison group.

†SAPS: Short Assessment of Patient Satisfaction.

GRSI: government response stringency index, NOS: Newcastle-Ottawa Scale.

LIC: lower income country. HIC: high income country. LMIC: lower middle income country. UMIC: upper middle income country. CS: caesarean section. VD: vaginal delivery. WHC: women's health clinic. MFM: maternal foetal medicine, NA: not applicable: .

disorders of pregnancy [44], diabetes [40,45,72], women at risk of preterm birth [65] and women with inflammatory bowel disease [46]. For example, in Nanjing, China, strict screening protocols were introduced for face-to-face antenatal care while telemedicine appointments were offered as an alternative for routine visits, and home monitoring of blood glucose and blood pressure was utilised [28]. For 2458 women studied, the number of in-person visits was significantly reduced from approximately 500 to 200 visits per week, without an associated change in maternal and neonatal outcomes or hospital acquired infections for women with diabetes or high blood pressure.

In one obstetric service in New York, telemedicine via either audio or video link was introduced for most high-risk prenatal care, including gestational diabetes education, genetic counselling and maternal-foetal medicine consultations. The telemedicine protocol included self-monitoring of blood pressure via the provision of automated sphygmomanometers but not foetal heart auscultation, and the majority of contacts were conducted via video link. For 91 women studied, 29% of visits were conducted using telemedicine; patient non-attendance was decreased and both patients (869%) and providers (878%) reported satisfaction with the service [70]. Another New York centre reported conducting 318% (1354 of 4248) of prenatal care visits using video communication within the electronic record accessed by the patients on smart phones or other devices [66], with 92% provider satisfaction when appointments were scheduled appropriately. This group identified that Medicaid patients had higher rates of non-attendance than patients with private insurance. A further New York study reported similar findings: patients with public insurance were less likely to have had at least one telehealth visit (609 vs. 873%, $P < 0001$), although it is not clear if this was patient or insurer driven [43]. One unit in Texas offered audio-only virtual appointments because they predicted that most of their patients lacked access to high-speed internet [29]. By the end of the study period of two weeks, around 25% of prenatal visits were conducted remotely. Average waiting times for women attending in person were reduced (21 min) and a greater proportion of prenatal visits were completed virtually than in person (88% vs 82%, $P < 0001$). The benefits of virtual appointments cited by patients included reduced

requirement to use public transport during the pandemic, less time away from work, and less need to arrange childcare assistance. In a relatively more privileged population in Michigan, a prenatal care schedule utilising virtual appointments via either audio or video link was implemented and supported with the distribution of home sphygmomanometers to patients in the third trimester [42]; average weekly clinic visit volume decreased by 332 (316%), and virtual visits increased from 101 to 239 (1366%). Around two-thirds of respondents felt that virtual visits were as safe as in-person visits (648% of patients and 65% of providers), but only 371% of patients and 455% of providers felt that the overall quality of virtual appointments was equivalent to face-to-face visits [42]. Interestingly, there was a discrepancy between patient and provider enthusiasm for continuing virtual visits after the pandemic, with only 403% of patients in favour compared to 922% of providers.

A specialised preterm birth clinic in the UK reported that it reduced face-to-face appointments by 54% from 341 to 157, by increasing their telephone consultations from 0 to 221 (64%) and changing definitions of high- and intermediate-risk referral criteria. By questionnaire, 625% of women indicated they 'did not mind' having remote consultations, and 75% were happy or had no preference for telephone over video consultations [65]. A Spanish clinic caring for women with gestational diabetes reduced their face-to-face visits by 886% by using a smartphone app to monitor blood glucose remotely [72].

Five studies reported the number of virtual or over the phone visits per week during, compared to before, the pandemic [29,42,47,66,70]. There was an almost 46-fold increase in the number of virtual appointments during the pandemic period (pooled IRR 4656, 95% CI 7762–2794, $P < 00001$) (Table 4), which balanced the reduction in the number of in-person appointments. There was significant heterogeneity amongst included studies ($I^2=906%$).

3.2.2. Intrapartum analgesia

Quantitative synthesis of two studies found no change in epidural analgesia use during labour (pooled RR 1044, 95% CI 0993–1098, $P = 00896$) (Table 3) [30,50].

Table 2

Quality Assessment of the included studies using the Newcastle-Ottawa Scale (NOS).

Author	Selection Represent-ativeness of exposed cohort (max score: *)	Comparability Selection of non-exposed (max score: *)	Outcome Ascertain-ment of exposure (max score: *)	Total (max score: 9*) Demonstration that outcome of interest was not present at start of study (max score: *)	Comparability of cohorts on the basis of design or analysis (max score: ‡)	Assessment of outcome (max score: *)	Was follow-up long enough for outcomes to occur (max score: *)	Adequacy of follow up (max score: *)	
Abdela (2020) [37]	*	*	*	*	—	*	*	*	7*
Abel (2021) [23]	*	*	*	—	—	*	*	*	6*
Ahmed (2021) [56]	*	*	*	*	—	*	*	*	7*
Albert (2020) [72]	*	—	*	—	—	*	*	*	5*
Anderson (2020) [77]	*	*	*	*	—	*	*	*	7*
Baptiste (2021) [32]	*	*	*	*	**	*	*	*	9*
Bernstein (2021) [33]	*	*	*	*	—	*	*	*	7*
Bertozzi-Villa (2021) [34]	*	*	*	*	—	*	*	*	7*
Bhatia (2020) [48]	*	*	*	*	—	*	*	*	7*
Biviá-Rovig (2020) [69]	—	*	*	—	—	—	*	*	4*
Bornstein (2020) [52]	*	*	*	*	—	*	*	*	7*
Chen (2020) [59]	*	*	*	*	—	*	*	*	7*
Dell'Utri (2020) [24]	*	*	*	*	—	*	*	*	7*
Duryea (2021) [35]	*	*	*	*	***	*	*	*	9*
Facco (2021) [36]	*	*	*	*	***	*	*	*	8*
Filice (2020) [25]	*	*	*	*	—	*	*	*	7*
Futtermann (2020) [26]	*	*	*	*	***	*	*	*	9*
Gildner (2020) [73]	—	—	*	*	***	—	*	*	5*
Goyal (2020) [13]	—	*	*	*	—	*	*	*	6*
Greco (2021) [44]	*	*	*	*	***	*	*	*	9*
Greene (2020) [27]	*	*	*	*	***	*	*	*	8*
Gu (2020) [28]	*	*	*	*	**	*	*	*	9*
Holcomb (2020) [29]	*	*	*	*	—	*	*	*	7*
Hui (2020) [30]	*	*	*	—	—	—	*	*	5*
Hussain (2021) [45]	*	*	*	*	***	*	*	*	9*
Jeganathan (2020) [70]	*	*	*	*	***	*	*	*	8*
Justman (2020) [50]	*	*	*	*	***	*	*	*	9*
KC (2020) [64]	*	*	*	*	***	*	*	*	9*
Khalil (2020) [31]	*	*	*	*	—	*	*	*	7*
Krishnamurti (2021) [81]	*	*	*	—	*	*	*	*	7*
Kugelman (2020) [71]	*	*	*	*	***	*	*	*	9*
Kumari (2020) [63]	*	*	*	*	***	*	*	*	9*
Limaye (2020) [43]	*	*	*	*	***	*	*	*	8*
Liu (2020) [62]	—	*	*	—	***	—	*	*	6*
Madden (2020) [66]	—	*	*	*	***	*	*	*	7*
McDonnell (2020) [38]	*	*	*	*	*	*	*	*	8*
Meyer (2020) [39]	*	*	*	*	—	*	*	*	7*
Monni (2020) [53]	*	*	*	*	—	*	*	*	7*
Moyer (2020) [60]	—	—	*	*	*	*	*	*	6*
Ozalp (2020) [54]	*	*	*	*	**	*	*	*	9*
Patkar-Kattimani (2021) [49]	*	*	*	*	—	*	*	*	7*
Peahl (2020) [42]	*	*	*	*	***	—	*	*	8*
Racine (2021) [51]	*	*	*	*	—	*	*	*	7*
Sarkar (2021) [58]	*	*	*	*	—	*	*	*	7*
Sakowicz (2021) [74]	*	*	*	—	*	*	*	*	7*
Sakowicz (2) (2021) [75]	*	*	*	—	***	*	*	*	8*
Salsi (2020) [68]	*	*	*	*	***	*	*	*	8*
Selinger (2021) [46]	*	—	*	*	**	*	*	*	8*

(continued on next page)

Table 2 (Continued)

Author	Selection Representativeness of exposed cohort (max score: *)	Comparability Selection of non-exposed (max score: *)	Outcome Ascertainment of exposure (max score: *)	Total (max score: 9*) Demonstration that outcome of interest was not present at start of study (max score: *)	Comparability of cohorts on the basis of design or analysis (max score: #)	Assessment of outcome (max score: *)	Was follow-up long enough for outcomes to occur (max score: *)	Adequacy of follow up (max score: *)
Shields (2020) [41]	*	*	*	*		*	*	7*
Silverman (2020) [67]	*	*	*	-		*	*	6*
Soffer (2021) [47]	*	*	*	*		*	*	7*
Sun (2020) [55]	-	*	*	*		*	*	6*
Tadesse (2020) [61]	*	-	*	*	**	*	*	8*
Wanyana (2021) [57]	*	*	*	*		*	*	7*
Weingarten (2021) [40]	*	*	*	*	**	*	*	9*
Zaravand (2020) [65]	*	*	*	*		*	*	7*

In a study of six UK hospitals with over 17,000 births collectively, the rate of general anaesthesia for caesarean section was reduced from 77% in 2019 to 37% during the equivalent period in 2020, an RR of 0.50 (95% CI 0.39 – 0.63) [48]. A similar proportional reduction in intra-operative conversion from regional to general anaesthesia was observed from 16% (n = 39) to 0.8% (n = 19). This finding was supported by a second UK based study [49], and pooled analysis showed that general anaesthesia use for category I (the most urgent) caesarean sections were significantly reduced during the pandemic period (pooled RR 0.529, 95% CI 0.407–0.690, P<0.0001) while general anaesthesia use for elective caesarean was unchanged (pooled RR 0.831, 95% CI 0.205–3.356, P = 0.79) (Table 4).

3.2.3. Companionship in labour

Three papers reported changes in the proportion of women having personal companionship in labour. In Nepal (20,354 women), the reduction was small (89.4% to 83.4%, P = 0.0014) [64], while in Hong Kong (2138 women) the reduction was large (88.8% to 21.8%, P<0.005) [30]. One paper reported on the number of women anticipating a reduction in support persons present in labour and an associated increase in planned home births [73]. The data could not be pooled as the definition of companionship varied amongst studies – in some contexts family members take an active role in personal care and physical support of the labouring person while in others the role is primarily the provision of social and emotional support.

3.2.4. Hospital length of stay after birth

Four studies reported length of hospital stay after admission for birth using varying metrics; in all cases, length of stay was reduced after both vaginal births and caesarean sections during the pandemic [27,36,51,52]. In California, the proportion of women (n = 1339) discharged fewer than three nights after caesarean section increased from 11.8% prior to COVID-19 practice alterations to 40.9% afterwards (P<0.0001); after vaginal birth, the proportion of 597 women (n = 597) who stayed only one night in hospital increased from 24.9% to 48.5% (P<0.0001) [27]. In New York, as the number of hospitalised patients with COVID-19 increased, the median postpartum length of stay decreased from a median of 48 to 34 h after vaginal birth (P<0.0001) and from a median of 74 to 51 h after caesarean section (P<0.0001) [52].

3.3. Postnatal care

Postnatal care is critical to the long term health of both mother and child – several studies highlighted reduced postpartum visit attendance [58,74], which was in some cases associated with reduced uptake of postnatal long acting contraception (OR 0.67 (95% CI 0.53–0.84)) [75] or probability of receiving screening for postpartum depression (86.2% vs 45.5%, P <0.01) [74]. This was not universal; in Rwanda postnatal care attendance was unaffected even as antenatal care contacts were reduced [57], while for women receiving remote postnatal follow up for hypertension in pregnancy care contacts were actually increased [44].

4. Discussion

This review has provided evidence that pregnant women have altered their healthcare-seeking behaviour during the COVID-19 pandemic, in a variety of contexts, and there has also been rapid and substantive change to maternity care provision globally. There has been a substantial decrease in the number of scheduled and unscheduled antenatal care visits, hospitalisations when urgent care has been sought, a reduction in antenatal care screening uptake (including but not limited to ultrasound and prenatal genetic testing), and delayed attendance at the planned place of care when labour starts. Maternity healthcare provision has also been affected as evidenced by a clear

Table 3

Characteristics and summary of the findings of studies and outcomes included in the meta-analysis.

Author (Year)	Pandemic Period	Comparison Period	Events in exposed cohort	Total number in exposed cohort	Events in comparison cohort	Total number in exposed cohort
Abdela (2020) [37]	23/3/2020–19/4/2020	24/02/2020–22/3/2020	ANC per week: 60	n/a	ANC per week: 86	n/a
Abel (2021) [23]	4/3/2020–1/9/2020	01/01/2020–03/03/2020	Unscheduled visits per week: 554	n/a	Unscheduled visits per week: 778	n/a
Baptiste (2021) [32]	01/01/2020–31/07/2020	01/01/2019–31/07/2019	Prenatal diagnosis procedures per year: 377	n/a	Prenatal diagnosis procedures per year: 464	n/a
Bhatia (2020) [48]	01/04/2020–01/07/2020	01/04/2019–01/07/2019	GA for ELCS: 14	1083	GA for ELCS: 28	1059
Dell'Utri (2020) [24]	23/02/2020 – 24/06/2020	23/02/2019 – 24/06/2019	Unscheduled visits per week: 260	n/a	Unscheduled visits per week: 403	n/a
Facco (2020) [36]	01/04/2020–01/07/2020	01/04/2019–01/07/2019	Unscheduled visits per week: 18	n/a	Unscheduled visits per week: 22	n/a
Holcomb (2020) [29]	22/3/2020–31/5/2020	23/2/2020–21/3/2020	ANC per week 1888 Virtual per week 399	n/a	ANC per week 2409 Virtual per week 0	n/a
Hui (2020) [30]	5/01/2020 – 30/04/2020	1/01/2019 – 4/01/2020	Epidural use: 126	954	Epidural use: 461	3577
Jeganathan (2020) [70]	1/3/2020–30/5/2019	1/3/2019–30/5/2019	ANC per week 42 Virtual per week 21	n/a	ANC per week 79 Virtual per week 0	n/a
Justman (2020) [50]	01/03/2020 – 30/04/2020	01/03/2019 – 30/04/2019	Epidural use: 507	610	Epidural use: 572	742
Khalil (2020) [31]	1/2/2020–14/6/2020	1/10/2019–31/1/2020	Unscheduled visits per week: 96	n/a	Unscheduled visits per week: 119	n/a
Kugelman (2020) [71]	15/03/2020 – 12/04/2020	15/03/2019 – 12/04/2019	Unscheduled visits per week: 136 Hospital admissions: 257	398	Unscheduled visits per week: 136 Hospital admissions: 279	544
Madden (2020) [66]	17/3/2020 – 12/4/2020	9/3/2020–16/3/2020	ANC per week 378 Virtual per week 187	n/a	ANC per week 417 Virtual per week 4	n/a
Meyer (2020) [39]	1/2/2020–28/3/2020	1/2/2019 – 28/3/2019	Unscheduled visits per week: 462	n/a	Unscheduled visits per week: 483	n/a
Monni (2020) [53]	10/03/2020–18/05/2020	10/03/2019–18/05/2019	Prenatal diagnosis procedures per year: 793	n/a	Prenatal diagnosis procedures per year: 772	n/a
Ozalp (2020) [54]	11/03/2020–30/06/2020	11/03/2019–30/06/2019	Prenatal diagnosis procedures per year: 56	129	Prenatal diagnosis procedures per year: 88	138
Patkar-Kamminati (2021) [49]	12/03/2020–11/06/2020	01/10/2019–31/12/2019	GA for ELCS: 4 GA for EMCS: 8	172 58	GA for ELCS: 2 GA for EMCS: 18	186 67
Peahl (2020) [42]	20/3/2020 – 28/6/2020	16/12/2019–20/3/2020	ANC per week 430 Virtual per week 332	n/a	ANC per week 805 Virtual per week 97	n/a
Salsi (2020) [68]	1/3/2020 – 31/3/2020	1/3/2019 – 31/3/2019	Unscheduled visits per week: 57 Hospital admissions: 164	254	Unscheduled visits per week: 90 Hospital admissions: 223	400
Sarkar (2021) [58]	23/02/2020–31/05/2020	12/01/2020–22/03/2020	ANC per week 99	n/a	ANC per week 321	n/a
Soffer (2021) [47]	01/04/2020–31/07/2020	01/04/2019–31/07/2019	ANC per week 594 Virtual per week 251	1296	ANC per week 1156 Virtual per week 0	1345

increase in virtual or remote consultations, decrease in face-to-face appointments, and reduction in waiting times; however, people with fewer resources within a population group may have had less access to telehealth, based on data from the USA. Moreover, there was a reduction in companionship allowed during birth, and a reduction in postpartum hospital length of stay, regardless of mode of birth.

The strengths of this review include the comprehensive literature search and rigorous methodology. However, the findings are limited

by the heterogeneity of the included studies and the variety of outcomes reported, which frequently precluded meta-analysis. Where substantial heterogeneity was identified in the quantitative synthesis, we must advise caution in reliance on the meta-analysis outcomes. First, there was no information about public health and local healthcare messaging to which people were exposed, although we are unaware of women in any jurisdiction being advised against healthcare-seeking when concerned. Second, although every hospital




Fig. 2. Forest plots for antenatal care visits per week (2a), unscheduled care attendance per week (2b), hospital admissions amongst unscheduled attendance (2c) and prenatal diagnostic procedures per year (2d).

and care provider will certainly have made changes in response to the COVID-19 pandemic, only a small proportion will have published their experience and outcomes, and not all of these can be guaranteed to have been identified from this search. In particular, despite evidence highlighting the disproportionate impact of the pandemic response on women and children in low resource settings [76], fewer than a third of the included papers originated from low or middle income countries. Third, patient and provider experience of remote antenatal care during the pandemic may not be generalisable to the future post-pandemic world. Patients frequently cited fear of COVID-19 as a key driver for avoiding face-to-face appointments, whether or not virtual care was available, so when this is no longer a concern, the perceived benefits of in-person consultation may vary. Fourth, potential advantages of remote consultation technology to increase flexibility and efficiency in pregnancy care must not compromise patient safety or the development of the essential therapeutic relationship that is core to safety in maternity care. Finally, we do not know the cost implications of the changes observed.

As each individual study reported on specific and highly variable protocols, they are unlikely to be generalisable but, taken together,

these studies demonstrate that significant changes in patient and provider behaviour and care provision occurred during the pandemic response. Where positive developments (e.g. increased access to antenatal services via hybrid face-to-face and remote monitoring care pathways or expedited postnatal discharge pathways) have been identified, some of these rapidly developed innovations are likely to result in permanent change.

The altered patient maternity care-seeking and maternity health-care provision demonstrated in this review must be considered as potentially contributing to worsening of pregnancy outcomes observed during the pandemic [9]. At this point, it is not possible to establish a causal link; where studies did report clinical outcomes for their included cohorts, the findings were mixed. For example, although it would seem plausible that delayed presentation in labour might be associated with worse perinatal outcomes, the small studies that reported on this outcome reported no differences in maternal or neonatal mortality [38,55,71]. One group used their detailed patient records to identify specific complications potentially attributable to, or exacerbated by, delay in seeking care, including anaemia, post-term pregnancy and pregnancy induced hypertension as the most




Fig. 2. Continued.

common, and suggested that these relate to an observed increase in ICU admission and maternal mortality during the pandemic in their small cohort [13]. One study explicitly assessed the changes in antenatal care provision in tandem with perinatal outcomes in the same period and found no difference in their high resource setting where a large proportion of antenatal contacts were delivered remotely [35], but this is not applicable to the contexts where antenatal visits were reduced and not mitigated by increased provision of remote or community based services.

This review provides evidence of reduced attendance for antenatal care and reduced uptake of antenatal screening for infection and foetal anomaly [25,32,54]. In addition, there is clear evidence that in some contexts women have avoided seeking urgent care for pregnancy concerns or attending the planned place of birth when labour occurs. All of these changes may introduce additional risk to mothers and babies, and are plausibly linked to the observed worsening of pregnancy outcomes during the pandemic, including an increase in stillbirths [9]. The significant increase in proportion of hospitalisations of those presenting for urgent care could reflect the proportionate increase in mandatory emergency visits, such as labour or rupture of membranes, compared to less urgent presentations, such as emesis or cramps [68,71,77].

Multiple reports considered the changes that occurred in antenatal care provision during the pandemic response. The introduction or upscaling of remote access technologies was a common feature of the

pandemic mitigation strategies implemented worldwide, particularly in high and middle income countries. Potential benefits identified included high levels of both patient and provider satisfaction, and a reduction in 'no shows' or 'did not attend' [29,42,66,69,70]. There are concerns that virtual consultations carried out in effect in the patient's home may make it less possible for pregnant people to disclose concerns for their own or their children's safety, and impair the development of the therapeutic relationship between woman and care providers. There has, for example, been an increase in intimate partner violence (IPV) during the pandemic, primarily directed against women [78–80]. Krishnamurthi et al. report increased uptake during the pandemic period of an app developed to support IPV reporting [81] but clearly this pathway is only available to women with access to a smartphone. Several groups noted the danger of reducing access for low income or vulnerable women who might be less able to access high speed internet or video capable personal devices.

This review has also identified evidence of change in practice in relation to obstetric anaesthesia and analgesia in labour. The initial concerns that women might be denied access to epidural analgesia in labour because of redeployment of limited anaesthetic staff were not borne out in the studies reporting on this outcome. We did find evidence that the use of general anaesthesia for intrapartum caesarean section was reduced, which must be examined further in relation to outcomes. Both the studies reporting this outcome originated from

Table 4
Results of the quantitative synthesis.

Outcomes	Studies	Pre-pandemic	Post-pandemic	Pooled estimate †(95% CI)	P	I[2]				
Number of antenatal care appointments per week	Holcomb [29]	2409	1888	IRR 0•614 (0•486 – 0•776)	<0•0001	54•6%				
	Jeganathan [70]	79	42							
	Abdela [37]	86	60							
	Madden [66]	417	378							
	Peahl [42]	805	430							
	Sarkar [58]	321	99							
	Soffer [47]	1156	594							
Number of virtual or over the phone visits per week	Holcomb [29]	0	399	IRR 46•56 (7•762 – 279•4)	<0•0001	90•6%				
	Jeganathan [70]	0	21							
	Madden [66]	4	187							
	Peahl [42]	97	332							
	Soffer [47]	0	251							
	Number of unscheduled care visits per week	Dell'Utri [24]	403				260	IRR 0•741 (0•602 – 0•911)	0•0046	0•0%
		Khalil [31]	119				96			
Kugelman [71]		136	99							
Meyer [39]		483	462							
Salsi [68]		90	57							
Abel [23]		778	554							
Facco [36]		22	18							
Baptiste [32]		464	377							
Number of prenatal diagnosis procedures per year	Baptiste [32]	464	377	IRR 0•880 (0•645 – 1•199)	0•41	0•0%				
	Ozalp [54]	88	56							
Hospital admissions amongst unscheduled care visits	Monni [53]	772	793	RR 1•214 (1•118 – 1•319)	<0•0001	0•0%				
	Salsi [68]	55•7% (223/400)	64•6% (164/254)							
	Kugelman [71]	51•3% (279/544)	64•6% (257/398)							
General anaesthesia for ELCS	Bhatia [48]	2•6% (28/1059)	1•3% (14/1083)	RR 0•831 (0•205 – 3•356)	0•79	61•8%				
General anaesthesia for EMCS	Patkar-Kamminati [49]	1•1% (2/186)	2•3% (4/172)	RR 0•529 (0•407 – 0•690)	<0•0001	0•0%				
	Bhatia [48]	24•3% (118/486)	12•9% (61/472)							
Epidural use	Patkar-Kamminati [49]	26•9% (18/67)	13•8% (8/58)	RR 1•044 (0•993 – 1•098)	0•0896	37•4%				
	Hui [30]	12•9% (461/3577)	13•2% (126/954)							
	Justman [50]	77•1% (572/742)	83•1% (507/610)							

†Individual patient data quantitative synthesis using generalised Poisson regression with random intercepts for studies reporting events per week. Mantel-Haenszel random effects meta-analysis for outcomes reported given a number of dependant events.
IRR: incidence rate ratio, RR: risk ratio, CI: confidence interval.

the UK where full aerosol PPE was recommended by Public Health England for emergency caesarean section under GA but not under spinal. This may not be observed in other settings, although the increased risk of GA to staff would have been known in every health-care facility. It may be that general anaesthesia was in fact overused in the pre-pandemic times, without benefits for mothers and babies, but it is also possible that reluctance to initiate general anaesthesia in the context of a pandemic respiratory virus could have contributed to delays in time critical emergency deliveries.

Parents and midwives have repeatedly expressed real concern about the impact of reduction in labour companionship on maternal experience and intrapartum outcomes, and this review has shown that labour companionship has significantly reduced in settings as disparate as Nepal and Hong Kong [30,64]. Labour companionship has been shown to affect both birth experience and outcome, and restrictions on companions should be carefully considered. Throughout the pandemic, the NHS in England has issued guidance emphasising the importance of supporting women to have a birth partner of their choice, although there was anecdotal reporting of local restrictions initially [82]. In high income settings fear of birthing alone has

been identified as a driver of increased planned home births, in lower resource settings without skilled attendance provision in the community and robust transfer pathways, this could potentially increase unattended birth. In Dessie region in Ethiopia, for example, 393% of women giving birth reported that their carers and attendants were not permitted to enter the hospital with them for the birth [83].

The COVID-19 pandemic has shone a harsh light of racial and social inequality, both within and between societies and regions. While we have identified some potential positive alterations in maternity care provision, it is likely that the majority of these benefits will be available to financially secure women in high income countries able to benefit from digital innovations in care provision. While in high income countries antenatal care has shifted to a hybrid model without sacrificing the number of contacts [29,33,35,36,44,47,65,66] in low resource settings without recourse to alternative models, steep reductions in antenatal care attendance were observed [13,28,37,56–58,60,61]. The development of pandemic response and recovery strategies must be sensitive to the needs of the most vulnerable women in their population – whether considering the loan of smart devices to vulnerable women in high income settings or

provision of alternate pathways to care. In low resource settings antenatal care attendance must be facilitated by ensuring women have confidence in the safety of healthcare facilities and the means to reach care where transport and mobility restrictions are in place.

Key themes identified by this review – maintaining key preventive care such as antenatal screening and routine care, the importance of clear communication, and considering the needs of those in lower socio-economic groups and lower income countries – are of relevance to all providers of maternity care. These must be considered as locally-responsive and culturally-appropriate care pathways are re-developed during the evolving pandemic response and into the future. They also provide the opportunity to challenge the established norms of maternity care and consider whether ‘returning to normal’ should be our goal.

One of the greatest healthcare lessons of this global pandemic has been that large structural change in maternity services in an extremely short time frame is possible. If length of postpartum stay can be safely and swiftly reduced, why did it take a pandemic to make this happen? Enhanced recovery protocols for obstetric care exist and could be more widely implemented, and innovative models of care make it possible to provide continuity of carer across hospital, clinic and community sites, and help to achieve the recommended 8 antenatal visits as recommended by the World Health Organization [84].

The COVID-19 pandemic has posed an unprecedented challenge to individuals, society and healthcare systems. This systematic review comprises a detailed and rigorous global assessment of changes in maternity healthcare provision, as well as use by pregnant women. It confirms that reduced maternity care-seeking and healthcare provision have occurred globally. These changes must be considered when evaluating whether demonstrable harms to both mothers and babies could have been avoided. We now have an opportunity to examine in depth the effects of this pandemic on maternity healthcare systems and outcomes, harness and refine the examples of excellent practice that have been implemented at pace, and discard or mitigate those that may have increased the risk of adverse outcomes. The strategic choices made now could either reverse or entrench the harms of this pandemic and their disproportionate effects on the poorest and most vulnerable women globally. It is imperative that we put in place mitigation strategies to minimise the collateral harm to mothers and babies in future health system shocks.

Funding

There was no funding received for this study. Therefore, no funder had any involvement in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Contribution and the data sharing statement

BC, IB, RT, EK and AK participated in conceptualisation, data curation, formal analysis, validation, visualisation, writing – original draft, as well as writing, review & editing of the manuscript.

LAM, JvdM, IGU EM, TD, ST, KL and SL participated in the conceptualisation, investigation, visualisation, and writing (original draft, as well as reviewing and editing).

PvD and POB participated in the conceptualisation, investigation, validation, visualisation, and writing (original draft, as well as reviewing and editing).

All authors have read and agreed to the published version of the manuscript.

Data collected for this meta-analysis have already been published in other studies. Data extracted from these published articles will be made available to others upon request. There are no individual

participant data due to the nature of this meta-analysis. The protocol is already published and is included as supplementary material. The data will be made available 3 months beginning 3 months and ending 5 years following article publication. The data will be made available to researchers who provide a methodologically sound proposal. Proposals should be directed to akhaliil@sgul.ac.uk; to gain access, data requestors will need to sign a data access agreement. Data are available for 5 years following the publication date

Declaration of Competing Interest

Dr Morris reports grants and other from Gedeon Richter, grants and other from Chugai Pharma, personal fees from Pfizer, personal fees from Gedeon Richter, other from Kebomed, from null, outside the submitted work; and President and Trustee, RCOG Trustee, British Menopause Society Chair of Trustees, Group B Strep Support.

Supplementary materials

Supplementary material associated with this article can be found in the online version at doi:[10.1016/j.eclinm.2021.100947](https://doi.org/10.1016/j.eclinm.2021.100947).

References

- [1] Khalil A, Kalafat E, Benlioglu C, et al. SARS-CoV-2 infection in pregnancy: a systematic review and meta-analysis of clinical features and pregnancy outcomes. *EClinicalMedicine* 2020;25:100446.
- [2] Allotey J, Stallings E, Bonet M, et al. Clinical manifestations, risk factors, and maternal and perinatal outcomes of coronavirus disease 2019 in pregnancy: living systematic review and meta-analysis. *BMJ* 2020;370:m3320.
- [3] Knight M, Bunch K, Cairns A, et al. Saving lives, improving mothers' care rapid report: learning from SARS-CoV-2-related and associated maternal deaths in the UK March – May 2020. Vol. 31, MBRRACE-UK. 2020. <https://www.npeu.ox.ac.uk/news/2013-mbrance-uk-release-saving-lives-improving-mothers-care-rapid-report> [Accessed 28.03.2021]
- [4] Khalil A, von Daddelsen P, Draycott T, Ugwumadu A, O'Brien P, Magee L. Change in the incidence of stillbirth and preterm delivery during the COVID-19 pandemic. *JAMA* 2020;324:705–6.
- [5] De Curtis M, Villani L, Polo A. Increase of stillbirth and decrease of late preterm infants during the COVID-19 pandemic lockdown. *Arch Dis Child Fetal Neonatal* editor 2020. doi: [10.1136/archdischild-2020-320682](https://doi.org/10.1136/archdischild-2020-320682).
- [6] Been JV, Burgos Ochoa L, Bertens LCM, Schoenmakers S, Steegers EAP, Reiss IKM. Impact of COVID-19 mitigation measures on the incidence of preterm birth: a national quasi-experimental study. *Lancet Public Health* 2020;5:e604–e11.
- [7] Philip RK, Purtill H, Reidy E, et al. Unprecedented reduction in births of very low birthweight (VLBW) and extremely low birthweight (ELBW) infants during the COVID-19 lockdown in Ireland: a ‘natural experiment’ allowing analysis of data from the prior two decades. *BMJ Glob Health* 2020;5:e003075.
- [8] Hedermann G, Hedley PL, Baekvad-Hansen M, et al. Danish premature birth rates during the COVID-19 lockdown. *Arch Dis Child Fetal Neonatal* editor 2021;106:93–5.
- [9] Chmielewska B, Barratt I, Townsend R, et al. Effects of the COVID-19 pandemic on maternal and perinatal outcomes: a systematic review and meta-analysis. *Lancet Glob Health* 2021. doi: [10.1016/S2214-109X\(21\)00079-6](https://doi.org/10.1016/S2214-109X(21)00079-6).
- [10] Jardine J, Relph S, Magee LA, et al. Maternity services in the UK during the coronavirus disease 2019 pandemic: a national survey of modifications to standard care. *BJOG* 2021;128:880–9.
- [11] Coxon K, Turienzo CF, Kweekel L, et al. The impact of the coronavirus (COVID-19) pandemic on maternity care in Europe Midwifery. 2020;88:102779.
- [12] Royal College of Obstetrics and Gynaecology. Coronavirus (COVID-19) Infection in Pregnancy. 2021. <https://www.rcog.org.uk/globalassets/documents/guidelines/2021-02-19-coronavirus-covid-19-infection-in-pregnancy-v13.pdf> [Accessed 28.03.2021]
- [13] Goyal M, Singh P, Singh K, Shekhar S, Agrawal N, Misra S. The effect of the COVID-19 pandemic on maternal health due to delay in seeking health care: experience from a tertiary center. *Int J Gynaecol Obstet* 2021;152:231–5.
- [14] Czeisler MÉ, Marynak K, Clarke KEN, et al. Delay or Avoidance of medical care because of COVID-19–related concerns – United States, June 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:1250–7.
- [15] Yerger P, Jalloh M, Coltart CEM, King C. Barriers to maternal health services during the Ebola outbreak in three West African countries: a literature review. *BMJ Global Health* 2020;5:e002974.
- [16] Moher D, Liberati A, Tetzlaff J, Altman DG. PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 2009;62:1006–12.
- [17] Hale T, Angrist N, Goldszmidt R, et al. A global panel database of pandemic policies (Oxford COVID-19 government response tracker). *Nat Hum Behav* 2021. doi: [10.1038/s41562-021-01079-8](https://doi.org/10.1038/s41562-021-01079-8).

- [18] Wells G., Shea B., O'Connell D., Peterson J., Welch V., Losos M., et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses [Internet]. Available from: http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp Accessed 01 January 2021.
- [19] Deeks JJ, Higgins JP, Altman DG, obotCSM Group. Analysing data and undertaking meta-analyses. *Cochrane handbook for systematic reviews of interventions*; 2019:241–84.
- [20] Hunter JP, Saratzis A, Sutton AJ, Boucher RH, Sayers RD, Bown MJ. In meta-analyses of proportion studies, funnel plots were found to be an inaccurate method of assessing publication bias. *J Clin Epidemiol* 2014;67:897–903.
- [21] Manzoli L, Flacco ME, D'Addario M, et al. Non-publication and delayed publication of randomized trials on vaccines: survey. *BMJ* 2014;348:g3058.
- [22] Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.
- [23] Abel MK, Alavi MX, Tierney C, Weintraub MR, Avins A, Zaritsky E. Coronavirus disease 2019 (COVID-19) and the incidence of obstetric and gynecologic emergency department visits in an integrated health care system. *Obstet Gynecol* 2021;137:581–3.
- [24] Dell'Utri C, Manzoni E, Cipriani S, et al. Effects of SARS Cov-2 epidemic on the obstetrical and gynecological emergency service accesses. What happened and what shall we expect now? *Eur J Obstet Gynecol Reprod Biol* 2020;254:64–8.
- [25] Filice C, Poma GL, Above E, et al. Is the fear of COVID-19 infection the same in all subjects? *Int J Infect Dis* 2020;97:331–3.
- [26] Futterman I, Rosenfeld E, Toaff M, et al. Addressing disparities in prenatal care via telehealth during COVID-19: prenatal satisfaction survey in East Harlem. *Am J Perinatol* 2021;38:88–92.
- [27] Greene NH, Kilpatrick SJ, Wong MS, Ozimek JA, Naqvi M. Impact of labor and delivery unit policy modifications on maternal and neonatal outcomes during the coronavirus disease 2019 pandemic. *Am J Obstet Gynecol MFM* 2020;2:100234.
- [28] Gu X-X, Chen K, Yu H, Liang G-Y, Chen H, Shen Y. How to prevent in-hospital COVID-19 infection and reassure women about the safety of pregnancy: experience from an obstetric center in China. *J Int Med Res* 2020;48:0300060520939337.
- [29] Holcomb D, Faucher MA, Bouzid J, Quint-Bouzid M, Nelson DB, Duryea E. Patient perspectives on audio-only virtual prenatal visits amidst the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic. *Obstet Gynecol* 2020;136:317–22.
- [30] Hui PW, Ma G, Seto MTY, Cheung KW. Effect of COVID-19 on delivery plans and postnatal depression scores of pregnant women. *Hong Kong Med* 2020. doi: 10.12809/hkmj208774.
- [31] Khalil A, von Dadelszen P, Kalafat E, et al. Change in obstetric attendance and activities during the COVID-19 pandemic. *Lancet Infect Dis* 2021. doi: 10.1016/S1473-3099(20)30779-9.
- [32] Baptiste C, Sutton D, Jacob T, et al. 904 Prenatal genetic diagnostic testing during the COVID-19 pandemic. *Am J Obstet Gynecol* 2021;224:S560–1.
- [33] Bernstein M, Mei JY, Patton E, Negi M. 787 Evaluating standard of care in a reduced prenatal care model in the COVID-19 pandemic. *Am J Obstet Gynecol* 2021;224:S490–1.
- [34] Bertozzi-Villa C, Arditi B, Syeda SK, et al. 1050 Effect of the COVID-19 pandemic on initial obstetric ultrasounds in New York City. *Am J Obstet Gynecol* 2021;224:S650–1.
- [35] Duryea EL, Adhikari EH, Ambia A, Spong C, McIntire D, Nelson DB. Comparison between in-person and audio-only virtual prenatal visits and perinatal outcomes. *JAMA Netw Open* 2021;4(4):e215854.
- [36] Facco F, Himes K. 993 did health care delivery changes driven by COVID alter pregnancy and postpartum visits? *Am J Obstet Gynecol* 2021;224 S615–6.
- [37] Abdela SG, Berhanu AB, Ferede LM, van Griensven J. Essential healthcare services in the face of COVID-19 prevention: experiences from a referral hospital in Ethiopia. *Am J Trop Med Hyg* 2020;103:1198–200.
- [38] McDonnell S, McNamee E, Lindow SW, O'Connell MP. The impact of the Covid-19 pandemic on maternity services: a review of maternal and neonatal outcomes before, during and after the pandemic. *Eur J Obstet Gynecol Reprod Biol* 2020;255:172–6.
- [39] Meyer R, Levin G, Hendin N, Katorza E. Impact of the COVID-19 outbreak on routine obstetrical management. *Isr Med Assoc J* 2020;22:483–8.
- [40] Weingarten SJ, Clare CA. 150 Management of diabetes in pregnancy during the COVID-19 pandemic at a New York City hospital. *Am J Obstet Gynecol* 2021;224:S103.
- [41] Shields AD, Wagner RK, Knutzen D, Deering S, Nielsen PE. Maintaining access to maternal fetal medicine care by telemedicine during a global pandemic. *J Telemed Telecare* 2020 10.1177/1357633 × 20957468.
- [42] Peahl AF, Powell A, Berlin H, et al. Patient and provider perspectives of a new prenatal care model introduced in response to the coronavirus disease 2019 pandemic. *Am J Obstet Gynecol* 2021;224 384.e1–e11.
- [43] Limaye MA, Lantigua-Martinez M, Trostle ME, et al. Differential uptake of telehealth for prenatal care in a large New York City academic obstetrical practice during the COVID-19 pandemic. *Am J Perinatol* 2021;38:304–6.
- [44] Greco P, Hesson A, Clifford C, Sangtani A, Stout MJ, Langen ES. 801 Hypertensive disorders of pregnancy in a pandemic: care delivery, outcomes, and lessons learned. *Am J Obstet Gynecol* 2021;224:S499.
- [45] Hussain FN, Garvey KL, Karotkin L, et al. 1028 Managing gestational diabetes mellitus with telemedicine during COVID-19: was there an impact on pregnancy outcomes? *Am J Obstet Gynecol* 2021;224:S636–7.
- [46] Selinger CP, Fraser A, Collins P, et al. Impact of the coronavirus infectious disease (COVID-19) pandemic on the provision of inflammatory bowel disease (IBD) antenatal care and outcomes of pregnancies in women with IBD. *BMJ Open Gastroenterol* 2021;8:e006063. doi: 10.1136/bmjgast-2021-000603.
- [47] Soffer MD, Sinnott C, Clapp MA, Bernstein SN. 514 Impact of a hybrid model of prenatal care on the diagnosis of fetal growth restriction. *Am J Obstet Gynecol* 2021;224:S324–5.
- [48] Bhatia K, Columb M, Bewlay A, et al. The effect of COVID-19 on general anaesthesia rates for caesarean section. A cross-sectional analysis of six hospitals in the north-west of England. *Anaesthesia* 2021;76:312–9.
- [49] Patkar-Kattimani C, Athod R, Sangtani D. COVID-19 and obstetric anaesthetic services in a tertiary maternity care unit. *Int J Obstet Anesth* 2021;45:152–3.
- [50] Justman N, Shahak G, Gutzeit O, et al. Lockdown with a Price: the impact of the COVID-19 Pandemic on Prenatal Care and Perinatal Outcomes in a Tertiary Care Center. *Isr Med Assoc J* 2020;22:533–7.
- [51] Racine JL, Hetzel S, Iruretagoyena JJ, Hoppe KK. 889 Did institutional changes and patient behaviors surrounding COVID-19 affect perinatal outcomes? *Am J Obstet Gynecol* 2021;224:S551–2.
- [52] Bornstein E, Guleren M, Husk G, et al. Early postpartum discharge during the COVID-19 pandemic. *J Perinat Med* 2020;48:1008–12.
- [53] Monni G, Corda V, Iuculano A. Prenatal screening diagnosis and management in the era of coronavirus: the Sardinian experience. *J Perinat Med* 2020;48:943–9.
- [54] Ozalp M, Demir O, Akbas H, Kaya E, Celik C, Osmanagaoglu MA. Effect of COVID-19 pandemic process on prenatal diagnostic procedures. *J Matern Fetal Neonatal Med* 2020;1–6. doi: 10.1080/14767058.2020.1815190.
- [55] Sun SY, Guazzelli CAF, de Moraes LR, et al. Effect of delayed obstetric labor care during the COVID-19 pandemic on perinatal outcomes. *Int J Gynaecol Obstet* 2020;151:287–9.
- [56] Ahmed T, Rahman AE, Amole TG, et al. The effect of COVID-19 on maternal newborn and child health (MNCH) services in Bangladesh, Nigeria and South Africa: call for a contextualised pandemic response in LMICs. *Int J Equity Health* 2021;20:77.
- [57] Wanyana D, Wong R, Hakizimana D. Rapid assessment on the utilization of maternal and child health services during COVID-19 in Rwanda. *Public Heal action* 2021;11:12–21.
- [58] Sarkar S, Chowdhury RROY, Mukherji J, Samanta M, Bera G. Comparison of attendance of patients pre-lockdown and during lockdown in gynaecology and antenatal outpatient department in a tertiary care hospital of Nadia, West Bengal, India. *J Clin Diagnostic Res* 2021;15(2) QC05–8.
- [59] Chen M, Liu X, Zhang J, et al. Characteristics of online medical care consultation for pregnant women during the COVID-19 outbreak: cross-sectional study. *BMJ Open* 2020;10:e043461.
- [60] Moyer CA, Sakyi KS, Sacks E, Compton SD, Lori JR, Williams JEO. COVID-19 is increasing Ghanaian pregnant women's anxiety and reducing healthcare seeking. *Int J Gynecol Obstet* 2021;152:444–5.
- [61] Tadesse E. Antenatal care service utilization of pregnant women attending antenatal care in public hospitals during the COVID-19 pandemic period. *Int J Womens Health* 2020;12:1181–8.
- [62] Liu X, Chen M, Wang Y, et al. Prenatal anxiety and obstetric decisions among pregnant women in Wuhan and Chongqing during the COVID-19 outbreak: a cross-sectional study. *BJOG* 2020;127:1229–40.
- [63] Kumari V, Mehta K, Choudhary R. COVID-19 outbreak and decreased hospitalisation of pregnant women in labour. *Lancet Glob Heal* 2020;8:e1116–7.
- [64] KC A, Gurung R, Kinney M V, et al. Effect of the COVID-19 pandemic response on intrapartum care, stillbirth, and neonatal mortality outcomes in Nepal: a prospective observational study. *Lancet Glob Heal* 2020;8:e1273–e81.
- [65] Zarasvand S, Bayar E, Adan M, et al. Rapid quality improvement in a preterm birth clinic care pathway during the COVID-19 pandemic. *BMJ Open Qual* 2020;9:e001049.
- [66] Madden N, Emeruwa UN, Friedman AM, et al. Telehealth uptake into prenatal care and provider attitudes during the COVID-19 pandemic in New York City: a quantitative and qualitative analysis. *Am J Perinatol* 2020;37:1005–14.
- [67] Silverman ME, Medeiros C, Burgos L. Early pregnancy mood before and during COVID-19 community restrictions among women of low socioeconomic status in New York City: a preliminary study. *Arch Womens Ment Health* 2020;23:779–82.
- [68] Salsi G, Seidenari A, Diglio J, Bellussi F, Pilu G, Bellussi F. Obstetrics and gynecology emergency services during the coronavirus disease 2019 pandemic. *Am J Obstet Gynecol MFM* 2020;2:100214.
- [69] Biviá-Roig G, La Rosa VL, Gómez-Tébar M, et al. Analysis of the impact of the confinement resulting from COVID-19 on the lifestyle and psychological wellbeing of Spanish pregnant women: an internet-based cross-sectional survey. *Int J Environ Res Public Health* 2020;17:5933.
- [70] Jeganathan S, Prasannan L, Blitz MJ, Vohra N, Rochelson B, Meirowitz N. Adherence and acceptability of telehealth appointments for high-risk obstetrical patients during the coronavirus disease 2019 pandemic. *Am J Obstet Gynecol MFM* 2020;2:100233.
- [71] Kugelmann N, Lavie O, Assaf W, et al. Changes in the obstetrical emergency department profile during the COVID-19 pandemic. *J Matern Neonatal Med* 2020. doi: 10.1080/14767058.2020.1847072.
- [72] Albert L, Capel I, García-Sáez G, Martín-Redondo P, Hernando ME, Rigla M. Managing gestational diabetes mellitus using a smartphone application with artificial intelligence (SineDie) during the COVID-19 pandemic: much more than just telemedicine. *Diabetes Res Clin Pract* 2020;169:108396. doi: 10.1016/j.diabres.2020.108396.
- [73] Gildner TE, Thayer ZM. Birth plan alterations among American women in response to COVID-19. *Heal Expect* 2020;23:969–71.

- [74] Sakowicz A, Imeroni S, Matovina C, Daiter M, Grobman WA, Miller ES. 590 Postpartum depression screening during the COVID-19 pandemic. *Am J Obstet Gynecol* 2021;224:S371–2.
- [75] Sakowicz A, Matovina C, Imeroni S, et al. 591 The association between COVID-19 related health services changes and postpartum contraception. *Am J Obstet Gynecol* 2021;224:S372.
- [76] Kumar J, Kumar P. COVID-19 pandemic and health-care disruptions: count the most vulnerable. *Lancet Glob Heal* 2021. doi: 10.1016/S2214-109X(21)00098-X.
- [77] Anderson TS, Stevens JP, Pinheiro A, Li S, Herzig SJ. Hospitalizations for emergent medical, surgical, and obstetric conditions in Boston during the COVID-19 pandemic. *J Gen Intern Med* 2020;35:3129–32.
- [78] Hamadani JD, Hasan MI, Baldi AJ, et al. Immediate impact of stay-at-home orders to control COVID-19 transmission on socioeconomic conditions, food insecurity, mental health, and intimate partner violence in Bangladeshi women and their families: an interrupted time series. *Lancet Glob Heal* 2020;8:e1380–9.
- [79] Feder G, Lucas d'Oliveira AF, Rishal P, Johnson M. Domestic violence during the pandemic. *BMJ* 2021;372:n722.
- [80] Evans ML, Lindauer M, Farrell ME. A pandemic within a pandemic – intimate partner violence during Covid-19. *N Engl J Med* 2020;383:2302–4.
- [81] Krishnamurti T, Davis AL, Quinn B, Castillo AF, Martin KL, Simhan HN. Mobile remote monitoring of intimate partner violence among pregnant patients during the COVID-19 shelter-in-place order: quality improvement pilot study. *J Med Internet Res* 2021;23:e22790.
- [82] England NHS. Supporting pregnant women using maternity services during the coronavirus pandemic. London, UK: Actions for NHS providers; 2020 <https://www.england.nhs.uk/coronavirus/publication/supporting-pregnant-women-using-maternity-services-during-the-coronavirus-pandemic-actions-for-nhs-providers/> [Accessed 28.03.2021].
- [83] Assefa KT, Gashu AW, Mulualem TD. The impact of COVID-19 infection on maternal and reproductive health care services in governmental health institutions of Dessie town. North-East Ethiopia; 2020 G.C. *medRxiv* 2020; 2020.09.20.20198259.
- [84] World Health Organisation (WHO). WHO recommendations on antenatal care for a positive pregnancy experience. Geneva, 2016.