



Contents lists available at ScienceDirect

Journal of Traditional and Complementary Medicine

journal homepage: <http://www.elsevier.com/locate/jtcme>



Original Article

Therapeutic potential of active components of saffron in post-surgical adhesion band formation



Mohammad-Hassan Arjmand ^{a,1}, Milad Hashemzehi ^{b,1}, Atena Soleimani ^{c,1}, Fereshteh Asgharzadeh ^{d,1}, Amir Avan ^{e,f}, Saeedeh Mehraban ^g, Maryam Fakhraei ^d, Gordon A. Ferns ^h, Mikhail Ryzhikov ⁱ, Masoumeh Gharib ^j, Roshanak Salari ^k, Sayyed Hadi Sayyed Hoseini ^l, Mohammad Reza Parizadeh ^c, Majid Khazaei ^{d,e,**}, Seyed Mahdi Hassanian ^{c,e,*}

^a Medical Plants Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

^b Iranshahr University of Medical Sciences, Iranshahr, Iran

^c Department of Clinical Biochemistry, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

^d Department of Physiology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

^e Metabolic Syndrome Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

^f Student Research Committee and Department of Medical Genetics, Faculty of Medicine, Mashhad University of Medical Science, Mashhad, Iran

^g Immunology Research Center, Inflammation and Inflammatory Diseases Division, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

^h Brighton & Sussex Medical School, Division of Medical Education, Falmer, Brighton, BN1 9PH, UK

ⁱ Division of Pulmonary and Critical Care Medicine, Washington University, School of Medicine, Saint Louis, MO, USA

^j Department of Pathology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

^k Department of Pharmaceutical Sciences in Persian Medicine, School of Persian and Complementary Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

^l Orthopedic Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

ARTICLE INFO

Article history:

Received 15 June 2020

Received in revised form

26 October 2020

Accepted 4 January 2021

Available online 20 January 2021

Keywords:

Saffron

Crocin

Crocetin

Post-surgical adhesion band formation

Fibrosis

Inflammation

ABSTRACT

Background: Abdominal adhesions are common and often develop after abdominal surgery. There are currently no useful targeted pharmacotherapies for adhesive disease. Saffron and its active constituents, Crocin and Crocetin, are widely used in traditional medicine for alleviating the severity of inflammatory or malignant disease.

Purpose: The aim of this study was to investigate the therapeutic potential of the pharmacological active component of saffron in attenuating the formation of post-operative adhesion bands using different administration methods in a murine model.

Material method: saffron extract (100 mg/kg), Crocin (100 mg/kg), and Crocetin (100 mg/kg) were administered intraperitoneally and by gavage in various groups of male Wistar rat post-surgery. Also three groups were first treated intra-peritoneally by saffron extract, Crocin, and Crocetin (100 mg/kg) for 10 days and then had surgery. At the end of the experiments, animals sacrificed for biological assessment.

Result: A hydro-alcoholic extract of saffron and crocin but not crocetin potently reduced the adhesion band frequency in treatment and pre-treatment groups in the mice given intra-peritoneal (i.p) injections. Following the saffron or crocin administration, histological evaluation and quantitative analysis represented less inflammatory cell infiltration and less collagen composition, compared to control group. Moreover, the oxidative stress was significantly reduced in treatment groups.

Abbreviations: APC, activated protein C; DSS, dextran sodium sulfate; HE, Hematoxylin & Eosin; IP, intera-peritoneal; MDA, malondialdehyde; PSAB, post-surgical adhesion band; PDGF, platelet-derived growth factor; TGF- β , transforming growth factor-beta; α -SMA, α -smooth muscle actin; SOD, superoxidase dismutase; TAA, thioacetamide.

* Corresponding author. Department of Clinical Biochemistry, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

** Corresponding author. Department of Medical Physiology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

E-mail addresses: khazaeim@mums.ac.ir (M. Khazaei), hasanianmehrm@mums.ac.ir (S.M. Hassanian).

Peer review under responsibility of The Center for Food and Biomolecules, National Taiwan University.

¹ Made equal contribution to this study. This study was supported by grants awarded by the National Institute for Medical Research Development (Grant No.977089).

Conclusion: These findings suggest that a hydro-alcoholic extract of saffron or its active compound, crocin, is a potentially novel therapeutic strategy for the prevention of adhesions formation and might be used as beneficial anti-inflammatory or anti-fibrosis agents in clinical trials.

Taxonomy: Abdominal surgeries/post-surgical adhesions.

© 2021 Center for Food and Biomolecules, National Taiwan University. Production and hosting by Elsevier Taiwan LLC. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Intraperitoneal adhesion formation is a serious worldwide complication of surgery developing between abdominal wall and intra-abdominal surfaces following peritoneal irritation. This condition is highly prevalent (up to 90%) in patients undergoing open gynecological pelvic or open abdominal surgery.^{1,2} Intestinal obstruction,³ female infertility,⁴ chronic abdominal and pelvic pain⁵ are the most important consequences of post-operative adhesion bands. Placement of some solid barriers such as Sepra-film® is a standard strategy for adhesion bands prevention which needs to accurately recognize the damage sites.⁶ Predicting where these fibrotic bands may develop is a barriers usage limitation encouraging the researchers to find an efficient therapeutic method.⁶ Therefore, identifying the exact mechanisms involved in adhesion band formation would help to progress the treatment process.

It is generally accepted that pathological adhesions generation is the end-result of improper healing process in peritoneal cavity.⁷ In post-operative physiological responses, immune cell-released inflammatory cytokines, as well as platelet-derived growth factor (PDGF), and fibroblast-induced transforming growth factor-beta (TGF-β) are increased in peritoneum.^{8–11} It has been shown that inflammation and fibrosis play key roles in the pathogenesis of adhesion band formation.^{11–13} We previously investigated the effects of anti-fibrotic or anti-inflammatory compounds, EW-7197¹⁴ and activated protein C (APC)¹⁵ on post-surgical adhesions, representing the hopeful results in reduction of these fibrotic bands. Although there are extensive studies on post-surgical adhesion band formation, our knowledge of pathogenic mechanism of this surgery-associated complication is still limited.

The plant-based therapies to improve pathologic conditions have been used for centuries and counted as potential treatments due to low complications and minimal toxicities. The huge parts of FDA-approved components are contained the therapeutics agents derived from natural components.^{16,17} The natural products continue to play an important role in drug discovery programs for different diseases. Using different technical approaches,^{18,19} studies showed the effectiveness of these components for various disorders such as carcinomas,^{20–26} diabetes,²⁷ various inflammatory and fibrotic conditions.^{28–32} One of these natural products was isolated from *Crocus sativus*, a flowering plant in the Iridaceae family and is commonly known as saffron.³³

Saffron, the dried stigmas of *Crocus sativus*, is cultivated in Iran and other countries such as Spain and Turkey³⁴ and has a wide range of activities including oxytocic, anti-carcinogenic,^{35,36} and etc. . There are several active pharmacological components in saffron including crocin and crocetin that are responsible for the biological effects of saffron.^{37,38} Saffron contains essential mineral and various important vitamins. Generally, due to anti-aging and anti-oxidant function of saffron, its daily consumption is very high in many parts of the world. Also, the ability of saffron, crocin or crocetin to reducing adverse effects of chemotherapeutic components showed their drug modulator activities.³⁹ Animal studies

showed low or non-toxicity of saffron and its extract.⁴⁰ A temporary immunomodulatory activity with no renal, hepatic, hematological, or any side effects were reported in sub-chronic use of 100 mg/day saffron in a randomized controlled trial study.⁴¹ Studies revealed that these compounds elicit anti-inflammatory and anti-fibrotic responses by decreasing the inflammatory mediators in different animal models^{42–44} which is consistent with our previous work, showing anti-inflammatory effects of crocin in mice model.⁴⁵ In line with this, saffron potently decreased the level of Malondialdehyde (MDA) and inflammatory cytokines in bleomycin-induced pulmonary fibrosis.⁴⁶ Similarly, crocin significantly reduced the over-expression of fibrotic molecules such as collagen 1a, α-smooth muscle actin (α-SMA), and TGF-β in hepatic fibrosis condition.⁴⁷

In this study, we investigated the protective effects of saffron, crocin, or crocetin in preventing post-surgical adhesion band formation in animal models. Our results showed that saffron or crocin (i.p) could suppress the formation of adhesion bands through anti-inflammatory, anti-oxidant and anti-fibrosis function in rat model, suggesting the therapeutic potential of this non-toxic compounds against post-surgery-associated complications.

2. Material and methods

2.1. Animal

Animal experiments were conducted in according to the guideline for Care and Use of Laboratory Animals from Mashhad University of Medical Sciences (MUMS). Animals were purchased from pasture institute (Tehran, Iran) and were subjected to controlled conditions of temperature ($20 \pm 2^\circ\text{C}$) and humidity (50%). The animals were housed in a standard cage in a 12 h light-dark cycle room with free access to both food and water. This study was approved by ethical committee of Mashhad University of Medical Sciences, ethical approval number: 960917.

2.2. Reagents

Pure crocin and crocetin were purchased from BuAli Pharmaceutical Research Center (MUMS, Mashhad, Iran). Hydro-alcoholic extract of Saffron was isolated by Maceration method as described.^{48,49} Reagents need for hematoxylin and eosin, malonyl dialdehyde (MDA), total thiol, and catalase were purchased from Sigma Co (Saint Louis, MO).

2.3. Surgical abrasion model

Surgical method for adhesion band formation was performed as described.^{50,51} Briefly, the animals were anesthetized by intraperitoneal injection of ketamine.⁵² After shaving and preparation of the site of surgery with 1% antiseptic povidone-iodine solution, the abdominal cavity was opened with 1.5 cm incision. The anterior cecal surface was gently rubbed until partial petechial hemorrhages were generated and the interior surface of abdomen was damaged using a medical electric scalpel. The cecum placed in front of

damaged surface of abdomen and the muscular layer was sutured. Intraperitoneal adhesion formation occurs with this procedure probability above 90% between damaged surfaces.⁵³ The evaluation of post-surgical adhesion bands was performed according to the previously explained adhesions grade criteria by Nair et al.⁵⁴ in a blinded fashion, assessing the incidence and frequency of fibrotic bands. The detailed of Nair scoring system is shown in Table 1.

2.4. Experimental design

Rats were randomly divided into the following groups of 6 rats in each group: 1) Control group (animals with surgical abrasion received saline as vehicle intraperitoneally (i.p)); 2) animals with surgical abrasion treated intraperitoneally with 100 mg/kg/day saffron extract; 3) animals with surgical abrasion treated intraperitoneally with 100 mg/kg/day crocin; 4) animals with surgical abrasion treated intraperitoneally with 100 mg/kg/day crocetin.^{55–58}

To investigate whether type of administration, i. p or gavage could affect protective effects of these compounds, we also add the following groups: 5) animals with surgical abrasion treated by oral gavage with 100 mg/kg/day saffron extract; 6) animals with surgical abrasion treated by oral gavage with 100 mg/kg/day crocin; 7) animals with surgical abrasion treated by oral gavage with 100 mg/kg/day crocetin.

Next, we were interested in evaluating the preventive potential of these compounds in post-surgical adhesion band formation. In this case, before the surgery, we consider more groups: 8) animals were first treated intra-peritoneally with 100 mg/kg/day saffron extract for 10 days and then had surgery; 9) animals were first treated intra-peritoneally with 100 mg/kg/day crocin for 10 days and then had surgery; 10) animals were first treated intra-peritoneally with 100 mg/kg/day crocetin for 10 days and then had surgery.

All animals weighted before surgical abrasion and during the experiment. At the end of the experiments, animals sacrificed for biological assessment. The excised tissue samples were quickly frozen in liquid nitrogen or fixed in 10% formalin solution.

2.5. Histological analysis

Tissue sections were fixed in formalin 10% for 24–72 h. After processing and paraffin embedding, staining was done with either Hematoxylin & Eosin (HE) or Masson's Trichrome for assessing inflammation and collagen deposition, respectively.^{59,60} Quantifying the tissue staining presented histological grading of inflammation and fibrosis scores.^{61,62} The slides were seen under light microscope (magnification $\times 400$).

2.6. Preparation tissue homogenates

Adhesion bands-linked cecum was collected for making tissue homogenate. The tissue sample (100 mg tissue sample for each animal) was weighted and homogenized with PBS as the homogenization medium. The supernatant of the tissue homogenate was

used for the assay of stress oxidative markers.⁶³

2.7. Measurement of stress oxidative markers

The oxidative stress was evaluated in each subject by measuring the concentration levels of malondiladehyde (MDA) and thiol, as well as catalase activity in the tissue homogenates. All procedures were conducted according to the kit protocol.^{64–66}

2.8. Statistical analysis

Quantitative variables were described as mean \pm SEM. One-way ANOVA test was used for comparison between different groups. The collected data were imported into GraphPad Prism for statistical analysis. The statistical significance was set to 5%.

3. Results

3.1. Saffron and crocin significantly reduced adhesion scores

We investigated the therapeutic effect of hydro-alcoholic extracted of saffron, crocin, and crocetin in different administration in adhesion rat model. No animal death occurred during the experiment, and the body weight did not significantly differ after second surgery. There was no infection or bleeding in post-operative time. Schematic representation of the study protocol is shown in Fig. 1A. According to clinical observations, pre-treatment with crocin is the treatment of choice, offering more recovery and reducing the adhesion formation probability (Fig. 1B). We quantitatively analyzed the incidence of adhesion band in different groups, using Nair⁵⁴ scoring scheme (Fig. 1C). Compared to the rat control group, pre-treatment or treatment (i.p) with saffron or crocin significantly improved the frequency of post-operative intra-abdominal adhesion bands (P value < 0.05). Moreover, decrease in adhesion bands frequency (Nair grade) was not significant in oral administration groups. No reduction of adhesion bands were found in the case of crocetin treatment (Fig. 1B and C), compared to control cases. We did not continue experiments on crocetin due to adverse results.

3.2. Saffron extract and crocin decreased inflammation in post-surgical adhesion bands

Since inflammatory responses are increased during adhesion bands formation,⁶⁷ we evaluated the inhibitory effects of saffron and its pharmacological active components on inflammation in adhesion bands tissues, using Hematoxylin & eosin (HE) staining. Histological analysis and quantitative evaluation showed a significant reduction of inflammation in treatment as well as pre-treatment groups (Fig. 2A–C). As expected, oral treatment with hydro-alcoholic extract of Saffron showed no significant differences, compared to control tissues (Fig. 2C). The detailed of inflammation scoring system is shown in Table 2.

Table 1
Adhesion score system for macroscopic evaluation (Nair's et al.).

0 to 4	Adhesion grade
0	Complete absence of adhesion
1	Single band of adhesion, between viscera or from viscera to abdominal wall
2	Two bands, either or from viscera to abdominal wall
3	More than two bands, between viscera or viscera to abdominal wall
4	Viscera directly adherent to abdominal wall, irrespective of number and extent of adhesive bands




Fig. 1. Saffron and Crocin significantly attenuate adhesion band formation in rat model. (A) Schematic representation of the experimental protocol. (B) Macroscopic illustrations of different groups of treatment (Identified adhesion bands are shown with arrows). (C) Compared to control group, intraperitoneal treatment or pre-treatment with Saffron (100 mg/kg/day) or Crocin (100 mg/kg/day) significantly reduced the frequency of adhesion bands in rat. Treated mice showed no response to intraperitoneal injection of Crocetin (100 mg/kg/day). The sample size For all groups was n = 6 mice/group. *p < 0.05; **p < 0.01; ***p < 0.001.

3.3. Saffron and crocin decreased inflammatory responses via reducing oxidative stress markers in post-surgical adhesion band

To further determine the anti-inflammatory mechanisms of Crocin and Saffron in adhesion bands, we measured concentration of oxidant marker, Malon deladehyde (MDA), as well as antioxidants agents including total Thiol concentration and Catalase activity in adhesive tissue homogenates. Our findings showed significant reduction of MDA concentration in all treatment group, compared to control group (Fig. 3A). Consistently, the level of total Thiol (Fig. 3B) and catalase activity (Fig. 3C) were higher in treated groups than control tissues. These results supported the hypothesis that saffron and crocin elicited their protective functions at least partially by attenuating oxidative stress reactions in post-operative adhesion rat model.

3.4. Inhibitory effects of saffron and crocin on tissue fibrosis in adhesion rat model

Next, we investigated the regulatory effects of saffron and crocin on fibrosis as key factors in the pathogenesis of adhesion band formation. Masson's trichrome staining accompanied with quantitative analysis demonstrated that crocin or hydro-alcoholic extracted of Saffron could significantly decrease the areas of fibrosis and collagen deposition in adhesive tissues (Fig. 4A–C). The therapeutic effect was more potent in crocin administration group. The fibroblast activity quantified according to the scoring system presented in Table 2. Similar to previous results, oral treatment of Saffron elicited no protective effect in adhesion tissues (Fig. 4B) (P value < 0.05).

4. Discussion

The present study investigated the protective effects of hydro-

alcoholic extract of saffron, crocin and crocetin on intra-abdominal adhesion models, using different administration methods. Macroscopic results demonstrated a significant reduction of adhesion bands frequency in intraperitoneal administration of saffron or crocin as well as the pre-treatment group. However, neither crocetin, nor oral administration of saffron could prevent post-surgical adhesion band formation in rat model. We also showed that decrease in the oxidative stress responses as well as attenuating fibrosis and collagen depositions are some of the mechanisms by which saffron and crocin (i.p.) exert their protective responses in adhesion model, supporting the therapeutic potential of these low-toxic compounds against post-surgical adhesion band formation.

Damage to peritoneum leading to the deposition of peritoneal fibrin and local inflammation are the key steps of adhesiogenesis.⁹ Studies have shown that using fibrinolytic compounds prevent the progress of adhesion formation via stimulation of the intraperitoneal fibrinolytic system.^{68–70} Moreover, it has been found that adhesion formation is accompanied by lower tissue oxygenation and free oxygen radical generation.^{71,72} Ezberci et al.⁷³ showed that decrease in oxidant agent including MDA whereas increase in catalase activity attenuated adhesion score in Bacterial peritonitis rat model. Consistent with these results, our previous finding revealed that decrease in oxidative stress markers, as well as collagen deposition and infiltration of inflammatory cells to injured site attenuated the severity of adhesions in animal model.¹⁴ Also, we showed that human APC significantly reduced the formation of adhesion bands which is correlated with lower concentration of pro-inflammatory cytokines and higher tissues plasminogen activator (TPA) *in vivo*.¹⁵

Furthermore, there are several studies supporting the anti-oxidative, anti-inflammatory, and anti-fibrotic properties of saffron and crocin in different models. For instance, Hemshekhar et al.⁷⁴ showed that crocin enhanced anti-oxidant status by




Fig. 2. The inhibitory effects of Saffron and Crocin on inflammation in adhesive tissues. (A, C) Intraperitoneal injection of Crocin (100 mg/kg/day) or Saffron (100 mg/kg/day) reduced inflammatory cell infiltration (black arrows) in (A) treatment and (C) pre-treatment groups. (B) Despite crocin, oral administration of Saffron showed no significant protective responses in adhesion rat model. ***p < 0.001.

Table 2

Infiltration of inflammatory cells and fibrosis scoring according to Swolin.

Grade	Inflammatory cell infiltrate	Fibrosis
0	Absent or normal in number	None
1	Slight increase	Slight
2	Moderate infiltration	Moderate
3	Dense	Dense

increasing catalase, and superoxidase dismutase (SOD) activities as well as attenuating serum level of inflammatory factors in arteritis rat model. Moreover, Samarghandian et al.⁷⁵ reported that crocin improved aged rat kidney functions by reducing oxidative stress and inflammatory responses in rat. Consistently, Hashemi et al. indicated the effects of saffron carotenoids, crocin and crocetin on oxidative stress in breast tumor. They showed that crocin and crocetin increase the catalase⁷⁶ and superoxide dismutase⁷⁷ activities in BALB/c mice, after 28 days of treatment. Since crocin treatment has been shown antioxidant activities under various conditions, Nasimian et al. showed that crocin elevated the apoptotic death of human breast cancer cell lines, partially via ROS-activated FOXO3a axis.⁷⁸ Moreover, clinical trial studies evaluated the effect of saffron aqueous extract, crocin⁷⁹ and crocetin⁸⁰ in coronary artery disease (CAD). It has been shown that crocin and crocetin treatment resulted in a significant reduction in lectin-like oxidized LDL receptor 1 (LOX1), nuclear factor kappa-B (NF- κ B) and Serum ox-LDL. Also, the levels of monocyte chemoattractant

protein 1 (MCP-1) was reduced in all treatment groups in CAD patients.^{79,80} Similarly, we recently showed that crocin significantly inhibited oxidative stress and histopathological scores, representing a reduction of inflammation and fibrosis in dextran sodium sulfate (DSS)-induced colitis model.⁴⁵ Another study showed that 100 mg/kg of saffron reduced MDA, Myeloperoxidase, and tumor-necrosis factor-alpha (TNF- α) in pulmonary fibrosis.⁸¹

Consistent with the fibrinolysis effect of crocin in post-surgical adhesion model, it has been shown that saffron could significantly prevent the thickness of alveolar septa and collagen deposition in bleomycin-induced pulmonary fibrosis.⁸¹ In line with these results, Mehrabani et al. demonstrated the protective effects of crocin against fibrosis and hydroxyproline content of lungs.⁸² Chhimwal et al. showed that crocin decreased the hepatic fibrosis via expression of peroxisome proliferator-activated receptor γ (PPAR- γ), modulating the inflammatory and fibrogenic pathways.⁸³ These results are consistent with another study showing the anti-fibrotic and anti-inflammatory properties of crocin in thioacetamide (TAA)-induced liver fibrosis.⁴⁷

In conclusion, this study suggested that saffron and crocin have potential therapeutic value in preventing intra-abdominal adhesion band formation. The mechanism underlying abdominal adhesions has not yet been completely understood. Further animal and clinical studies are required to clarify this issue and to assess the exact mechanism of action for Saffron and its pharmacologically active component, crocin, for preventing adhesion bands formation.




Fig. 3. Oxidative stress is attenuated following Saffron- or Crocin-treatment in adhesive tissues. (A–C) The tissue concentration of (A) MDA, (B) total Thiol, and (C) the activity of Catalase were compared between different groups. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.




Fig. 4. Saffron and Crocin suppressed fibrosis and Collagenesis in adhesive tissues. (A, C) Masson's trichrome staining showed a significant reduction of deposition of Collagen (asterix) in Crocin- or Saffron-treated rat in different groups (B) The decrease in collagen thickening was not significant in the case of oral administration of Saffron, compared to control. *** $p < 0.001$.

Data availability statement

Research Data are not shared.

Declaration of competing interest

The authors have no conflicts of interest.

References

- Menzies D, Ellis H. Intestinal obstruction from adhesions—how big is the problem? *Ann R Coll Surg Engl.* 1990;72(1):60.
- Okabayashi K, Ashrafi H, Zacharakis E, et al. Adhesions after abdominal surgery: a systematic review of the incidence, distribution and severity. *Surg Today.* 2014;44(3):405–420.
- Menzies D. Peritoneal adhesions. Incidence, cause, and prevention. *Surg Annu.* 1992;24:27.
- Kavic SM, Kavic SM. Adhesions and adhesiolysis: the role of laparoscopy. *J Soc Laparoendosc Surg; J Soc Laparoendosc Surg.* 2002;6(2):99.
- Sulaiman H, Gabella G, Davis C, et al. Growth of nerve fibres into murine peritoneal adhesions. *J Pathol.* 2000;192(3):396–403.
- Cai X, Hu S, Yu B, et al. Transglutaminase-catalyzed preparation of crosslinked carboxymethyl chitosan/carboxymethyl cellulose/collagen composite membrane for postsurgical peritoneal adhesion prevention. *Carbohydr Polym.* 2018;201:201–210.
- diZerega GS, Campeau JD. Peritoneal repair and post-surgical adhesion formation. *Hum Reprod Update.* 2001;7(6):547–555.
- Surgeons PCoASRMicwtSoR. Pathogenesis, consequences, and control of peritoneal adhesions in gynecologic surgery: a committee opinion. *Fertil Steril.* 2013;99(6):1550–1555.
- Saed GM, Zhang W, Diamond MP. Molecular characterization of fibroblasts isolated from human peritoneum and adhesions. *Fertil Steril.* 2001;75(4):763–768.
- Saed GM, Diamond MP. Modulation of the expression of tissue plasminogen activator and its inhibitor by hypoxia in human peritoneal and adhesion fibroblasts. *Fertil Steril.* 2003;79(1):164–168.
- Rout UK, Saed GM, Diamond MP. Transforming growth factor- β 1 modulates expression of adhesion and cytoskeletal proteins in human peritoneal fibroblasts. *Fertil Steril.* 2002;78(1):154–161.
- Chegini N. TGF- β system: the principal profibrotic mediator of peritoneal adhesion formation. Paper presented at: Seminars in reproductive medicine 2008.
- White JC, Jiang ZL, Diamond MP, Saed GM. Macrophages induce the adhesion phenotype in normal peritoneal fibroblasts. *Fertil Steril.* 2011;96(3):758–763. e753.
- Soleimani A, Asgharzadeh F, Rahmani F, et al. Novel oral transforming growth factor- β signaling inhibitor potently inhibits postsurgical adhesion band formation. *J Cell Physiol.* 0(0).
- Dinavand P, Hassanian SM, Weiler H, Rezaie AR. Intraperitoneal administration of activated protein C prevents postsurgical adhesion band formation. *Blood.* 2015;125(8):1339–1348.
- Baig B, Hilal-Alnaqbi A, Amin A. Cancer and biotechnology: a matchup that should never slowdown. In: *Biotechnology and Production of Anti-cancer Compounds*; Springer. 2017:73–97.
- Al-Dabbagh B, Elhaty IA, Al-Hrout A, et al. Antioxidant and anticancer activities of Trigonella foenum-graecum, Cassia acutifolia and Rhazya stricta. *BMC Compl Alternative Med.* 2018;18(1):240.
- Hanna EM, Zaki N, Amin A. Detecting protein complexes in protein interaction networks modeled as gene expression biclusters. *Plos One.* 2015;10(12), e0144163.
- Mahmoud-Ghoneim D, Amin A, Corr P. MRI-based texture analysis: a potential technique to assess protectors against induced-liver fibrosis in rats. *Radiol Oncol.* 2009;43(1):30.
- Kamal H, Jafar S, Mudgil P, Murali C, Amin A, Maqsood S. Inhibitory properties of camel whey protein hydrolysates toward liver cancer cells, dipeptidyl peptidase-IV, and inflammation. *J Dairy Sci.* 2018;101(10):8711–8720.
- Hamza AA, Heeba GH, Elwey HM, Murali C, El-Awady R, Amin A. Molecular characterization of the grape seeds extract's effect against chemically induced liver cancer: in vivo and in vitro analyses. *Sci Rep.* 2018;8(1):1270.
- Amin A, Bashir A, Zaki N, McCarthy D, Ahmed S, Lotfy M. Insights into glycan biosynthesis in chemically-induced hepatocellular carcinoma in rats: a glycomic analysis. *World J Gastroenterol.* 2015;21(20):6167–6179.
- Baig B, Halim SA, Farrukh A, Greish Y, Amin A. Current status of nanomaterial-based treatment for hepatocellular carcinoma. *Biomed Pharmacother.* *Biomed Pharmacother.* 2019;116:108852.
- Al-Dabbagh B, Elhaty IA, Elhaw M, et al. Antioxidant and anticancer activities of chamomile (Matricaria recutita L.). *BMC Res Notes.* 2019;12(1):3.
- Al-Akhras MA, Aljarrah K, Al-Khateeb H, et al. Introducing Cichorium Pumilum as a potential therapeutic agent against drug-induced benign breast tumor in rats. *Electromagn Biol Med.* 2012;31(4):299–309.
- Hamza AA, Ahmed MM, Elwey HM, Amin A. Melissa officinalis protects against doxorubicin-induced cardiotoxicity in rats and potentiates its anticancer activity on MCF-7 cells. *Plos One.* 2016;11, e0167049, 11.
- Amin A, Lotfy M, Mahmoud-Ghoneim D, et al. Pancreas-protective effects of chlorella in STZ-induced diabetic animal model: insights into the mechanism. *J Diabetes Mellitus.* 2011;1:36, 03.
- Amin A. Ketoconazole-induced testicular damage in rats reduced by Gentiana extract. *Exp Toxicol Pathol : Off J Gesellschaft fur Toxikologische Pathol.* 2008;59(6):377–384.
- Al-Dabbagh B, Elhaty IA, Murali C, Al Madhoon A, Amin A. Salvadoria persica (Miswak): antioxidant and promising antiangiogenic insights. *Am J Plant Sci.* 2018;9:1228, 06.
- El-Dakhly SM, Salama AAA, Hassanin SOM, Yassen NN, Hamza AA, Amin A. Aescin and diosmin each alone or in low dose- combination ameliorate liver damage induced by carbon tetrachloride in rats. *BMC Res Notes.* 2020;13(1):259.
- Amin A, Mahmoud-Ghoneim D. Texture analysis of liver fibrosis microscopic images: a study on the effect of biomarkers. *Acta Biochim Biophys Sin.* 2011;43(3):193–203.
- Hamza AA, Mohamed MG, Lashin FM, Amin A. Dandelion prevents liver fibrosis, inflammatory response, and oxidative stress in rats. *J Basic Applied Zool.* 2020;81(1):1–13.
- Ashktorab H, Soleimani A, Singh G, et al. Saffron: the golden spice with therapeutic properties on digestive diseases. *Nutrients.* 2019;11(5).
- Hosseinzadeh H, Modaghegh MH, Saffari Z, Crocus sativus L (Saffron) extract and its active constituents (crocin and safranal) on ischemia-reperfusion in rat skeletal muscle. *Evid Base Compl Alternative Med.* 2009;6(3):343–350.
- Khorasanchi Z, Shafee M, Kermanshahi F, et al. Crocus sativus a natural food coloring and flavoring has potent anti-tumor properties. *Phytomedicine.* 2018;43:21–27.
- Naeimi M, Shafee M, Kermanshahi F, et al. Saffron (*Crocus sativus*) in the treatment of gastrointestinal cancers: current findings and potential mechanisms of action. *J Cell Biochem.* 0(0).
- Amin A, Bajbouj K, Koch A, Gandesiri M, Schneider-Stock R. Defective autophosome formation in p53-null colorectal cancer reinforces crocin-induced apoptosis. *Int J Mol Sci.* 2015;16(1):1544–1561.
- Al-Hrout A, Chaiboonchoe A, Khraiwesh B, et al. Safranal induces DNA double-strand breakage and ER-stress-mediated cell death in hepatocellular carcinoma cells. *Sci Rep.* 2018;8(1):16951.
- Vahedi M, Govil S, Kumar S, Shrivastava D, Karimi R, S Bisen P. Therapeutic applications of *Crocus sativus* L. (saffron): a review. *Nat Prod J.* 2016;6(3):162–171.
- Aghaei Z, Jafari SM, Dehnad D. Effect of different drying methods on the physicochemical properties and bioactive components of saffron powder. *Plant Foods Hum Nutr.* 2019;74(2):171–178.
- Kianbakhsh S, Ghazavi A. Immunomodulatory effects of saffron: a randomized double-blind placebo-controlled clinical trial. *Phytotherapy research. PTR.* 2011;25(12):1801–1805.
- Nam KN, Park Y-M, Jung H-J, et al. Anti-inflammatory effects of crocin and crocetin in rat brain microglial cells. *Eur J Pharmacol.* 2010;648(1–3):110–116.
- Yarjani ZM, Pourmotabbed A, Pourmotabbed T, Najafi H. Crocin has anti-inflammatory and protective effects in ischemia-reperfusion induced renal injuries. *Iranian J. Basic Med. Sci.* 2017;20(7):753.
- Hosseinzadeh H, Younesi HM. Antinociceptive and anti-inflammatory effects of *Crocus sativus* L. stigma and petal extracts in mice. *BMC Pharmacol.* 2002;2(1):7.
- Rezaei N, Avan A, Fashirzad M, et al. Crocin as a novel therapeutic agent against colitis. *Drug Chem Toxicol.* 2019;1–8.
- Bahtouee M, Fatemikia H, Pourkhalili K, et al. Saffron protection against bleomycin-induced pulmonary fibrosis in rats. *Iranian J. Toxicol.* 2017;11(6):7–12.
- Algandaby MM. Antifibrotic effects of crocin on thioacetamide-induced liver fibrosis in mice. *Saudi J Biol Sci.* 2018;25(4):747–754.
- Makhlouf H, Saksouk M, Habib J, Chahine R. Determination of antioxidant activity of saffron taken from the flower of *Crocus sativus* grown in Lebanon. *Afr J Biotechnol.* 2011;10(41):8093–8100.
- Ghadroost B, Vafei AA, Rashidy-Pour A, et al. Protective effects of saffron extract and its active constituent crocin against oxidative stress and spatial learning and memory deficits induced by chronic stress in rats. *Eur J Pharmacol.* 2011;667(1–3):222–229.
- Hemadeh O, Chilukuri S, Bonet V, Hussein S, Chaudry IH. Prevention of peritoneal adhesions by administration of sodium carboxymethyl cellulose and oral vitamin E. *Surgery.* 1993;114(5):907–910.
- Wei G, Chen X, Wang G, et al. Inhibition of cyclooxygenase-2 prevents intra-abdominal adhesions by decreasing activity of peritoneal fibroblasts. *Drug Des Dev Ther.* 2015;9:3083.
- Green CJ, Knight J, Precious S, Simpkin S. Ketamine alone and combined with diazepam or xylazine in laboratory animals: a 10 year experience. *Lab Anim.* 1981;15(2):163–170.
- Hemadeh O, Chilukuri S, Bonet V, Hussein S, Chaudry IH. Prevention of peritoneal adhesions by administration of sodium carboxymethyl cellulose and oral vitamin E. *Surgery.* 1993;114(5):907–910.
- Nair SK, Bhat IK, Aurora AL. Role of proteolytic enzyme in the prevention of postoperative intraperitoneal adhesions. *Arch Surg.* 1974;108(6):849–853.
- Bostan HB, Mehri S, Hosseinzadeh H. Toxicology effects of saffron and its constituents: a review. *Iranian J. Basic Med. Sci.* 2017;20(2):110.
- Shirali S, Zahra Bathaei S, Nakhjavani M. Effect of crocin on the insulin resistance and lipid profile of streptozotocin-induced diabetic rats. *Phytother Res.* 2013;27(7):1042–1047.
- Bathaei SZ, Hoshyar R, Miri H, Sadeghzadeh M. Anticancer effects of crocetin in both human adenocarcinoma gastric cancer cells and rat model of gastric cancer. *Biochem Cell Biol.* 2013;91(6):397–403.
- Zhou C, Bai W, Chen Q, et al. Protective effect of crocetin against burn-induced intestinal injury. *J Surg Res.* 2015;198(1):99–107.
- Srayva T, Rao GV, Kumari MG, Sagar YV, Sivarajanji Y, Sudheeranthy K. Evaluation of biosafe alternatives as xylene substitutes in hematoxylin and eosin staining procedure: a comparative pilot study. *J Oral Maxillofac Pathol: JOMFP.*

- 2018;22(1):148.
60. O'connor WN, Valle S. A combination Verhoeff's elastic and Masson's trichrome stain for routine histology. *Stain Technol.* 1982;57(4):207–210.
 61. Swolin K. Experimentell studien zur prophylaxe von intraabdominalen verwachsungen: versuche an der Ratte mit einer Emulsion aus Lipid und Prednisolon. *Acta Obstet Gynecol Scand.* 1966;45(4):473–498.
 62. Giusto G, Vercelli C, Iussich S, et al. A pectin-honey hydrogel prevents post-operative intraperitoneal adhesions in a rat model. *BMC Vet Res.* 2017;13(1):55.
 63. Roghani M, Baluchnejadmojarad T, Roghani DF. The Sesame Lignan Sesamin Attenuates Vascular Permeability in Rats with Streptozotocin-Induced Diabetes: Involvement of Oxidative Stress. 2011.
 64. Marjaneh RM, Rahmani F, Hassanian SM, et al. Phytosomal curcumin inhibits tumor growth in colitis-associated colorectal cancer. *J Cell Physiol.* 2018;233(10):6785–6798.
 65. Hashemzehi M, Behnam-Rassouli R, Hassanian SM, et al. Phytosomal-curcumin antagonizes cell growth and migration, induced by thrombin through AMP-Kinase in breast cancer. *J Cell Biochem.* 2018;119(7):5996–6007.
 66. Asgharzadeh F, Bargi R, Beheshti F, Hosseini M, Farzadnia M, Khazaei M. Thymoquinone restores liver fibrosis and improves oxidative stress status in a lipopolysaccharide-induced inflammation model in rats. *Avicenna J. Phytomed.* 2017;7(6):502.
 67. Chiorescu S, Andercou OA, Grad NO, Mironiu IA. Intraperitoneal administration of rosuvastatin prevents postoperative peritoneal adhesions by decreasing the release of tumor necrosis factor. *Cujul Med.* 2018;91(1):79.
 68. Holmdahl L, Ivarsson ML. The role of cytokines, coagulation, and fibrinolysis in peritoneal tissue repair. *European J. Surg. Acta Chirurgica.* 1999;165(11):1012–1019.
 69. Raftery AT. Regeneration of peritoneum: a fibrinolytic study. *J Anat.* 1979;129(Pt 3):659–664.
 70. Menzies D. Postoperative adhesions: their treatment and relevance in clinical practice. *Ann R Coll Surg Engl.* 1993;75(3):147–153.
 71. Gutmann J, Penzias A, Diamond M. *Adhesions in Reproductive Surgery. Reproductive Medicine and Surgery.* St. Louis: Mosby; 1995:681, 193.
 72. Bertuglia S, Colantuoni A, Intaglietta M. Effect of leukocyte adhesion and microvascular permeability on capillary perfusion during ischemia-reperfusion injury in hamster cheek pouch. *Int J Microcirc Clin Exp.* 1993;13(1):13–26.
 73. Ezberci F, Bulbuloglu E, Ciragil P, et al. Intraperitoneal tenoxicam to prevent abdominal adhesion formation in a rat peritonitis model. *Surg Today.* 2006;36(4):361–366.
 74. Hemshekhar M, Santhosh MS, Sunitha K, et al. A dietary colorant crocin mitigates arthritis and associated secondary complications by modulating cartilage deteriorating enzymes, inflammatory mediators and antioxidant status. *Biochimie.* 2012;94(12):2723–2733.
 75. Samarghandian S, Azimi-Nezhad M, Borji A, Farkhondeh T. Effect of crocin on aged rat kidney through inhibition of oxidative stress and proinflammatory state. *Phytother Res.* 2016;30(8):1345–1353.
 76. Hashemi SA, Bathaei SZ, Mohagheghi MA. Interaction of saffron carotenoids with catalase: in vitro, in vivo and molecular docking studies. *J Biomol Struct Dynam.* 2020;38(13):3916–3926.
 77. Hashemi SA, Karami M, Bathaei SZ. Saffron carotenoids change the superoxide dismutase activity in breast cancer: in vitro, in vivo and in silico studies. *Int J Biol Macromol.* 2020;158:845–853.
 78. Nasimian A, Farzaneh P, Tamanoi F, Bathaei SZ. Cytosolic and mitochondrial ROS production resulted in apoptosis induction in breast cancer cells treated with Crocin: the role of FOXO3a, PTEN and AKT signaling. *Biochem Pharmacol.* 2020;177:113999.
 79. Abedimanesh N, Motlagh B, Abedimanesh S, Bathaei SZ, Separham A, Ostadrakimani A. Effects of crocin and saffron aqueous extract on gene expression of SIRT1, AMPK, LOX1, NF- κ B, and MCP-1 in patients with coronary artery disease: a randomized placebo-controlled clinical trial. *Phytotherapy research : PTR.* 2020;34(5):1114–1122.
 80. Abedimanesh S, Bathaei SZ, Ostadrakimani A, Jafarabadi MA, Sadeghi MT. The effect of crocetin supplementation on markers of atherogenic risk in patients with coronary artery disease: a pilot, randomized, double-blind, placebo-controlled clinical trial. *Food Funct.* 2019;10(11):7461–7475.
 81. Bahtouee M, Fatemikia H, Movahed A, et al. A comparative analysis of saffron and methylprednisolone on bleomycin-induced pulmonary fibrosis in rats. *Iranian J. Toxicol.* 2018;12(3):9–13.
 82. Mehrabani M, Goudarzi M, Mehrzadi S, et al. Crocin: a protective natural antioxidant against pulmonary fibrosis induced by bleomycin. *Pharmacol Rep.* 2020:1–10.
 83. Chhimwal J, Sharma S, Kulurkar P, Patial V. Crocin attenuates CCl4-induced liver fibrosis via PPAR- γ mediated modulation of inflammation and fibrogenesis in rats. *Hum Exp Toxicol.* 2020;39(12):1639–1649.