


CORRIGENDUM: The effects and costs of home-based rehabilitation for heart failure with reduced ejection fraction: The REACH-HF multicentre randomized controlled trial

European Journal of Preventive Cardiology
2020, Vol. 27(18) NP17
© The European Society of Cardiology 2020
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/2047487320974256
journals.sagepub.com/home/cpr



The effects and costs of home-based rehabilitation for heart failure with reduced ejection fraction: The REACH-HF multicentre randomized controlled trial by Dalal HM, Taylor RS, Jolly K, et al. European Journal of Preventive Cardiology, 26(3) pp. 262–272, DOI: 10.1177/2047487318806358

In Table 3 of the above article the values in the 4 months column were printed incorrectly. The correct version of this table is below:

Table 3. Secondary objective outcomes at baseline and follow up. Data are mean (standard deviation (SD), n) unless otherwise indicated.

Outcome	Baseline		Follow up				Between-group difference	p value
			4 months		12 months			
	REACH-HF	Control	REACH-HF	Control	REACH-HF	Control		
ISWT (meters)	262.3 (153.4, 99)	239.7 (152.4, 103)	319.2 (174.5, 85)	282.8 (166.8, 85)	328.5 (181.3, 66)	294.3 (215.5, 75)	0.1 (–33.3 to 33.5)	0.995
Number of days/week with at least 10 minutes/day activity >100 milli-g*	5.8 (2.3, 99)	5.9 (1.9, 103)	5.5 (2.4, 88)	5.8 (2.1, 93)	5.6 (2.4, 78)	5.5 (2.6, 84)	0.2 (–0.4 to 0.7)	0.601
Average time/day (minutes)								
≤20 milli-g*	1104 (102, 99)	1106 (114, 103)	1107 (110, 88)	1092 (116, 93)	1092 (124, 78)	1103 (118, 84)	–7 (–29 to 15)	0.534
21–40 milli-g*	141 (35, 99)	136 (35, 103)	140 (35, 88)	138 (30, 93)	142 (39, 78)	138 (34, 84)	–1 (–9 to 8)	0.880
41–60 milli-g*	80 (25, 99)	80 (27, 103)	80 (27, 88)	82 (26, 93)	81 (30, 78)	81 (28, 84)	0 (–6 to 6)	0.901
61–80 milli-g*	45 (21, 99)	46 (21, 103)	45 (22, 88)	48 (22, 93)	48 (23, 78)	46 (22, 84)	2 (–2 to 5)	0.372
81–100 milli-g*	26 (16, 99)	27 (16, 103)	26 (16, 88)	28 (17, 93)				
>100 milli-g*	42 (34, 99)	46 (40, 103)	43 (37, 88)	51 (46, 93)				

*1000 milli-g = 1 g = 9.81 m/s², <40 milli-g is approximately equivalent to sedentary activities such as sitting, lying and ≥100 milli-g is approximately equivalent to activities undertaken at a moderate to vigorous intensity.

EQ-5D-3L = three-level version of five-dimension EuroQol scale; HADS = Hospital Anxiety and Depression Scale; ISWT = incremental shuttle walk test; milli-g = milli-gravity unit; MLHFQ = Minnesota Living with Heart Failure Questionnaire; REACH-HF = Rehabilitation EnAblement in CHronic Heart Failure.