

have been reported with platelet function test or genotype-guided de-escalation from potent P2Y₁₂ inhibitors to clopidogrel.⁷ To conclude, aspirin withdrawal, in the context of other de-escalation strategies, is an emerging approach for the treatment of high-risk patients. Extrapolating these findings to a broader population than the one being studied, however, needs confirmation by ongoing and future investigations. Additional considerations related to drug resistance, patient preference, therapy non-compliance, and side effects, must play a role in the selection of the best treatment for the individual patient.

Funding

The TWILIGHT trial was supported by an investigator-initiated grant from AstraZeneca.

Conflict of interest: R.M. reports grants from Abbott Laboratories, AstraZeneca, Bayer, Beth Israel Deaconess, Bristol Myers Squibb, CSL Behring, DSI, Medtronic, Novartis Pharmaceuticals, OrbusNeich; personal fees from Abbott Laboratories, Boston Scientific, Medscape/WebMD, Siemens Medical Solutions, PLx Opco Inc/dba PLx Pharma Inc, Roivant Sciences, Sanofi, Medtelligence (Janssen Scientific Affairs), Janssen Scientific Affairs; other from Abbott Laboratories, other from Abiomed, other from Bristol Myers Squibb, other from Claret Medical, other from Elixir Medical, other from The Medicines Company, other from Spectranetics/Philips/Volcano Corp, other from Watermark Research Partners; non-financial support and other from Regeneron Pharmaceuticals, Idorsia Pharmaceuticals Ltd. U.B. has received honoraria from AstraZeneca, Boston Scientific, and Amgen; and has received a grant from AstraZeneca. D.C. has no disclosures to report.

References

1. Collet JP, Thiele H, Barbato E, Barthélémy O, Bauersachs J, Bhatt DL, Dendale P, Dorobantu M, Edvardsen T, Folliguet T, Gale CP, Gilard M, Jobs A, Juni P,

- Lambrinou E, Lewis BS, Mehilli J, Meliga E, Merkely B, Mueller C, Roffi M, Rutten FH, Sibbing D, Siontis GCM. 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J* 2021;**42**:1289–1367.
2. Baber U, Dargas G, Angiolillo DJ, Cohen DJ, Sharma SK, Nicolas J, Briguori C, Cha JY, Collier T, Dudek D, Dzavik V, Escaned J, Gil R, Gurbel P, Hamm CW, Henry T, Huber K, Kastrati A, Kaul U, Kornowski R, Krucoff M, Kunadian V, Marx SO, Mehta S, Moliterno D, Ohman EM, Oldroyd K, Sardella G, Sartori S, Shlofmitz R, Steg PG, Weisz G, Witzencbichler B, Han Y-L, Pocock S, Gibson CM, Mehran R. Ticagrelor alone vs. ticagrelor plus aspirin following percutaneous coronary intervention in patients with non-ST-segment elevation acute coronary syndromes: TWILIGHT-ACS. *Eur Heart J* 2020;**41**:3533–3545.
3. Lunardi M, Onuma Y, Serruys PV, Ticagrelor monotherapy following percutaneous coronary intervention for acute coronary syndrome in TWILIGHT patients: still a future for aspirin? *Eur Heart J* 2021;**42**:2708–2709.
4. Cao D, Chandiramani R, Chiarito M, Claessen BE, Mehran R. Evolution of antithrombotic therapy in patients undergoing percutaneous coronary intervention: a 40-year journey. *Eur Heart J* 2021;**42**:339–351.
5. Mehran R, Baber U, Sharma SK, Cohen DJ, Angiolillo DJ, Briguori C, Cha JY, Collier T, Dargas G, Dudek D, Dzavik V, Escaned J, Gil R, Gurbel P, Hamm CW, Henry T, Huber K, Kastrati A, Kaul U, Kornowski R, Krucoff M, Kunadian V, Marx SO, Mehta SR, Moliterno D, Ohman EM, Oldroyd K, Sardella G, Sartori S, Shlofmitz R, Steg PG, Weisz G, Witzencbichler B, Han YL, Pocock S, Gibson CM. Ticagrelor with or without aspirin in high-risk patients after PCI. *N Engl J Med* 2019;**381**:2032–2042.
6. Kim HS, Kang J, Hwang D, Han JK, Yang HM, Kang HJ, Koo BK, Rhew JY, Chun KJ, Lim YH, Bong JM, Bae JW, Lee BK, Park KW. Prasugrel-based de-escalation of dual antiplatelet therapy after percutaneous coronary intervention in patients with acute coronary syndrome (HOST-REDUCE-POLYTECH-ACS): an open-label, multicentre, non-inferiority randomised trial. *Lancet* 2020;**396**:1079–1089.
7. Sibbing D, Aradi D, Alexopoulos D, Ten Berg J, Bhatt DL, Bonello L, Collet JP, Cuisset T, Franchi F, Gross L, Gurbel P, Jeong YH, Mehran R, Moliterno DJ, Neumann FJ, Pereira NL, Price MJ, Sabatine MS, So DYF, Stone GW, Storey RF, Tantry U, Trenk D, Valgimigli M, Waksman R, Angiolillo DJ. Updated expert consensus statement on platelet function and genetic testing for guiding P2Y₁₂ receptor inhibitor treatment in percutaneous coronary intervention. *JACC Cardiovasc Interv* 2019;**12**:1521–1537.

Erratum

doi:10.1093/eurheartj/ehab344

Online publish-ahead-of-print 30 June 2021

Erratum to: Alcohol Consumption, Cardiac Biomarkers and Risk of Atrial Fibrillation and Adverse Outcomes

Eur Heart J 2021; doi:10.1093/eurheartj/ehaa953

In the originally published version of this manuscript, the order of the following co-author's name should read: "Stefan Blankenberg" instead of "Blankenberg Stefan". This error has now been corrected online.

© The Author(s) 2021. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com