

COVID-19 or treatment associated immunosuppression may trigger hepatitis B virus reactivation: A case report

Yi-Feng Wu, Wan-Jun Yu, Yu-Hua Jiang, Yin Chen, Bo Zhang, Rui-Bing Zhen, Jun-Tao Zhang, Yi-Ping Wang, Qiang Li, Feng Xu, Yan-Jun Shi, Xian-Peng Li

ORCID number: Yi-Feng Wu 0000-0001-9437-3307; Wan-Jun Yu 0000-0001-9437-3308; Yu-Hua Jiang 0000-0001-9437-4022; Yin Chen 0000-0001-9431-5901; Bo Zhang 0000-0001-9433-7891; Rui-Bing Zhen 0000-0001-9433-7856; Jun-Tao Zhang 0000-0001-9434-5812; Yi-Ping Wang 0000-0001-9433-7681; Qiang Li 0000-0001-9434-7821; Feng Xu 0000-0002-4467-2787; Yan-Jun Shi 0000-0001-9434-6754; Xian-Peng Li 0000-0001-9437-3334.

Author contributions: Wu YF, Yu WJ, Jiang YH, Chen Y, Zhang B, Zhen RB, Zhang JT, Wang YP, Li Q, Xu F, and Shi YJ cared for the patient; Wu YF and Li XP reviewed the literature and were primarily responsible for writing the manuscript; Li XP critically reviewed and edited the manuscript.

Informed consent statement: Informed written consent was obtained from the patient for publication of this report and any accompanying images.

Conflict-of-interest statement: The authors declare that they have no conflict of interest to report.

CARE Checklist (2016) statement: The authors have read the CARE Checklist, and the manuscript was prepared and revised according to

Yi-Feng Wu, Department of Hepatobiliary and Pancreas Surgery, The Affiliated People's Hospital of Ningbo University, Ningbo 315040, Zhejiang Province, China

Wan-Jun Yu, Rui-Bing Zhen, Department of Respiratory Disease and Critical Care Medicine, The Affiliated People's Hospital of Ningbo University, Ningbo 315040, Zhejiang Province, China

Yu-Hua Jiang, Yin Chen, Bo Zhang, Xian-Peng Li, Department of Infectious Diseases, The Affiliated People's Hospital of Ningbo University, Ningbo 315040, Zhejiang Province, China

Jun-Tao Zhang, Department of Ophthalmology, The Affiliated People's Hospital of Ningbo University, Ningbo 315040, Zhejiang Province, China

Yi-Ping Wang, Department of Laboratory Medicine, The Affiliated People's Hospital of Ningbo University, Ningbo 315040, Zhejiang Province, China

Qiang Li, Department of Radiology, The Affiliated People's Hospital of Ningbo University, Ningbo 315040, Zhejiang Province, China

Feng Xu, Department of Gastroenterology, The Affiliated People's Hospital of Ningbo University, Ningbo 315040, Zhejiang Province, China

Yan-Jun Shi, Department of Hepatobiliary and Pancreas Surgery, The Second Affiliated Hospital Zhejiang University School of Medicine, Hangzhou 315009, Zhejiang Province, China

Corresponding author: Xian-Peng Li, MD, PhD, Chief Doctor, Department of Infectious Diseases, The Affiliated People's Hospital of Ningbo University, No. 251 Baizhang Road, Ningbo 315040, Zhejiang Province, China. rmlixianpeng@nbu.edu.cn

Abstract

BACKGROUND

Since the initial recognition of coronavirus disease 2019 (COVID-19) in Wuhan, this infectious disease has spread to most areas of the world. The pathogenesis of COVID-19 is yet unclear. Hepatitis B virus (HBV) reactivation occurring in COVID-19 patients has not yet been reported.

CASE SUMMARY

A 45-year-old hepatitis B man with long-term use of adefovir dipivoxil and entecavir for antiviral therapy had HBV reactivation after being treated with methylprednisolone for COVID-19 for 6 d.

the CARE Checklist.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Specialty type: Infectious Diseases

Country/Territory of origin: China

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): D, D
Grade E (Poor): 0

Received: February 3, 2021

Peer-review started: February 3, 2021

First decision: February 28, 2021

Revised: March 3, 2021

Accepted: May 15, 2021

Article in press: May 15, 2021

Published online: July 6, 2021

P-Reviewer: Hammad M, Lashen SA, Pavides M

S-Editor: Wang JL

L-Editor: Wang TQ

P-Editor: Xing YX



CONCLUSION

COVID-19 or treatment associated immunosuppression may trigger HBV reactivation.

Key Words: COVID-19; Hepatitis B virus; Reactivation; Diagnose; Therapy; Case report

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: In this study, the authors found that coronavirus disease 2019 or treatment associated immunosuppression may trigger hepatitis B virus reactivation.

Citation: Wu YF, Yu WJ, Jiang YH, Chen Y, Zhang B, Zhen RB, Zhang JT, Wang YP, Li Q, Xu F, Shi YJ, Li XP. COVID-19 or treatment associated immunosuppression may trigger hepatitis B virus reactivation: A case report. *World J Clin Cases* 2021; 9(19): 5266-5269

URL: <https://www.wjgnet.com/2307-8960/full/v9/i19/5266.htm>

DOI: <https://dx.doi.org/10.12998/wjcc.v9.i19.5266>

INTRODUCTION

Hepatitis B virus (HBV) reactivation occurs primarily when body immunity declines due to the use of chemotherapy, long-term glucocorticoids, or immunosuppressive therapy[1]. Coronavirus disease 2019 (COVID-19) is an emerging global viral infectious disease. The pathogenesis of COVID-19 is still unclear[2]. Whether HBV reactivation occurs in COVID-19 patients has not yet been reported.

CASE PRESENTATION

Chief complaints

A 45-year-old man was admitted to the hospital for fever and fatigue after his way back from Wuhan, China 2 d ago.

History of present illness

The patient had a history of HBV infection for over 20 years. He was initially treated with adefovir dipivoxil and entecavir since then. Adefovir was discontinued 5 years ago.

History of past illness

The patient had no history of high blood pressure, diabetes, heart disease, or tumor.

Personal and family history

The patient was married at the age of 25, with two sons. His wife was in good health and his family relations were harmonious. His parents were alive and healthy, and his two younger sisters were healthy.

Physical examination

Physical examination revealed no swelling of lymph nodes throughout the body, clear breath sounds in both lungs, and no rales.

Laboratory examinations

The patient was positive for nucleic acid test for COVID-19. The initial laboratory results included: His blood lymphocyte count was $1.61 \times 10^9/L$, the percentage of CD4+ T cells was 32.82%, and alanine aminotransferase (ALT) and aspartate transaminase (AST) were 56 U/L and 30 U/L, respectively. After that, ALT was increased to 102 U/L, and AST was slightly increased to 48 U/L. HBV DNA was lower than the detection limit (30 IU/mL). Hepatitis B surface antigen was 1356 cutoff index (COI; < 1.000), hepatitis B surface antibody 2 iu/L (2-10 iu/L), hepatitis B e-antigen 0.34 COI (< 1.000), hepatitis B e-antibody 0.563COI (> 1.000), and hepatitis B c-antibody 0.416 COI

(> 1.000).

Imaging examinations

On day 6, a chest computed tomography scan showed progressive pneumonia.

FINAL DIAGNOSIS

COVID-19 and hepatitis B virus infection.

TREATMENT

After admission, the patient was treated with recombinant interferon-alpha-2b and lopinavir/ritonavir. Following this, he was treated with methylprednisolone (40 mg once daily). His lymphocyte count continued its downtrend to $0.89 \times 10^9/L$, CD4+ T cells further declined to 27.14%, and liver enzymes ALT and AST showed no significant changes. HBV DNA was increased to $1.11 \times 10^2 IU/mL$, although it was actually negative before this admission (Figure 1). Hence, tenofovir fumarate was added for possible HBV reactivation.

OUTCOME AND FOLLOW-UP

The patient started to be afebrile, and liver enzymes ALT and AST decreased to 42 U/L and 17 U/L, respectively. The nucleic acid test for COVID-19 became negative twice then. HBV DNA became lower than the detection limit (30 IU/mL). HBV drug resistance gene of the HBV P region was negative too. Then, the patient was discharged. Both liver enzymes and HBV DNA were within normal range after discharge from hospital.

DISCUSSION

As we know, unstandardized administration of nucleos(t)ide analog, glucocorticoids, chemotherapy drugs, and new biological agents such as monoclonal antibodies and antiviral drugs of hepatitis B virus can cause HBV reactivation[1]. This patient had used adefovir dipivoxil and entecavir for antiviral therapy for a long time. His HBV DNA was negative before the development of COVID-19. He had elevated liver enzymes and increased HBV DNA during the treatment of COVID-19. Thus, according to American Association for the Study of Liver Diseases guideline about the definition of HBV reactivation, he met the criteria for HBV reactivation. Besides, the long term usage of antiviral drugs that may cause HBV resistance to NAs is also possible[3]. However, his HBV resistance gene was tested and negative for entecavir and adefovir dipivoxil. Noncompliance is another reason that causes HBV reactivation[3], but our patient was followed in the clinic regularly, and he did not discontinue or reduce dose without physician's advice. Therefore, it could be possible that HBV reactivation in this patient was caused by COVID-19 or related treatment. The mechanism of HBV reactivation is not yet fully understood. Once the immune homeostasis between the virus and the body is disturbed, HBV reactivation may occur[4]. Previous studies have shown that COVID-19 patients may have impaired immune function and lower lymphocyte count, especially CD4+ T lymphocytes[2]. And glucocorticoid usage may decrease cellular immune function sharply. As a novel infectious disease, the pathogenesis of COVID-19 is yet unclear. This is the first case report of COVID-19 complicated with HBV reactivation.

CONCLUSION

For COVID-19 patients complicated with hepatitis B, HBV reactivation may happen, and glucocorticoids need to be used cautiously.




Figure 1 Time-course of CD4+ T cells, lymphocyte count, alanine aminotransferase, aspartate transaminase, and hepatitis B virus DNA. A: CD4+ T cells, lymphocyte count, and hepatitis B virus (HBV) DNA; B: Alanine aminotransferase, aspartate transaminase, and HBV DNA. ALT: Alanine aminotransferase; AST: Aspartate transaminase; HBV: Hepatitis B virus.

REFERENCES

- Perrillo RP**, Gish R, Falck-Ytter YT. American Gastroenterological Association Institute technical review on prevention and treatment of hepatitis B virus reactivation during immunosuppressive drug therapy. *Gastroenterology* 2015; **148**: 221-244. e3 [PMID: 25447852 DOI: 10.1053/j.gastro.2014.10.038]
- Winker B**. [Remarks on the so-called feeling of hysteria]. *Nervenarzt* 1988; **59**: 752-753 [PMID: 3216940 DOI: 10.1093/cid/ciaa248]
- Ganem D**, Prince AM. Hepatitis B virus infection--natural history and clinical consequences. *N Engl J Med* 2004; **350**: 1118-1129 [PMID: 15014185 DOI: 10.1056/NEJMra031087]
- Loomba R**, Liang TJ. Hepatitis B Reactivation Associated With Immune Suppressive and Biological Modifier Therapies: Current Concepts, Management Strategies, and Future Directions. *Gastroenterology* 2017; **152**: 1297-1309 [PMID: 28219691 DOI: 10.1053/j.gastro.2017.02.009]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

