



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

## Spike-antibody waning after second dose of BNT162b2 or ChAdOx1

Vaccines based on the spike glycoprotein of SARS-CoV-2 are being rolled out globally to control transmission and limit morbidity and mortality due to COVID-19. Current evidence indicates strong immunogenicity and high short-term efficacy for BNT162b2 (Pfizer–BioNTech) and ChAdOx1 nCoV-19 (Oxford–AstraZeneca).<sup>1–3</sup> Both vaccines are delivered through a prime-boost strategy, and many countries, including the UK, have used dose intervals longer than 3–4 weeks, expecting to maximise first-dose coverage and immunogenicity. With continued high global incidence, and potential for more transmissible SARS-CoV-2 variants, data on longer-term vaccine efficacy and antibody dynamics in infection-naïve individuals are essential for clarifying the need for further booster doses.

To identify early indications of waning antibody levels to the spike protein (S-antibody) after complete two-dose vaccination, we did a cross-sectional analysis of fully vaccinated adults (aged  $\geq 18$  years) who submitted capillary blood samples for Virus Watch, a longitudinal community cohort study in England and Wales.<sup>4</sup> The study received ethical approval from the Hampstead NHS Health Research Authority Ethics Committee (20/HRA/2320). Sera were tested using Elecsys Anti-SARS-CoV-2 S and N electro-chemiluminescent immunoassays (Roche Diagnostics, Basel, Switzerland); the S assay targets total antibodies to the S1 subunit of the spike protein (range 0.4–25 000 units per mL [U/mL]), whereas the N assay targets total antibodies to the full-length nucleocapsid protein, which we took as a proxy for previous SARS-CoV-2 infection (specificity 99.8% [99.3–100]).<sup>5</sup> Serological results were linked with demographic

and clinical information collected at enrolment and with weekly self-reported vaccination status.

605 adults submitted a valid sample on June 14–15, 2021. 321 (53%) of 605 participants were women, and the median age was 63 years (IQR 58–67). Of 605 participants, 186 (31%) were categorised as clinically vulnerable, 117 (19%) as clinically extremely vulnerable, and 302 (50%) as not clinically vulnerable (additional participant characteristics and definitions of clinical vulnerability are available in the appendix). Participants contributed a single sample, taken 14–154 days after their second vaccine dose (median 42 days [IQR 30–53]). 197 (33%) of 605 samples were from BNT162b2 vaccinees and 405 (67%) samples were from ChAdOx1 vaccinees; vaccine type was missing for three (<1%) participants. The median interval between first and second doses was 77 days (IQR 70–78).

Participants with previous infection (N-seropositive; n=47) had a median S-antibody level of 9091 U/mL (IQR 3143 to 16 135), with 2.5-fold lower median levels for ChAdOx1 (median 5179 [IQR 2432.5 to 9513.5]) than BNT162b2 (median 13 025 [9091 to  $\geq 25 000$ ]). N-seronegative individuals had seven-fold lower average S-antibody levels than N-seropositive individuals (median 1257 U/mL [616 to 3526]) and six-fold lower median levels were seen after ChAdOx1 (median 864 [IQR 481 to 1395]) compared to BNT162b2 (median 5311 [3133 to 8829]) within this infection-naïve group.

We examined the distribution of S-antibody levels for confirmed N-seronegative samples 14–20 days, 21–41 days, 42–55 days, 56–69 days, and 70 days or more after second vaccination to infer the general trend in antibody levels with time, stratified by vaccine type, with p values derived from non-parametric tests for trend. We excluded two individuals with shorter dose intervals of 21–28 days

(and assumed those missing first dose date had a longer dose interval) as this has been demonstrated (in part, through preliminary data) to be less immunogenic than longer intervals for both ChAdOx1 and BNT162b2,<sup>6,7</sup> giving a total of 552 individuals included in the analysis.


A significant trend of declining S-antibody levels was seen with time for both ChAdOx1 ( $p < 0.001$ ) and BNT162b2 ( $p < 0.001$ ; figure; appendix), with levels reducing by about five-fold for ChAdOx1, and by about two-fold for BNT162b2, between 21–41 days and 70 days or more after the second dose. This trend remained consistent when results were stratified by sex, age, and clinical vulnerability (appendix). For BNT162b2, S-antibody levels reduced from a median of 7506 U/mL (IQR 4925–11 950) at 21–41 days, to 3320 U/mL (1566–4433) at 70 or more days. For ChAdOx1, S-antibody levels reduced from a median of 1201 U/mL (IQR 609–1865) at 0–20 days to 190 U/mL (67–644) at 70 or more days.

Across both vaccine types, women had higher initial S-antibody levels than men at 21–42 days after complete



Published Online  
July 15, 2021  
[https://doi.org/10.1016/S0140-6736\(21\)01642-1](https://doi.org/10.1016/S0140-6736(21)01642-1)

See Online for appendix



**Figure:** Levels of antibody against the spike glycoprotein of SARS-CoV-2 (S-antibody) at defined timepoints after second dose of vaccination (with extended dose intervals) in individuals with no previous infection, stratified by vaccine type p values derived from non-parametric tests for trend for each vaccine subgroup are given in parentheses in the key.

vaccination; also ending with higher levels at 70 days or more (appendix). Similarly, those aged 18–64 years had higher levels at 21–42 days compared to those aged 65 years and older, with correspondingly higher levels at 70 or more days (appendix).

For BNT162b2 vaccinees, some disparity was noted by clinical vulnerability status in peak antibody levels at 21–41 days, although this pattern was not observed with ChAdOx1 (appendix). At 70 days or more, the pattern of disparities was different, with higher antibody levels in vulnerable groups for BNT162b2 and the reverse for ChAdOx1. These data suggest substantial underlying heterogeneity within clinical vulnerability groupings and are also limited by small numbers in the clinically extremely vulnerable strata. However, the trend for declining S-antibody levels with time remains consistent, and the low levels in clinically vulnerable ChAdOx1 vaccinees at 70 days or more might be cause for concern.

Our data suggest waning of S-antibody levels in infection-naïve individuals over a 3–10-week period after a second dose of either ChAdOx1 or BNT162b2. These data are consistent with the decline in S-antibody and neutralising antibody levels observed after infection, although memory B-cell populations appear to be maintained.<sup>8,9</sup> As such, the clinical implications of waning antibody levels post-vaccination are not yet clear, and it remains crucial to establish S-antibody thresholds associated with protection against clinical outcomes.

Although trends were consistent after stratification by key variables that are likely to affect the immune response, there might be residual confounding due to age and dosing interval as small numbers precluded more precise strata. These findings are also limited by the cross-sectional nature of the data. This analysis should be repeated with a larger number of participants to allow better adjustment for potential confounding,

and with longitudinal follow-up of antibody dynamics in individuals over 6–12 months to establish plateau levels, or time to seroreversion.

Higher antibody levels are possibly associated with greater protection against variants that can partially evade immunity, which could explain the observed higher efficacy (partly preliminary) of BNT162b2 compared to ChAdOx1 against the Delta variant (B.1.617.2).<sup>10,11</sup> Disparity in peak antibody levels between vaccine types, and to a lesser extent between population groups, might therefore be important if antibody levels in some groups drop below (as yet undefined) thresholds of protection earlier than in others. There is, however, accumulating evidence suggesting the importance of T-cell-mediated immunity, particularly in individuals with weak or absent antibody responses,<sup>12</sup> so it is possible that T-cell responses compensate to some extent as antibody responses wane.

In the context of recent advice in support of booster vaccinations from the UK's Joint Committee on Vaccination and Immunisation,<sup>13</sup> and given the potentially rapid S-antibody decline suggested by these data, heterologous regimens, which preliminary data suggest elicit stronger antibody and T-cell responses,<sup>14,15</sup> might provide more durable immunity and greater protection against emerging variants. However, the ultimate effect of different dose intervals and various heterologous combinations on clinical outcomes remain important unanswered questions. Principally, the ethical basis for universal booster dose deployment in high-income settings should be carefully considered in the context of widening global vaccine inequities. Data on disparities in peak antibody levels and rates of decline might therefore inform targeted and equitable booster deployment.

ACH serves on the UK New and Emerging Respiratory Virus Threats Advisory Group.

All other authors declare no competing interests. The research costs for the study have been supported by the Medical Research Council Grant awarded to University College London. The study also received US\$15 000 of Facebook advertising credit to support a pilot social media recruitment campaign on Aug 18, 2020. Virus Watch received funding via the UK Government Department of Health and Social Care's Vaccine Evaluation Programme to provide monthly Thruva antibody tests to adult participants. This work was supported by the Wellcome Trust through a Wellcome Clinical Research Career Development Fellowship to RWA. Author contributions and members of the Virus Watch Collaborative are listed in the appendix.

**Madhumita Shrotri, Annalan M D Navaratnam, Vincent Nguyen, Thomas Byrne, Cyril Geismar, Ellen Fragaszy, Sarah Beale, Wing Lam Erica Fong, Parth Patel, Jana Kovar, Andrew C Hayward, \*Robert W Aldridge, on behalf of the Virus Watch Collaborative**  
r.aldridge@ucl.ac.uk

Institute of Health Informatics, University College London, London WC1E 6BT, UK

- 1 Polack FP, Thomas SJ, Kitchin N, et al. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *N Engl J Med* 2020; **383**: 2603–15.
- 2 Voysey M, Clemens SAC, Madhi SA, et al. Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *Lancet* 2021; **397**: 99–111.
- 3 Lopez Bernal J, Andrews N, Gower C, et al. Effectiveness of the Pfizer-BioNTech and Oxford-AstraZeneca vaccines on covid-19 related symptoms, hospital admissions, and mortality in older adults in England: test negative case-control study. *BMJ* 2021; **13**: 373.
- 4 Hayward A, Fragaszy E, Kovar J, et al. Risk factors, symptom reporting, healthcare-seeking behaviour and adherence to public health guidance: protocol for Virus Watch, a prospective community cohort study. *BMJ Open* 2021; **11**: e048042.
- 5 The National SARS-CoV-2 Serology Assay Evaluation Group. Performance characteristics of five immunoassays for SARS-CoV-2: a head-to-head benchmark comparison. *Lancet Infect Dis* 2020; **20**: 1390–400.
- 6 Voysey M, Costa Clemens SA, Madhi SA, et al. Single-dose administration and the influence of the timing of the booster dose on immunogenicity and efficacy of ChAdOx1 nCoV-19 (AZD1222) vaccine: a pooled analysis of four randomised trials. *Lancet* 2021; **397**: 881–91.
- 7 Parry H, Tut G, Faustini S. BNT162b2 vaccination in people over 80 years of age induces strong humoral immune responses with cross neutralisation of P.1 Brazilian variant. *SSRN* 2021; published online March 31. <https://dx.doi.org/10.2139/ssrn.3816840> (preprint).

- 8 Wheatley AK, Juno JA, Wang JJ, et al. Evolution of immune responses to SARS-CoV-2 in mild-moderate COVID-19. *Nat Commun* 2021; **12**: 1–11.
- 9 Gaebler C, Wang Z, Lorenzi JCC, et al. Evolution of antibody immunity to SARS-CoV-2. *Nature* 2021; **591**: 639–44.
- 10 Bernal JL, Andrews N, Gower C, et al. Effectiveness of COVID-19 vaccines against the B.1.617.2 variant. *medRxiv* 2021; published online May 24 (preprint).
- 11 Sheikh A, McMenamin J, Taylor B, Robertson C, on behalf of Public Health Scotland and the EAVE II Collaborators. SARS-CoV-2 Delta VOC in Scotland: demographics, risk of hospital admission, and vaccine effectiveness. *Lancet* 2021; **397**: 2461–62.
- 12 Wyllie D, Jones HE, Mulchandani R, et al. SARS-CoV-2 responsive T cell numbers and anti-Spike IgG levels are both associated with protection from COVID-19: a prospective cohort study in keyworkers. *medRxiv* 2020; published online Nov 4. <https://doi.org/10.1101/2020.11.02.20222778> (preprint).
- 13 Joint Committee on Vaccination and Immunisation, Department of Health and Social Care. JCVI interim advice on a potential coronavirus (COVID-19) booster vaccine programme for winter 2021 to 2022. June 30, 2021. <https://www.gov.uk/government/publications/jcvi-interim-advice-on-a-potential-coronavirus-covid-19-booster-vaccine-programme-for-winter-2021-to-2022> (accessed July 6, 2021).
- 14 Hillus D, Schwarz T, Tober-lau P, et al. Safety, reactogenicity, and immunogenicity of homologous and heterologous prime-boost immunisation with ChAdOx1-nCoV19 and BNT162b2: a prospective cohort study. *medRxiv* 2021; published online June 2. <https://doi.org/10.1101/2021.05.19.21257334> (preprint).
- 15 Barros-Martins J, Ramos GM, Dopfer-Jablunka A, et al. Humoral and cellular immune response against SARS-CoV-2 variants following heterologous and homologous ChAdOx1 nCoV-19 / BNT162b2 vaccination. *medRxiv* 2021; published online June 3. <https://doi.org/10.1101/2021.06.01.21258172> (preprint).

## Education and mental health: good reasons to vaccinate children

With the elevated transmissibility of circulating SARS-CoV-2 variants, vaccination coverages as high as 90% in adults might be necessary to fully relax control measures towards the end of 2021.<sup>1</sup> Such targets might be hard to reach because of vaccine hesitancy. Therefore, there is a risk that COVID-19 might cause substantial stress on health care in the winter months at the end of 2021 and

beginning of 2022. Modelling data suggest that vaccination of children and adolescents could help mitigate this risk of SARS-CoV-2 dissemination by ensuring they do not act as a reservoir.<sup>1</sup> However, since COVID-19 is mild in children,<sup>2</sup> such intervention might be ethically problematic if the population benefits come without individual benefits for children. Here, we argue that vaccinating children and adolescents is important to secure their continued access to education and protect their mental health.

In the event of a COVID-19 epidemic rebound during the winter months, we anticipate that control strategies will evolve to preferably target unvaccinated individuals, accounting for the reduced contribution of vaccinated individuals to disease spread. Living with children aged 11–17 years increases the risk of SARS-CoV-2 infection by 18–30%.<sup>3</sup> This contribution to disease spread could substantially increase once children are the only unvaccinated group, leading to a larger proportion of infections and clusters occurring in schools. Although such clusters might be tolerated if the rate of admission to hospital remains low, there is a point beyond which class closures might be reinstated. These closures would be highly detrimental to the education and wellbeing of children and adolescents who have had their schooling increasingly disrupted.<sup>4</sup> School closure can affect learning, lead to anxiety and depressive symptoms, exacerbate tensions or even intrafamily violence, and deepen social inequalities.

Early data from clinical trials suggest that the BNT162b2 mRNA COVID-19 vaccine (Pfizer-BioNTech) is safe and highly immunogenic in adolescents aged 12–15 years.<sup>5</sup> On May 10, 2021, the US Food and Drug Administration, followed by the European Medicines Agency on May 28, 2021, extended the use of this vaccine to include adolescents aged 12–15 years. Side-effects in vaccinated adolescents

should be carefully monitored at population level to make sure that rare but severe side-effects will not go unnoticed. As data from ongoing trials in children younger than 12 years become available, vaccination in younger age groups could be considered.

At a time when we all want to return to normal life, we cannot ignore the fact that children share the same aspirations. The vaccination of children against COVID-19 would be the best way to insulate them from the risk of class closures, secure their continued access to education, and protect their mental health.

We declare funding from Investissement d'Avenir programme, the Laboratoire d'Excellence Integrative Biology of Emerging Infectious Diseases programme, and the EU's Horizon 2020 research and innovation programme. The funders had no role in the writing of or decision to submit this Correspondence.

\*Simon Cauchemez, Paolo Bosetti, Cécile Tran Kiem, Violette Mourou, Angèle Consoli, Arnaud Fontanet [simon.cauchemez@pasteur.fr](mailto:simon.cauchemez@pasteur.fr)

Mathematical Modelling of Infectious Diseases Unit, Institut Pasteur, UMR2000, CNRS, Paris 75015, France (SC, PB, CTK); Collège Doctoral, Sorbonne Université, Paris, France (CTK); French COVID-19 Vaccine Strategy Committee, Sorbonne Université, Paris, France (VM); Department of Child and Adolescent Psychiatry, Sorbonne Université, Hôpital Pitié-Salpêtrière, AP-HP, Paris, France (AC); GRC 15 PSYDEV, Troubles psychiatriques et développement, Sorbonne Université, Paris, France (AC); Emerging Diseases Epidemiology Unit, Institut Pasteur, Paris, France (AF); PACRI Unit, Conservatoire National des Arts et Métiers, Paris, France (AF)

- 1 Tran Kiem C, Massonnaud C, Levy-Bruhl D, et al. Short and medium-term challenges for COVID-19 vaccination: from prioritisation to the relaxation of measures. *EClinicalMedicine* (in press).
- 2 Hoang A, Chorath K, Moreira A, et al. COVID-19 in 7780 pediatric patients: a systematic review. *EClinicalMedicine* 2020; **24**: 100433.
- 3 Galmiche S, Charmet T, Schaeffer L, et al. Exposures associated with SARS-CoV-2 infection in France: a nationwide online case-control study. *Lancet Reg Health Eur* 2021; **7**: 100148.
- 4 YoungMinds. Coronavirus: impact on young people with mental health needs. London: YoungMinds, 2021.
- 5 Frencik RW, Klein NP, Kitchin N, et al. Safety, immunogenicity, and efficacy of the BNT162b2 COVID-19 vaccine in adolescents. *New Engl J Med* 2021; published online May 27. <https://doi.org/10.1056/NEJMoa2107456>.



Europa Press News/Getty Images

Published Online  
July 14, 2021  
[https://doi.org/10.1016/S0140-6736\(21\)01453-7](https://doi.org/10.1016/S0140-6736(21)01453-7)