



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Contents lists available at ScienceDirect

# Multiple Sclerosis and Related Disorders

journal homepage: [www.elsevier.com/locate/msard](http://www.elsevier.com/locate/msard)

## Editorial

### COVID-19 vaccines and multiple sclerosis disease-modifying therapies



Much more evidence has emerged since our first editorial in April 2020 on the use of multiple sclerosis (MS) disease-modifying therapies (DMTs) during the pandemic (Giovannoni et al., 2020) and a further publication in MSARDs by Baker and colleagues on the biology underpinning the interaction between DMTs and SARs-CoV-2 (David Baker, Amor, et al., 2020). It is now clear that apart from anti-CD20 therapies (rituximab and ocrelizumab) MS DMTs do not appear to increase the risk of COVID-19 or severe COVID-19 (Maria Pia Sormani et al., 2021; Peeters et al., 2020; Salter et al., 2021). Importantly, people with MS (pwMS) treated with interferon-beta are less likely to get severe COVID-19 (Maria Pia Sormani et al., 2021), presumably due to its antiviral effects (Monk et al., 2021). The risks of COVID-19 associated with anti-CD20 therapy are relatively small with an approximate doubling of liability compared to people on other DMTs (Maria Pia Sormani et al., 2021; Peeters et al., 2020). The most important drivers of severe COVID-19 and death are age, disability and the presence of comorbidities (Maria P. Sormani et al., 2021; Louapre et al., 2020). Male sex and other social determinants of health such as deprivation and ethnicity also play an important role in outcomes from COVID-19 (Bhaskaran et al., 2021). When counselling individual patients about the doubling of the risk of getting severe COVID-19 when on an anti-CD20 therapy it should be stressed that a doubling of a low risk, particularly a low absolute risk, remains still a low absolute risk. People with MS could use one of the online risk calculators, for example the online QCovid® risk calculator developed by the University of Oxford (<https://qcovid.org/>) to assess their personal risk.

As we hopefully approach the final wave of the COVID-19 pandemic, some countries anticipate herd immunity towards the end of the summer in the northern hemisphere. The question of vaccine readiness in relation to DMTs (D. Baker et al., 2020; Reyes et al., 2021) and vaccine hesitancy concerning the long-term safety of the COVID-19 vaccines remain very topical (Reyes et al., 2021; Xiang et al., 2021; Moniz Diónísio et al., 2021).

A recent publication from Israel shows blunted antibody response to the Pfizer-BionTech mRNA-COVID-19 vaccine in ocrelizumab and fingolimod treated patients, compared to those treated with cladribine or no DMT (Achiron et al., 2021). The suboptimal ocrelizumab and fingolimod vaccine response results have been substantiated by a small French study (Bigaut et al., 2021) and in part refuted by an Italian real-life study that examined 32 patients with MS. In 10 out of 16 (62.5%) patients managed with fingolimod and 6 out of 16 (37.5%) treated with ocrelizumab there was a positive serological response after vaccination (Guerrieri et al., 2021). We suspect the discrepancy may relate to sensitivity of the assay used to detect the anti-SARS-CoV-2 spike protein antibodies.

These results are not surprising as there is good evidence that both anti-CD20 therapies (Cioc et al., 2008) and fingolimod (Han et al., 2004), a S1P modulator, disrupt germinal centre (GC) functioning in lymphoid tissue where naive B-cells are educated with the help of follicular T-helper cells. In the GCs, antibody gene rearrangement results in class switching, for example from IgM to one of the IgG subtypes, followed by affinity maturation and selection of high-affinity B-cell receptors or membrane-bound antibodies, prior to formation of memory B-cells and plasmablast clones that exit the GCs to produce high-affinity soluble IgG that can be detected in the peripheral blood (Lu and Craft, 2021).

It is important to stress that vaccine immunity is not all about antibody responses (B-cell mediated) and that T-cell responses are also important (Tejaro and Farber, 2021).

A Swiss study, that has yet to be peer-reviewed, evaluated 96 anti-CD20 treated patients and 29 immunocompetent controls. Anti-spike SARS-CoV-2 IgG antibodies were detected in 49% of patients after the second mRNA-COVID-19 vaccine dose, compared to 100% of controls. SARS-CoV2 specific interferon- $\gamma$  release, a T-cell assay, was detected in 17% of patients and 86% of healthy controls (Moor et al., 2021). Only 5% of patients, but 86% of healthy controls showed positive results in both the antibody (B cell) and T cell assays. Importantly, the time elapsed since the last dose of an anti-CD20 therapy (>7.6 months), peripheral blood B-cell reconstitution (CD19+ cells >27/ $\mu$ l) and CD4+ lymphocyte count above 653/ $\mu$ l predicted an antibody vaccine response (Moor et al., 2021). In contrast, a US group of investigators confirmed that anti-CD20 therapy significantly reduced an anti-SARS-CoV-2 spike and receptor-binding domain (RBD) specific antibody and memory B cell responses in most patients. They also demonstrated that the effect was reduced with longer duration from the last infusion of anti-CD20 treatment and extent of B cell reconstitution (Apostolidis et al., 2021). In contrast, all patients treated with anti-CD20 therapies generated antigen-specific CD4+ and CD8+ T-cell responses following vaccination with mRNA vaccines (Apostolidis et al., 2021). In their group of patients, anti-CD20 therapy tended to skew the immune response away from so-called T-follicular helper cells and augmented the induction of antigen-specific CD8+ T cells (Apostolidis et al., 2021). These conflicting results are probably due to methodological issues. It is generally agreed that a reasonable IgG antibody response to a vaccine indicates a good CD4+ T cell response, particularly a T-follicular helper cell reaction, which is required to help educate and produce antigen-specific memory B-cells and plasmablasts in GCs (Lu and Craft, 2021). In the Swiss study above, 49% of anti-CD20 treated patients made an antibody response compared to 17% with the interferon- $\gamma$  or T-cell assay. This suggests a problem with the sensitivity of their T-cell assay (Moor et al.,

<https://doi.org/10.1016/j.msard.2021.103155>

Available online 18 July 2021

2211-0348/© 2021 Published by Elsevier B.V.

2021).

It is important to note that the majority of patients with MS on anti-CD20 therapy or a S1P-modulator make an unremarkable recovery from COVID-19. This implies that innate and T-cell responses which are critical in clearing SARS-CoV-2, are functional. In comparison, B-cell and antibody responses are not vital for eliminating the primary infection (Maria Pia Sormani et al., 2021) (Hughes et al., 2021), but are likely to play an important role in secondary immune responses, particularly sterilizing immunity, i.e. preventing repeat infection in people infected in the past or infection after vaccination (Sette and Crotty, 2021).

Putting all of this in context, it seems highly likely that patients with MS placed on anti-CD20 therapy or a S1P-modulator will have blunted, but not necessarily absent, antibody responses to the COVID-19 vaccines. These patients are likely to benefit from protective or at least partially protective T-cell responses to the vaccine, in keeping with what happens when they experience naturally occurring SARS-CoV-2 infection. This immunity is unlikely to be sterilizing, particularly because the new emerging SARS-CoV-2 varieties, e.g. the delta variant, are partially immune escape strains. Despite this, this immunity may be sufficient to prevent symptomatic infection, severe disease or death from COVID-19. The message should therefore remain the same; patients on anti-CD20 therapy or a S1P modulator (fingolimod, siponimod, ozanimod, ponemod) should receive one of the licensed COVID-19 vaccines as soon as possible on the principle that during the pandemic some immunity, particularly T-cell immunity, is better than no immunity.

At this point in the pandemic we would not recommend withdrawal of anti-CD20 or S1P modulator therapy, to optimise vaccine responses. The risk of rebound disease activity from stopping and washing out a S1P modulator is particularly hazardous (Barry et al., 2019). Conversely, the longer duration of action of anti-CD20 therapies makes rebound disease activity after withdrawal less of an issue (David Baker, Pryce, et al., 2020). Despite this, we do not know whether this is necessary and if needed, what level of B-cell reconstitution is required to optimise vaccine responses. We anticipate through studies such as those highlighted by Pedotti and colleagues (Pedotti et al., 2021) in this journal, that we will find out if pwMS on these therapies, who are vaccinated against COVID-19, albeit suboptimally from an immunological perspective, have enough immunity to prevent COVID-19 in any shape or form.

We would like to conclude by addressing the recommendation for people with MS to have seasonal influenza vaccines, 5-year pneumococcal vaccines or travel vaccines as required and perhaps a seasonal/booster COVID-19 vaccine. At some point people with MS on anti-CD20 and S1P modulators, and potentially other emerging MS DMTs, such as the Bruton Tyrosine Kinase (BTK) inhibitors, must accept that they are immunocompromised and that this immunosuppression extends to blunted, but not necessarily absent, vaccine responses as well. Through accumulation of global data we will learn if and how optimization of vaccine responses may be achieved without compromising effective management of MS. It is vital that MS-treating clinicians balance the small absolute risks of suffering severe COVID-19 and potentially death and the blunted vaccine responses on anti-CD20 therapies and the blunted vaccine responses on S1P modulators against the risks of under-treating or not treating MS.

#### Declaration of Competing Interest

In the last 5 years, Gavin Giovannoni has received compensation for serving as a consultant or speaker for or has received research support from AbbVie, Aslan, Atara Bio, Biogen, BMS-Celgene, GlaxoSmithKline, GW Pharma, Janssens/Actelion, Japanese Tobacco, Jazz Pharmaceuticals, LifNano, Merck & Co, Merck KGaA/EMD Serono, Novartis, Sanofi-Genzyme, Roche/Genentech and Teva.

#### References

- Achiron, A.n.a.t, Mandel, M.a.thilda, Dreyer-Alster, S.a.pir, Harari, G.i.l, Magalashvili, Dav.id, Sonis, P.o.lina, Dolev, M.ark, et al., 2021. Humoral immune response to COVID-19 mRNA vaccine in patients with multiple sclerosis treated with high-efficacy disease-modifying therapies. *Therap. Adv. Neurol. Disord.* 14 (April), 17562864211012835.
- Apostolidis, S.A., Kakara, M., Painter, M.M., Goel, R.R., Mathew, D., Lenzi, K., Rezk, A., et al. n.d. "Altered Cellular and Humoral Immune Responses Following SARS-CoV-2 mRNA Vaccination in Patients with Multiple Sclerosis on Anti-CD20 Therapy." <https://doi.org/10.1101/2021.06.23.21259389>.
- Baker, D.a.vid, Amor, S.andra, Kang, Angr.ay S., Schmierer, K.l.a.us, Giovannoni, Gav.in, 2020. The underpinning biology relating to multiple sclerosis disease modifying treatments during the COVID-19 pandemic. *Mult. Scler. Relat. Disord.* 43 (August), 102174.
- Baker, D.a.vid, Pryce, G.areth, James, L.ouisa K., Marta, M.onica, Schmierer, K.l.a.us, 2020. The ocrelizumab phase II extension trial suggests the potential to improve the risk: benefit balance in multiple sclerosis. *Mult. Scler. Relat. Disord.* 44 (September), 102279.
- Baker, D., Roberts, C.A.K., Pryce, G., Kang, A.S., Marta, M., Reyes, S., Schmierer, K., Giovannoni, G., Amor, S., 2020. COVID-19 vaccine-readiness for anti-CD20-depleting therapy in autoimmune diseases. *Clin. Exp. Immunol.* 202 (2), 149–161.
- Barry, Br.ian, Erwin, April.A., Stevens, Je.ssica, Tornatore, C.arlo, 2019. Fingolimod rebound: a review of the clinical experience and management considerations. *Neurol. Therapy* 8 (2), 241–250.
- Bhaskaran, Kri.shnan, Bacon, Seb.astian, Evans, Ste.phen J.w, Bates, Ch.ris J., Rentsch, Christ.opher T., MacKenna, Br.ian, Tomlinson, Lau.rie, et al., 2021. Factors associated with deaths due to COVID-19 versus other causes: population-based cohort analysis of UK primary care data and linked national death registrations within the OpenSAFELY platform. *Lancet Regional Health. Europe* 6 (July), 100109.
- Bigaut, K., Kremer, L., Fleury, M., Lanotte, L., Collongues, N., de Seze, J., 2021. Impact of disease-modifying treatments on humoral response after COVID-19 vaccination: a mirror of the response after SARS-CoV-2 infection. *Rev. Neurol. (Paris)*. <https://doi.org/10.1016/j.neuro.2021.05.001>. June.
- Cioc, Adina.M., Vanderwerf, Steve.n M., Peterson, Bruce.A., Robu, Va.lentin G., Forster, Collee.n L., Pambuccian, Ste.fan E., 2008. Rituximab-induced changes in hematology tissues found at autopsy. *Am. J. Clin. Pathol.* 130 (4), 604–612.
- Giovannoni, Ga.vin, Hawkes, Ch.ris, Lechner-Scott, Jean.nette, Levy, M.ichael, Waubant, Emmanu.elle, Gold, J.ulian, 2020. The COVID-19 pandemic and the use of MS disease-modifying therapies. *Mult. Scler. Relat. Disord.* 39 (April), 102073.
- Han, Shuh.ua, Zhang, Xu.ejun, Wang, Ga.ngdou, Guan, H.ui, Garcia, Ga.briela, Li, Pi.ng, Feng, L.ili, Zheng, B.iao, 2004. FTY720 Suppresses humoral immunity by inhibiting germinal center reaction. *Blood* 104 (13), 4129–4133.
- Hughes, Rich.ard, Whitley, Lo.aise, Fitovski, Koc.ho, Schneble, H.ans-M.artin, Muros, Er.wan, Sauter, A.nette, Craveiro, Lici.nio, et al., 2021. COVID-19 in ocrelizumab-treated people with multiple sclerosis. *Mult. Scler. Relat. Disord.* 49 (April), 102725.
- Louapre, Cé.line, Collongues, N.icolas, Stankoff, B.runo, Giannesi, Cla.ire, Papeix, Caro.line, Bensa, Car.oline, Deschamps, Rom.ain, et al., 2020. Clinical characteristics and outcomes in patients with coronavirus disease 2019 and multiple sclerosis. *JAMA Neurol.* 77 (9), 1079–1088.
- Lu, Yi.si, Craft, J.oe, 2021. T Follicular regulatory cells: choreographers of productive germinal center responses. *Front. Immunol.* 12 (June), 679909.
- Moniz Dionisio, Jo.ana, Santos, Mar.iana, Rodrigues, An.a Mar.tins, Régo, A.ndré, Vítor, Jo.ana, Delgado, S.ofia, Pinheiro, Ri.ta, et al., 2021. Opinions, beliefs and knowledge of people with multiple sclerosis on COVID-19 pandemic and vaccine. *Mult. Scler. Relat. Disord.* 54 (June), 103113.
- Monk, Philli.p D., Marsden, Rich.ard J., Tear, Vict.oria J., Brookes, Jod.y, Batten, To.by N., Mankowski, M.arcin, Gabbay, Felicity J., et al., 2021. Safety and efficacy of inhaled nebulised interferon beta-1a (SNG001) for treatment of SARS-CoV-2 infection: a randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet. Respiratory Med.* 9 (2), 196–206.
- Moor, M.B., Suter-Riniker, F., Horn, M.P., Aeberli, D., Amsler, J., Möller, B., Njue, L.M., et al. n.d. "Humoral and cellular responses to mRNA vaccines against SARS-CoV2 in patients with a history of CD20-B-Cell depleting therapy." <https://doi.org/10.1101/2021.07.04.21259848>.
- Pedotti, Rosetta, Muros-Le Rouzic, Erwan, Raposo, Catarina, Schippling, Sven, Jessop, Nikki, 2021. Understanding the Impacts of COVID-19 Pandemic in People with Multiple Sclerosis Treated with Ocrelizumab. *Multiple Sclerosis and Related Disorders*.
- Reyes, Sa.úl, Cunningham, Anth.ony L., Kalincik, To.mas, Havrdová, E.va Ku.bala, Isobe, Nori.ko, Pakpoor, Juli.a, Airas, La.ura, et al., 2021. Update on the management of multiple sclerosis during the COVID-19 pandemic and post pandemic: an international consensus statement. *J. Neuroimmunol.* 357 (August), 577627.
- Salter, Am.ber, Fox, Ro.ber.t J., Newsome, Sc.ott D., Halper, Ju.ne, Li, Dav.id K.B., Kanellis, Pa.mela, Costello, Kathle.en, et al., 2021. Outcomes and risk factors associated with SARS-CoV-2 infection in a North American registry of patients with multiple sclerosis. *JAMA Neurol.* 78 (6), 699–708.
- Sette, Alessa.ndro, Crotty, Sh.ane, 2021. Adaptive immunity to SARS-CoV-2 and COVID-19. *Cell* 184 (4), 861–880.
- Guerrieri, S., Lazzarin, S., Zanetta, C., Nozzolillo, A., Filippi, M., Moiola, L., 2021. Serological Response to SARS-CoV-2 Vaccination in Multiple Sclerosis Patients Treated with Fingolimod or Ocrelizumab: An Initial Real-Life Experience. *J. Neurol.* <https://doi.org/10.1007/s00415-021-10663-x>. June.
- Peeters, LM, Parciak, T, Walton, C, Geys, L, Moreau, Y, De Brouwer, E, Raimondi, D, Pirmani, A, Kalincik, T, Edan, G, Simpson-Yap, S, De Raedt, L, Dauxais, Y, Gauthrais, C, Rodrigues, PR, McKenna, L, Lazovski, N, Hillert, J, Foursberg, L, Spelman, T, McBurney, R, Schmidt, H, Bergmann, A, Braune, S, Stahmann, A,

- Middleton, R, Salter, A, Bebo, BF, Rojas, JI, van der Walt, A, Butzkueven, H, van der Mei, I, Ivanov, R, Hellwig, K, Sciascia do Olival, G, Cohen, JA, Van Hecke, W, Dobson, R, Magyari, M, Brum, DG, Alonso, R, Nicholas, R, Bauer, J, Chertcoff, A, de Sèze, J, Louapre, C, Comi, G, Rijke, N, 2020. COVID-19 in people with multiple sclerosis: A global data sharing initiative. *Mult Scler.* 26 (10), 1157–1162. <https://doi.org/10.1177/1352458520941485>. Epub 2020 Jul 14. PMID: 32662757; PMCID: PMC7361123.
- Sormani, Maria P., Rossi, Nico, La D.e, Schiavetti, Irene, Carmisciano, Luca, Cordioli, Cinzia, Moiola, Lucia, Radaelli, Marta, et al., 2021. Disease-modifying therapies and coronavirus disease 2019 severity in multiple sclerosis. *Ann. Neurol.* 89 (4), 780–789.
- Sormani, Maria P., Salvetti, Marco, Labauge, Pierre., Schiavetti, Irene, Zephir, Helene, Carmisciano, Luca, Bensa, Caroline, et al., 2021. DMTs and Covid-19 severity in MS: a pooled analysis from Italy and France. *Ann. Clin. Transl. Neurol.* <https://doi.org/10.1002/acn3.51408>. July.
- Tejaro, John R., Farber, Donna L., 2021. COVID-19 vaccines: modes of immune activation and future challenges. *Nat. Rev. Immunol.* 21 (4), 195–197.
- Xiang, Xinran M., Hollen, Chris, Yang, Qian, Brumbach, Barbara H., Spain, Rebecca L., Wooliscroft, Lindsey, 2021. COVID-19 Vaccination willingness among people with multiple sclerosis. *Mult. Scler. J. - Exp., Transl. Clin.* 7 (2), 20552173211017159.
- Gavin Giovannoni\*, Christopher H. Hawkes, Jeannette Lechner-Scott, Michael Levy, E. Ann Yeh, David Baker\*,<sup>1</sup>  
*Blizard Institute, Barts and The London School of Medicine and Dentistry, 4 Newark Street, London E1 2AT, UK*

\* Corresponding authors.

E-mail address: [g.giovannoni@qmul.ac.uk](mailto:g.giovannoni@qmul.ac.uk) (G. Giovannoni).

---

<sup>1</sup> Guest editor