



# Antimicrobial Resistance Hidden within Multiserovar *Salmonella* Populations

Amy T. Sicheloff,<sup>a</sup> Naomi Ohta,<sup>b\*</sup> Keri N. Norman,<sup>c</sup> Guy H. Loneragan,<sup>d</sup> Bo Norby,<sup>e</sup>  H. Morgan Scott,<sup>b</sup>  Nikki W. Shariat<sup>a</sup>

<sup>a</sup>Department of Population Health, College of Veterinary Medicine, University of Georgia, Athens, Georgia, USA

<sup>b</sup>Department of Veterinary Pathobiology, College of Veterinary Medicine and Biomedical Sciences, Texas A&M University, College Station, Texas, USA

<sup>c</sup>Department of Veterinary Integrative Biosciences, College of Veterinary Medicine and Biomedical Sciences, Texas A&M University, College Station, Texas, USA

<sup>d</sup>School of Veterinary Medicine, Texas Tech University, Amarillo, Texas, USA

<sup>e</sup>Department of Large Animal Clinical Sciences, College of Veterinary Medicine, Michigan State University, East Lansing, Michigan, USA

**ABSTRACT** *Salmonella enterica* can exist in food animals as multiserovar populations, and different serovars can harbor diverse antimicrobial resistance (AMR) profiles. Conventional *Salmonella* isolation assesses AMR only in the most abundant members of a multiserovar population, which typically reflects their relative abundance in the initial sample. Therefore, AMR in underlying serovars is an undetected reservoir that can readily be expanded upon antimicrobial use. CRISPR-SeroSeq profiling demonstrated that 60% of cattle fecal samples harbored multiple serovars, including low levels of *Salmonella* serovar Reading in 11% of samples, which were not found by culture-based *Salmonella* isolation. An *in vitro* challenge revealed that *Salmonella* serovar Reading was tetracycline resistant, while more abundant serovars were susceptible. This study highlights the importance of AMR surveillance in multiserovar populations.

**KEYWORDS** *Salmonella*, tetracycline, CRISPR-SeroSeq, cattle, serovar Reading, antimicrobial resistance

*Salmonella enterica* is responsible for more than a million human salmonellosis cases each year in the United States, with 212,500 cases attributed to antimicrobial-resistant *Salmonella* (1–3). *Salmonella* surveillance and isolation rely on culture methodology that typically concludes with serotyping one or a few colonies per sample (4). These colonies usually reflect those serovars that were most abundant in the original sample or were favored by the chosen culture methods. A previous study reported that in order to detect two serovars that are in equal proportion in a population with a 95% probability, six colonies must be selected per sample (5). Since it is not feasible to routinely pick several colonies, in samples with mixed serovars of unequal relative frequencies (or where some serovars or strains are outcompeted during *Salmonella* isolation), the serovar(s) present at lower frequencies remains undetected due to effective masking by more abundant serovars. As a result, the antimicrobial resistance (AMR) profiles of these low-abundance serovars also remain unknown. Since *Salmonella* serovars can exhibit different AMR profiles (6), it is possible that multiserovar populations contribute to a more diverse AMR reservoir. Cattle can harbor multiserovar *Salmonella* populations (7, 8); however, high-resolution analysis of these populations is almost impossible to discern using conventional *Salmonella* culture methodology.

*Salmonella* clustered regularly interspaced short palindromic repeat (CRISPR) spacer content is tractable with serovar identity, and these sequences have been employed effectively for molecular serotyping (9–13). CRISPR-SeroSeq is an amplicon-based sequencing tool that uses *Salmonella* CRISPR identities to quantify the relative frequency of multiple serovars in a single sample, down to serovars comprising as little as 0.003% of the

**Citation** Sicheloff AT, Ohta N, Norman KN, Loneragan GH, Norby B, Scott HM, Shariat NW. 2021. Antimicrobial resistance hidden within multiserovar *Salmonella* populations. *Antimicrob Agents Chemother* 65:e00048-21. <https://doi.org/10.1128/AAC.00048-21>.

**Copyright** © 2021 American Society for Microbiology. All Rights Reserved.

Address correspondence to Nikki W. Shariat, [nikki.shariat@uga.edu](mailto:nikki.shariat@uga.edu).

\* Present address: Naomi Ohta, Faculty of Veterinary Medicine, Okayama University of Science, Imabari, Ehime, Japan.

**Received** 11 January 2021

**Returned for modification** 14 March 2021

**Accepted** 23 March 2021

**Accepted manuscript posted online** 29 March 2021

**Published** 18 May 2021

population (14, 15). In a complementary but targeted approach, quantitative PCR (qPCR) assays can detect a single serovar at low quantities within a mixed *Salmonella* culture (13).

In a previous study, the effects of injectable ceftiofur crystalline-free acid (CCFA) and in-feed chlortetracycline (CTC) administration on *Salmonella* in feedlot cattle were explored (16). Culturing of fecal samples identified selection of multidrug-resistant (MDR) (including resistance to tetracycline) *Salmonella* serovar Reading within 4 days of CTC treatment. Importantly, *Salmonella* serovar Reading was never detected in fecal samples collected prior to antibiotic administration. We hypothesized that *Salmonella* serovar Reading was present in the pretreatment samples at lower levels than other serovars and was not detected in the initial study because of the low resolution provided by culture-based methods. Given the profiling capabilities and high-resolution detection of underlying serovars in a sample, we hypothesized that CRISPR-SeroSeq could be used to reveal the presence of *Salmonella* serovar Reading in pretreated cattle. This hypothesis is further supported by a subsequent study that showed most resistant *Salmonella* were below the limit of quantification prior to antibiotic treatment (17). Using that study as a framework, we used CRISPR-SeroSeq analysis to retrospectively reveal a high prevalence of multiserovar *Salmonella* populations in cattle feces, including the presence of *Salmonella* serovar Reading at low levels in fecal samples collected prior to antibiotic treatment. A subsequent *in vitro* challenge showed these *Salmonella* serovar Reading bacteria were resistant to tetracycline, while other more abundant serovars were tetracycline susceptible.

Cattle fecal samples from our previous study were stored at  $-80^{\circ}\text{C}$  in glycerol. Since this was a retrospective study, *Salmonella* was reisolated for this study from the *Salmonella*-positive fecal samples that were collected on day 0 and day 20 of the original study (16) by preenrichment and subsequent selective enrichment and plating, as previously described. Total genomic DNA was isolated from 1 ml of the enriched broth culture as described previously (14). An additional 5 ml of the enrichment culture was centrifuged at 5,000 rpm for 5 min, and the bacterial pellet was resuspended in tryptic soy broth (TSB) with 20% glycerol and stored at  $-80^{\circ}\text{C}$  for the tetracycline challenge experiments. CRISPR-SeroSeq was performed using  $2\ \mu\text{l}$  of DNA as the template as described previously (14). Thirty-six samples were multiplexed on a single MiSeq (Illumina Inc., San Diego, CA) run, including one positive (*S. enterica* serovar Enteritidis) and two negative controls. A CRISPR-SeroSeq Python script that scans sequence reads and then uses BLAST to match sequence reads to a database of more than 130 serovars was used to profile serovars, and the output was written directly to a spreadsheet (14, 15). Serovars were counted if they contained multiple CRISPR spacers unique to that serovar and if the cumulative number of reads for all the spacers in that serovar constituted a relative frequency of at least 0.02% of the population. CRISPR-SeroSeq data were visualized via graphs and Sankey plots built using SankeyMATIC ([www.sankeymatic.com](http://www.sankeymatic.com)).

A  $200\text{-}\mu\text{l}$  aliquot of thawed enriched cattle fecal sample was inoculated into 10 ml LB broth and *Salmonella* were allowed to recover for 5 h at  $37^{\circ}\text{C}$ . Two 4-ml aliquots were removed from the culture, and  $16\ \mu\text{g/ml}$  tetracycline (MilliporeSigma, Burlington, MA) was added to one aliquot. Following incubation at  $37^{\circ}\text{C}$  for 19 h, the samples were subcultured 1:100 into fresh medium, maintaining tetracycline selection in the one culture, and incubated for an additional 24 h at  $37^{\circ}\text{C}$ . Genomic DNA was isolated as described above from cultures directly after the 5-h recovery incubation (but before antibiotic addition) and again after 48 h. The tetracycline challenge experiments were performed in biological triplicates on separate days with fresh aliquots of the frozen enrichment culture. Each qPCR was performed in triplicate with  $2\ \mu\text{l}$  genomic DNA as a template as described previously (13, 18) on the qTower3 platform and analyzed using qPCRsoft 4.0 software (Analytik Jena, Jena, Germany). The primer and probe sequences are shown in Table 1. The fold change in target DNA between 48-h-treated and untreated samples was calculated as the  $\log_2$  difference in threshold cycle ( $C_T$ ) values.

To establish whether cattle fecal samples from the Ohta et al. (16) study harbored multiple *Salmonella* serovars, we performed CRISPR-SeroSeq on 55 enriched fecal samples collected before CCFA and CTC were administered (day 0) and 11 enriched fecal samples collected after antibiotic treatment was completed (day 20). The samples from

**TABLE 1** qPCR primer and probe sequences used in this study


Serovar	Forward primer (5'–3')	Reverse primer (5'–3')	Probe (5'–3') <sup>a</sup>	CRISPR locus <sup>b</sup>
Give	GCGGCAGCGGTGGCTAATATA	GCGGGGAACACATGGTCTGAAA	CGGATCATGTCCATGTGCGGTTTATCCCC	CRISPR2, sp 17-18
Mbandaka	ACCGGTACGGAAATTTGTGTCAGA	GGGAACACTATCCTGC GCAATTC	CGAACTGTGGGCACGGTTTATCCCC	CRISPR1, sp 8-9
Montevideo	CCCTGGTTAATGATGGTTGTCAGCTT	CGGGGAACACCACCGGATA	CCGGGTTCTCAGTGCACC	CRISPR1, sp 34-35
Reading	GCTAACAGAAACATAGCTGATAGTTGGCG	CGGGGAACACACTGGTCTG	ACGGTCAGTCTGCAAACGGTTTATCCCC	CRISPR1, sp 31-32

<sup>a</sup>Probes were labeled with a 6-carboxyfluorescein (FAM) fluorophore and contained an IOWA-Black quencher.


<sup>b</sup>sp, spacers; refers to spacer location within the designated CRISPR array.

this study are listed in Table S1 in the supplemental material and show which cattle were positive for *Salmonella* (and the serovar) at three different time points in the original study. Day 20 samples were selected as they represented the completion of both antimicrobial treatment regimens (i.e., CCFA and CTC) and because data from Ohta et al. (16) suggested that by the end of the experiment (day 26), the effects of the antimicrobials were waning, and *Salmonella* populations were reverting back to “normal” (Table S1). Overall, 60% (44/60) fecal samples contained two or more serovars (Fig. 1). We found up to five serovars in a single sample (sample 2366, day 0), though there was no difference in the number of serovars found between day 0 and day 20 samples (two-tailed *t* test, *P* = 0.65). There was, however, a drastic difference in the serovar identity between the two sets of fecal samples: on day 0, *Salmonella* serovars Give and Mbandaka were the most frequently detected serovars, each found in 62% of samples, while in day 20 samples, *Salmonella* serovar Reading was found in all samples (100%). These data are consistent with earlier work (16) and with serotyping performed in that original study, which was not previously published (Table S1). Importantly, we detected *Salmonella* serovar Reading in six day 0 fecal samples, with a relative frequency ranging from 0.2% of the population (sample 2313) to 37% (sample 2326) (Fig. 2A). In every instance, other serovars had a greater relative frequency than *Salmonella* serovar Reading, explaining why it was not detected in our earlier work, which relied on characterizing individual colonies.

There is some discordance between *Salmonella* serovars that have the greatest relative frequency as defined by CRISPR-SeroSeq (Fig. 1) and the colonies that were serotyped in the first study (Table S1). In previous studies where we have directly compared CRISPR-SeroSeq frequencies to cultured isolates that are serotyped, we find



**FIG 1** *Salmonella* populations in feedlot cattle feces are diverse and consist of multiple serovars. CRISPR-SeroSeq identified multiple serovars in 60% and 45% of *Salmonella* enriched fecal samples from day 0 (55 total samples) and day 20 (11 total samples), respectively. Day 0 samples that contain serovar Reading (yellow) are noted with an asterisk above the bar. Each color represents a different serovar as indicated, and the colors used are the same colors used by Ohta et al. (16). At the bottom of the figure, the gray numbers show the numbers of different serovars per sample. The cattle identification numbers are shown as black numbers; each unique number corresponds to a single steer. All cattle in this study were treated with in-feed CTC.



**FIG 2** *Salmonella* serovar Reading that is present at low levels in untreated cattle is resistant to tetracycline. (A) CRISPR-SeroSeq identified six fecal samples collected before CCFA and CTC treatment that contained low levels of *Salmonella* serovar Reading. The relative *Salmonella* serovar frequency is represented by the thickness of the line in the Sankey plots, and each serovar is represented by a different color as labeled on the right and matches the color used by Ohta et al. (16). Each node on the left side of the Sankey plots represents an enriched fecal sample from a single steer; the cattle identification number is shown in bold type, and each number is unique to a single steer. The percentage values represent the percentages of *Salmonella* serovar Reading found in the samples. (B) Fecal samples enriched from cattle at day 0 were recovered from frozen glycerol stocks by culturing for 5 h in LB broth (white bars) and then cultured in the presence (black bars) or absence (diagonal bars) of 16  $\mu\text{g}/\text{ml}$  tetracycline for 48 h. TaqMan-based qPCR assays targeting CRISPR sequences unique to each serovar were used to assess relative changes in serovar abundance. The numbers above the black bars represent the fold differences in DNA between treated and untreated samples at 48 h, calculated as the  $\log_2$  of the difference in  $C_T$  values. This experiment is representative of three independent experiments performed on separate days with different aliquots of the frozen culture. Sample 2313 was not included in these experiments as we could not reproducibly detect *Salmonella* serovar Reading at the 5-h time point. We suspect this is because its relative frequency was very low (0.2%), and the necessitated repeated freezing and thawing of the glycerol cultures damaged the integrity of that sample. For sample 2392, *Salmonella* serovar Mbandaka was undetectable after 40 cycles when cultured in tetracycline.

extremely high concordance (14, 15). We expect that the discrepancies found here are due to differences that occurred as a result of reenriching a different aliquot of the same fecal sample, which may have had an unequal distribution of *Salmonella*. Nonetheless, the serovars found by CRISPR-SeroSeq were concordant with those enriched in a particular pen (Table S1).

The population analysis presented here provides evidence that *Salmonella* serovar Reading was present in a small number of cattle on day 0. Of the six cattle that were positive for *Salmonella* serovar Reading at the beginning of the study, only two were positive for *Salmonella* on day 20 in the earlier study (samples 2244 and 2366; Table S1). Given the intermittent nature of *Salmonella* shedding in cattle, this is expected. Of

these two samples, only fecal sample 2244 yielded *Salmonella* when the enrichment was repeated for this study; therefore, we have only one paired sample. CRISPR-SeroSeq provides a high resolution of *Salmonella* serovar frequency, and in this study, we were able to detect serovars contributing to as low as 0.02% of the *Salmonella* population. Therefore, the possibility that other cattle at day 0 carried *Salmonella* serovar Reading but that we did not capture it is low. Rather, the transient nature of *Salmonella* transmission within the dense feedlot pen environments and between different cattle within and between pens likely contributed to identifying *Salmonella* serovar Reading in multiple different cattle by the end of the study.

To determine whether the *Salmonella* serovar Reading detected at day 0 was in fact resistant to tetracycline, we treated the enriched day 0 cultures that contained *Salmonella* serovar Reading with or without tetracycline for 48 h and analyzed changes in *Salmonella* serovar Reading levels by qPCR. The  $C_T$  values from samples collected after a 5-h recovery were congruent with the relative serovar frequencies detected using CRISPR-SeroSeq in day 0 fecal samples (Fig. 2B). In all five tetracycline-treated cultures, there was an increase of *Salmonella* serovar Reading after 48 h in comparison to the untreated samples, with a maximum 43-fold increase in sample 2244. The smallest change was in sample 2326, which was expected as it had a higher initial relative amount of *Salmonella* serovar Reading. Consistent with an increase in *Salmonella* serovar Reading, we observed a decrease of *Salmonella* serovars Give, Mbandaka, and Montevideo (when present), confirming their susceptibility to tetracycline.

Cattle are an important *Salmonella* reservoir, and using a high-resolution amplicon sequencing approach, we have revealed that nearly two thirds of the *Salmonella*-positive cattle fecal samples analyzed contained multiple serovars. The population analysis presented here provides evidence that *Salmonella* serovar Reading was present in a small number of cattle before CTC treatment. The transient nature of *Salmonella* transmission within feedlots and the selection pressure of antibiotic administration, as observed in the first study (16), likely contributed to identifying *Salmonella* serovar Reading in multiple cattle by the end of the study, including those where *Salmonella* serovar Reading was not present at the beginning of the study. The contracted list of *Salmonella*-positive cattle on day 20 (Fig. 1 and Table S1) reflects not only the antibiotic-driven expansion of resistant *Salmonella* serovar Reading but also the elimination of susceptible *Salmonella*. Our data also suggest that in the absence of selective pressure, *Salmonella* serovar Reading is outcompeted by other serovars, including *Salmonella* serovars Mbandaka and Give. This is supported by other studies showing that MDR carriage can incur a fitness cost in *Salmonella* and other members of the *Enterobacteriaceae* family (19, 20). This is the first study to precisely reveal the composition of multiseroovar *Salmonella* populations in cattle, demonstrating how they can shift in response to antimicrobial treatment, and to use the population information to detect less abundant AMR serovars. With respect to AMR in low-abundance serovars, this study highlights the importance of a high-resolution surveillance platform that can detect multiseroovar populations. This study was designed based on the hindsight provided by our earlier work, and we strongly suspected that we would find *Salmonella* serovar Reading present in the pretreated cattle. Future applications would be performing this blind, where the presence of an AMR phenotype is unknown. Here, we expect that this approach would work in two steps. First, CRISPR-SeroSeq would be used to reveal serovar profiles in a sample. Second, enriched, mixed cultures would be treated with an antibiotic. Using the information from CRISPR-SeroSeq, serovar-specific qPCR assays could then be used to rapidly screen the antibiotic-treated aliquots to determine AMR serovars. This approach would be faster than the current alternative, which involves streaking multiple samples onto agar and then serotyping individual colonies. Additionally, further characterization of isolates would need to be conducted to identify AMR. The two-step approach suggested could be scaled up to treat aliquots of a sample with a panel of different antibiotics and could potentially be used to define the antimicrobial profile of all serovars in a population. Collectively, the work presented here underscores the importance of being able to analyze phenotypes of clinical importance, including AMR, within entire *Salmonella* serovar populations and provides a

powerful framework with which to assess the dynamics of antimicrobial resistance among bacterial populations.

## SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

**SUPPLEMENTAL FILE 1**, PDF file, 0.1 MB.

## ACKNOWLEDGMENTS

We thank David Medina and Cameron Thompson for their assistance in preparing CRISPR-SeroSeq libraries, Erin Schroeder for helping to design the qPCR assays, Javier Vinasco for curating and preparing samples and isolates from previous work, and Edgar Morales for his help in isolating *Salmonella* from fecal samples.

We are grateful for funding support from USDA-NIFA (award number 2016-69003-24615 to N.W.S. for development of CRISPR-SeroSeq and qPCR assays), USDA-NIFA-AFRI (formerly CSREES; award numbers 2008-35201-30235 and 2008-35201-04682 to B.N., G.H.L., and H.M.S. for the cattle field trial), and USDA-NIFA-NIFSI (award number 2010-51110-21083 to G.H.L. and H.M.S. for phenotypic analysis). Any recommendations, opinions, findings or conclusions expressed in this publication are those of the publishing authors and do not necessarily represent those of the U.S. Department of Agriculture.

## REFERENCES

- Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, Jones JL, Griffin PM. 2011. Foodborne illness acquired in the United States—major pathogens. *Emerg Infect Dis* 17:7–15. <https://doi.org/10.3201/eid1701.p11101>.
- Tack DM, Marder EP, Griffin PM, Cieslak PR, Dunn J, Hurd S, Scallan E, Lathrop S, Muse A, Ryan P, Smith K, Tobin-D'Angelo M, Vugia DJ, Holt KG, Wolpert BJ, Tauxe R, Geissler AL. 2019. Preliminary incidence and trends of infections with pathogens transmitted commonly through food – Foodborne Diseases Active Surveillance Network, 10 U.S. sites, 2015–2018. *MMWR Morb Mortal Wkly Rep* 68:369–373. <https://doi.org/10.15585/mmwr.mm6816a2>.
- Centers for Disease Control and Prevention. 2019. Antibiotic resistance threats in the United States, 2019. Centers for Disease Control and Prevention, Atlanta, GA.
- US Department of Agriculture. 2017. Laboratory Guidebook Notice of Change Title: Isolation and identification of *Salmonella* from meat, poultry, pasteurized egg, and siluriformes (fish) products and carcass and environmental sponges. US Department of Agriculture, Washington, DC.
- Cason J, Cox N, Buhr R, Bourassa D, Richardson L. 2011. Probability of identifying different *Salmonella* serotypes in poultry samples, abstr 262P, p 75–76. In International poultry scientific forum. Southern Poultry Science Society, Mississippi State, MS.
- Shah DH, Paul NC, Sischo WC, Crespo R, Guard J. 2017. Population dynamics and antimicrobial resistance of the most prevalent poultry-associated *Salmonella* serotypes. *Poult Sci* 96:687–702. <https://doi.org/10.3382/ps/pew342>.
- Gragg SE, Loneragan GH, Nightingale KK, Brichta-Harhay DM, Ruiz H, Elder JR, Garcia LG, Miller MF, Echeverry A, Ramirez PR, Brashears MM. 2013. Substantial within-animal diversity of *Salmonella* isolates from lymph nodes, feces, and hides of cattle at slaughter. *Appl Environ Microbiol* 79:4744–4750. <https://doi.org/10.1128/AEM.01020-13>.
- Vohra P, Bugarel M, Turner F, Loneragan GH, Hope JC, Hopkins J, Stevens MP. 2017. Quantifying the survival of multiple *Salmonella enterica* serovars *in vivo* via massively parallel whole-genome sequencing to predict zoonotic risk. *Appl Environ Microbiol* 84:e02262-17. <https://doi.org/10.1128/AEM.02262-17>.
- Touchon M, Rocha EPC. 2010. The small, slow and specialized CRISPR and anti-CRISPR of *Escherichia* and *Salmonella*. *PLoS One* 5:e11126. <https://doi.org/10.1371/journal.pone.0011126>.
- Fabre P-H, Hautier L, Dimitrov D, Douzery EJ. 2012. A glimpse on the pattern of rodent diversification: a phylogenetic approach. *BMC Evol Biol* 12:88. <https://doi.org/10.1186/1471-2148-12-88>.
- Shariat N, Timme RE, Pettengill JB, Barrangou R, Dudley EG. 2015. Characterization and evolution of *Salmonella* CRISPR-Cas systems. *Microbiology (Reading)* 161:374–386. <https://doi.org/10.1099/mic.0.000005>.
- Bugarel M, den Bakker H, Grout J, Vignaud M-L, Loneragan GH, Fach P, Brisabois A. 2018. CRISPR-based assay for the molecular identification of highly prevalent *Salmonella* serotypes. *Food Microbiol* 71:8–16. <https://doi.org/10.1016/j.fm.2017.03.016>.
- Richards AK, Hopkins BA, Shariat NW. 2020. Conserved CRISPR arrays in *Salmonella enterica* serovar Infantis can serve as qPCR targets to detect Infantis in mixed serovar populations. *Lett Appl Microbiol* 71:138–145. <https://doi.org/10.1111/lam.13296>.
- Thompson CP, Doak AN, Amirani N, Schroeder EA, Wright J, Kariyawasam S, Lamendella R, Shariat NW. 2018. High-resolution identification of multiple *Salmonella* serovars in a single sample by using CRISPR-SeroSeq. *Appl Environ Microbiol* 84:e01859-18. <https://doi.org/10.1128/AEM.01859-18>.
- Cox N, Berrang M, House S, Medina D, Cook K, Shariat N. 2019. Population analyses reveal preenrichment method and selective enrichment media affect *Salmonella* serovars detected on broiler carcasses. *J Food Prot* 82:1688–1696. <https://doi.org/10.4315/0362-028X.JFP-19-166>.
- Ohta N, Norman KN, Norby B, Lawhon SD, Vinasco J, den Bakker H, Loneragan GH, Scott HM. 2017. Population dynamics of enteric *Salmonella* in response to antimicrobial use in beef feedlot cattle. *Sci Rep* 7:14310. <https://doi.org/10.1038/s41598-017-14751-9>.
- Ohta N, Norby B, Loneragan GH, Vinasco J, den Bakker HC, Lawhon SD, Norman KN, Scott HM. 2019. Quantitative dynamics of *Salmonella* and *E. coli* in feces of feedlot cattle treated with ceftiofur and chlortetracycline. *PLoS One* 14:e0225697. <https://doi.org/10.1371/journal.pone.0225697>.
- Tracy LM, Hicks JA, Grogan KB, Nichols JA, Morningstar-Shaw BR, Shariat NW. 2020. Molecular detection of *Salmonella enterica* subsp. *arizonae* by quantitative PCR. *Avian Dis* 64:305–309. <https://doi.org/10.1637/aviandiseases-D-19-00197>.
- Bythwood TN, Soni V, Lyons K, Hurley-Bacon A, Lee MD, Hofacre C, Sanchez S, Maurer JJ. 2019. Antimicrobial resistant *Salmonella enterica* Typhimurium colonizing chickens: the impact of plasmids, genotype, bacterial communities, and antibiotic administration on resistance. *Front Sustain Food Syst* 3:20. <https://doi.org/10.3389/fsufs.2019.00020>.
- Johnson T, Singer R, Isaacson R, Danzeisen J, Lang K, Kobluk K, Rivet B, Borewicz K, Frye J, Englen M, Anderson J, Davies P. 2015. *In vivo* transmission of an IncA/C plasmid in *Escherichia coli* depends on tetracycline concentration, and acquisition of the plasmid results in a variable cost of fitness. *Appl Environ Microbiol* 81:3561–3570. <https://doi.org/10.1128/AEM.04193-14>.