

Article

Cytoplasmic Localization of RXR α Determines Outcome in Breast Cancer

Alaleh Zati zehni ^{1,†}, Falk Batz ^{1,†}, Vincent Cavailles ², Sophie Sixou ^{3,4}, Till Kaltofen ¹, Simon Keckstein ¹, Helene Hildegard Heidegger ¹, Nina Ditsch ⁵, Sven Mahner ¹, Udo Jeschke ^{1,5,*} and Theresa Vilsmaier ¹

¹ Department of Obstetrics and Gynecology, University Hospital Munich LMU, 81377 Munich, Germany; Alaleh.Zati@med.uni-muenchen.de (A.Z.z.); falk.batz@med.uni-muenchen.de (F.B.); till.kaltofen@med.uni-muenchen.de (T.K.); simon.keckstein@med.uni-muenchen.de (S.K.); helene.heidegger@med.uni-muenchen.de (H.H.H.); Sven.Mahner@med.uni-muenchen.de (S.M.); Theresa.Vilsmaier@med.uni-muenchen.de (T.V.)

² IRCM—Institut de Recherche en Cancérologie de Montpellier, INSERM U1194, Université Montpellier, Parc Euromédecine, 208 rue des Apothicaires, CEDEX 5, F-34298 Montpellier, France; vincent.cavailles@inserm.fr

³ Faculté des Sciences Pharmaceutiques, Université Paul Sabatier Toulouse III, CEDEX 09, 31062 Toulouse, France; sophie.sixou@inserm.fr

⁴ Cholesterol Metabolism and Therapeutic Innovations, Cancer Research Center of Toulouse (CRCT), UMR 1037, Université de Toulouse, CNRS, Inserm, UPS, 31037 Toulouse, France

⁵ Department of Obstetrics and Gynecology, University Hospital, 86156 Augsburg, Germany; Nina.Ditsch@uk-augsburg.de

* Correspondence: Udo.jeschke@med.uni-muenchen.de; Tel.: +49-8214-0016-5505

† These authors equally contributed to this work.



Citation: Zati zehni, A.; Batz, F.; Cavailles, V.; Sixou, S.; Kaltofen, T.; Keckstein, S.; Heidegger, H.H.; Ditsch, N.; Mahner, S.; Jeschke, U.; et al. Cytoplasmic Localization of RXR α Determines Outcome in Breast Cancer. *Cancers* **2021**, *13*, 3756. <https://doi.org/10.3390/cancers13153756>

Academic Editors: Tommaso Susini and Nicoletta Biglia

Received: 26 May 2021

Accepted: 21 July 2021

Published: 26 July 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Simple Summary: Considering the immense development of today's therapeutic approaches in oncology towards customized therapy, this study aimed to assess the prognostic value of nuclear versus cytoplasmic retinoid X receptor α (RXR α) expression in breast cancer. Our results demonstrate that RXR α expression may have different roles in tumorigenesis according to its subcellular localization. This study strengthens the need for further research on the behavior of RXR α , depending on its intracellular localization.

Abstract: The aim of this retrospective study was to assess the prognostic value of cytoplasmic versus nuclear RXR α expression in breast cancer (BC) tissue samples and to correlate the results with clinicopathological parameters. In 319 BC patients, the expression of RXR α was evaluated via immunohistochemistry. Prognosis-determining aspects were calculated through uni- and multi-variate analyses. Correlation analysis revealed a trend association with nuclear RXR α expression regarding an improved overall survival (OS) ($p = 0.078$), whereas cytoplasmic RXR α expression was significantly correlated with a poor outcomes in terms of both OS ($p = 0.038$) and disease-free survival (DFS) ($p = 0.037$). Strengthening these results, cytoplasmic RXR α was found to be an independent marker for DFS ($p = 0.023$), when adjusted to clinicopathological parameters, whereas nuclear RXR α expression was positively associated with lower TNM-staging, i.e., pT ($p = 0.01$), pN ($p = 0.029$) and pM ($p = 0.001$). Additionally, cytoplasmic RXR α expression was positively associated with a higher histopathological tumor grading ($p = 0.02$). Cytoplasmic RXR α was also found to be a negative prognosticator for Her-2neu-negative and triple-negative patients. Altogether, these findings support the hypothesis that the subcellular localization of RXR α plays an important role in carcinogenesis and the prognosis of BC. The expression of cytoplasmic RXR α is correlated with a more aggressive course of the disease, whereas nuclear RXR α expression appears to be a protective factor. These data may help to identify high-risk BC subgroups in order to find possible specific options in targeted tumor therapy.

Keywords: breast cancer; retinoid X receptor; subcellular localization; steroid hormone receptor; prognosis; overall survival; disease-free survival; immunohistochemistry

1. Introduction

Breast cancer (BC) is the world's most prevalent cancer and the most frequent cause of cancer death worldwide [1,2]. According to the World Health Organization (WHO) in 2020, 2.3 million women have been diagnosed with BC and 685,000 deaths were BC-related [3]. As a highly heterogeneous disease, BC diagnostics and treatment are complex and differ based on clinical tumor subtypes [4,5]. Options for BC therapies have advanced immensely over the past decades, offering a variety of therapeutic approaches depending on the therapy intention, i.e., in adjuvant, neoadjuvant or metastatic settings. Therapies include surgical interventions, radiation and systematic regimes such as chemotherapy and endocrine therapy [6–8]. Lately, novel therapeutic options have been introduced and implemented into international therapeutic guidelines in BC treatment, including, for instance, monoclonal antibodies targeting human epidermal growth factor receptor 2 (HER2). Therapies targeting nuclear receptors (NRs) such as the estrogen receptor (ER) and the progesterone receptor (PR) are very successful treatment options. Endocrine therapy regimes led to a decline in the BC-associated mortality rate of approximately 30%, which makes them indispensable for the treatment of hormone-receptor-positive (HR+) BC [9–11]. Furthermore, clinical trials indicate a strong correlation between the expression of “classical steroid hormone receptors”, such as ER and PR, and the progression of the disease [12–16]. Nevertheless, some tumors are resistant to those established therapeutic options, so the identification of new therapeutic targets is consequently central to research interests [17].

Currently, individual personalized BC therapy already entails NR-specific targeted therapies for both prevention and treatment [18]. NRs are activated by binding to lipophilic metabolites and mainly function as transcription factors in the nucleus [19,20]. Novel literature demonstrates that, besides the well-established NR, other receptors, including the retinoid X receptor (RXR), thyroid hormone receptors (THRs) and vitamin D receptor (VDR), play a significant role in the pathophysiology not only of BC but also of other cancer entities [21–23]. Studies on the role of NR in distinct intracellular compartments indicate a specific prognostic value, depending on the subcellular localization [24]. Czogalla et al. demonstrated a direct link between the cytoplasmic localization of VDR and impaired overall survival (OS) in ovarian cancer [25]. In the case of THR, high nuclear localization was reported to be a positive predictive factor for OS in epithelial ovarian cancer [22]. In contrast, nuclear THR has been identified to have cancer-promoting activities in BC development [24].

As a promising protective and antitumoral factor, RXRs appear to be regulatory factors with important roles in various processes during the initiation and progression of BC [26]. RXR α plays an important role in innate and adaptive immune responses by coordinating cell metabolism and the mononuclear phagocyte system [27,28]. Previous studies revealed that the activation of RXR α regulates levels of cytokines and chemokines, coordinates phagocytosis after apoptosis and attenuates antiviral immune reactions in myelocytes [27,29]. RXR α controls cell differentiation, is a crucial factor for physiological cell homeostasis and operates as a potent regulator of pathogenesis in diverse diseases, including BC [30,31]. Retinoids derived from vitamin A and co-activator molecules bind and activate RXRs, which then regulate the transcriptional activity by means of heterodimers with other nuclear receptors such as THR and VDR and translocates into the nucleus to eventually promote its transcriptional activity [31]. Since retinoids have been defined as inducers of cancer cell differentiation and cell proliferation arrest, they are key components in the tumorigenesis process and cancer cell metabolism [20,32]. After activation and heterodimerization, RXR subsequently translocate from the cytoplasm to the nucleus [33] and binds to promoters of the target genes [34]. So far, three different RXRs have been described: RXR α (NR2B1), RXR β (NR2B2) and RXR γ (NR2B3) [35].

To date, specific pathways, such as the nuclear import of RXR-VDR heterodimers, have been investigated. Yet the exact mechanisms that determine the translocation of RXR-containing heterodimers into the nucleus remain insufficiently described [36]. Zhang et al. reported that RXR α modulated important biological processes in the cytoplasm, such

as mitochondria-dependent apoptosis, inflammation and phosphatidylinositol 3-kinase (PI3K)/AKT-mediated cell survival [37]. Interestingly, new small-molecule-binding sites on the surface of RXR α have been identified recently, which are described to mediate the regulation of the nongenomic actions of RXR α by means of a class of small molecules derived from a nonsteroidal anti-inflammatory drug [37]. RXR α expression is described to be increased in miscarriage in the endometrial glands, a process linked to inflammation [38]. RXR consists of an amino-terminal domain, which plays a role in the basal transcriptional activity, a zinc finger containing a DNA binding domain, a connecting hinge region and a carboxyl-terminal region (ligand binding domain, LBD). The LBD binds the ligand, regulates interactions with transcriptional co-regulators and coordinates the formation of dimers or (in the case of RXR) of tetramers. Thus, the LBD regulates the ligand-dependent transcriptional activities [39]. In the absence of a ligand, RXRs are transcriptionally silent. After being activated by its ligand, RXRs form homotetramers, which dissociate immediately, or RXRs heterodimerize with other nuclear receptors such as VDR and initiate transcriptional activity [39,40]. Since RXR-containing oligomers play a significant role in the nucleus, it is crucial to understand the mechanisms that underlie the nuclear translocation that for now have not been investigated sufficiently. According to the Human Protein Atlas (<https://www.proteinatlas.org/ENSG00000186350-RXRA/tissue/breast>, accessed on 19 July 2021) there is an intense expression of RXR α exclusively in the nucleus of normal glandular breast tissue and 40% of these cells show a positive RXRA expression in premenopausal women, and 15% show positive expression in postmenopausal women [41].

In cell studies, Raffo et al. identified RXRs as specific mediators of proliferation inhibition and apoptosis induction in BC cell lines [42]. Furthermore, in comparison with benign breast tissue, Friedrich et al. found higher expression levels of RXR in breast cancer cells [43]. As suppressors of cell proliferation, these receptors inhibit tumor cell growth [32]. Studies on BC cell lines and in vivo approaches demonstrated a proapoptotic effect of RXR in BC cells, which is likely to harm BC growth [44–46]. Moreover, there is evidence of a cooperative interaction between RXR and ER in this antiproliferative process [47,48]. Several studies suggest that RXR might represent a marker or even a therapeutic target in BC. Furthermore, the literature indicates that high expression of RXR is an antitumoral mediator and thus a favorable prognostic factor in BC [49–51]. On the contrary, our previous study identified RXR expression as a risk factor with a significant worse DFS in patients with multifocal or multicentric BC [52].

Due to its alleged contradictory role in BC prognosis, it appeared necessary to further investigate the behavior of RXR in BC. No study so far has identified the subcellular localization of RXR α as a prognostic factor in human breast cancer specimens. New insights could potentially be promising in regard to individualized targeted BC therapy. In the present study, we define the prognostic role of cytoplasmic versus nuclear expression of RXR α in BC and correlate the results with clinicopathological criteria.

2. Materials and Methods

2.1. Patient Collective

The cohort for this study includes 319 formalin-fixed paraffin-embedded primary BC tissues that were collected from patients who underwent surgery from the year 2000 to 2002 at the Department of Gynecology and Obstetrics of the Ludwig Maximilian University in Munich, Germany.

The total collective (TC) included 319 patients (Table 1). After an observation period of up to 10 years, DFS and OS were statistically analyzed, and these follow-up data were retrieved from the Munich Cancer Registry. The Union for International Cancer Control (UICC) TNM classification was completed to assess the size of the primary tumor (pT) [53,54], lymph node involvement (pN), and distant metastasis (pM). An experienced pathologist of the LMU Department of Pathology determined the tumor grade and histological status. Tumor grade was defined according to the Bloom and Richardson grading system [55]. The hormone receptor status was determined by immunohistochemistry

on paraffin-embedded material. Cells were regarded as hormone-receptor-positive with positive staining in $\geq 10\%$ of the tumor cell nuclei. The immune-reactive scoring system (IRS) of Remmele and Stegner (IRS) was used [56].

Table 1. Patient characteristics of the total collective.

Patient Characteristics	n (%)
Age (years)	Median 59.09 \pm 13.1 Standard Deviation
Tumor foci	Unifocal 173 (54.2) Multifocal 146 (45.7)
Histology	No special type (NST) 188 (61.4) Non-NST 118 (38.5)
Estrogen Receptor	Negative 45 (18.3) Positive 201 (81.7)
Progesterone Receptor	Negative 92 (37.4) Positive 154 (62.6)
Tumor grade	G1 or G2 165 (52.2) G3 151 (47.7)
pT	pT1 197 (64.3) pT2–pT4 109 (35.6)
pN	pN0 166 (54.2) pN1–pN3 140 (45.7)
pM	pM0 239 (78.1) pM1 67 (21.8)
Nuclear RXR α	Negative 124 (42.5) Positive 168 (57.5)
Cytoplasmic RXR α	Negative 122 (40.0) Positive 183 (60.0)

2.2. Patient Treatment

As described previously [57,58], the primary surgical treatment involved either breast conservation or modified radical mastectomy. Routine axillary dissections were performed on level I and II lymph nodes, whereas level III lymph nodes were only excised in the expression of macroscopic metastatic lesions of the lower levels was observed. Single embedded lymph nodes were screened at up to three levels for the diagnosis of lymph node metastasis.

According to the guidelines of the Cancer Treatment Center of Munich, the patients in this study received chemotherapy in cases of lymph node involvement. Hormone receptor-positive postmenopausal patients received adjuvant endocrine therapy with tamoxifen (20–30 mg/day). Premenopausal woman received GnRH analogues in the later years of the follow-up period. Aromatase inhibitors were used in the case of contraindications.

However, the guidelines for surgical, radiation oncology and chemotherapy treatment options changed substantially within the observation time of the study. Therefore, the authors did not include oncological treatment details. Nevertheless, in our patient cohort, the most common type of chemotherapy consisted of six cycles of CMF, cyclophosphamide (600 mg/m² body-surface area), methotrexate (40 mg/m²) and 5-fluoruracil (600 mg/m²) every 21 days.

2.3. Immunohistochemistry

According to the earlier published and well described methods [52,59–61], the immunohistochemistry of RXR α on formalin fixed paraffin-embedded sections was assessed. Concisely, a combination of pressure cooker heating and the standard streptavidin-biotin-peroxidase complex with a mouse/rabbit-IgG-Vectastain Elite ABC kit (Vector Laboratories, Burlingame, CA, USA) were used. The staining procedure was carried out with commercially available kits with a monoclonal mouse antibody to detect RXR α expression (Perseus Proteomics Inc., Tokyo, Japan).

Therefore, the paraffin-embedded tissue sections were dewaxed in xylol for 15 min and rehydrated twice for 15 min in a solution containing 100% alcohol. Endogenous peroxidase activity was quenched by immersion in 3% hydrogen peroxide (H_2O_2) (Merck; Darmstadt, Germany) in methanol for 20 min. Again, sections were put into a 96% and then a 70% alcohol solution. After washing in phosphate-buffered saline (PBS), sections were subjected to 10 min in a pressure cooker using sodium citrate buffer with a pH of 6.0 for epitope retrieval. To create the pH of 6.0, 0.1 citric acid was diluted in 1 L distilled water (solution A) and 0.1 M sodium citrate was diluted in 1 L distilled water (solution B). The solution contained 18 mL of solution A and 82 mL of solution B, diluted with 900 mL distilled water. The steps involved in washing the sections in distilled water and PBS were then carried out. To prevent the nonspecific binding of the primary antibodies (Anti-RXR α), sections were incubated with diluted normal serum (10 mL PBS containing 150 μ L horse serum, Vector Laboratories). Afterwards, incubation of the tissue sections with the primary antibodies diluted in PBS (1:1000) was carried out for one hour at room temperature. Sections were washed twice for 2 min in PBS. Incubation with the secondary antibody binding the streptavidin-biotin-peroxidase complex (ABC-Complex), diluted in 10 mL PBS for 30 min, was performed. This was followed by repeated PBS washing steps and incubation with the ABC complex. Coloration of the substrate was achieved with the use of chromogen 3,3'-diaminobenzidine (DAB; Dako, Glostrup, Denmark) for one minute. After washing in PBS, sections were counterstained with Mayer's acidic hematoxylin for 2 min. Finally, the sections were rehydrated in an ascending alcohol series and covered with Eukitt.

Serving for negative controls were human placenta tissue sections incubated with pre-immune IgGs (supersensitive rabbit negative control, BioGenex, Fremont, CA, USA), instead of the primary antibody. As positive controls, we used human placenta samples for RXR α detection (Figure 1). Pictures were taken with a digital charged coupled device (CCD) camera system (JVC, Tokyo, Japan).




Figure 1. Immunohistochemical staining of retinoid X receptor (RXR) in human placenta tissue, serving as negative and positive control; (a) RXR α -negative control and (b) positive control. (a,b) shows 10 \times (scale bar = 200 μ m) magnification.

2.4. Staining Evaluation (Immunoreactive Score)

To quantify the specific RXR α immunoreactivity in the nuclei and the cytoplasm, meaning the distribution and intensity patterns, the well-established semi-quantitative immune-reactive scoring system (IRS) of Remmele and Stegner (IRS) [56] was used. Two independent blinded observers evaluated the intensity and distribution pattern of the staining reaction. In six circumstances ($n = 1.9\%$), the assessment of the two independent observers varied. Both observers re-evaluated these cases together, eventually agreeing upon the same result. The concordance prior to the re-evaluation was stated at 98.1%. The scoring method has previously been described and used in numerous studies conducted by our study group [52,59–61]. Analyzing the staining, a light microscope by Leitz (Immuno-

histochemistry Type 307–148.001 512 686) (Wetzlar, Germany) and a 3CCD color camera (JVC, Victor company of Japan, Japan) were used.

The IRS scoring system ranges from 0 to 12. To obtain the IR score results, the staining intensity (score 0 = no staining, score 1 = weak staining, score 2 = moderate staining, score 3 = strong staining) and the percentage of positively stained cells (0: no staining, 1: $\leq 10\%$ of the cells, 2: 11–50% of the cells, 3: 51–80% of the cells and 4: $\geq 81\%$ of the cells) were multiplied.

Nuclear and cytoplasmic stainings of RXR α were evaluated in parallel, with the separate determination of nuclear and cytoplasmic IRS. Cut-off scores for the IRS were defined as follows: tissue samples that had been assigned an IRS greater than 1, for either nuclear or cytoplasmic RXR α expression, were scored as positive.

2.5. Ethical Approval

The tissue samples used in this study were left over material after all diagnostics had been completed and were retrieved from the archive of Gynecology and Obstetrics, Ludwig Maximilian University, Munich, Germany. All patients gave their consent to participate in the study. All patient data and clinical information from the Munich Cancer Registry were fully anonymized and encoded for statistical analysis. The study was performed according to the standards set in the Declaration of Helsinki, 1975. The current study was approved by the Ethics Committee of Ludwig Maximilian University, Munich, Germany (approval number 048–08). The authors were blinded from the clinical information during the experimental analysis.

2.6. Statistical Analysis

Statistical analysis was performed with the IBM Statistical Package for the Social Sciences (IBM SPSS Statistic v26.0 Inc., Chicago, IL, USA). The gathered results were inserted into the SPSS database in the implied manner, building the TC. The Chi-squared test was used to assess the distribution of clinical-pathological variables. Correlations between findings of immunohistochemical staining were determined with Spearman's analysis. The nonparametric Kruskal–Wallis test was used to test for differences in cytoplasmic and nuclear RXR α expression regarding the set prognostic markers. OS (in years) and disease-free survival (DFS) (in years) was compared using Kaplan–Meier graphics and differences in patient survival times were tested for significance using the Chi-squared statistics of the log rank test. For multivariate analyses, the Cox regression model for survival was used and the following factors were included: pT and pN of the TNM staging system, grading, histology type, focality, and estrogen and progesterone receptors. Each parameter to be considered showed significance at the level of $p < 0.05$. The p -value and the number of patients analyzed in each group are given for each chart.

3. Results

3.1. Patient Characteristics

Table 1 presents a summary of the detailed patient characteristics from the TC. The relatively large TC, including 319 patients, and the fairly equal distribution amongst the subgroups, strengthen the statistical power of our study. The mean age (\pm STDV) of the cohort at the time of initial diagnosis was 59 ± 13.1 years. In terms of BC focality, 173 patients were diagnosed with unifocal and 146 with multifocal and/or multicentric BC. A total of 61.4% of the patient collective had histological invasive carcinoma of no special type (NST); 81.7% of BC patients were ER-positive, whereas 62.6% of patients were PR-positive; 52.2% of patients had a low-grade carcinoma (G1–G2 = 52.2%, G3 = 47.7%) and 64.3% were staged with a tumor size smaller than 2 cm (pT1: 64.3%, pT2–pT4: 35.6). Furthermore, 54.2% of all patients were staged pN0. The majority of the patient collective was staged with no present metastasis at initial diagnosis (pM0 = 78.1%, pM1 = 21.8%). A total of 12 patients showed a high cytoplasmic expression and 37 patients showed a high nuclear expression. The cut-off for high expression was IRS scores > 4 . The negligible

different total patient numbers (n) in the subgroups may be explained by the lack of a limited number of input variables, which could not be obtained due to the retrospective character of the study.

3.2. RXR α Expression Correlates with Clinicopathological Data

The distribution of tumors, with negative or positive nuclear or cytoplasmic RXR α staining, was analyzed for all tumors. Figure 2 shows immunohistochemically stained RXR α images. Positive stained tissue appeared in a brownish color (Figure 2a–c) and negative or unstained cells appeared blue (Figure 2d). Human placenta tissue was used for RXR α negative controls (Figure 1a) and positive controls (Figure 1b). Tumor staining either appeared negative or positive for both nuclear and cytoplasmic RXR α localizations (Table 1). In cases of RXR α expression in both localizations, a nucleo-cytoplasmic IRS ratio was given (Figure 1b). Correlation analyses show the results of any nuclear or cytoplasmic staining present.




Figure 2. Immunohistochemical staining of retinoid X receptor (RXR). Immunohistochemical staining of RXR α in human breast cancer samples is illustrated in (a–d): (a) positive cytoplasmic RXR α expression only, (b) positive nuclear and cytoplasmic RXR α expression with a nucleo-cytoplasmic IRS ratio of 8:4, (c) positive nuclear RXR α expression only, (d) negative nuclear and cytoplasmic RXR α expression; (a–d) show 25 \times (scale bar = 100 μ m) magnification.

RXR α expression displayed correlations to clinical and pathological data (Table 2) in univariate analysis, using the Spearman correlation coefficient. A positive correlation was observed between high cytoplasmic RXR α expression and higher histopathological tumor grading ($p = 0.029$; $Cc = 0.125$). In the nucleus, RXR α was significantly and negatively associated with tumor size (pT) ($p = 0.012$; $Cc = -0.143$).

Table 2. High cytoplasmic versus nuclear RXR α expression in correlation to clinicopathological data.

Variables	Cytoplasmic RXR α		Nuclear RXR α	
	<i>p</i>	Correlation Coefficient	<i>p</i>	Correlation Coefficient
pT	0.497	−0.039	0.012 *	−0.143
pN	0.630	0.028	0.063	−0.107
Histology	0.610	0.029	0.558	−0.034
Grading	0.029 *	0.125	0.227	−0.069

Clinicopathologic data and RXR α expression were correlated to each other in pT and grading status using Spearman's correlation analysis. Significant correlations are in bold and indicated by asterisks (* $p < 0.05$) (p : two-tailed significance).

In addition, Kruskal–Wallis analysis revealed a statistical positive association for nuclear RXR α expression in BC tissue samples with lower pN ($p = 0.029$) and lower pM ($p = 0.001$) staging cases in BC patients. No further significant association between nuclear or cytoplasmic RXR α expression and clinicopathological characteristics could be found.


3.3. Nuclear RXR α Expression Correlates with Clinicopathological Parameters Linked with Good Prognosis

Nuclear RXR α expression in BC tissue samples revealed a trend association with improved OS. The Kaplan–Meier curve visualized a positive association of OS (Figure 3) with the expression of nuclear RXR α . The log rank test yielded a p -value of 0.078 for the OS, which was not statistically significant per our definition. Nevertheless, an obvious trend in terms of a positive correlation of nuclear RXR α expression with OS can be observed. Nuclear RXR α expression showed no significant effect on the DFS ($p = 0.914$), calculated using the Kaplan–Meier curve and the log rank test. Finally, multivariate Cox regression identified nuclear RXR α as a dependent prognostic factor for OS (HR 0.733, 95%CI 0.487–1.103, $p = 0.136$) (Table 3).

Table 3. Multivariate Cox regression analysis of nuclear RXR α expression in relation to OS.

Variable	Coefficient	HR (95% CI)	<i>p</i> Value
RXRn > 0	−0.310	0.733 (0.487–1.103)	0.136
Histology	−0.003	0.997 (0.970–1.024)	0.821
Grading	0.004	1.004 (1.000–1.009)	0.056
pT	0.551	1.736 (1.500–2.008)	0.001
pN	0.015	1.015 (1.006–1.024)	0.001
Focality	−0.240	0.787 (0.599–1.033)	0.085
Estrogen Receptor	0.084	1.088 (0.643–1.841)	0.754
Progesterone Receptor	−0.082	0.921 (0.545–1.557)	0.759

Significant results are shown in bold ($p < 0.05$); HR: hazard ratio; CI: confidence interval.




Nuclear RXR α	Mean	SEM	Mean OS Time (Years)	
			95% Confidence Interval (CI) Lower border	Higher border
Negative ($n = 124$)	7.954	0.303	7.360	8.548
Positive ($n = 168$)	8.549	0.221	8.117	8.982
Overall ($n = 292$)	8.358	0.184	7.997	8.720

Figure 3. Kaplan–Meier survival analysis of nuclear RXR α positive and negative expression in relation to OS. The risk table demonstrates the mean survival time, SEM and 95% confidence interval (CI) for univariate analyses.

3.4. Cytoplasmic RXR α Is an Independent Prognostic Factor for DFS

Cytoplasmic RXR α expression in BC tissue samples was associated with impaired OS and DFS. The Kaplan–Meier curve visualized a significant negative association of the OS (Figure 4) and DFS (Figure 5) when expressing cytoplasmic RXR α . A statistically negative significant correlation was observed for the OS ($p = 0.032$) and for the DFS ($p = 0.037$), as calculated by the log rank test.

Multivariate Cox regression identified cytoplasmic RXR α as a dependent prognostic factor for OS (HR 1.603, 95%CI 0.964–2.665, $p = 0.069$) (Table 4) but an independent factor for DFS (HR 1.696, 95%CI 1.077–2.671, $p = 0.023$) (Table 5).




Cytoplasmic RXR α	Mean OS Time (Years)			
	Mean	SEM	95% Confidence Interval (CI) Lower Border	Higher Border
Negative ($n = 116$)	8.655	0.261	8.144	9.166
Positive ($n = 176$)	8.080	0.248	7.594	8.567
Overall ($n = 292$)	8.358	0.184	7.997	8.720

Figure 4. Kaplan–Meier survival analysis of cytoplasmic RXR α positive and negative expression in relation to OS. Statistical significance is shown as p -value determined using the log-rank-test ($p < 0.05$). The risk table demonstrates the mean survival time, SEM and 95% confidence interval (CI) for univariate analyses.

Table 4. Multivariate Cox regression analysis of cytoplasmic RXR α expression in relation to OS.

Variable	Coefficient	HR (95% CI)	p Value
RXR α > 0	0.472	1.603 (0.964–2.665)	0.069
Histology	−0.003	0.997 (0.973–1.023)	0.837
Grading	0.004	1.004 (0.999–1.009)	0.132
pT	0.572	1.772 (1.514–2.074)	0.001
pN	0.013	1.013 (1.003–1.023)	0.010
Focality	−0.172	0.842 (0.623–1.139)	0.265
Estrogen Receptor	0.253	1.288 (0.725–2.291)	0.253
Progesterone Receptor	−0.254	0.776 (0.437–1.377)	0.386

Significant results are shown in bold ($p < 0.05$); HR: hazard ratio; CI: confidence interval.



Cytoplasmic RXRα	Mean DFS Time (Years)			
	Mean	SEM	95% Confidence Interval (CI)	
			Lower Border	Higher Border
Negative ($n = 122$)	8.392	0.470	7.471	9.313
Positive ($n = 181$)	7.154	0.417	6.336	7.972
Overall ($n = 303$)	7.722	0.317	7.101	8.344

Figure 5. Kaplan–Meier survival analysis of cytoplasmic RXRα positive and negative expression in relation to DFS. Statistical significance is shown as p -values determined using the log-rank-test ($p < 0.05$). The risk table demonstrates the mean survival time, SEM and 95% confidence interval (CI) for univariate analyses.

Table 5. Multivariate Cox regression analysis of cytoplasmic RXRα expression in relation to DFS.

Variable	Coefficient	HR (95% CI)	p Value
RXRc > 0	0.528	1.696 (1.077–2.671)	0.023
Histology	−0.059	0.942 (0.873–1.017)	0.127
Grading	−0.001	0.999 (0.995–1.004)	0.811
pT	0.353	1.423 (1.190–1.701)	0.001
pN	0.000	1.000 (0.986–1.014)	0.996
Focality	0.040	1.041 (0.782–1.386)	0.783
Estrogen Receptor	−0.173	0.841 (0.488–1.451)	0.534
Progesterone Receptor	0.172	1.188 (0.689–2.047)	0.535

Significant results are shown in bold ($p < 0.05$); HR: hazard ratio; CI: confidence interval.


3.5. Subcellular Localisation of RXRα

Breast cancer cases were furthermore classified into four combinatorial phenotypic groups as follows: (1) cytoplasmic and nuclear RXRα expression negative ($n = 47$); (2) cytoplasmic RXRα expression positive only ($n = 83$); (3) nuclear RXRα expression positive only ($n = 75$); (4) cytoplasmic and nuclear RXRα expression positive ($n = 98$). As described

previously, cut-off scores for the IRS were defined as follows: tissue samples that had been assigned an IRS greater than 1, for either nuclear or cytoplasmic RXR α expression, were scored as positive. Tumor staining results appeared either negative or positive for both nuclear and cytoplasmic RXR α localizations. In cases of RXR α expression in both localizations, a nucleo-cytoplasmic IRS ratio was given.


In the patient cohort with a positive nuclear RXR α expression only, this group revealed a significant correlation with improved OS and DFS when compared to the other combinatorial RXR α expression cohorts.

The Kaplan–Meier curve visualized a positive correlation of OS (Figure 6) and a positive correlation of DFS (Figure 7) with the expression of nuclear RXR α only in contrast to the combinatorial RXR α expression groups. The log rank test yielded a significant correlation both for OS ($p = 0.030$) and for DFS ($p = 0.010$).



Combined RXR α Expression	Mean OS Time (Years)			
	Mean	SEM	95% Confidence Interval (CI) Lower Border	95% Confidence Interval (CI) Higher Border
Cytoplasmic and nuclear RXR α negative ($n = 47$)	8.591	0.634	7.348	9.834
Cytoplasmic RXR α positive ($n = 83$)	9.399	0.491	8.437	10.361
nuclear RXR α positive ($n = 75$)	10.389	0.388	9.629	11.149
Cytoplasmic and nuclear RXR α positive ($n = 100$)	9.286	0.429	8.445	10.128
Total ($n = 305$)	9.704	0.251	9.212	10.196

Figure 6. Kaplan–Meier survival analysis of the four combinatorial phenotypic RXR α groups and their expression in relation to OS. Statistical significance is shown as p -values determined using the log-rank-test ($p < 0.05$). The risk table demonstrates the mean survival time, SEM and 95% confidence interval (CI) for univariate analyses.



Combined RXRα Expression	Mean DFS Time (Years)			
	Mean	SEM	95% Confidence Interval (CI) Lower Border	95% Confidence Interval (CI) Higher Border
Cytoplasmic and nuclear RXRα negativ ($n = 47$)	6.837	0.763	5.343	8.332
Cytoplasmic RXRα positiv ($n = 83$)	7.902	0.599	6.727	9.077
Nuclear RXRα positiv ($n = 75$)	9.132	0.527	8.099	10.166
Cytoplasmic and nuclear RXRα positiv ($n = 98$)	6.361	0.548	5.288	7.435
Total ($n = 303$)	7.722	0.317	7.101	8.344


Figure 7. Kaplan–Meier survival analysis of the four combinatorial phenotypic RXRα groups and their expression in relation to DFS. Statistical significance is shown as p -values determined using the log-rank-test ($p < 0.05$). The risk table demonstrates the mean survival time, SEM and 95% confidence interval (CI) for univariate analyses.

3.6. Cytoplasmic RXRα Is a Negative Prognosticator for Her-2neu-Negative and Triple-Negative Patients

Breast cancer cases were furthermore divided into the known subgroups; ER, PR, Her-2neu and triple-negative cases. The prognostic value of RXRα expression in the different subcohorts was analyzed and revealed a significant correlation in Her-2neu negative and triple-negative patients and RXRα expression. No further BC subtype group displayed a prognostic association with RXRα expression.

The Her-2neu-negative patient cohort with positive cytoplasmic RXRα BC tissue expression revealed a significant correlation with impaired OS. The TC included 94 patients with a Her-2neu-negative status and 95 patients with a Her-2neu-positive status. The Kaplan–Meier curve visualized a negative correlation of the OS (Figure 8) with the expression of cytoplasmic RXRα in Her-2neu-negative patients. The log rank test yielded a significant correlation for the OS ($p = 0.035$). Her-2neu-positive patients revealed no

significant correlation in relation to survival ($p = 0.471$). Her-2neu-positive or negative patients revealed no statistical correlation in relation to nuclear RXR α -positive ($p = 0.415$) or negative ($p = 0.351$) expression and survival.




Mean OS Time (Years) in Her-2neu-Negative BC Patients

Cytoplasmic RXR α	Mean	SEM	95% Confidence Interval (CI)	
			Lower Border	Higher Border
Negative ($n = 45$)	8.969	0.369	8.245	9.692
Positive ($n = 47$)	7.883	0.443	7.015	8.752
Overall ($n = 92$)	8.406	0.296	7.825	8.986

Figure 8. Kaplan–Meier survival analysis assessing cytoplasmic RXR α -positive expression in the Her-2neu-negative patient cohort in relation to OS. Statistical significance is shown as p -values determined using the log-rank-test ($p < 0.05$). The risk table demonstrates the mean survival.

The triple-negative patient cohort with a positive cytoplasmic RXR α BC tissue expression revealed a significant correlation with worse OS. The TC included 20 patients with a triple-negative BC status. The Kaplan–Meier curve visualized a negative correlation of the OS (Figure 9) with the expression of cytoplasmic RXR α in triple-negative patients. The log rank test yielded a significant correlation for the OS ($p = 0.044$).



Mean OS Time (Years) in Triple-Negative BC Patients

Cytoplasmic RXR α	Mean	SEM	95% Confidence Interval (CI)	
			Lower Border	Higher Border
Negative ($n = 8$)	9.554	1.398	6.813	12.295
Positive ($n = 12$)	5.681	1.031	3.661	7.701
Overall ($n = 20$)	7.360	0.963	5.473	9.248

Figure 9. Kaplan–Meier survival analysis assessing cytoplasmic RXR α positive expression in the triple-negative patient cohort in relation to OS. Statistical significance is shown as p -values determined using the log-rank test ($p < 0.05$). The risk table demonstrates the mean survival time in triple-negative patients, SEM and the 95% confidence interval (CI) for univariate analyses.

4. Discussion

The aim of this study was to evaluate the prognostic impact of the subcellular expression of RXR α in a large cohort of BC tissues, and to correlate the results with clinicopathological criteria. To date, the role of RXRs in BC patients has not been sufficiently investigated [32]. This is the first study to define the prognostic role of cytoplasmic versus nuclear expression of RXR α in BC using a relatively large patient cohort that did not receive any treatment before surgery and a long-term follow-up. Results from the present study provide evidence that the expression of cytoplasmic RXR α is a significant negative prognostic marker, whereas nuclear RXR α expression appears to be a protective factor.

To understand better the molecular function of RXR α in the pathogenesis of BC, this study focused separately on nuclear versus cytoplasmic RXR α expression in BC. Our study confirmed that RXR α is expressed with a nuclear and cytoplasmic localization. Interestingly, nuclear and cytoplasmic forms of RXR α may exhibit opposite roles in mammary carcinogenesis. This is possibly due to the activation status of RXR. Inactivated cytoplasmic RXR α does not show anticancerogenic potential, in contrast to the activated nuclear form of RXR α . Furthermore, the presence of inactive cytoplasmic RXR α may also be due to the

absence of a corresponding ligand or decreased receptor response to the ligand. This could cause lower levels of activated nuclear RXR α . Indeed, considering the correlation with the cytoplasmic expression of RXR α , DFS and OS were significantly lower, whereas the nuclear expression of RXR α revealed a trend association with improved OS. Similarly, the patient cohort with positive nuclear RXR α expression only revealed a significant correlation with improved OS and DFS when compared to the other combinatorial RXR α expression cohorts. These correlations are strengthened by the fact that, in the multivariate analysis, cytoplasmic RXR α was found to be an independent prognostic marker for poorer outcomes in DFS and a dependent prognostic factor in OS. However, nuclear RXR α was found to be a dependent prognostic marker for DFS and OS.

The role of RXR α in DNA binding and transactivation is known; however, several studies indicate that RXR α also has extranuclear functions. RXR α is reported to reside in the cytoplasm at different stages of development and can migrate from the nucleus to the cytoplasm in response to differentiation, survival, apoptosis and inflammation [37,62–65]. In our study, cytoplasmic RXR α in BC tissues was negatively associated with patient survival, whereas nuclear RXR α expression appeared to be a protective factor. Numerous studies have confirmed that modification of the subcellular localization of RXR α is connected to the development of malignant diseases and inflammation processes, thus offering potential explanations for this difference at the cellular level [37]. Ghose et al. described how inflammation processes reduced the nuclear localization of RXR α in the liver and how inflammation-induced signaling can lead to fast, diverse and multiple alterations in hepatic gene expression [62]. Mey et al. confirmed RXR α translocation from the nucleus to the cytoplasm in response to endotoxin and other inflammatory mediators and described how it inhibits its transactivation function [63]. In highly malignant human breast cancer cells, an altered localization of RXR α to the splicing factor compartments was confirmed [64], whereas Zhou et al. reported that an N-terminally truncated form of RXR α produced in cancer cells resides in the cytoplasm to promote tumor cell growth [65].

To date, no studies have described the specific role of RXR α in different cell compartments in BC tissue. However, recent data on other nuclear receptors in cancer research indicate that the prognostic value of nuclear receptors depends on their subcellular localization [24]. Consistently with our findings, Ditsch et al. demonstrated that there is a direct link between the nuclear localization of THR and increased OS in epithelial ovarian cancer [22]. However, THR has been identified to show cancer-promoting activities during BC development [24]. In the case of VDR, a direct link between the cytoplasmic localization of VDR and impaired OS in ovarian cancer has been reported [25]. Furthermore, higher cytoplasmic VDR expression in colon and vulvar cancer, as well as in malignant melanoma, had an impact on tumor progression and prognosis [66–68].

In order to gain further insights on potential individualized targeted BC treatments, we evaluated the subcellular RXR α expression in correlation with clinicopathological characteristics. In our study, cytoplasmic RXR α expression was strongly positively correlated with higher histopathological tumor grading. Nuclear RXR α expression, on the other hand, was significantly positively associated with a decreased tumor size, as well as lower pathological staging for distant metastases and regional lymph node involvement. Our data suggest a specific role of the subcellular localization of RXR α . In this study, cytoplasmic RXR α was significantly correlated with a worse prognosis in BC. Furthermore, a trend towards favorable prognostic outcomes could be seen in the case of nuclear RXR α in BC tissue. Interestingly, when looking at the different BC subcohorts in our patient collective, triple-negative and Her-2neu-negative BC tissue that stained positive for cytoplasmic RXR α showed a significant decrease in OS. These results further support the hypothesis that the shuttling of RXR α out of the nucleus, or RXR α knock-out, may lead to a deterioration of survival. No studies have so far examined the prognostic value of the intracellular expression of RXR α in BC subcohorts. Previously, Joseph et al. reported that high-nuclear retinoid X receptor gamma (RXR γ) expression in ER-positive BC tissue samples was associated with a better prognostic impact [69]. This is consistent with previous reports indicating a

positive prognostic value of nuclear RXR γ in ER- and PR-positive BC [70]. Nevertheless, the mechanisms by which the intracellular localization of RXRs exert their effects in BC subtypes remain incompletely understood [32]. The more detailed investigation of the intracellular localization of the RXR protein in BC in triple-negative breast cancer is of particular interest, as this BC subtype is characterized by worse OS, DFS and increased metastatic potential compared with other major BC subtypes. The identification of reliable predictive biomarkers is fundamental in finding new therapeutic regimes.

In cancer research, RXR has been found to be a rather protective factor. Lee et al. documented impaired OS in the case of the epigenetic inactivation of RXR genes in non-small cell lung cancer [71]. Furthermore, both the activation of RXR and the higher expression of RXR in epithelial ovarian cancer have been found to foster pro-apoptotic mechanisms [72]. In BC cells and other tumor entities, RXR is able to execute responses such as cell proliferation, cellular differentiation and programmed cell death [45,69]. Consequently, RXR is a potential therapeutic target in breast cancer cell lines [44,45]. Earlier research indicates the high expression of RXR to be an antitumoral mediator and thus a favorable prognostic factor in BC [49–51]. Consistently, cell studies on BC cells demonstrated upregulated apoptosis in BCL2-positive human cancer cells after the activation of RXR [44].

Furthermore, *in vivo* studies on transgenic mice demonstrated that RXR ligands decreased vascularization in BC [73]. Additionally, Wu et al. indicated that RXR-selective retinoids in BC tumors in transgenic mice prevented carcinogenesis and that receptor-selective retinoids may be an option in the molecular-based chemoprevention of BC [74]. For instance, LGD1069, an RXR-selective retinoid, was described to be a promising therapeutic agent. After the binding and activation of RXR, LGD1069 leads to a downregulation of cyclooxygenase-2 expression and induces a temporary G1 cell cycle arrest in human breast cells [75,76]. In relation to hereditary breast cancer, RXR has been described to be overexpressed in BRCA1-mutated BC cells. Consequently, RXR may potentially serve as a marker or even a therapeutic target in hereditary breast cancer [49].

In contrast to previous findings, which attribute a protective and antineoplastic function to RXR, our previous study identified RXR expression to be a risk factor, correlated with a significantly worse DFS in patients with multifocal or multicentric BC [52]. To explain the tumorigenic function of RXR in multifocal or multicentric BC, we suggest an increased interaction between RXR and other NRs such as VDR with elevated levels of heterodimers. Ditsch et al. previously identified RXR, THR and VDR to form functional homodimers and heterodimers with many other NRs in human BC cell lines [31]. Depending on their activation status and possibly their corresponding NRs, specific responses such as growth arrest and apoptosis may be induced. Thus, we assume a protective role of these NRs in breast cancer development [45]. Contrary to this assumption, a multicenter phase II study of oral bexarotene, a third-generation retinoid which is indicated for the treatment of cutaneous T-cell lymphoma, did not show an inhibitory effect on cancer growth. This therapeutic approach was ineffective for patients with metastatic BC [77,78]. Due to the abovementioned contradictory data on the role of RXR in BC development and progression, further scientific research on RXR is needed.

There are some factors limiting our study. First, it was a retrospective analysis based on a single dataset. Second, the sample size was comparatively low and may thus be possibly insufficient to elucidate all the heterogenous entities in breast cancer. Furthermore, specific information on possible toxic environmental aspects or a history of endocrine therapy and other patient characteristics could enrich the investigation of how additional factors interact with RXR α . Hormone treatment with estrogen might influence breast cancer growth, since a cooperative antiproliferative interaction between RXR and ER has been described [47,48]. In addition, the guidelines for surgical, radiation oncology and chemotherapy treatment options changed substantially within the observation time of the study. Therefore, the authors did not include oncological treatment details.

However, aside from these limitations, our data demonstrate that the RXR α pathway could represent a promising therapeutic target in BC. This study might provide an impetus

to further investigate the crosstalk between potential NR ligands and the RXR α pathway in regard to its therapeutic potential in BC.

In summary, our findings demonstrate the complexity of the links between nuclear and cytoplasmic RXR expression and their impact on patient outcome. Our results emphasize the need for more detailed investigations of the intracellular localization of the RXR protein in mammary carcinoma in order to understand its biomolecular function and role as a possible biomarker in BC diagnostics.

5. Conclusions

In this study, we investigated the prognostic value of the nuclear localization of the RXR α receptor versus its cytoplasmic expression in human BC specimens. Furthermore, we investigated the correlation between clinicopathological criteria as well as patient outcomes and the subcellular localization of RXR α . This is the first retrospective cohort study to define the prognostic role of cytoplasmic versus nuclear expression of RXR α in sporadic mammary cancer using a large clinical patient cohort and a long-term follow-up. RXR α expression was observed to play an incongruous role for BC prognosis depending on its intracellular localization: RXR α expressed in the cytoplasm of BC tissues was negatively associated with prognostic factors, such as patient survival, and the opposite result was observed in nucleus-localized RXR α .

In summary, nuclear receptors, such as RXR α , and possible targeted treatments should become the subjects of future research. Our findings confirm the need of further examinations on the subcellular expression of RXR α and other members of the NR family. Further investigations studying the biomolecular role of RXR α in BC would be of major interest.

Author Contributions: U.J., V.C. and N.D. conceived and designed the project. T.V., A.Z.z. and F.B. wrote the paper. A.Z.z. and T.V. carried out the statistical evaluation. A.Z.z. and T.V. carried out the method. T.V. and U.J. provided the concept, substantially contributed to the manuscript and supervised the research. V.C., S.S., H.H.H., S.K. and T.K. revised the manuscript for critical content and helped with statistical evaluation. Funding acquisition, S.M. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by internal departmental sources and has no funding grant number and name.

Institutional Review Board Statement: The tissue samples used in this study were left over material after all diagnostics had been completed and were retrieved from the archive of Gynecology and Obstetrics, Ludwig Maximilian University, Munich, Germany. All patients gave their consent to participate in the study. All patient data were fully anonymized; the study was performed according to the standards set out in the Declaration of Helsinki, 1975. The current study was approved by the Ethics Committee of Ludwig Maximilian University, Munich, Germany (approval number 048–08) 05 February 2018. The authors were blinded from the clinical information during the experimental analysis.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The data presented in this study are available on request from the corresponding author. The data are not publicly available due to ethical issues.

Conflicts of Interest: Sven Mahner: research support, advisory board, honoraria and travel expenses from AbbVie, AstraZeneca, Clovis, Eisai, GlaxoSmithKline, Medac, MSD, Novartis, Olympus, PharmaMar, Roche, Sensor Kinesis, Teva and Tesaro. All other authors declare no conflict of interest.

Abbreviations

BC	Breast cancer
DFS	Disease-free survival
NR	Nuclear receptor
OS	Overall survival
IRS	Immunoreactive score
ER	Estrogen receptor
PBS	Phosphate-buffered saline
PR	Progesterone receptor
RXR	Retinoid X receptor
RXR α	Retinoid X receptor alpha
RXR γ	Retinoid X receptor gamma
TC	Total collective
VDR	Vitamin D receptor
THR	Thyroid hormone receptor

References

- Bray, F.; Ferlay, J.; Soerjomataram, I.; Siegel, R.L.; Torre, L.A.; Jemal, A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J. Clin.* **2018**, *68*, 394–424. [[CrossRef](#)]
- Harbeck, N.; Gnant, M. Breast Cancer. *Lancet* **2017**, *389*, 1134–1150. [[CrossRef](#)]
- WHO. *Breast Cancer*; World Health Organization: Geneva, Switzerland, 2021.
- Tao, Z.; Shi, A.; Lu, C.; Song, T.; Zhang, Z.; Zhao, J. Breast Cancer: Epidemiology and Etiology. *Cell Biophys.* **2015**, *72*, 333–338. [[CrossRef](#)] [[PubMed](#)]
- Harbeck, N.; Penault-Llorca, F.; Cortes, J.; Gnant, M.; Houssami, N.; Poortmans, P.; Ruddy, K.; Tsang, J.; Cardoso, F. Breast Cancer. *Nat. Rev. Dis. Primers* **2019**, *5*, 66. [[CrossRef](#)] [[PubMed](#)]
- Fisher, B.; Costantino, J.P.; Wickerham, D.L.; Cecchini, R.; Cronin, W.M.; Robidoux, A.; Bevers, T.B.; Kavanah, M.T.; Atkins, J.N.; Margolese, R.G.; et al. Tamoxifen for the Prevention of Breast Cancer: Current Status of the National Surgical Adjuvant Breast and Bowel Project P-1 Study. *J. Natl. Cancer Inst.* **2005**, *97*, 1652–1662. [[CrossRef](#)] [[PubMed](#)]
- Cauley, J.A.; Norton, L.; Lippman, M.E.; Eckert, S.; Krueger, K.A.; Purdie, D.W.; Farrerons, J.; Karasik, A.; Mellstrom, D.; Ng, K.W.; et al. Continued Breast Cancer Risk Reduction in Postmenopausal Women Treated with Raloxifene: 4-Year Results from the MORE Trial. *Breast Cancer Res. Treat.* **2001**, *65*, 125–134. [[CrossRef](#)]
- Goss, P.E.; Ingle, J.N.; Alés-Martínez, J.E.; Cheung, A.M.; Chlebowski, R.T.; Wactawski-Wende, J.; McTiernan, A.; Robbins, J.; Johnson, K.C.; Martin, L.W.; et al. Exemestane for Breast-Cancer Prevention in Postmenopausal Women. *N. Engl. J. Med.* **2011**, *364*, 2381–2391. [[CrossRef](#)]
- Giordano, S.H.; Elias, A.D.; Gradishar, W.J. Gradishar. Nccn Guidelines Updates: Breast Cancer. *J. Natl. Compr. Cancer Netw.* **2018**, *16*, 605–610. [[CrossRef](#)]
- Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften e.V. (AWMF). Interdisziplinäre S3-Leitlinie Für Die Früherkennung, Diagnostik, Therapie Und Nachsorge Des Mammakarzinoms. Available online: https://www.awmf.org/uploads/tx_szleitlinien/032-045OLI_S3_Mammakarzinom_2017-12.pdf (accessed on 19 July 2021). (In German).
- Shaikh, T.; Tam, T.Y.; Li, T.; Hayes, S.B.; Goldstein, L.; Bleicher, R.; Boraas, M.; Sigurdson, E.; Ryan, P.D.; Anderson, P. Multifocal and Multicentric Breast Cancer is Associated with Increased Local Recurrence Regardless of Surgery Type. *Breast J.* **2015**, *21*, 121–126. [[CrossRef](#)]
- Zhang, X.; Hofmann, S.; Rack, B.; Harbeck, N.; Jeschke, U.; Sixou, S. Fluorescence Analysis of Vitamin D Receptor Status of Circulating Tumor Cells (CTCS) in Breast Cancer: From Cell Models to Metastatic Patients. *Int. J. Mol. Sci.* **2017**, *18*, 1318. [[CrossRef](#)]
- Ditsch, N.; Toth, B.; Mayr, D.; Lenhard, M.; Gallwas, J.; Weissenbacher, T.; Dannecker, C.; Friese, K.; Jeschke, U. The Association between Vitamin D Receptor Expression and Prolonged Overall Survival in Breast Cancer. *J. Histochem. Cytochem.* **2011**, *60*, 121–129. [[CrossRef](#)]
- Lang, Z.; Wu, Y.; Li, C.; Li, X.; Wang, X.; Qu, G. Multifocal and Multicentric Breast Carcinoma: A Significantly More Aggressive Tumor Than Unifocal Breast Cancer. *Anticancer Res.* **2017**, *37*, 4593–4598. [[PubMed](#)]
- Reinert, T.; De Paula, B.; Shafae, M.N.; Souza, P.H.; Ellis, M.J.; Bines, J. Endocrine therapy for ER-positive/HER2-negative metastatic breast cancer. *Chin. Clin. Oncol.* **2018**, *7*, 25. [[CrossRef](#)]
- Welsh, J. Function of the vitamin D endocrine system in mammary gland and breast cancer. *Mol. Cell. Endocrinol.* **2017**, *453*, 88–95. [[CrossRef](#)]
- Liu, C.-Y.; Wu, C.-Y.; Petrossian, K.; Huang, T.-T.; Tseng, L.-M.; Chen, S. Treatment for the endocrine resistant breast cancer: Current options and future perspectives. *J. Steroid Biochem. Mol. Biol.* **2017**, *172*, 166–175. [[CrossRef](#)]

18. Muscat, G.E.O.; Eriksson, N.A.; Byth, K.; Loi, S.; Graham, D.; Jindal, S.; Davis, M.J.; Clyne, C.; Funder, J.W.; Simpson, E.R.; et al. Research Resource: Nuclear Receptors as Transcriptome: Discriminant and Prognostic Value in Breast Cancer. *Mol. Endocrinol.* **2013**, *27*, 350–365. [CrossRef]
19. Escriva, H.; Bertrand, S.; Laudet, V. The evolution of the nuclear receptor superfamily. *Essays Biochem.* **2004**, *40*, 11–26.
20. Dawson, M.I.; Xia, Z. The retinoid X receptors and their ligands. *Biochim. Biophys. Acta (BBA) Mol. Cell Biol. Lipids* **2012**, *1821*, 21–56. [CrossRef] [PubMed]
21. Hua, S.; Kittler, R.; White, K.P. Genomic Antagonism between Retinoic Acid and Estrogen Signaling in Breast Cancer. *Cell* **2009**, *137*, 1259–1271. [CrossRef]
22. Ditsch, N.; Heublein, S.; Jeschke, U.; Sattler, C.; Kuhn, C.; Hester, A.; Czogalla, B.; Trillsch, F.; Mahner, S.; Engel, J.; et al. Cytoplasmic versus nuclear THR alpha expression determines survival of ovarian cancer patients. *J. Cancer Res. Clin. Oncol.* **2020**, *146*, 1923–1932. [CrossRef] [PubMed]
23. Zehni, A.Z.; Jacob, S.-N.; Mumm, J.-N.; Heidegger, H.H.; Ditsch, N.; Mahner, S.; Jeschke, U.; Vilsmaier, T. Hormone Receptor Expression in Multicentric/Multifocal versus Unifocal Breast Cancer: Especially the VDR Determines the Outcome Related to Focality. *Int. J. Mol. Sci.* **2019**, *20*, 5740. [CrossRef]
24. Shao, W.; Kuhn, C.; Mayr, D.; Ditsch, N.; Kailuweit, M.; Wolf, V.; Harbeck, N.; Mahner, S.; Jeschke, U.; Cavailles, V.; et al. Cytoplasmic and Nuclear Forms of Thyroid Hormone Receptor Beta1 Are Inversely Associated with Survival in Primary Breast Cancer. *Int. J. Mol. Sci.* **2020**, *21*, 330. [CrossRef] [PubMed]
25. Czogalla, B.; Deuster, E.; Liao, Y.; Mayr, D.; Schmoeckel, E.; Sattler, C.; Kolben, T.; Hester, A.; Fürst, S.; Burges, A.; et al. Cytoplasmic VDR expression as an independent risk factor for ovarian cancer. *Histochem. Cell Biol.* **2020**, *154*, 421–429. [CrossRef] [PubMed]
26. Doan, T.B.; Graham, J.D.; Clarke, C. Emerging functional roles of nuclear receptors in breast cancer. *J. Mol. Endocrinol.* **2017**, *58*, R169–R190. [CrossRef]
27. Röszer, T.; Menéndez-Gutiérrez, M.P.; Cedenilla, M.; Ricote, M. Retinoid X receptors in macrophage biology. *Trends Endocrinol. Metab.* **2013**, *24*, 460–468. [CrossRef]
28. Leal, A.S.M.; Zydeck, K.; Carapellucci, S.; Reich, L.A.; Zhang, D.; Moerland, J.A.; Sporn, M.B.; Liby, K.T. Retinoid X receptor agonist LG100268 modulates the immune microenvironment in preclinical breast cancer models. *NPJ Breast Cancer* **2019**, *5*, 1–15. [CrossRef] [PubMed]
29. Nunez, V.; Alameda, D.; Rico, D.; Mota, R.; Gonzalo, P.; Cedenilla, M.; Fischer, T.; Bosca, L.; Glass, C.; Arroyo, K.A.G.; et al. Retinoid X Receptor Alpha Controls Innate Inflammatory Responses through the up-Regulation of Chemokine Expression. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 10626–10631. [CrossRef] [PubMed]
30. Han, S.; Roman, J. Peroxisome Proliferator-Activated Receptor Gamma: A Novel Target for Cancer Therapeutics? *Anticancer Drugs* **2007**, *18*, 237–244. [CrossRef]
31. Ditsch, N.; Vrekoussis, T.; Lenhard, M.; Ruhl, I.; Gallwas, J.; Weissenbacher, T.; Friese, K.; Mayr, D.; Makriganakis, A.; Jeschke, U. Retinoid X Receptor Alpha (Rxralpha) and Peroxisome Proliferator-Activated Receptor Gamma (Ppargamma) Expression in Breast Cancer: An Immunohistochemical Study. *In Vivo* **2012**, *26*, 87–92.
32. Tang, X.-H.; Gudas, L.J. Retinoids, Retinoic Acid Receptors, and Cancer. *Annu. Rev. Pathol. Mech. Dis.* **2011**, *6*, 345–364. [CrossRef]
33. Forman, B.M.; Umesono, K.; Chen, J.; Evans, R. Unique response pathways are established by allosteric interactions among nuclear hormone receptors. *Cell* **1995**, *81*, 541–550. [CrossRef]
34. Koeffler, H.P. Peroxisome Proliferator-Activated Receptor Gamma and Cancers. *Clin. Cancer Res.* **2003**, *9*, 1–9.
35. Nuclear Receptors Nomenclature Committee. A Unified Nomenclature System for the Nuclear Receptor Superfamily. *Cell* **1999**, *97*, 161–163. [CrossRef]
36. Yasmin, R.; Williams, R.; Xu, M.; Noy, N. Nuclear Import of the Retinoid X Receptor, the Vitamin D Receptor, and Their Mutual Heterodimer. *J. Biol. Chem.* **2005**, *280*, 40152–40160. [CrossRef] [PubMed]
37. Zhang, X.-K.; Su, Y.; Chen, L.; Chen, F.; Liu, J.; Zhou, H. Regulation of the nongenomic actions of retinoid X receptor- α by targeting the coregulator-binding sites. *Acta Pharmacol. Sin.* **2015**, *36*, 102–112. [CrossRef] [PubMed]
38. Knabl, J.; Vattai, A.; Hüttenbrenner, R.; Hutter, S.; Karsten, M.; Jeschke, U. RXR α is upregulated in first trimester endometrial glands of spontaneous abortions unlike LXR and PPAR γ . *Eur. J. Histochem.* **2016**, *60*, 2665. [CrossRef] [PubMed]
39. Laudet, V.; Gronemeyer, H. *The Nuclear Receptor FactsBook*; Academic Press: San Diego, CA, USA, 2002.
40. Kersten, S.; Dong, D.; Lee, W.-Y.; Reczek, P.R.; Noy, N. Auto-silencing by the retinoid X receptor 1 Edited by M. Yaniv. *J. Mol. Biol.* **1998**, *284*, 21–32. [CrossRef]
41. The Human Protein Atlas. 2021. Available online: <http://www.proteinatlas.org> (accessed on 19 July 2021).
42. Raffo, P.; Emionite, L.; Colucci, L.; Belmonto, F.; Moro, M.G.; Bollag, W.; Toma, S. Retinoid Receptors: Pathways of Proliferation Inhibition and Apoptosis Induction in Breast Cancer Cell Lines. *Anticancer Res.* **2000**, *20*, 1535–1543.
43. Friedrich, M.; Axt-Fliedner, R.; Villena-Heinsen, C.; Tilgen, W.; Schmidt, W.; Reichrath, J. Analysis of Vitamin D-Receptor (Vdr) and Retinoid X-Receptor Alpha in Breast Cancer. *Histochem. J.* **2002**, *34*, 35–40. [CrossRef] [PubMed]
44. Elstner, E.; Williamson, E.A.; Zang, C.; Fritz, J.; Heber, D.; Fenner, M.; Possinger, K.; Koeffler, H.P. Novel Therapeutic Approach: Ligands for Ppargamma and Retinoid Receptors Induce Apoptosis in Bcl-2-Positive Human Breast Cancer Cells. *Breast Cancer Res. Treat* **2002**, *74*, 155–165. [CrossRef]

45. Crowe, D.L.; Chandraratna, R.A. A Retinoid X Receptor (Rxr)-Selective Retinoid Reveals That Rxr-Alpha Is Potentially a Therapeutic Target in Breast Cancer Cell Lines, and That It Potentiates Antiproliferative and Apoptotic Responses to Peroxisome Proliferator-Activated Receptor Ligands. *Breast Cancer Res.* **2004**, *6*, R546–R555. [[CrossRef](#)] [[PubMed](#)]
46. Suh, N.; Wang, Y.; Williams, C.R.; Risingsong, R.; Gilmer, T.; Willson, T.M.; Sporn, M.B. A new ligand for the peroxisome proliferator-activated receptor-gamma (PPAR-gamma), GW7845, inhibits rat mammary carcinogenesis. *Cancer Res.* **1999**, *59*, 5671–5673.
47. Zanardi, S.; Serrano, D.; Argusti, A.; Barile, M.; Puntoni, M.; DeCensi, A. Clinical trials with retinoids for breast cancer chemoprevention. *Endocr. Relat. Cancer* **2006**, *13*, 51–68. [[CrossRef](#)] [[PubMed](#)]
48. Ross-Innes, C.S.; Stark, R.; Holmes, K.A.; Schmidt, D.; Spyrou, C.; Russell, R.; Massie, C.E.; Vowler, S.L.; Eldridge, M.; Carroll, J.S. Cooperative Interaction between Retinoic Acid Receptor-Alpha and Estrogen Receptor in Breast Cancer. *Genes Dev.* **2010**, *24*, 171–182. [[CrossRef](#)] [[PubMed](#)]
49. Heublein, S.; Mayr, D.; Meindl, A.; Kircher, A.; Jeschke, U.; Ditsch, N. Vitamin D Receptor, Retinoid X Receptor and Peroxisome Proliferator-Activated Receptor Gamma Are Overexpressed in Brca1 Mutated Breast Cancer and Predict Prognosis. *J. Exp. Clin. Cancer Res.* **2017**, *36*, 57. [[CrossRef](#)]
50. Bonofiglio, D.; Cione, E.; Vizza, D.; Perri, M.; Pingitore, A.; Qi, H.; Catalano, S.; Rovito, D.; Genchi, G.; Ando, S. Bid as a Potential Target of Apoptotic Effects Exerted by Low Doses of Ppargamma and Rxr Ligands in Breast Cancer Cells. *Cell Cycle* **2011**, *10*, 2344–2354. [[CrossRef](#)]
51. Yasmin, R.; Kannan-Thulasiraman, P.; Kagechika, H.; Dawson, M.I.; Noy, N. Inhibition of Mammary Carcinoma Cell Growth by Rxr Is Mediated by the Receptor's Oligomeric Switch. *J. Mol. Biol.* **2010**, *397*, 1121–1131. [[CrossRef](#)]
52. Zehni, A.Z.; Batz, F.; Vattai, A.; Kaltofen, T.; Schrader, S.; Jacob, S.N.; Mumm, J.N.; Heidegger, H.H.; Ditsch, N.; Mahner, S.; et al. The Prognostic Impact of Retinoid X Receptor and Thyroid Hormone Receptor Alpha in Unifocal vs. Multifocal/Multicentric Breast Cancer. *Int. J. Mol. Sci.* **2021**, *22*, 957. [[CrossRef](#)] [[PubMed](#)]
53. Cserni, G.; Chmielik, E.; Cserni, B.; Tot, T. The new TNM-based staging of breast cancer. *Virchows Archiv.* **2018**, *472*, 697–703. [[CrossRef](#)] [[PubMed](#)]
54. Hortobagyi, G.N.; Edge, S.B.; Giuliano, A. New and Important Changes in the TNM Staging System for Breast Cancer. *Am. Soc. Clin. Oncol. Educ. Book* **2018**, *38*, 457–467. [[CrossRef](#)]
55. Elston, C.; Ellis, I. Pathological prognostic factors in breast cancer. I. The value of histological grade in breast cancer: Experience from a large study with long-term follow-up. *Histopathology* **1991**, *19*, 403–410. [[CrossRef](#)]
56. Remmele, W.; Stegner, H.E. Recommendation for Uniform Definition of an Immunoreactive Score (Irs) for Immunohistochemical Estrogen Receptor Detection (Er-Ica) in Breast Cancer Tissue. *Pathologe* **1987**, *8*, 138–140.
57. Weissenbacher, T.; Hirte, E.; Kuhn, C.; Janni, W.; Mayr, D.; Karsten, U.; Rack, B.; Friese, K.; Jeschke, U.; Heublein, S.; et al. Multicentric and Multifocal Versus Unifocal Breast Cancer: Differences in the Expression of E-Cadherin Suggest Differences in Tumor Biology. *BMC Cancer* **2013**, *13*, 361. [[CrossRef](#)]
58. Weissenbacher, T.M.; Zschage, M.; Janni, W.; Jeschke, U.; Dimpfl, T.; Mayr, D.; Rack, B.; Schindlbeck, C.; Friese, K.; Dian, D. Multicentric and Multifocal Versus Unifocal Breast Cancer: Is the Tumor-Node-Metastasis Classification Justified? *Breast Cancer Res. Treat* **2010**, *122*, 27–34. [[CrossRef](#)]
59. Ditsch, N.; Toth, B.; Himsl, I.; Lenhard, M.; Ochsenkühn, R.; Friese, K.; Mayr, D.; Jeschke, U. Thyroid hormone receptor (TR)alpha and TRbeta expression in breast cancer. *Histol. Histopathol.* **2013**, *28*, 227–237. [[PubMed](#)]
60. Heublein, S.; Mayr, R.; Meindl, A.; Angele, M.; Gallwas, J.; Jeschke, U.; Ditsch, N. Thyroid Hormone Receptors Predict Prognosis in BRCA1 Associated Breast Cancer in Opposing Ways. *PLoS ONE* **2015**, *10*, e0127072. [[CrossRef](#)] [[PubMed](#)]
61. Ditsch, N.; Mayr, D.; Lenhard, M.; Strauss, C.; Vodermaier, A.; Gallwas, J.; Stoeckl, D.; Graeser, M.; Weissenbacher, T.; Friese, K.; et al. Correlation of Thyroid Hormone, Retinoid X, Peroxisome Proliferator-Activated, Vitamin D and Oestrogen/Progesterone Receptors in Breast Carcinoma. *Oncol. Lett.* **2012**, *4*, 665–671. [[CrossRef](#)]
62. Ghose, R.; Zimmerman, T.L.; Thevananther, S.; Karpen, S.J. Endotoxin Leads to Rapid Subcellular Re-Localization of Hepatic Rxralpha: A Novel Mechanism for Reduced Hepatic Gene Expression in Inflammation. *Nucl. Recept.* **2004**, *2*, 4. [[CrossRef](#)] [[PubMed](#)]
63. Mey, J.; Schrage, K.; Wessels, I.; Vollpracht-Crijns, I. Effects of Inflammatory Cytokines Il-1beta, Il-6, and Tnfalpha on the Intracellular Localization of Retinoid Receptors in Schwann Cells. *Glia* **2007**, *55*, 152–164. [[CrossRef](#)]
64. Tanaka, T.; Dancheck, B.L.; Trifiletti, L.C.; Birnkrant, R.E.; Taylor, B.J.; Garfield, S.H.; Thorgeirsson, U.; De Luca, L.M. Altered Localization of Retinoid X Receptor Alpha Coincides with Loss of Retinoid Responsiveness in Human Breast Cancer Mda-Mb-231 Cells. *Mol. Cell. Biol.* **2004**, *24*, 3972–3982. [[CrossRef](#)] [[PubMed](#)]
65. Zhou, H.; Liu, W.; Su, Y.; Wei, Z.; Liu, J.; Kolluri, S.K.; Wu, H.; Cao, Y.; Chen, J.Y.; Wu, T.; et al. Nsaid Sulindac and Its Analog Bind Rxralpha and Inhibit Rxralpha-Dependent Akt Signaling. *Cancer Cell* **2010**, *17*, 560–573. [[CrossRef](#)]
66. Salehin, D.; Haugk, C.; Thill, M.; Cordes, T.; William, M.; Hemmerlein, B.; Friedrich, M. Vitamin D receptor expression in patients with vulvar cancer. *Anticancer. Res.* **2012**, *32*, 283–289. [[PubMed](#)]
67. Matusiak, D.; Murillo, G.; Carroll, R.E.; Mehta, R.G.; Benya, R.V. Expression of Vitamin D Receptor and 25-Hydroxyvitamin D3-1[1]-Hydroxylase in Normal and Malignant Human Colon. *Cancer Epidemiol. Biomark. Prev.* **2005**, *14*, 2370–2376. [[CrossRef](#)] [[PubMed](#)]

68. Hutchinson, P.E.; Halsall, J.A.; Popovici, S.; Papadogeorgakis, E.; Osborne, J.E.; Powley, I.R.; Dasari, D.; Saldanha, G.; Pringle, J.H. Compromised vitamin D receptor signalling in malignant melanoma is associated with tumour progression and mitogen-activated protein kinase activity. *Melanoma Res.* **2018**, *28*, 410–422. [[CrossRef](#)] [[PubMed](#)]
69. Joseph, C.; Al-Izzi, S.; Alsaleem, M.; Kurozumi, S.; Toss, M.S.; Arshad, M.; Goh, F.Q.; Alshankyty, I.M.; Aleskandarany, M.; Ali, S.; et al. Retinoid X receptor gamma (RXRG) is an independent prognostic biomarker in ER-positive invasive breast cancer. *Br. J. Cancer* **2019**, *121*, 776–785. [[CrossRef](#)]
70. Hennigs, A.; Riedel, F.; Gondos, A.; Sinn, P.; Schirmacher, P.; Marmé, F.; Jäger, D.; Kauczor, H.U.; Stieber, A.; Lindel, K.; et al. Prognosis of Breast Cancer Molecular Subtypes in Routine Clinical Care: A Large Prospective Cohort Study. *BMC Cancer* **2016**, *16*, 734. [[CrossRef](#)]
71. Lee, S.M.; Lee, J.Y.; Choi, J.E.; Lee, S.Y.; Park, J.Y.; Kim, D.S. Epigenetic inactivation of retinoid X receptor genes in non-small cell lung cancer and the relationship with clinicopathologic features. *Cancer Genet. Cytogenet.* **2010**, *197*, 39–45. [[CrossRef](#)] [[PubMed](#)]
72. Kalra, R.S.; Bapat, S.A. Expression Proteomics Predicts Loss of Rxr-Gamma During Progression of Epithelial Ovarian Cancer. *PLoS ONE* **2013**, *8*, e70398. [[CrossRef](#)]
73. Liby, K.; Rendi, M.; Suh, N.; Royce, D.B.; Risingsong, R.C.; Williams, R.; Lamph, W.; Labrie, F.; Krajewski, S.; Xu, X.; et al. The Combination of the Retinoid, Lg100268, and a Selective Estrogen Receptor Modulator, Either Arzoxifene or Acolbifene, Synergizes in the Prevention and Treatment of Mammary Tumors in an Estrogen Receptor-Negative Model of Breast Cancer. *Clin. Cancer Res.* **2006**, *12*, 5902–5909. [[CrossRef](#)]
74. Wu, K.; Kim, H.T.; Rodriguez, J.L.; Hilsenbeck, S.G.; Mohsin, S.K.; Xu, X.C.; Lamph, W.W.; Kuhn, J.G.; Green, J.E.; Brown, P.H. Suppression of Mammary Tumorigenesis in Transgenic Mice by the Rxr-Selective Retinoid, Lgd1069. *Cancer Epidemiol. Biomark. Prev.* **2002**, *11*, 467–474.
75. Wu, K.; Dupré, E.; Kim, H.; Tin-U, C.K.; Bissonnette, R.P.; Lamph, W.W.; Brown, P.H. Receptor-selective retinoids inhibit the growth of normal and malignant breast cells by inducing G1 cell cycle blockade. *Breast Cancer Res. Treat.* **2005**, *96*, 147–157. [[CrossRef](#)] [[PubMed](#)]
76. Kong, G.; Kim, H.-T.; Wu, K.; De Nardo, D.; Hilsenbeck, S.G.; Xu, X.-C.; Lamph, W.W.; Bissonnette, R.; Dannenberg, A.J.; Brown, P.H. The Retinoid X Receptor-Selective Retinoid, LGD1069, Down-regulates Cyclooxygenase-2 Expression in Human Breast Cells through Transcription Factor Crosstalk: Implications for Molecular-Based Chemoprevention. *Cancer Res.* **2005**, *65*, 3462–3469. [[CrossRef](#)] [[PubMed](#)]
77. Gniadecki, R.; Assaf, C.; Bagot, M.; Dummer, R.; Duvic, M.; Knobler, R.; Ranki, A.; Schwandt, P.; Whittaker, S. The optimal use of bexarotene in cutaneous T-cell lymphoma. *Br. J. Dermatol.* **2007**, *157*, 433–440. [[CrossRef](#)]
78. Esteva, F.J.; Glaspy, J.; Baidas, S.; Laufman, L.; Hutchins, L.; Dickler, M.; Tripathy, D.; Cohen, R.; De Michele, A.; Yocum, R.C.; et al. Multicenter Phase II Study of Oral Bexarotene for Patients with Metastatic Breast Cancer. *J. Clin. Oncol.* **2003**, *21*, 999–1006. [[CrossRef](#)] [[PubMed](#)]