



# HHS Public Access

Author manuscript

*J Am Chem Soc.* Author manuscript; available in PMC 2021 October 14.

Published in final edited form as:

*J Am Chem Soc.* 2020 October 14; 142(41): 17236–17242. doi:10.1021/jacs.0c05595.

## Serine-Selective Bioconjugation

Julien C. Vantourout<sup>1</sup>, Srinivasa Rao Adusumalli<sup>2</sup>, Kyle W. Knouse<sup>1</sup>, Dillon Flood<sup>1</sup>, Antonio Ramirez<sup>4</sup>, Natalia M. Padial<sup>1</sup>, Alena Istrate<sup>2</sup>, Katarzyna Maziarz<sup>1</sup>, Justine N. deGruyter<sup>1</sup>, Rohan R. Merchant<sup>1</sup>, Jennifer X. Qiao<sup>4</sup>, Michael A. Schmidt<sup>4</sup>, Michael J. Deery<sup>5</sup>, Martin D. Eastgate<sup>\*,4</sup>, Philip E. Dawson<sup>\*,1</sup>, Gonçalo J. L. Bernardes<sup>\*,2,3</sup>, Phil S. Baran<sup>\*,1</sup>

<sup>1</sup>Department of Chemistry, Scripps Research, 10550 North Torrey Pines Road, La Jolla, CA 92037, United States.

<sup>2</sup>Department of Chemistry, University of Cambridge, Lensfield Road, CB2 1EW Cambridge, United Kingdom.

<sup>3</sup>Instituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, Avenida Professor Egas Moniz, 1649-028 Lisboa, Portugal.

<sup>4</sup>Chemical Process Development, Bristol-Myers Squibb, One Squibb Drive, New Brunswick, NJ 08903, United States.

<sup>5</sup>Cambridge Centre for Proteomics, Milner Therapeutics Institute, Jeffrey Cheah Biomedical Centre, University of Cambridge, Puddicombe Way, CB2 0AW Cambridge, United Kingdom.

### Abstract

The first general method for the rapid, chemoselective, and modular functionalization of serine residues in native polypeptides is reported using a reagent platform based on P(V) oxidation state. This redox-economic approach can be used to append nearly any kind of cargo onto serine, generating a stable, benign, and hydrophilic phosphorothioate linkage. The method tolerates all other known nucleophilic functional groups of naturally occurring proteinogenic amino acids. A variety of applications can be envisaged by this expansion of the toolbox of site-selective bioconjugation methods.

### Graphical Abstract


---

\*Corresponding Authors: martin.eastgate@bms.com, dawson@scripps.edu, gb453@cam.ac.uk, pbaran@scripps.edu.  
Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website. The Supporting Information contains all experimental procedures, analysis, the detail of DFT calculations and compound characterization data.



It is no surprise that of the twenty naturally-occurring amino acid residues bearing reactive groups, serine is rarely chosen as a handle for selective derivatization (Figure 1A).<sup>1–4</sup> This lack of serine functionalization clearly stems not from a lack of desire, but rather the unmet challenge of differentiating the nucleophilicity of serine residues from other nucleophilic sidechains and water (chemoselectivity) and single residue among its multiple copies (site-selectivity).<sup>1–7</sup> A process that could achieve this differentiation would be inherently valuable, as it would add a new dimension to the toolbox of options available for chemoselective modification of peptides and the site-selective labeling of proteins. Whereas nature has pointed to a possible solution to this problem using enzymatic machinery and phosphorus(V) reactivity,<sup>8–16</sup> there is currently no general method (chemoenzymatic or otherwise) to site-selectively attach a precise cargo to a serine residue in a native biomolecule.<sup>1–3,17</sup> Kinases are typically responsible for the phosphorylation of various amino acid residues such as serine (Figure 1B).<sup>8–16</sup> This process suggests that P(V)-based electrophiles could have an innate preference for alcohol-based nucleophiles.<sup>18</sup> In parallel with this biological precedent, chemical inspiration for P(V)-based bioconjugation stems from the new class of phosphorous reagents ( $\Psi$  and  $\Pi$ ) which react with both oxygen- and carbon-based nucleophiles in a rapid and predictable way.<sup>19,20</sup> In this Communication,  $\Psi$ -loaded reagents ( $\Psi$ -modules) are shown to exhibit strikingly selective reactivity towards serine. This method enables both selective post-translational labeling and bioconjugation of serine residues.

In 2018, a set of reagents based upon a limonene-scaffold fused to an oxathiophospholane heterocycle were introduced for the purpose of simplifying stereocontrolled access to phosphorothioate linkages (Phosphorous-Sulfur Incorporation, abbreviated PSI or  $\Psi$ ).<sup>19</sup> This redox-economic approach avoided the extraneous steps associated with a P(III)-based manifold, yet retained the high reactivity associated with phosphoramidites and related systems.<sup>19</sup> Shortly thereafter, related limonene-based reagents were introduced to access both phosphines and methylphosphonates through stereocontrolled P–C bond formation (Phosphorous Incorporation, abbreviated PI or  $\Pi$ ).<sup>20</sup>

During the studies using PSI reagent, it was sporadically observed that oxygen-based nucleophiles would react in preference to other heteroatoms such as sulfur and nitrogen.<sup>19,20</sup> After optimization of the coupling step (see supporting information, Figure S23), the generality of these observations was explored systematically in the context of amino-acid functionalization as depicted in Figure 2A. Thus, competitive coupling experiments between amino acids revealed a striking selectivity for serine functionalization using  $\Psi$ -module

**P(V)-1** in the presence of cysteine, lysine, tyrosine and selenocysteine, even at high concentration (0.1 M). These preliminary studies established the viability of such an approach, and represent selectivities on par with traditional protein labeling methods (e.g. NHS esters for lysine, iodoacetamide for cysteine, and MTAD for tyrosine/tryptophan).<sup>1-4</sup> Interestingly, reagent **P(V)-1** under similar conditions displayed a respectable selectivity of 7:1 favoring serine over threonine. Transitioning to more realistic models for chemoselective labeling, competitive coupling experiments were then pursued using small peptides with  $\Psi$ -modules **P(V)-1**, **P(V)-2a**, and **P(V)-3** (Figure 2B). Good-to-excellent conversions were observed within 15 minutes, and unless otherwise noted, only one product was formed with no epimerization. Exquisite selectivity was obtained for serine at both 50 mM and 100 mM reagent concentration (10 mM peptide concentration) when competing with cysteine, lysine, histidine, glutamine, methionine, tryptophan, and tyrosine. When employing cysteine containing peptides, excess EtSH was added to avoid the formation of adducts between cysteine and the thiirane by-product of the  $\Psi$ -module. In addition, a reductive quench (DTT) is used to reduce any S-S bond formed during the reaction.<sup>21,22</sup> When competing with threonine, a useful level of selectivity (determined by H-coupled <sup>31</sup>P NMR) was observed at 50 mM of reagents **P(V)-1** (8:1), **P(V)-2a** (6:1), or reagent **P(V)-3** (7:1) delivering compounds **15**, **16** and **17** in 95%, 72% and 94%, respectively.<sup>1-4</sup>

More realistic applications of this method were then explored in the context of longer peptides containing a variety of nucleophilic amino acids (Figure 3A). Excellent selectivity for serine and satisfactory conversions were always observed using a variety of  $\Psi$ -modules (Figure 3A). As an example, peptide **SI-14** was selectively labelled at the serine position to afford peptide **31**, **32**, and **33** in 23%, 36% and 17% conversions using reagents **P(V)-1**, **P(V)-2b**, and **P(V)-3**, respectively. In addition to these linear examples, the functionalization of cyclic peptides was explored due to their importance and relevance in the context of medicinal chemistry (Figure 3B).<sup>23-25</sup> The developed conditions were first applied to the diversification of a cyclic peptide containing several nucleophilic functionalities, affording 61% yield of the desired product **34** within 10 minutes. An oxytocin-derivative, bearing a sensitive disulfide bridge and a tyrosine, successfully underwent coupling in 42% yield, with complete selectivity for the serine residue.<sup>26-28</sup> Finally, vancomycin was selected as a key example with an impressive functional group array: one carboxylic acid, one primary amine, one primary amide, three phenolic alcohols, five secondary alcohols, and a single primary alcohol.<sup>29-33</sup> Nevertheless, both reagents **P(V)-1** and **P(V)-3** demonstrated remarkable selectivity for the primary alcohol site affording compounds **36** and **37** in 19% and 21% yields, respectively. The labeled position was confirmed by H-coupled <sup>31</sup>P NMR; assignment was simple as the molecule bears only one primary alcohol.

Next, we challenged this chemical method for the installation of various  $\Psi$ -loaded reagents on proteins. The protein ubiquitin was primarily chosen for its relevance in proteasome targeting.<sup>34-36</sup> When ubiquitin **38** was vortexed with **P(V)-2b** reagent and DBU, it resulted in mono-labeled ubiquitin **39** in 37% conversion (Figure 4A). Labeled ubiquitin **39** was digested with trypsin and analyzed by LC-MS/MS (see SI for details). The resulting MS/MS data were searched using the Mascot search algorithm using a variable modification of 176 Da and a database containing the ubiquitin sequence. MS/MS of P(V) tagged peptide

(ESTLHLVLR, E64-R72,  $m/z$  622.31  $[M+2H]^{2+}$ ) confirmed the site-selective labeling of Ser residue (S65; kinetic labeling). Subsequently, other  $\Psi$ -modules were screened for the labeling of ubiquitin. **P(V)-1**, **P(V)-3**, and **P(V)-4** reagents afforded the mono-labeling of ubiquitin in 32% (**40**), 40% (**41**), and 20% (**42**) conversions respectively. In addition, the method was efficiently translated to the labeling of the DNA binding protein 434 repressor<sup>37–39</sup> **43** using **P(V)-2b**; reaction of these components delivered **44** in 35% conversion (Figure 4B). Further advances in the design of  $\Psi$ -reagents that are more reactive, yet more stable in aqueous conditions, and cysteine compatible will facilitate further use of this new approach toward the chemo- and regioselective modification of serine residues on proteins.

DFT studies were undertaken to rationalize the selectivity observed for serine in the coupling step (Figure 5).<sup>40</sup> These studies support a stepwise  $A_N+D_N$  phosphorylation mechanism (Figure 5A) with limiting barriers for the formation (**TS1**) or collapse (**TS2**) of a trigonal bipyramidal pentacoordinated (TBP, **INT1**) intermediate (Figure 5B). The remaining steps of the reaction coordinate involve a ring flip of the limonene skeleton (**TS3**) followed by  $S_N2$  displacement of the phosphorylated amino acid and simultaneous release of the cyclohexene sulfide (**TS4**) with an exothermic balance ( $G \approx -37$  kcal/mol).<sup>41</sup>

Transition structures **TS3** represent the highest barrier within the sequence of conformational distortions<sup>42</sup> required to transform chair **INT2** into chair **INT3** (See SI for details).<sup>43</sup> The transition structures leading to the TBP, **TS1**, display the backside attack of the nucleophile to the P-O ring bond in a late, intermediate-like arrangement. **TS1** shows a large extent of RX-P (where X = O, or N) bond formation and concomitant proton transfer to the nitrogen atom of DBU acting as a general base. Amino acids with alcohol and amine side chains display a rate-limiting formation of the trigonal bipyramidal intermediate with barriers that increase in the order Ser < Thr << Tyr << Lys (**TS1**, Figure 5B). A  $G^\ddagger$  of 0.6 kcal/mol between **TS1**-Ser and **TS1**-Thr is in qualitative agreement with the lower but competitive reactivity of threonine, whereas a  $G^\ddagger$  of 4.9 kcal/mol between **TS1**-Ser and **TS1**-Tyr is consistent with the limited reactivity of tyrosine. Similarly, **TS1**-Ser is favored relative to **TS1**-Lys by  $\approx 9$  kcal/mol in agreement with the lower acidity of the primary amine and greater steric congestion during the partial deprotonation by DBU. A search for the transition structures corresponding to the collapse of the TBP intermediates led to structures **TS2**, which depict the dissociation of the ring S-P bond accompanied by pseudorotation<sup>44</sup> and reduction of the O-P-O angle as the tetrahedral phosphorylated P(V) product forms (Figure 5A and SI for details). The formation of a strong O-P bond and the axial arrangement of the methoxy group facilitate the dissociation of the weaker S-P bond (See SI for details).<sup>45</sup> For amino acids with alcohol side chains, transition structures **TS2** leading to the collapse of the TBP intermediates are  $\approx 8$  kcal/mol more stable than the transition structures **TS1** forming the TBP intermediates. In contrast with the reaction coordinate for the alcohol, the collapse of the **TBP**-Cys intermediate is disfavored relative to its formation (cf. **TS1**-Cys and **TS2**-Cys, Figure 5B). The existence of a low barrier for decomposition of the TBP intermediate back to reactants has been used to explain the slow thiolysis of phosphate triesters.<sup>46</sup> In agreement with experiment, the limiting barrier for the phosphorylation of cysteine (**TS2**-Cys,  $G^\ddagger = +22.2$  kcal/mol) is estimated to be 2.7 kcal/mol higher than the limiting barrier for the phosphorylation of serine (**TS1**-Ser,  $G^\ddagger =$

+19.5 kcal/mol). A comparison of **TS2**-Cys and **TS2**-Ser geometries shows the reluctance of cysteine to pseudorotate ( $S-P-O$ ring  $147.0^\circ$  in **TS2**-Cys versus  $111.2^\circ$  in **TS2**-Ser), become equatorial, and stabilize ring P-S dissociation via the axial-axial arrangement of the OMe group and the ring P-S bond (See SI). The higher pseudorotation barrier of cysteine relative to serine is consistent with the preference of good leaving groups (e.g. cysteine) to occupy an axial position.<sup>47</sup> It worth noting that the selectivity for serine versus cysteine and tyrosine may simply be associated with a higher degree of reversibility. Taken all together, these results rationalize the serine selectivity observed for the coupling step.

Expanding on our previous work on the chemistry of the  $\Psi$ -reagent system, a chemoselective, rapid, and robust method for the direct functionalization of serine residues in the context of both peptides and proteins has been developed. The presented methodology displays excellent chemoselectivity for serine residues under the optimized solvent conditions and will enable practitioners in this field to append virtually any kind of cargo desired in an orthogonal fashion. Applications of these findings to bioconjugation, chemical biology, as well as the pursuit of novel materials can all be envisaged.<sup>48</sup> Development of new PSI-reagents that are compatible to aqueous conditions and cysteine residues will also facilitate further use of this new approach toward the chemo- and regioselective modification of serine residues on proteins.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## ACKNOWLEDGMENT

Financial support for this work was provided by Bristol-Myers Squibb, NIH (GM-118176), the Marie Skłodowska-Curie Global Fellowships (749359-EnanSET, N.M.P) within the European Union research and innovation framework programme (2014-2020), FCT Portugal (IF/00624/2015) and the Royal Society (URF/R\180019). DTF was supported by the National Center for Advancing Translational Sciences, National Institutes of Health, through grant number UL1 TR002551 and linked award TL1 TR002551. Authors are grateful to Dr. Dee-Hua Huang and Dr. Laura Pasternack (Scripps Research) for assistance with nuclear magnetic resonance (NMR) spectroscopy, to Dr. Jason Chen, Brittany Sanchez and Emily Sturgell (Scripps Automated Synthesis Facility) for assistance with HPLC, HRMS and LCMS. We finally thank Dr Victor L. Ayora and Lavinia Dunsmore for 434 repressor protein sample.

## REFERENCES

- (1). Algar W. Russ, D. P, Medintz Igor L., Chemoselective and Bioorthogonal Ligation Reactions: Concepts and Applications. John Wiley & Sons: 2017.
- (2). deGruyter JN; Malins LR; Baran PS Residue-Specific Peptide Modification: A Chemist's Guide. *Biochemistry* 2017, 56, 3863–3873. [PubMed: 28653834]
- (3). (a) Hoyt EA; Cal PMSD; Oliveira BL; Bernardes GJL Contemporary approaches to site-selective protein modification. *Nat. Rev. Chem* 2019, 3, 147–171. (b) Rawale DG; Thakur K; Adusumalli SR; Rai V Chemical methods for selective labeling of proteins. *Eur. J. Org. Chem* 2019, 6749–6763.
- (4). Boutureira O; Bernardes GJL Advances in chemical protein modification. *Chem. Rev* 2015, 115, 2174–2195. [PubMed: 25700113]
- (5). Bruice TC; Fife TH; Bruno JJ; Brandon NE Hydroxyl group catalysis. II. The reactivity of the hydroxyl group of serine. The nucleophilicity of alcohols and the ease of hydrolysis of their acetyl esters as related to their pKa. *Biochemistry* 1962, 1, 7–12. [PubMed: 13873927]

- (6). Neet KE; Koshland DE The conversion of serine at the active site of subtilisin to cysteine: a “chemical mutation”. *Proc. Natl. Acad. Sci. USA* 1966, 56, 1606–1611. [PubMed: 5230319]
- (7). Polgar L; Bender ML A new enzyme containing a synthetically formed active site. thiol-subtilisin. *J. Am. Chem. Soc* 1966, 88, 3153–3154.
- (8). Jackson MD; Denu JM Molecular reactions of protein phosphatases-insights from structure and chemistry. *Chem. Rev* 2001, 101, 2313–2340. [PubMed: 11749375]
- (9). Manning G; Whyte DB; Martinez R; Hunter T; Sudarsanam S The protein kinase complement of the human genome. *Science* 2002, 298, 1912–1934. [PubMed: 12471243]
- (10). Kennelly PJ Protein phosphatases-a phylogenetic perspective. *Chem. Rev* 2001, 101, 2291–2312. [PubMed: 11749374]
- (11). Cohen P The regulation of protein function by multisite phosphorylation – a 25-year update. *Trends Biochem. Sci* 2000, 25, 596–601. [PubMed: 11116185]
- (12). Johnson LN; Lewis RJ Structural basis for control by phosphorylation. *Chem. Rev* 2001, 101, 2209–2242. [PubMed: 11749371]
- (13). Chen C; Ha BH; Thevenin AG; Lou HJ; Zhang R; Yip KY; Peterson JR; Gerstein M; Kim PM; Filippakopoulos P; Knapp S; Boggon TJ; Turk BE Identification of a major determinant for serine-threonine kinase phosphoacceptor specificity. *Mol. Cell* 2014, 53, 140–147. [PubMed: 24374310]
- (14). Tagliabracci VS; Wiley SE; Guo X; Kinch LN; Durrant E; Wen J; Xiao J; Cui J; Nguyen KB; Engel JL; Coon JJ; Grishin N; Pinna LA; Pagliarini DJ; Dixon JE A single kinase generates the majority of the secreted phosphoproteome. *Cell* 2015, 161, 1619–632. [PubMed: 26091039]
- (15). Miller CJ; Turk BE Homing in: mechanisms of substrate targeting by protein kinases. *Trends Biochem. Sci* 2018, 43, 380–394. [PubMed: 29544874]
- (16). Westheimer FH Why nature chose phosphates. *Science* 1987, 235, 1173–1178. [PubMed: 2434996]
- (17). Liu H; Li X Serine/Threonine Ligation: Origin, Mechanistic Aspects, and Applications. *Acc. Chem. Res* 2018, 51, 1643–1655. [PubMed: 29979577]
- (18). For a recent report on selective phosphorylation of histidine residue, see: Jia S; He D; Chang CJ Bioinspired Thiophosphorodichloridate Reagents for Chemoselective Histidine Bioconjugation. *J. Am. Chem. Soc* 2019, 141, 7294–7301. [PubMed: 31017395]
- (19). Knouse KW; deGruyter JN; Schmidt MA; Zheng B; Vantourout JC; Kingston C; Mercer SE; McDonald IM; Olson RE; Zhu Y; Hang C; Zhu J; Yuan C; Wang Q; Park P; Eastgate MD; Baran PS Unlocking P(V): Reagents for chiral phosphorothioate synthesis. *Science* 2018, 361, 1234–1238. [PubMed: 30072577]
- (20). Xu D; Rivas-Bascón N; Padial, N. M; Knouse KW; Zheng B; Vantourout JC; Schmidt MA; Eastgate MD; Baran PS Enantiodivergent Formation of C-P Bonds: Synthesis of P-Chiral Phosphines and Methylphosphonate Oligonucleotides. *J. Am. Chem. Soc* 2020, 142, 5785–5792. [PubMed: 32109356]
- (21). Lu BY; Chang JY Rapid and irreversible reduction of protein disulfide bonds. *Anal. Biochem* 2010, 405, 67–72. [PubMed: 20488157]
- (22). Singh R; Whitesides GM Reagents for Rapid Reduction of Native Disulfide Bonds in Proteins. *Bioorg. Chem* 1994, 22, 109–115.
- (23). Zorzi A; Deyle K; Heinis C Cyclic peptide therapeutics: past, present and future. *Curr. Opin. Chem. Biol* 2017, 38, 24–29. [PubMed: 28249193]
- (24). Jing X; Jin K A gold mine for drug discovery: Strategies to develop cyclic peptides into therapies. *Med. Res. Rev* 2020, 40, 753–810. [PubMed: 31599007]
- (25). Lee AC; Harris JL; Khanna KK; Hong JH A Comprehensive Review on Current Advances in Peptide Drug Development and Design. *Int. J. Mol. Sci* 2019, 20, 2383–2403.
- (26). Ichinose W; Cherepanov SM; Shabalova AA; Yokoyama S; Yuhi T; Yamaguchi H; Watanabe A; Yamamoto Y; Okamoto H; Horike S; Terakawa J; Daikoku T; Watanabe M; Mano N; Higashida H; Shuto S Development of a Highly Potent Analogue and a Long-Acting Analogue of Oxytocin for the Treatment of Social Impairment-Like Behaviors. *J. Med. Chem* 2019, 62, 3297–3310. [PubMed: 30896946]

- (27). Gimpl G; Fahrenholz F The oxytocin receptor system: structure, function, and regulation. *Physiol. Rev* 2001, 81, 629–683. [PubMed: 11274341]
- (28). Bakermans-Kranenburg MJ; van IJMH Sniffing around oxytocin: review and meta-analyses of trials in healthy and clinical groups with implications for pharmacotherapy. *Transl. Psychiatry* 2013, 3, e258. [PubMed: 23695233]
- (29). Okano A; Isley NA; Boger DL Peripheral modifications of [ $\Psi$ [CH<sub>2</sub>NH]Tpg<sup>4</sup>]vancomycin with added synergistic mechanisms of action provide durable and potent antibiotics. *Proc. Natl. Acad. Sci. U.S.A* 2017, 114, E5052–E5061. [PubMed: 28559345]
- (30). Boger DL; Miyazaki S; Kim SH; Wu JH; Castle SL; Loiseleur O; Jin Q Total Synthesis of the Vancomycin Aglycon. *J. Am. Chem. Soc* 1999, 121, 10004–10011.
- (31). Nicolaou KC; Mitchell HJ; Jain NF; Winssinger N; Hughes R; Bando T Total Synthesis of Vancomycin. *Angew. Chem. Int. Ed* 1999, 38, 240–244.
- (32). Monteiro JF; Hahn SR; Goncalves J; Fresco P Vancomycin therapeutic drug monitoring and population pharmacokinetic models in special patient subpopulations. *Pharmacol Res. Perspect* 2018, 6, e00420. [PubMed: 30156005]
- (33). Boger DL Vancomycin, teicoplanin, and ramoplanin: synthetic and mechanistic studies. *Med. Res. Rev* 2001, 21, 356–381. [PubMed: 11579438]
- (34). Pickart CM Ubiquitin in chains. *Trends in Biochem. Sci* 2000, 25, 544–548. [PubMed: 11084366]
- (35). Swatek KN; Komander D Ubiquitin modifications. *Cell Res.* 2016, 26, 399–422. [PubMed: 27012465]
- (36). Glickman MH; Ciechanover A The ubiquitin-proteasome proteolytic pathway: destruction for the sake of construction. *Physiol. Rev* 2002, 82, 373–428. [PubMed: 11917093]
- (37). Bushman FD; Ptashne M Activation of transcription by the bacteriophage 434 repressor. *Proc. Natl. Acad. Sci. U.S.A* 1986, 83, 9353–9357. [PubMed: 3467310]
- (38). Rodgers DW; Harrison SC The complex between phage 434 repressor DNA-binding domain and operator site OR3: structural differences between consensus and non-consensus half-sites. *Structure* 1993, 1, 227–240. [PubMed: 8081737]
- (39). Koudelka GB Recognition of DNA structure by 434 repressor. *Nucleic Acids Res.* 1998, 26, 669–675. [PubMed: 9421532]
- (40). See SI for references regarding the DFT calculations process. Based on reported pKa values of the side chain functional groups, Ser, Thr, and Tyr were modeled following a general base catalysis path, whereas Cys was modeled as a fully deprotonated nucleophile.
- (41). (a)Uchimaru T; Stec WJ; Taira K Mechanism of the Chemoselective and Stereoselective Ring Opening of Oxathiaphospholanes: An Ab Initio Study. *J. Org. Chem* 1997, 62, 5793–5800. (b)Uchimaru T; Stec WJ; Tsuzuki S; Hirose T; Tanabe K; Taira K Ab Initio Investigation on Nucleophilic Ring opening of 1, 3, 2-Oxathiaphospholane: Nucleophilic Substitution at Phosphorus Coupled with Pseudorotation. *Chem. Phys. Lett* 1996, 263, 691–696.
- (42). Dixon DA; Komornicki A Ab Initio Conformational Analysis of Cyclohexane. *J. Phys. Chem* 1990, 94, 5630–5636.
- (43). The S–C–O dihedral angle was scanned at fixed intervals and the highest energy geometry of the resulting potential energy was fully optimized to afford a twisted half-chair transition structure.
- (44). (a)Westheimer FH Pseudo-Rotation in the Hydrolysis of Phosphate Esters. *Acc. Chem. Res* 1968, 1, 70–78. (b)Boyd DB Mechanism of Hydrolysis of Cyclic Phosphate Esters. *J. Am. Chem. Soc* 1969, 91, 1200–1205.
- (45). Park K-H; Kim M-H; Um I-H Kinetic Study on Nucleophilic Displacement Reactions of Phenyl Y-Substituted Phenyl Carbonates with 1,8-Diazabicyclo [5.4.0] undec-7-ene: Effects of Amine Nature on Reaction Mechanism. *Bull. Korean Chem. Soc* 2016, 37, 77–81.
- (46). Arantes GM; Chaimovich H Thiolytic and Alcoholysis of Phosphate Tri- and Monoesters with Alkyl and Aryl Leaving Groups. An ab Initio Study in the Gas Phase. *J. Phys. Chem. A* 2005, 109, 5625–5635. [PubMed: 16833895]

- (47). DeBruin KE; Petersen JR Steric and Electronic Effects on the Stereochemistry of the Alkaline Hydrolysis of Acyclic Dialkoxyphosphonium Salts. Pseudorotation of Intermediates in Phosphorus Ester Reactions. *J. Org. Chem* 1972, 37, 2272–2278.
- (48). Flood DT; Knouse KW; Vantourout JC; Sanchez BB; Sturgell EJ; Chen JS; Baran PS; Dawson PE Synthetic Elaboration of Native DNA by RASS (SENDR) ChemRxiv Preprint 2020.


Author Manuscript

Author Manuscript


Author Manuscript

Author Manuscript







**FIGURE 1.** (A) Serine: Unanswered Challenge for Selective Amino Acid Functionalization. (B) PSI Reagents Mimic Natural P(V)-based Phosphorylation of Serine.

**FIGURE 2.**


(A) Competitive Experiments Between Amino Acids. (B) Competitive Experiments Using Small Tetrapeptides (10 mM). <sup>a</sup> Using EtSH (5 equiv).

**FIGURE 3.**

(A) Peptide Scope. (B) Functionalization of Cyclic Peptide.



**FIGURE 4.** Functionalization of Proteins (A) Functionalization of Ubiquitin. (B) Functionalization of 434 Repressor.



**FIGURE 5.** (A) General Stepwise Mechanism of the Coupling Step. (B) DFT Calculations to Rationalize the Serine Selectivity.