



Published in final edited form as:

*AIDS Care*. 2020 August ; 32(8): 1061–1068. doi:10.1080/09540121.2019.1698703.

## Enhancers and Barriers to Uptake of Male Circumcision Services in Northern Uganda: A qualitative study

Barbara M Nanteza<sup>a,b</sup>, Fredrick E Makumbi<sup>b</sup>, Ronald H Gray<sup>c</sup>, David Serwadda<sup>d</sup>, Ping Teresa Yeh<sup>e</sup>, Caitlin E Kennedy<sup>e</sup>

<sup>a</sup>Ministry of Health, AIDS Control Program, Uganda <sup>b</sup>Makerere University, College of Health Sciences, School of Public Health, Department of Epidemiology and Biostatistics, Uganda <sup>c</sup>Johns Hopkins Bloomberg School of Public Health, Department of Epidemiology, Baltimore, MD, USA <sup>d</sup>Makerere University, College of Health Sciences, School of Public Health, Department of Disease Control and Environmental Health, Uganda <sup>e</sup>Johns Hopkins Bloomberg School of Public Health, Department of International Health, Baltimore, MD, USA

### Abstract

Uganda adopted voluntary medical male circumcision (VMMC) in 2010, but uptake remains disproportionately low in the Northern region despite implementing several demand creation strategies. This study explored the socio-cultural and structural enhancers and barriers to uptake of VMMC services in Gulu, a district in Northern Uganda where uptake is lowest. In September 2016, we conducted 19 focus group discussions, 9 in-depth interviews, and 11 key informant interviews with 149 total participants. Data were collected and analyzed thematically using both inductive and deductive approaches, then framed in four levels of the social ecological model. Enhancers included adequate knowledge about VMMC services, being young and single, partner involvement, peer influence, perceived increased libido after circumcision, and availability of free and high-quality VMMC services. Barriers included sexual abstinence during wound healing, penile appearance after circumcision, religion, culture, and misconceptions. Optimizing enhancers and addressing barriers could increase VMMC service uptake in northern Uganda.

---

Corresponding author: Dr. Barbara M Nanteza, National Safe Male Circumcision (SMC) Coordinator, AIDS Control Program, Ministry of Health, Uganda, P.O. Box 7272 K'la Plot 6 Lourdel Rd, Nakasero, Tel: +256 701 -657-704, dr\_nanteza@yahoo.co.uk. Contributions

Barbara M Nanteza

Generated the research question, conducted the data collection, analysis and write up of the draft manuscript

Fredrick E Makumbi

Supervised the protocol development, data collection, and reviewed the various drafts and final version of the manuscript

Ronald H Gray

Reviewed all the various drafts and final manuscripts and guided on the analysis and interpretation of findings and the policy and program benefits

David Serwadda

Provided technical support for the analysis methodology, interpretation of findings, reviewed the drafts and final manuscript

Ping Teresa Yeh

Provided technical support in the analysis, interpretation of the findings and reviewed the manuscripts and

Caitlin E Kennedy

Supervised and provided technical support for the analysis methodology, reviewed the drafts and final manuscript

Disclosure statement

The authors disclose no financial interest or benefit that has arisen from the direct applications of our research.

Geographical information

This work took place in Gulu District, Northern Uganda.

## Keywords

male circumcision; uptake; facilitators; challenges

---

## Introduction

Voluntary medical male circumcision (VMMC) reduces female-to-male transmission of HIV by approximately 60% (Auvert, Taljaard et al. 2005, Bailey, Moses et al. 2007, Gray, Kigozi et al. 2007). Since 2010, 14 priority countries in Eastern and Southern Africa have scaled up VMMC as part of combination HIV prevention following recommendations by the World Health Organization (WHO) and the United Nations Joint Programme on HIV/AIDS (UNAIDS) (WHO and UNAIDS 2007, WHO and UNAIDS 2011). A modelling study in Uganda suggested that 6.9 million 10–49 year old males had to be circumcised by 2020 to achieve 80% coverage needed for public health benefit (Kripke, Vazzano et al. 2016).

The Uganda Ministry of Health (MoH), with support from the U.S. President's Emergency Fund for AIDS Relief (PEPFAR), increased availability of VMMC services and intensified mobilization in communities. VMMC prevalence among Ugandan men aged 15–49 years increased from 26% in 2011 to 43% in 2016, but wide regional variations persist, ranging from 69% in the Mid-Eastern region to 14% in the Mid-Northern Region (MoH 2019). Several strategies have been suggested to increase uptake including mobile outreaches, dedicated teams, dedicated VMMC days, devices, education entertainment, and implementing VMMC as a vertical program (Gourlay, Birdthistle et al. 2013, Kitara, Lagoro et al. 2013, WHO Regional Office for Africa 2013, Menon, Gold et al. 2014, Mati, Adegoke et al. 2016). Anecdotal reports suggest specific socio-cultural reasons and inadequate knowledge of key VMMC messages may explain the disproportionately low uptake in this region. In a complementary quantitative study, we found that over a quarter (27.5%) of male participants in northern Uganda had inadequate knowledge of key VMMC messages (Nanteza, Serwadda et al. 2018). More in-depth understanding of enhancers and barriers is needed in such a region with very low uptake of VMMC. However similar studies that have been done in the other 14 East and Southern priority countries (Hatzold K, Mavhu W et al. 2014, Abunah B, Onkoba R et al. 2016, Gavin George, Kaymarlin Govender et al. 2017, Maria A. Carrasco, Jessica Wilkinson et al. 2019)

## Methods

### Study design and setting

The study was conducted in Gulu district, northern Uganda, a region scarred from a recent twenty-year civil war that fueled HIV epidemic and internally displaced many people. The study was conducted in September 2016 in the catchment areas of the four public health facilities providing VMMC: Lacor Hospital, Gulu Regional Referral Hospital, Awach Health Center IV, and Lalogi Health Center IV.

Data were collected from 149 individuals aged 10 to 69 years who participated in 19 focus group discussions (FGDs), 11 key informant interviews (KIIs), 9 in-depth interviews (IDIs)

and a one day workshop with all district health team (DHT) members where we discussed the barriers and enhancers at district level. Each FGD consisted of 6–8 individuals of similar age, gender, and marital status (Table 1). After the 19 FGDs, IDIs and KIIs were conducted until saturation was attained. IDIs were conducted with men and women who held strong opinions for and against circumcision, as identified by village health teams. Cultural leaders from the Acholi cultural institution and religious leaders from the main religions (Catholic, Anglican, Pentecostal, Muslim, Seventh Day Adventist, and traditional) were selected for KIIs. Circumcision status was not considered in study recruitment because it was very low in this region. Study team members were trained persons recruited from northern Uganda who were fluent in both Acholi and English. The lead investigator was the national VMMC coordinator.

FGD, KII, and IDI guides covered factors that enhance or hinder uptake of VMMC services, including culture, religion, tribe, socioeconomic status, education, and occupation. Participants were informed of the purpose of the study, the extent of their involvement, and their rights before obtaining written consent and assent. Interviews were conducted in the participant's preferred language and digitally recorded. They lasted 60–90 minutes for FGDs and 45 minutes for IDIs and KIIs. Participants received refreshments and 7,000 Ugandan shillings (~US\$2) for their time.

Following a preliminary analysis of qualitative results, a dissemination workshop with VMMC major stakeholders in Gulu district was held in July 2017. The workshop generated additional information and discussion about the factors that may enhance or hinder uptake of VMMC services mainly at the district level.

### Data analysis

All FGDs, IDIs, and KIIs data were first transcribed in Acholi, then translated into English. MAXDQA software was used for data management and analysis. We used a socio-ecological model (SEM) comprising four levels (individual, household, community, and district) to conduct deductive analysis (Sweat and Denison 1995). A codebook was developed based on initial discussions using the SEM. The resulting codes were discussed, narratively summarized, and presented.

**Ethical Considerations**—The Makerere University School of Public Health Higher Degrees Research and Ethics Committee (HDREC) and the Uganda National Council for Science and Technology (UNCST) approved the protocol (# TR2015002451). Permission for data collection was obtained from the district health office, the residential district commissioner, and local community leaders.

### Results

FGD participants included a wide range of community members, including married and unmarried men and women of various ages and some boys; KIIs included both cultural and religious leaders, and IDIs explored pro- and anti-circumcision viewpoints (Table 1). Findings are presented below across the four levels of the SEM; themes are summarized with supportive quotes in Table 2.

## Individual Level

### Enhancers

**Knowledge of VMMC benefits among highly educated participants:** Participants (both male and female) with secondary or higher education levels demonstrated high knowledge about VMMC and its benefits.

**Peer influence among youth:** Young people (aged 10–29 years) considered circumcision fashionable; uptake in this age group was driven by peer influence. Older (aged 30 years and above) participants considered circumcision to be for young men who were still experimenting and making discoveries about their bodies.

**Perceived improved sexual performance:** Almost all circumcised participants reported improved sexual performance following circumcision with some spouses of circumcised men reporting “better quality” sexual performance after circumcision.

### Barriers

**Other health issues prioritized:** Due to the civil war, there many internally displaced persons. They felt they had other health issues that were more important than circumcision since they saw Muslims and other persons from traditionally circumcising communities die from HIV.

**Duration of sexual abstinence during wound healing:** Sexually active men said that the required six weeks of sexual abstinence after circumcision was too long. They preferred to get circumcised during the absence of their sexual partners to avoid arousal that might lead to prolonged healing.

**Penile appearance:** Participants expressed concern about the appearance of a circumcised penis, using the word “layom” (having a penis with an exposed glans like a baboon’s).

**Adverse events after circumcision:** Some participants reported that friends’ experiences of adverse events following circumcision made them shy away from circumcision.

**Non-involvement of women:** In the current communication strategy, women are considered a secondary audience yet they mothers and spouses to VMMC clients.

**Disinhibition:** Before getting circumcised, all clients are counselled about the benefits of VMMC. Some participants said individuals mistake approximately 60% reduction of HIV infection after VMMC for immunity, resulting in promiscuity.

## Household Level

### Enhancers

**Monogamous families with greater gender equality:** Although men make most family decisions, male participants seemed more likely to seek VMMC if they were in a monogamous relationship or had greater gender equality within the family.

## Barriers

**Low-income families:** Wound care is an important aspect of successful circumcision. Low-income families felt they already had enough health problems and were not willing for additional burden of wound care and the risk of potential infection.

**Low literacy levels:** Implementing partners (IPs) frequently mobilized men to seek VMMC services through written materials like billboards and brochures. These messages were often poorly understood by community members with low literacy levels.

**Both partners from Northern Uganda:** In families where both spouses were from the same tribe or from Northern Uganda, men seemed very reluctant to get circumcised.

## Community Level

### Enhancers

**Adequate comprehensive knowledge about VMMC:** Communities believed that adequate comprehensive knowledge about VMMC services would enhance uptake and combat HIV. Most community leaders were ready to promote VMMC only if IPs involved them.

**Availability and access to free VMMC services:** The communities were grateful that VMMC services were free of charge, readily available, and easily accessible at many health facilities.

**Safe and high quality VMMC services:** All IPs are required to provide quality VMMC services through trained and certified service providers.

### Barriers

**Religion:** The people in Acholi region held negative feelings towards Islam because of past experiences. Some associated circumcision with Islam, leading to negative feelings and perceptions about circumcision.

**Culture:** Cultural leaders informed the study team that their kingdom was a member of the district health committee and often attended district health meetings. However, they had not promoted VMMC because no one had ever approached their institution to promote circumcision.

**Misconceptions about circumcision:** Some of the misconceptions were; if you got circumcised part of you (the prepuce) dies and is buried, which would bring bad luck to you and your family. Also, any man who 'spilled' blood was considered ungodly while others thought that when you get circumcised you lose your libido. Some thought anesthesia makes a man sterile. These views were also elicited from women.

## District Level

### Enhancers

**Trained and certified service providers:** It was revealed that MoH and previous IPs had trained many service providers. However, they informed the team that all previous IPs did not allow these health care workers to offer VMMC services but would instead bring service providers from other regions whom they referred to as “Karuma crossers” (a derogatory term for outsiders).

**Availability of supplies and equipment:** All district health officers appreciated the availability of VMMC supplies, yet wondered how VMMC services could have emergency resuscitation kits while most of their maternity or theatres did not have these supplies.

### Barriers

**Lack of district ownership of the SMC program:** Almost all district health team (DHT) members agreed that they did not know how IPs were chosen to implement VMMC services. They also said they did not know how targets were set and allocated. This made them feel that they were not in control or did not own the VMMC program.

**Coordination and implementation of VMMC activities by IPs:** Many DHT members were not fully aware of VMMC activities conducted in their districts and attempts to get information about implementation plans were futile. This made government health care workers not want to encourage or support VMMC services.

**Non-involvement of districts in developing VMMC mobilization messages:** The DHT members complained that they were never involved in the development of VMMC mobilization messages. They said that some bill boards in their districts had VMMC messages written in other local languages which made them believe that these messages were not meant for their people.

## Discussion

Although VMMC coverage has improved from 4% in 2011, currently is just 13% in Mid-North region (MoH 2019). Uganda has not yet attained the recommended 80% coverage especially in northern Uganda (Carrasco, Nguyen et al. 2018). This study explored the factors that facilitate or hinder uptake of VMMC services in Gulu district. Some of the key enhancers were availability of safe VMMC services, adequate knowledge of benefits of VMMC, being young and single, perceived sexual improvement after VMMC, and partner’s involvement or interest in male circumcision. On the other hand, barriers were mainly social-cultural, especially peer influence, adverse events, appearance of a circumcised penis, duration of abstinence during wound healing, misconceptions, culture, and religion. Lack of district ownership or involvement during planning and implementation of the VMMC program was a major barrier at the district level.

Apart from a recent study that provided evidence about inadequate knowledge about VMMC messages, no studies have been conducted to understand socio-cultural factors that enhance

or hinder uptake of VMMC in northern Uganda (Nanteza, Serwadda et al. 2018). Almost all interviewed cultural and religious leaders wanted to be involved in sensitizing the communities about VMMC. Similar studies elsewhere have shown that when local leaders were involved in VMMC programs, uptake increased (WHO, UNAIDS et al. 2007, Ahmad, Jandu et al. 2013, Aizire, Fowler et al. 2013, Friedrich 2017).

Sexual behaviors after circumcision seemed to influence uptake of VMMC, especially among those who were sexually active and not in stable relationships (Andersson, Vardas et al. 2012, Abbott, Haberland et al. 2013, Wambura, Mahler et al. 2017). Another study showed that if peers were satisfied with VMMC services offered, they were more likely to encourage their friends to seek VMMC services (Kigozi, Musoke et al. 2014, Kibira, Daniel et al. 2017, Spense, Meller et al. 2017).

Our findings reinforce the important role that women play in VMMC uptake, as demonstrated in other studies (Lanham, L'Engle et al. 2012, Westercamp, Agot et al. 2012, Shacham, Godlonton et al. 2014, Maraux, Lissouba et al. 2017). In Uganda, regions where women are recruited as VHTs have slightly higher VMMC coverage as compared to regions that predominantly use male VHTs. Rate of adverse events seems to be lower because women usually provide post-surgical support and wound care to their family members (Gray, Kigozi et al. 2009, Friedland, Apicella et al. 2013). In many communities, women's perception about circumcision really mattered to unmarried men, so involving women should increase VMMC uptake (Sabapathy, Van den Bergh et al. 2012, Johnson, Chiu et al. 2016, Lilleston, Marcell et al. 2017).

We found that religion was considered a barrier to uptake of VMMC services. The majority of the population in this setting are Catholics. A recent study in Rakai district highlighted religion as a key barrier (Ssekubugu, Leontsini et al. 2013, Lilleston, Marcell et al. 2017). VMMC messaging should continue to emphasize that VMMC is not associated with a specific religious tradition.

Adverse events after circumcision were mentioned as potential barrier to VMMC uptake. However, the Safety Monitoring team at MoH has received reports of adverse events including at least sixteen deaths as of October 2019. Such occurrences of serious adverse and fatal events are significant barriers to VMMC uptake. VMMC IPs will need to ensure that their teams are well trained and certified, so as to prevent adverse events and build confidence within the population.

For VMMC uptake to increase in Gulu district, all the barriers must be addressed and the enhancers encouraged. VMMC IPs in the region should work with MoH, districts, and local leaders to address these issues at all levels. Since VMMC is for HIV prevention, future studies should focus on factors influencing men who are at higher risk of HIV infection. For policy makers, these findings can improve uptake of VMMC services if the IP in northern Uganda can address barriers while working with the district and regional stakeholders. Cultural and religious leaders should be involved in the implementation of VMMC programme.



## Strengths and Limitations

One strength of this study was its qualitative approach. Instead of counting, we tried to understand in detail the barriers and enhancers of VMMC uptake at four different levels. This was the first study in this region to include women with important perspectives on VMMC uptake.

The study was conducted in Gulu, a central district with urban, peri-urban, and rural communities. The study team majorly comprised of young men and women from the region so they freely dialogued with participants using both English and Acholi. VHTs who mobilize men for VMMC services were used to select participants since they were very conversant with the communities. The VHTs and the lead author were never part of the interviews or discussions, which limited bias and allowed the participants to discuss the real issues that affected uptake of VMMC services in Gulu.

Limitations of the study included the fact that it was conducted in a district that had the highest coverage of VMMC in the region, which may limit the transferability of our findings. Participant responses may have been influenced by social desirability, particularly as the study team may have been seen as being related to health care service provision in the setting. However, we tried to combat this by specifically including interviews with people who were both supportive and unsupportive of circumcision.

## Conclusion

The study revealed that more needs to be done in Gulu district if people are to benefit from VMMC. Although culture and religion were barriers, they were not the major concerns. Adequate knowledge and women involvement were the main enhancers.

Since VMMC is an HIV prevention intervention, barriers should be addressed and enhancers encouraged. Specifically, we suggest that women's involvement should be encouraged, since women act as both spouses and mothers to potential clients. Second, we suggest addressing misconceptions through improved community mobilization, sensitization and health education. Finally, a wider range of stakeholders should be engaged for program ownership.

## Acknowledgements

We hereby acknowledge support for this work from Fogarty through the Rakai Health Science Program-Uganda and African Doctoral Dissertation Research Fellowship (ADDRF). We thank the supervisors at the Ministry of Health, including Dr. Jane Ruth Acheng, Professor Anthony K. Mbonye, and Dr. Joshua Musinguzi. Dr Peter Simon Kibira who helped in reviewing the study protocol especially the methodology. Finally, we thank our research assistants led by Julius Kisenyi, David Makubuya, Brian Ssenoga, Okello James Stewart, Arop Stephen, Abala Julius, Ochan Jesse, Odong Nicky, Akello Irene Olara Emmanuel, Otto Moses Oyika and Lamunu Winnie. Finally, we thank Gulu district leaders and all our participants.

### Funding

This work was supported by the National Institutes of Health (USA) Fogarty International Center under grant award number 5D43TW009578 and the African Doctoral Dissertation Research Fellowship (ADDRF).



## References

- Abbott SA, Haberland NA, Mulenga DM and Hewett PC (2013). "Female sex workers, male circumcision and HIV: a qualitative study of their understanding, experience, and HIV risk in Zambia." *PLoS One* 8(1): e53809.
- Abunah B, Onkoba R, Nyagero J and e. al. (2016). "Motivators and barriers to uptake of post-operative voluntary medical male circumcision follow-up in Yala division, Siaya County, Kenya. ." *Pan Afr Med* 25(7).
- Ahmad F, Jandu B, Albagli A, Angus JE and Ginsburg O. (2013). "Exploring ways to overcome barriers to mammography uptake and retention among South Asian immigrant women." *Health Soc Care Community* 21(1): 88–97. [PubMed: 23057604]
- Aizire J, Fowler M. and Coovadia H. (2013). "Operational issues and barriers to implementation of prevention of mother-to-child transmission of HIV (PMTCT) interventions in Sub-Saharan Africa." *Curr HIV Res* 11(2): 144–159. [PubMed: 23432490]
- Andersson KM, Vardas E, Niccolai LM, Niekerk RM, Mogale MM and Holdsworth IM (2012). "Anticipated changes in sexual risk behaviour following vaccination with a low-efficacy HIV vaccine: survey results from a South African township." *Int J STD AIDS* 23(10): 736–741. [PubMed: 23104749]
- Auvert B, Taljaard D, Lagarde E, Sobngwi-Tambekou J, Sitta R. and Puren A. (2005). "Randomized, controlled intervention trial of male circumcision for reduction of HIV infection risk: the ANRS 1265 Trial." *PLoS Med* 2(11): e298.
- Bailey RC, Moses S, Parker CB, Agot K, Maclean I, Krieger JN, Williams CFM, Campbell RT and Ndinya-Achola JO (2007). "Male circumcision for HIV prevention in young men in Kisumu, Kenya: a randomised controlled trial." *Lancet* 369(9562): 643–656. [PubMed: 17321310]
- Carrasco MA, Nguyen TQ and Kaufman MR (2018). "Low uptake of voluntary medical male circumcision among high risk men in Malawi." *AIDS Behav* 22(2): 447–453. [PubMed: 27943000]
- Friedland BA, Apicella L, Schenk KD, Sheehy M. and Hewett PC (2013). "How informed are clients who consent? A mixed-method evaluation of comprehension among clients of male circumcision services in Zambia and Swaziland." *AIDS Behav* 17(6): 2269–2282. [PubMed: 23392912]
- Friedrich MJ (2017). "Educating religious leaders increases male circumcision rates in Tanzania." *JAMA* 317(15): 1517.
- George Gavin, Govender Kaymarlin, Beckett Sean, Montague Carl and Frohlich J. (2017). "Factors associated with the take-up of voluntary medical male circumcision amongst learners in rural KwaZulu-Natal. ." *African Journal of AIDS Research* 16(3): 251–256. [PubMed: 28978292]
- Gourlay A, Birdthistle I, Mburu G, Iorpenda K. and Wringe A. (2013). "Barriers and facilitating factors to the uptake of antiretroviral drugs for prevention of mother-to-child transmission of HIV in sub-Saharan Africa: a systematic review." *J Int AIDS Soc* 19(16): 18588.
- Gray RH, Kigozi G, Serwadda D, Makumbi F, Nalugoda F, Watya S, Moulton L, Chen MZ, Sewankambo NK, Kiwanuka N, Sempijja V, Lutalo T, Kagayii J, Wabwire-Mangen F, Ridzon R, Bacon M. and Wawer MJ (2009). "The effects of male circumcision on female partners' genital tract symptoms and vaginal infections in a randomized trial in Rakai, Uganda." *Am J Obstet Gynecol* 200(1): 42.e41–47. [PubMed: 18976733]
- Gray RH, Kigozi G, Serwadda D, Makumbi F, Watya S, Nalugoda F, Kiwanuka N, Moulton LH, Chaudhary MA, Chen MZ, Sewankambo NK, Wabwire-Mangen F, Bacon MC, Williams CFM, Opendi P, Reynolds SJ, Laeyendecker O, Quinn TC and Wawer MJ (2007). "Male circumcision for HIV prevention in men in Rakai, Uganda: a randomised trial." *Lancet* 369(9562): 657–666. [PubMed: 17321311]
- Hatzold K, Mavhu W, Jasi P, Chatora K, Cowan FM and Taruberekera N (2014). "Barriers and Motivators to Voluntary Medical Male Circumcision Uptake among Different Age Groups of Men in Zimbabwe: Results from a Mixed Methods Study." *Plos One* 9(5).
- Johnson LF, Chiu C, Myer L, Davies MA, Dorrington RE, Bekker LG, Boulle A. and Meyer-Rath G. (2016). "Prospects for HIV control in South Africa: a model-based analysis." *Glob Health Action* 9(1): 30314. [PubMed: 27282146]

- Kibira SP, Daniel M, Atuyambe LM, Makumbi FE and Sandøy IF (2017). “Exploring drivers for safe male circumcision: Experiences with health education and understanding of partial HIV protection among newly circumcised men in Wakiso, Uganda.” *PLoS One* 12(3): e0175228.
- Kigozi G, Musoke R, Watya S, Kighoma N, Nkale J, Nakafeero M, Namuguzi D, Serwadda D, Nalugoda F, Sewankambo NK, Wawer MJ and Gray RH (2014). “The safety and acceptance of the PrePex device for non-surgical adult male circumcision in Rakai, Uganda. A non-randomized observational study.” *PLoS One* 9(8): e100008.
- Kitara D, Lagoro L, Ozero A, Lanyero J. and Ocom F. (2013). “Roll-out of medical male circumcision (MMC) for HIV prevention in non-circumcising communities of northern Uganda.” *Pan Afr Med J* 15: 100. [PubMed: 24198894]
- Kripke K, Vazzano A, Kirungi W, Musinguzi J, Opio A, Ssempebwa R, Nakawunde S, Kyobutungi S, Akao JN, Magala F, Mwidu G, Castor D. and Njehumeli E. (2016). “Modeling the impact of Uganda’s safe male circumcision program: Implications for age and regional targeting.” *PLoS One* 11(7): e0158693.
- Lanham M, L’Engle KL, Loolpapit M. and Oguma IO (2012). “Women’s roles in voluntary medical male circumcision in Nyanza Province, Kenya.” *PLoS ONE* 7(9): e44825.
- Lilleston PS, Marcell AV, Nakyanjo N, Leonard L. and Wawer MJ (2017). “Multilevel influences on acceptance of medical male circumcision in Rakai District, Uganda.” *AIDS Care* 29(8): 1049–1055. [PubMed: 28278564]
- Maraux B, Lissouba P, Rain-Taljaard R, Taljaard D, Bouscaillou J, Lewis D, Puren A. and Auvert B. (2017). “Women’s knowledge and perception of male circumcision before and after its roll-out in the South African township of Orange Farm from community-based cross-sectional surveys.” *PLoS One* 12(3): e0173595.
- Carrasco Maria A., Wilkinson Jessica, Kasdan B and Flemming P. (2019). “Systematic review of barriers and facilitators to voluntary medical male circumcision in priority countries and programmatic implications for service uptake.” *Global Public Health* 14 (1): 91–111. [PubMed: 29695201]
- Mati K, Adegoke KK and Salihu HM (2016). “Factors associated with married women’s support of male circumcision for HIV prevention in Uganda: a population based cross-sectional study.” *BMC Public Health* 16(1): 696. [PubMed: 27484177]
- Menon V, Gold E, Godbole R, Castor D, Mahler H, Forsythe S, Ally M. and Njehumeli E. (2014). “Costs and impacts of scaling up voluntary medical male circumcision in Tanzania.” *PLoS One* 9(5): e83925.
- MoH (2019). Uganda Population-Based HIV Impact Assessment: UPHIA 2016–2017. Kampala, Uganda Ministry of Health.
- Nanteza B, Serwadda D, Kankaka E, Mongo G, Gray R. and Makumbi F. (2018). “Knowledge on voluntary medical male circumcision in a low uptake setting in northern Uganda.” *BMC Public Health* 18(1): 1278. [PubMed: 30453966]
- Sabapathy K, Van den Bergh R, Fidler S, Hayes R. and Ford N. (2012). “Uptake of home-based voluntary HIV testing in Sub-Saharan Africa: A systematic review and meta-analysis.” *PLoS Med* 9(12): e1001351.
- Shacham E, Godlonton S. and Thornton RL (2014). “Perceptions of male circumcision among married couples in rural Malawi.” *J Int Assoc Provid AIDS Care* 13(5): 443–449. [PubMed: 24162614]
- Spense J, Meller J, Abbey J, Foster K, Sirri C. and Naqvi M. (2017). “Why are we cutting? A survey of cultural views on circumcision in the Texas panhandle.” *Glob Pediatr Health* 4: 2333794×17711767.
- Ssekubugu R, Leontsini E, Wawer M, Serwadda D, Kigozi G, Kennedy C, Nalugoda F, Sekamwa R, Wagman J. and Gray R. (2013). “Contextual barriers and motivators to adult male medical circumcision in Rakai, Uganda.” *Qual Health Res* 23(6): 795–804. [PubMed: 23515302]
- Sweat MD and Denison JA (1995). “Reducing HIV incidence in developing countries with structural and environmental interventions.” *AIDS* 9 Suppl A: S251–257. [PubMed: 8819593]
- Wambura M, Mahler H, Grund JM, Larke N, Mshana G, Kuringe E, Plotkin M, Lija G, Makokha M, Terris-Prestholt F, Hayes RJ, Changalucha J. and Weiss HA (2017). “Increasing voluntary medical

male circumcision uptake among adult men in Tanzania.” *Aids* 31(7): 1025–1034. [PubMed: 28350578]

Westercamp M, Agot KE, Ndinya-Achola J. and Bailey RC (2012). “Circumcision preference among women and uncircumcised men prior to scale-up of male circumcision for HIV prevention in Kisumu, Kenya.” *AIDS Care* 24(2): 157–166. [PubMed: 21854351]

WHO and UNAIDS (2007). *New data on male circumcision and HIV prevention: policy and programme implications*. Geneva, Switzerland, World Health Organization.

WHO and UNAIDS (2011). *Progress in male circumcision scale-up: Country implementation and research update*. Geneva, Switzerland, World Health Organization.

WHO, UNAIDS and London School of Hygiene and Tropical Medicine (2007). *Male circumcision: global trends and determinants of prevalence, safety and acceptability*. Geneva, Switzerland, World Health Organization.

WHO Regional Office for Africa (2013). *Progress in scaling up voluntary medical male circumcision for HIV prevention in East and Southern Africa: January-December 2012*. Brazzaville, Republic of Congo, World Health Organization.

**Table 1:**

Summary of the IDIs, KIIs and FGDs conducted

	<b>Number conducted</b>	
	<b>No. FDGs (19)</b>	<b>No. Participants (129)</b>
<b>Focus Group Discussions (FGDs)</b>		
Children boys (10–12 years)	1	8
Adolescent boys (13–19 years)	3	19
Unmarried men (20–25, 26–34, 35+ years)	3 (one per subgroup)	21
Married men (20–25, 26–34 years)	2 (one per subgroup)	15
Married men (35+ years)	4	27
Unmarried women (18–24, 25–34, 35+ years)	3 (one per subgroup)	20
Unmarried women (18–24, 25–34, 35+ years)	3 (one per subgroup)	19
<b>Key Informant Interviews (KIIs)</b>	11 Participants total	
<b>Cultural leaders</b>	5	
<b>Religious leaders</b>	6	
<b>In-depth interviews (IDIs)</b>	9 Participants total	
<b>Pro-circumcision</b>	3	
<b>Anti-circumcision</b>	6	

**Table 2.**

Themes and supportive quotes

Category	Factors/ Theme	Quote
<i>Individual level</i>		
Enhancers	Knowledge about benefits of VMMC among high education levels	"Ever since I learned about the benefits of circumcision, I cannot date a man who is not circumcised. On every first date, that is the first question I ask." [Female, 26 years, tertiary education, ID1]
	Peer influence among youth	"I think being circumcised is cool. Some of my friends at school are circumcised and they are happy. So that is why I also decided to get circumcised." [Male, 21 years, secondary education, FGD]
	Perceived improved sexual performance	"Before circumcision I never used to satisfy my wife, but now she says I am the best. I would only take a few minutes then I ejaculate, but now I take close to an hour." [Male, 35 years, primary education, ID1]
	VMMC is not a priority health issue	"We have health priorities of which circumcision is not among. Instead of wasting our taxes on programs that don't work, why don't you instead supply medicines in all government hospitals?" [Male, 36 years, primary education, ID1]
	Duration of sexual abstinence during wound healing	"I tried to abstain from sex for 4 weeks but I could not resist the temptation. I love my wife very much and we continued sharing the same bed." [Newly circumcised male, 27 years, tertiary education, ID1]
	Penile appearance	"My friend was rejected by his girlfriend after he was circumcised on grounds that his penis looked like that of a baboon and behaved like he was taking drugs making love for a much longer than before. If he hadn't been circumcised, he would still have his girlfriend!" [Male, 18 years, secondary education, FGD]
	Adverse events after being circumcised	"It's very sad that even health care workers have started operating like taxi drivers. Before you are circumcised they promise you heaven on earth, but the moment you get circumcised they do not pick your calls." [Male, 34, primary education, FGD]
	Non-involvement of women	It's me who knows what's good or bad for my children, thus my opinion on SMC [safe male circumcision] really matters a lot if you are to circumcise them." [Female, 27 years, primary education, FGD]
	Disinhibition	"Most of my friends after getting circumcised they say that they were told that they are immune to HIV." [Uncircumcised male, 21 years, Secondary FGD]
<i>Household level</i>		
Enhancers	Monogamous families with greater gender equality	"After learning about the benefits of circumcision from my brother, I begged my husband to get circumcised. He requested for my support during wound healing and he got circumcised the following day. Since then we are very happy and all our sons are circumcised." [Female, 42 years, secondary education, KII]
	Low income families	"My neighbors suffered with their son after he got an infection after being circumcised. They spent all the little money they had." [Female, 32 years, primary education, FGD]
Barriers	Low literacy levels	"I have seen Muslims die. I think government wants to kill us and take our land. We survived the war but now they are trying to use VMMC. [Male, 57 years, no education, KII]
	Both partners from northern Uganda	"In our culture it's not good for a man to be circumcised. That's for men from other tribes who want to entice women. We are very happy as God created us." [Male, 46 years, primary education, FGD]
<i>Community level</i>		
Enhancers	Adequate comprehensive knowledge about VMMC	"Our youth are dying from HIV, so if circumcision is promoted by the government to reduce HIV among our youth, we should be involved by the service providers to promote it." [Male, 67 years, primary education, KII]

Category	Factors/ Theme	Quote
	Availability and easy access to free, safe, and high quality MC services	"We are very grateful that we can access these services free of charge and they are usually brought in our communities through outreaches." [Male, 52 years, primary education, KII]
Barriers	Religion	"Circumcision is a ritual for Muslims. This was for those greedy people who sold our people to whites. As long as I live I will never embrace anything that they stood for" [Male, 62 years, primary education, KII] "After all, the Bible recommends circumcision of the heart, not the body. There are many other ways you can reduce your chances of acquiring HIV other than circumcision." [Male, 43 years, secondary education, KII]
	Culture	"I respect my king so I only promote what my chief tells me to tell his people. No one has ever told us about circumcision so we cannot promote it" [Male, 54 years, primary education, KII]
	Misconceptions about circumcision	"When you allow to bury any part of your body while still alive, not only you get bad luck but even all your family members." [Male, 59 years, primary education, IDI] "Any man that spills blood intentionally is ungodly. Even he is rejected by traditional herbalists." [Male, 49 years, no education, FGD]
<i>District level</i>		
Enhancers	Trained and certified service providers	"We are very grateful that almost all the service providers are trained" [District Health Officer]
	Availability of supplies and equipment	"We wish PEPFAR could also fund other health services, VMMC programme has supplies that theatres and maternity centres that do not have." [Chief Administrative Officer]
	Regularly updated data collection and reporting tools	"Unlike other programmes, VMMC is a very special programme with all tools reviewed more regularly than other programmes." [District Biostatistician]
Barriers	Lack of district ownership of the SMC programme	"As district leaders we do not have a say in the VMMC programme, we do not set or allocate targets, which makes supervision very difficult." [District Health Officer]
	Coordination and implementation of VMMC activities by IPs	"All we do is to coordinate implementing partners, usually they are more than one and all are doing VMMC in the same location. This makes coordination a nightmare." [District Education Officer]
	Non-involvement of districts in developing VMMC mobilization messages	"The VMMC programme has a lot of resources, that you just wake up one morning and see a very big billboard which at times is not even the local language." [District Mobilizer]