



Published in final edited form as:

Wiley Interdiscip Rev Syst Biol Med. 2016 May ; 8(3): 227–241. doi:10.1002/wsbm.1331.

Signaling via the NF κ B system

Simon Mitchell, Jesse Vargas, Alexander Hoffmann*

Department of Microbiology, Immunology, and Molecular Genetics, and Institute for Quantitative and Computational Biosciences, University of California, Los Angeles, Los Angeles, CA, USA

Abstract

The nuclear factor kappa B (NF κ B) family of transcription factors is a key regulator of immune development, immune responses, inflammation, and cancer. The NF κ B signaling system (defined by the interactions between NF κ B dimers, I κ B regulators, and IKK complexes) is responsive to a number of stimuli, and upon ligand–receptor engagement, distinct cellular outcomes, appropriate to the specific signal received, are set into motion. After almost three decades of study, many signaling mechanisms are well understood, rendering them amenable to mathematical modeling, which can reveal deeper insights about the regulatory design principles. While other reviews have focused on upstream, receptor proximal signaling (Hayden MS, Ghosh S. Signaling to NF- κ B. *Genes Dev* 2004, 18:2195–2224; Verstrepen L, Bekaert T, Chau TL, Tavernier J, Chariot A, Beyaert R. TLR-4, IL-1R and TNF-R signaling to NF- κ B: variations on a common theme. *Cell Mol Life Sci* 2008, 65:2964–2978), and advances through computational modeling (Basak S, Behar M, Hoffmann A. Lessons from mathematically modeling the NF- κ B pathway. *Immunol Rev* 2012, 246:221–238; Williams R, Timmis J, Qwarnstrom E. Computational models of the NF-KB signalling pathway. *Computation* 2014, 2:131), in this review we aim to summarize the current understanding of the NF κ B signaling system itself, the molecular mechanisms, and systems properties that are key to its diverse biological functions, and we discuss remaining questions in the field.

INTRODUCTION

Nuclear factor kappa B (NF κ B) is a family of dimeric transcription factors central to coordinating inflammatory responses; innate and adaptive immunity; and cellular differentiation, proliferation, and survival in almost all multicellular organisms.^{1–4} The NF κ B system is tightly regulated, and misregulation of NF κ B has been implicated in a wide range of diseases ranging from cancers to inflammatory and immune disorders. As a result, the NF κ B regulatory network and its dynamics offer a multitude of promising therapeutic targets that remain to be fully explored and translated into clinical use.^{5–7} However, there continues to be an untapped potential for finer grained therapeutic targeting of the NF κ B signaling system that requires a quantitative understanding of dynamical control and the integration of various physiological and pathological signals and stimuli.^{8–10}

*Correspondence to: ahoffmann@ucla.edu.

Conflict of interest: The authors have declared no conflicts of interest for this article.

In *Mammalia*, the NF κ B network consists of five family member protein monomers (p65/RelA, RelB, cRel, p50, and p52) that form homodimers or heterodimers that bind DNA differentially^{11–14} and are regulated by two pathways: the canonical, NF κ B essential modulator (NEMO)-dependent pathway and the noncanonical, NEMO-independent pathway. These pathways tightly control the levels and dynamics of the transcriptionally active NF κ B dimer repertoire constitutively and in response to stimuli, and thus control broad gene expression programs^{15,16} via the recruitment of co-activators¹⁷ or interplay with other transcription factors.^{18,19} The activation pathways control NF κ B activity through multiple mechanisms: degradation of I κ B inhibitor proteins, processing of NF κ B precursor proteins, and expression of NF κ B monomer proteins.^{10,20} Signals from tumor necrosis factor receptor (TNFR), toll-like receptor (TLR) superfamilies, interleukin receptor (IL-1R) and metabolic genotoxic, and shear stresses are integrated by the I κ B/NF κ B signaling network to produce signal-specific, context-specific, and cell-type-specific transcriptional responses.^{21–23}

CANONICAL SIGNALING

Signaling via the NEMO-Associated IKK Complex

The canonical NF κ B signaling pathway (a.k.a. NEMO-dependent pathway) is mediated by kinase complexes consisting of the scaffold/adaptor protein NEMO (a.k.a. IKK γ) and two I κ B kinases (IKK1/2, a.k.a. IKK α and IKK β) (Figure 1(a)). This IKK complex is activated by mechanisms that are NEMO dependent. The IKK kinases are activated by phosphorylation of serines in the activation T-loop, characteristic of the MAPK superfamily, and three mechanisms for IKK activation have emerged: (1) NEMO multimerizes IKK subunits²⁷ to allow for activation via trans-autophosphorylation, (2) and/or brings them in proximity to upstream kinases such as TAK1,²⁸ which may also lead to mutual activation forming positive feedback and digital dose-response characteristics.^{29,30} These mechanisms are mediated by NEMO's ubiquitin-binding domain that allows for IKK's recruitment to nondegradative K63-linked ubiquitin chains, which are a hallmark of inflammatory signaling. Finally, (3) NEMO itself is a substrate of ubiquitination, particularly linear ubiquitin chains produced by the LUBAC enzyme,³¹ which also facilitates the formation of transient signalsomes.

A wide variety of inflammatory cytokines (such as TNF and IL-1), a wide variety of pathogen-associated molecular patterns (PAMPs), or antigen/immune stimulatory signals result in IKK phosphorylation-dependent activation of the NEMO-containing complex, by one or a subset of these mechanisms,²⁴ resulting in complex dynamical control.³² Once activated, the complex binds to and phosphorylates I κ B proteins on specific serines in the N-terminal, leading to ubiquitination and subsequent proteasomal degradation.²⁵ The degradation of inhibitors releases NF κ B dimers associated with them, freeing NF κ B dimers and allowing them to bind κ B site-containing DNA and thus rapidly accumulate in the nucleus. Following I κ B release the NF κ B subunits are subject to a variety of posttranslational modification that fine-tune gene-expression control.³³

Further, NEMO was shown to function as a scaffold between IKK and I κ B α , thereby directing IKK activity to I κ B α .³⁴ This mechanism ensures that the activation of NF κ B

dimers associated with I κ B α , which is RelA:p50 in most cells and conditions, is directly linked to signals propagating through the NEMO hub. Hence, canonical signaling is often thought to be synonymous with RelA activation, but this is not always the case. In inflammatory dendritic cells, I κ B α was also shown to be associated with RelB:p50 dimers, thus rendering RelB a key transcriptional effector of the canonical NF κ B pathway in that cell type.³⁵

While the activation mechanism of IKK is beginning to be elucidated with the aid of recent structural and biophysical characterizations^{36,37} and, for example, combined single-cell computational studies that have identified distinct pathways for robust digital responses and noisy sustained responses,³⁸ how IKK is inactivated remains unclear. One attractive proposal is that IKK is regulated in an autocatalytic cycle of at least three states in which activation occurs from a poised state and is followed by an inactive state. While the dynamic properties of such a kinase control cycle have been studied,³⁹ the biophysical evidence remains scant, but could involve trans-autophosphorylation of an inhibitory C-terminal domain in IKK β ⁴⁰ and/or conformational changes of the complex.⁴¹

I κ B α Negative Feedback

Among the target genes regulated by κ B sites are the I κ Bs that, upon transcriptional induction and resynthesis, are able to translocate to the nucleus, bind to and inhibit NF κ B activity, trafficking it back to the cytosol. This constitutes the primary component of the self-regulating negative feedback loop²⁵ (Figure 1(b)). This feedback loop not only prevents run-away NF κ B activity in response to transient inflammatory signals but also poises the system for reactivation when IKK activity is longer lasting. I κ B α negative feedback has been studied in some detail with a combined experimental and mathematical modeling approach, and interesting properties have emerged: (1) Given the delay intrinsic to I κ B α resynthesis, even very transient cytokine exposure (1 min TNF) results in almost a full hour of NF κ B activity.⁴² That 1 h of NF κ B activity is invariant to the duration of the signal unless the incoming signal lasts longer than approximately 45 min. (2) Transcriptional induction of I κ B α is necessary but not sufficient for mediating this effective negative feedback control of NF κ B activity.⁴³ I κ B β , when controlled by an NF κ B-inducible promoter, is unable to provide normal physiologically observed negative feedback⁴⁴; nuclear import, export, and protein degradation mechanisms specific to I κ B α are critically important to recapitulating proper negative feedback and the NF κ B dynamic responses characteristic of normal activity.^{45–49} In addition, I κ B α has been shown to be able to strip NF κ B off DNA (or associate with chromatin⁵⁰), a function that I κ B β does not possess.⁵¹ (3) Given effective I κ B α negative feedback, stimulation conditions that produce longer lasting IKK activities allow for repeated cycles of reactivation, leading to oscillatory NF κ B activity observed in biochemical bulk population assays²⁵ or single-cell assays.^{38,52–54} Intrinsic variability in the kinetic mechanisms that govern I κ B α feedback is thought to render the oscillatory behavior more robust to variations in dynamic signals.^{55,56} However, the potential function of NF κ B oscillations remains unclear, and no satisfying answer has been presented as to how such oscillations are interpreted by downstream gene regulatory networks.⁵⁷

I κ B δ Negative Feedback

Another transcriptional target gene that is induced by nuclear NF κ B activity is the *nfk2* gene, which produces the p100 protein.⁴ p100 was first known as the precursor for the p52 protein, a dimerization partner of RelB and potentially other Rel proteins. However, p100 that is not processed to p52 is able to form higher-molecular-weight complexes that are capable of binding to NF κ B, acting as an I κ B, termed I κ B δ .²⁶ As such, NF κ B control of *nfk2* forms another negative feedback loop that may terminate NF κ B signaling.⁵⁸ However, transcriptional induction and protein synthesis are slow, in part due to the length and half-life of the mRNA and protein, and the subsequent required oligomerization step also takes several hours.⁵⁹ Thus, in contrast to I κ B α that functions rapidly, I κ B δ 's role is primarily in attenuating persistent signals.⁵⁸ Indeed, while I κ B α negative feedback is reversible as it is a sensitive substrate for continued canonical IKK activity, I κ B δ is insensitive to canonical signals and thus attenuates the canonical pathway regardless of whether incoming signals persist. Further, because I κ B δ has a longer half-life than other I κ Bs, it contributes to a signaling memory in which sequential stimulation events are dampened.⁵⁸ However, as a substrate for noncanonical signaling (see below), I κ B δ is a signaling crosstalk node that integrates canonical and noncanonical signals that may result in noncanonical signals emanating from LT β R or BAFF to activate (or prolong the activation of) NF κ B RelA or cRel dimers.^{26,60,61}

Other Feedback Mechanisms

There are several other negative and positive feedback mechanisms that contribute to the complex and potentially oscillatory dynamics of nuclear NF κ B.^{58,62} I κ B ϵ was shown to provide delayed negative feedback (due to a transcriptional delay) that may partially compensate for the loss of I κ B α ,⁶³ and was suggested to dampen I κ B α -mediated oscillations by forming a dual, antiphase negative feedback system.⁶² There is also evidence that I κ B α and ϵ preferentially inhibit distinct NF κ B family members.⁶⁴ In B cells, I κ B ϵ has been shown to play a key role in regulating cRel containing NF κ B dimers, with loss of I κ B ϵ resulting in increased B-cell survival and proliferation.²¹

Within the TNF pathway, both negative and positive feedback has been reported. Expression of the de-ubiquitinase A20 is strongly NF κ B inducible. However, owing to a long protein half-life and enzymatic effector function, its negative feedback effects do not shape NF κ B dynamics acutely, but rather integrate the history of prior exposure to render the NF κ B pathway less sensitive to subsequent stimuli.⁴² TNF itself is NF κ B inducible, but for full activation, additional signaling mechanisms controlling splicing, mRNA half-life, pro-TNF processing, and secretion must be activated.^{65,66} Hence, TNF functions more like a feed-forward loop in response to PAMPs rather than a positive feedback loop *per se*.

NONCANONICAL NF κ B SIGNALING

Signaling via NIK and IKK α

Noncanonical signals are NF κ B activating signals that are transduced in a NEMO-independent, but NIK and IKK α -dependent manner. Noncanonical pathways activating signals are primarily developmental signals that activate TNF receptors (BAFFR,

CD40, LT β R, RANK, TNFR2, Fn14, etc.), some of which also activate the canonical NF κ B pathway.^{67–74} Noncanonical NF κ B signaling is known to control a wide variety of developmental phenotypes including B-cell survival and maturation, dendritic cell activation, and bone metabolism.⁷⁵ Several chemokines that regulate lymphoid organogenesis are induced specifically by noncanonical NF κ B activation.⁶⁹ Base pair differences in κ B sites may contribute to noncanonical pathway-specific gene expression.^{76,77}

While canonical signals transduced by NEMO require phosphorylation-dependent activation of the IKK kinase complex, noncanonical signals transduced by NIK require stabilization and accumulation of the kinase, which, in the absence of signal, is rapidly degraded by a TRAF–cIAP complex.⁷⁸ This ubiquitination-dependent degradation ensures very low basal NIK levels.⁷⁹ Noncanonical stimuli trigger TRAF2-dependent cIAP1–cIAP2 activation, which in turn leads to the proteasomal degradation of TRAF3; this disrupts the cIAP–TRAF complex reducing NIK degradation and leads to accumulation of NIK.^{79–81} NIK activity is dependent on IKK1, but independent of IKK2. Once activated, the NF κ B-inducing-kinase (NIK) complex has dual roles within the noncanonical pathway (Figure 2(a)). Originally identified as a MAP kinase kinase kinase (encoded by *Map3k14*), NIK activates IKK α by phosphorylation of Ser and Thr residues within an activation loop between subdomains VII and VIII of the kinase domain.^{84,85} Although NIK overexpression results in canonical NF κ B activation,^{85,86} NIK knockouts are not defective in the canonical activation of IKK and NF κ B in response to inflammatory cytokines.⁸²

I κ B δ Degradation to Release Preformed Dimers

NIK's first role is in targeting the oligomeric I κ B complex, causing I κ B δ degradation and release of preexisting NF κ B dimers for nuclear localization.²⁶ This initial response relies on phosphorylation events and occurs quickly, as I κ B δ bound to preexisting dimers can release NF κ B to localize to the nucleus as soon as it is modified by NIK (Figure 2(b)). While NIK is primarily considered the transducer of noncanonical NF κ B signaling and I κ B δ differs from other I κ Bs in its ability to inhibit RelB-containing dimers and respond to noncanonical stimuli, I κ B δ also inhibits RelA-containing dimers. Therefore, NIK-induced I κ B δ degradation may induce an inflammatory and/or developmental response depending on the existing NF κ B dimer repertoire poised within the cell prior to stimulation. Both RelB and RelA induction in response to LT β R (noncanonical only) are abrogated in NIK knockout.²⁶

P100 Processing to p52 to Generate New RelB:p52

The second role of NIK is in initiating the proteasome-mediated processing of p100 into p52. Phosphorylation of C-terminal serine residues leads to p100 being recognized by SCF/ β TRCP ubiquitin ligase and subsequent partial degradation of the ARD by the 26S proteasome.^{82,83} This processing produces a mature p52 monomer that is then able to dimerize to form transcriptionally active RelB:p52 and other NF κ B dimers.⁸⁷ It appears that only newly translated p100 is able to be processed into p52,⁸⁸ probably because p100 oligomerizes into a complex that renders it unavailable for processing.²⁶ The multidomain interactions between RelB and p52 (and also p100) co-stabilize both proteins, as RelB

protein levels are decreased in *Nfkb2*^{-/-} cells and p100 protein is decreased in *RelB*^{-/-} cells.^{89,90} While p52 is only produced at very low levels in most mammalian cells, certain cell types, including B cells, show active generation of p52.⁹¹ Processing is tightly controlled by a processing-suppressive region in the C-terminal portion of the p100 protein and disruption of this domain leads to constitutive p100 processing.^{82,92} p100 that does not form p52 is free to form higher-molecular-weight inhibitors of NFκB (IκBδ), and therefore tight regulation is important to ensure production of IκBδ in the absence of noncanonical stimuli to regulate late-phase NFκB activity. NIK-induced p100 processing, and subsequent dimerization with RelB, is a relatively slow process compared to the release of preexisting NFκB dimer from inhibitor, and therefore this results in a late-phase and sustained gene-expression response.

During B-cell maturation, BAFF activates non-canonical NFκB signaling and the consequence of this is dependent on the state of p100 synthesis within the NFκB network: at moderate p100 synthesis rates BAFF-induced p52 production fully depletes p100 and prevents the formation of IκBδ, whereas in the context of high p100 synthesis and resultant IκBδ formation BAFF causes IκBδ degradation altering cRel activity and affecting B-cell expansion.⁶¹ Constitutive P100 degradation also contributes to p100 homeostasis and hence is important for a variety of cellular functions.⁹³

ALTERNATIVE NFκB ACTIVATION MECHANISMS

Ribotoxic Stress

Ribotoxic stress, as induced by ultraviolet light (UV) exposure or by unfolded protein response (UPR) inducers, has been found to activate NFκB by inhibiting translation of IκBα.^{94,95} Thus, these stimuli engage in crosstalk with inflammatory signaling and amplify NFκB activity (Figure 3(a)). Indeed, UV-induced activation of NFκB was shown to occur in enucleated cells, eliminating UV-induced DNA damage as the primary transducer of this pathway. Instead, in response to UV, eukaryotic initiation factor 2 α (eIF2α) is phosphorylated through stress response kinases GCN2/PERK. Phosphorylation of eIF2α broadly inhibits transcription initiation, resulting in reduced synthesis of IκBα.⁹⁴ As free IκBα is constantly degraded and synthesized, the UV-induced reduction in synthesis results in decreased IκBα and amplified NFκB responses (Figure 3(b)). Interestingly, in response to chronic reactive oxygen species exposure, NFκB may repress prosurvival genes and induce prodeath genes.⁹⁷

Genotoxic Stress

Genotoxic stress also activates NFκB, although through distinct mechanisms.⁹⁶ In this context, the initiation signal for NFκB responses originates from within the nucleus and is propagated to the cytosol via mechanisms that involve the nuclear export of upstream signaling molecules. In response to DNA damage NEMO is localized to the nucleus as a result of SUMO-1 attachment.⁹⁸ DNA damage-activated ATM (ataxia telangiectasia mutated) then phosphorylates NEMO in the nucleus and triggers mono-ubiquitination of NEMO. ATM then binds to this modified NEMO, which is exported to the cytoplasm and activates IKK to result in degradation of IκBs. NFκB activation upon genotoxic stress is

markedly slower and lower in amplitude than that observed for immune receptor signaling, and its physiological role remains unclear.

Shear Stress

Mechanical forces exerted on cells can also activate NF κ B. Shear stress (mechanotransduction) activates NF κ B in osteoblasts through phospholipase pathways that release intracellular Ca²⁺. Activated NF κ B in response to shear stress upregulates COX-2, which plays an important role in the response of bone to mechanical loading.⁹⁹ Hemodynamic forces also exert shear stresses on vascular cells during development of atherosclerosis that result in activation of NF κ B through a mechanism independent from canonical I κ B α degradation.¹⁰⁰

NF κ B GENERATION MECHANISMS

NF κ B dimeric transcription factors are formed by five monomers (Figure 4). Of the 15 possible dimers, 12 are thought to bind the DNA κ B element, and 3 (RelB:RelB, RelB:RelA, and RelB:cRel) form low-affinity intertwined dimers that are unable to bind DNA.¹⁰¹ Of the 12 DNA-binding dimers, 9 contain at least one of the activator proteins, RelA, cRel, or RelB (RelA being the most potent and RelB the least), and generally function as transcriptional activators. The remaining three (the abundant p50:p50 homodimer, and the lesser p52:p52 and p50:p52 homodimers and heterodimers, respectively) may function as activators in conjunction with co-activators, including Bcl3 and I κ B ζ .

The mechanisms underlying NF κ B dimer generation (Figure 4) have only recently received attention, but are critical to understanding how different cell types produce different NF κ B dimer repertoires during cell differentiation and development. In some cases, a shift in the dimer repertoire has been documented: for example, B cells shift from a RelA:p50 predominant state at the pre-B stage to a cRel:p50 dominant state in mature B-lymphoid cells and then show strong upregulation of RelB and p52 in terminally differentiated B cells.¹⁰² Also, while most monocyte lineages rely on RelA:p50, GM-CSF-derived inflammatory dendritic cells show high levels of the unusual RelB:p50 dimer, whose generation was shown to be dependent on high constitutive RelB expression and NIK activity.³⁵ These examples show that the NF κ B dimer repertoire is responsive to stimuli, albeit at longer timescales (>8 h) than the activation of NF κ B from the latent dimer repertoire.

NF κ B family members are obligate dimers; as monomers they are unstable and are thought to be quickly degraded. Thus, the NF κ B dimer generation system is highly dynamic and homeostatic. Below, we summarize the key regulatory mechanisms that contribute to the NF κ B dimer repertoire. How these function together to determine the specific NF κ B signaling system of specific cell types ought to be a focus of future increasingly quantitative studies.

Dimerization Affinities

Key determinants of the NF κ B dimer repertoire are the interaction rate constants between the five monomers. Indirect evidence suggests that the 15 NF κ B dimers have dramatically

different dimerization affinities, yet remarkably little quantitative information has been published. Recently, analytical ultracentrifugation determined the affinities of the RelA:p50, p50:p50, and RelA:RelA dimers to be in the range of 1–5 nM, 20–50 nM, and 0.8–1.5 μ M, respectively.¹⁰³ It is not unreasonable to speculate that other dimers fall into these orders of magnitude, with large dimers (RelA:cRel, cRel:cRel and the DNA-binding incompetent dimers RelB:RelA, RelB; RelB, and RelB:cRel) having low affinities close to the μ M range, small dimers (p50:p52, p52:p52) in the high nM range, and dimers composed of one large and one small subunit (RelA:p52, RelB:p50, RelB:p52, cRel:p50, and cRel:p52) having the tightest affinities. However, even if these broad rules prove correct, differences between them are likely to be important in determining the dimer repertoire. Further, it is likely that association and dissociation rate constants, rather than steady-state affinities derived from ratios of these rates, may also be important, as a slow k_{on} rates will lead to a kinetic disadvantage within this potentially competitive dynamical system. If experimental pipelines are established, we expect that posttranslational modifications will likely be found that modulate these important parameters.

Expression of NF κ B Monomer Genes

In order to produce NF κ B dimers, a monomer must be expressed, yet given the interdependence of combinatorial dimerization, expression of all monomers must be considered in order to predict the abundance of a single dimer. NF κ B monomer genes are known to be expressed differentially. RelA is thought to be ubiquitously expressed at high levels, whereas cRel is largely restricted to lymphoid lineages (as well as inflammatory dendritic cells), and RelB is also known to be expressed at high levels in specific cell types. All NF κ B genes are to some degree NF κ B inducible, with cRel, RelB, *Nfkb1*, and *Nfkb2* being long-recognized targets of RelA,^{61,104} and RelA recently reported to be a RelA target as well.¹⁰⁵ However, it is unlikely that autoregulation alone accounts for the cell-type-specific expression patterns and little information is available about other transcription factors that control the expression of NF κ B genes.

Processing of NF κ B Monomer Precursors

Expression of *Nfkb1* and *Nfkb2* leads to the precursor proteins p105 and p100 that must be processed before the p50 and p52 dimerization partners are available. p105 is endoproteolytically cleaved into a mature protein, p50, that is able to dimerize with other NF κ B proteins (predominantly RelA).² p105 processing is generally a constitutive process that occurs in unstimulated cells.¹⁰⁶ It is unclear whether IKK2 may process p105 in a signal-responsive manner.¹⁰⁷

Nfkb2/p100 must be processed into p52 before it can dimerize, predominantly with RelB, to form an NF κ B dimer. In response to developmental signals p52 processing is increased as NIK is activated.⁸² p100 to p52 processing is not only significant as it generates p52 but also as it depletes p100 which would otherwise complex into the inhibitory complex I κ B δ . Competition for binding to RelA and RelB between p50 and p52 also contributes to p100 processing rate as RelB inhibits p100 to p52 processing.⁹⁰ The level of precursor protein controls the maximal signal-responsive induction of protein processing that can be achieved. When elevated, p100 processing depletes the cellular pool of p100 and

noncanonical signaling is unable to strongly induce further p52 production or RelB:p52 formation. Similarly, when all RelB is able to bind to p50 (for example, in *nfkb2*^{-/-} where no p52 is produced), the pool of precursor p105 is depleted.⁹⁰

Monomer Competition During Dimerization

The combinatorial nature of NFκB dimerization implies the potential for competition between monomers for the generation of specific dimers. This competition was first reported with regard to the formation of RelA:p50 versus RelB:p52, where a slightly higher affinity of p105/p50 for RelA reduces the opportunity for p100/p52 from complexing with RelA.¹⁰⁸ Conversely, a slightly higher affinity of p100/p52 for RelB diminishes the potential formation of RelA:p52 dimers. This model explains the appearance of such dimers in the respective knockouts.⁹⁰

More recently, a quantitative analysis of RelA homodimerization and heterodimerization (with p50) revealed the degree to which monomer competition reduces the abundance of low-affinity dimers¹⁰³: the high-affinity RelA:p50 dimer dramatically reduces the abundance of the low-affinity RelA homodimer, whose formation hence becomes entirely dependent on a chaperone. Only when p50 is genetically diminished does RelA homodimer formation become chaperone independent.¹⁰³

Dimer Stabilization and Chaperones

In addition to the combinatorial dimerization affinities and physiochemical properties of the monomers, dimer abundances may be enhanced by ‘third-party’ stabilizers, and dimer generation may be enhanced by dimerization chaperones. To date, one such example has been reported^{103,109}: IκBβ was identified as increasing the effective binding affinity of RelA homodimer.¹⁰³ As mentioned, RelA homodimer levels suffer not only from a poor affinity but also from competition from RelA:p50 dimerization. This effect is counteracted by IκBβ, which is able to increase the binding of the low-affinity RelA homodimer. A quantitative analysis arrives at such a high effective affinity that a two-step model for IκBβ function is plausible, i.e. that IκBβ binds one monomer first and then a second, enhancing the effective association rate constant and rendering it a *bona fide* ‘chaperone.’ However, direct evidence is currently outstanding.

But regardless of whether IκBs function as chaperones or merely stabilizers of specific dimers, their potential role adds a more dynamic and complex component to the control of the cell-type-specific NFκB dimer repertoire, as well as a redefinition of IκBs from being merely inhibitors to also being ‘licensing factors’ of NFκB activity.

Dimer Degradation

NFκB dimers are thought to be stable when associated with IκBs. They neither come apart nor are they degraded. However, while DNA interactions may also stabilize dimerizing interactions, NFκB dimers bound to DNA are thought to be subject to regulated degradation. Though an attractive hypothesis, the mechanisms remain less than clear. RelA was shown to be removed from specific chromatin sites even in the absence of new IκBα feedback synthesis¹¹⁰; RelA removal from chromatin was impaired in IKKα-deficient

macrophages¹¹¹; and the peptidyl-prolyl isomerase Pin1 and the E3 ligases SOCS1¹¹² and COMMD1/Cul2¹¹³ have been implicated in RelA degradation.

SIGNALING CROSSTALK MECHANISMS

While the canonical and noncanonical pathways are mediated by distinct kinases and immediate substrates, given the large number of shared components within the NF κ B system, there is a great potential for crosstalk between the two pathways.

Noncanonical Control of Canonical Signaling

Nfkb2/p100 is the primary signaling node at which canonical and noncanonical signals interact (Figure 5 (a)). That is because (1) *Nfkb2* expression is inducible by RelA, (2) any p100 that is not processed into p52 forms a higher-molecular-weight inhibitor of NF κ B, the I κ B δ -containing I κ Bsome, that may trap NF κ B dimers and thus diminish their association with canonical I κ Bs,^{26,114} and (3) p100 processing to p52 and I κ B δ degradation is triggered by noncanonical NIK activity. As a result, noncanonical signaling may extend the duration of canonical NF κ B activation,⁶¹ or in its absence may diminish canonical NF κ B activation.⁵⁸ Thus, noncanonical pathway activity tunes the potency of the canonical pathway in activating NF κ B (Figure 5(b)). Interestingly, in conditions with chronically elevated noncanonical activity, as in inflammatory dendritic cells, canonical pathway signaling is not only altered quantitatively but also qualitatively: in GM-CSF-derived dendritic cells, high noncanonical pathway activity diminishes p100 to such a degree that the RelB:p50 dimer forms, which then associates with I κ B α (and to some degree I κ B ϵ), rendering it responsive to noncanonical signals.³⁵ Hence, TLR-triggered maturation of these dendritic cells involves activation of RelB:p50, in addition to the expected RelA:p50 and cRel:p50 dimers.

Canonical Control of Noncanonical Signaling

In addition to *Nfkb2*, the *Relb* gene is also induced by NF κ B dimers in response to canonical pathway activity.^{104,115} Therefore, canonical pathway activity is essential for noncanonical activation of the RelB:p52 dimer.⁹⁰ The dependency of noncanonical signaling on canonical activity impacts lymph node development in RelA-deficient mice.⁸⁷

This potential for crosstalk may in principle also allow elevated canonical pathway activity to amplify noncanonical activation of RelB:p52 (Figure 5(b)), with potentially pathologic consequences.¹⁰⁸ However, this does not appear to be the case—yet the mechanism that insulates canonical signaling from high canonical activity remains to be elucidated.

Posttranslational Modifications

Several posttranslational modifications of RelA have been identified, including phosphorylation of serines and threonines in the DNA binding and activation domains, methylation, acetylation, and glycosylation.^{116,117} Mutational analysis of the modification acceptor residues has in many cases been shown to have detrimental effects on proper NF κ B control and target gene expression, and it appears that a variety of regulatory steps (e.g., nuclear localization, DNA binding, and co-activator recruitment) may be affected.

However, in many cases, it remains unclear whether the posttranslational modification occurs constitutively, is induced by the same stimulus as I κ B degradation, or may in fact be induced by a different stimulus or cell-type specifically. Evidence for the latter scenario would support the attractive hypothesis that NF κ B proteins function as integrators of distinct signals (to control nuclear localization and posttranslational modifications) to fine-tune NF κ B-responsive transcriptional programs. Such a scenario would suggest a signaling crosstalk that ought to be examined with a range of biochemical, cell biological, and computational research tools.

SUMMARY

NF κ B has been a key nexus of scientific interest over the past several decades. The breadth and depth of investigation, and the compiled knowledge of this key family of transcription factors are unparalleled. From early insights into its function as a transcription factor and immune response regulator to later studies on cellular signaling, dynamical responses, gene regulatory networks, and feedback mechanisms, it is clear that NF κ B has become a standard bearer for research into the inner workings of cellular communication and signal response paradigms. New work on signaling crosstalk and remaining questions and challenges in the field, an amenability to signaling and information transmission studies informed by computational modeling, as well as NF κ B's continued interest as a therapeutic target ensure a future of extensive interest and publication on the subject. Ongoing work to expand the scope of computational models and extend their applicability from the cellular scale to the tissue scale will require agent-based modeling techniques that may also leverage minimal model formulations.^{118–120} Identification and construction of appropriate minimal models for this purpose depend on identifying the physiologically relevant features within NF κ B's intricate and varied dynamic responses.

REFERENCES

1. Gerondakis S, Grumont R, Gugasyan R, Wong L, Isomura I, Ho W, Banerjee A. Unravelling the complexities of the NF- κ B signalling pathway using mouse knockout and transgenic models. *Oncogene*2006, 25:6781–6799. [PubMed: 17072328]
2. Vallabhapurapu S, Karin M. Regulation and function of NF- κ B transcription factors in the immune system. *Annu Rev Immunol*2009, 27:693–733. [PubMed: 19302050]
3. Hayden MS, Ghosh S. NF- κ B, the first quarter-century: remarkable progress and outstanding questions. *Genes Dev*2012, 26:203–234. [PubMed: 22302935]
4. Hoffmann A, Baltimore D. Circuitry of nuclear factor κ B signaling. *Immunol Rev*2006, 210:171–186. [PubMed: 16623771]
5. Baud V, Karin M. Is NF- κ B a good target for cancer therapy? Hopes and pitfalls. *Nat Rev Drug Discov*2009, 8:33–40. [PubMed: 19116625]
6. Cheong R, Hoffmann A, Levchenko A. Understanding NF- κ B signaling via mathematical modeling. *Mol Syst Biol*2008, 4:192. [PubMed: 18463616]
7. Kearns JD, Hoffmann A. Integrating computational and biochemical studies to explore mechanisms in NF- κ B signaling. *J Biol Chem*2009, 284:5439–5443. [PubMed: 18940809]
8. Basak S, Behar M, Hoffmann A. Lessons from mathematically modeling the NF- κ B pathway. *Immunol Rev*2012, 246:221–238. [PubMed: 22435558]
9. O'Dea E, Hoffmann A. NF- κ B signaling. *WIREs Syst Biol Med*2009, 1:107–115.
10. O'Dea E, Hoffmann A. The regulatory logic of the NF- κ B signaling system. *Cold Spring Harb Perspect Biol*2010, 2:a000216. [PubMed: 20182598]

11. Hoffmann A, Leung TH, Baltimore D. Genetic analysis of NF- κ B/Rel transcription factors defines functional specificities. *EMBO J*2003, 22:5530–5539. [PubMed: 14532125]
12. Leung TH, Hoffmann A, Baltimore D. One nucleotide in a κ B site can determine cofactor specificity for NF- κ B dimers. *Cell*2004, 118:453–464. [PubMed: 15315758]
13. Sanjabi S, Hoffmann A, Liou HC, Baltimore D, Smale ST. Selective requirement for c-Rel during IL-12 P40 gene induction in macrophages. *Proc Natl Acad Sci USA*2000, 97:12705–12710. [PubMed: 11058167]
14. Sanjabi S, Williams KJ, Saccani S, Zhou L, Hoffmann A, Ghosh G, Gerondakis S, Natoli G, Smale ST. A c-Rel subdomain responsible for enhanced DNA-binding affinity and selective gene activation. *Genes Dev*2005, 19:2138–2151. [PubMed: 16166378]
15. Hoffmann A, Natoli G, Ghosh G. Transcriptional regulation via the NF- κ B signaling module. *Oncogene*2006, 25:6706–6716. [PubMed: 17072323]
16. Shih VF, Tsui R, Caldwell A, Hoffmann A. A single NF κ B system for both canonical and non-canonical signaling. *Cell Res*2011, 21:86–102. [PubMed: 21102550]
17. Mukherjee SP, Behar M, Birnbaum HA, Hoffmann A, Wright PE, Ghosh G. Analysis of the RelA:CBP/p300 interaction reveals its involvement in NF- κ B-driven transcription. *PLoS Biol*2013, 11:e1001647. [PubMed: 24019758]
18. Ramirez-Carrozzi VR, Braas D, Bhatt DM, Cheng CS, Hong C, Doty KR, Black JC, Hoffmann A, Carey M, Smale ST. A unifying model for the selective regulation of inducible transcription by CpG islands and nucleosome remodeling. *Cell*2009, 138:114–128. [PubMed: 19596239]
19. Cheng CS, Feldman KE, Lee J, Verma S, Huang DB, Huynh K, Chang M, Ponomarenko JV, Sun SC, Benedict CA, et al. The specificity of innate immune responses is enforced by repression of interferon response elements by NF- κ B p50. *Sci Signal*2011, 4:ra11. [PubMed: 21343618]
20. Huxford T, Hoffmann A, Ghosh G. Understanding the logic of I κ B: NF- κ B regulation in structural terms. In: *NF- κ B in Health and Disease*. Berlin Heidelberg: Springer; 2011, 1–24.
21. Alves BN, Tsui R, Almaden J, Shokhirev MN, Davis-Turak J, Fujimoto J, Birnbaum H, Ponomarenko J, Hoffmann A. I κ B ϵ is a key regulator of B cell expansion by providing negative feedback on cRel and RelA in a stimulus-specific manner. *J Immunol*2014, 192:3121–3132. [PubMed: 24591377]
22. Escoubet-Lozach L, Benner C, Kaikkonen MU, Lozach J, Heinz S, Spann NJ, Crotti A, Stender J, Ghisletti S, Reichart D. Mechanisms establishing TLR4-responsive activation states of inflammatory response genes. *PLoS Genet*2011, 7:e1002401. [PubMed: 22174696]
23. Alkalay I, Yaron A, Hatzubai A, Orian A, Ciechanover A, Ben-Neriah Y. Stimulation-dependent I κ B α phosphorylation marks the NF- κ B inhibitor for degradation via the ubiquitin-proteasome pathway. *Proc Natl Acad Sci USA*1995, 92:10599–10603. [PubMed: 7479848]
24. Liu F, Xia Y, Parker AS, Verma IM. IKK biology. *Immunol Rev*2012, 246:239–253. [PubMed: 22435559]
25. Hoffmann A, Levchenko A, Scott ML, Baltimore D. The I κ B-NF- κ B signaling module: temporal control and selective gene activation. *Science*2002, 298:1241–1245. [PubMed: 12424381]
26. Basak S, Kim H, Kearns JD, Tergaonkar V, O’Dea E, Werner SL, Benedict CA, Ware CF, Ghosh G, Verma IM, et al. A fourth I κ B protein within the NF- κ B signaling module. *Cell*2007, 128:369–381. [PubMed: 17254973]
27. Iwai K. Diverse ubiquitin signaling in NF- κ B activation. *Trends Cell Biol*2012, 22:355–364. [PubMed: 22543051]
28. Adhikari A, Xu M, Chen Z. Ubiquitin-mediated activation of TAK1 and IKK. *Oncogene*2007, 26:3214–3226. [PubMed: 17496917]
29. Cheong R, Bergmann A, Werner SL, Regal J, Hoffmann A, Levchenko A. Transient I κ B kinase activity mediates temporal NF- κ B dynamics in response to a wide range of tumor necrosis factor- α doses. *J Biol Chem*2006, 281:2945–2950. [PubMed: 16321974]
30. Shinohara H, Behar M, Inoue K, Hiroshima M, Yasuda T, Nagashima T, Kimura S, Sanjo H, Maeda S, Yumoto N, et al. Positive feedback within a kinase signaling complex functions as a switch mechanism for NF- κ B activation. *Science*2014, 344:760–764. [PubMed: 24833394]
31. Niu J, Shi Y, Iwai K, Wu ZH. LUBAC regulates NF- κ B activation upon genotoxic stress by promoting linear ubiquitination of NEMO. *EMBO J*2011, 30:3741–3753. [PubMed: 21811235]

32. Werner SL, Barken D, Hoffmann A. Stimulus specificity of gene expression programs determined by temporal control of IKK activity. *Science*2005, 309:1857–1861. [PubMed: 16166517]
33. Chen L-F, Greene WC. Shaping the nuclear action of NF- κ B. *Nat Rev Mol Cell Biol*2004, 5:392–401. [PubMed: 15122352]
34. Schröfelbauer B, Polley S, Behar M, Ghosh G, Hoffmann A. NEMO ensures signaling specificity of the pleiotropic IKK β by directing its kinase activity toward I κ B α . *Mol Cell*2012, 47:111–121. [PubMed: 22633953]
35. Shih VF, Davis-Turak J, Macal M, Huang JQ, Ponomarenko J, Kearns JD, Yu T, Fagerlund R, Asagiri M, Zuniga EI. Control of RelB during dendritic cell activation integrates canonical and noncanonical NF- κ B pathways. *Nat Immunol*2012, 13:1162–1170. [PubMed: 23086447]
36. Xu G, Lo Y-C, Li Q, Napolitano G, Wu X, Jiang X, Dreano M, Karin M, Wu H. Crystal structure of inhibitor of κ B kinase β . *Nature*2011, 472:325–330. [PubMed: 21423167]
37. Polley S, Huang D-B, Hauenstein AV, Fusco AJ, Zhong X, Vu D, Schröfelbauer B, Kim Y, Hoffmann A, Verma IM. A structural basis for I κ B kinase 2 activation via oligomerization-dependent trans auto-phosphorylation. *PLoS Biol*2013, 11: e1001581. [PubMed: 23776406]
38. Cheng Z, Taylor B, Ourthiague DR, Hoffmann A. Distinct single-cell signaling characteristics are conferred by the MyD88 and TRIF pathways during TLR4 activation. *Sci Signal*2015, 8:ra69. [PubMed: 26175492]
39. Behar M, Hoffmann A. Tunable signal processing through a kinase control cycle: the IKK signaling node. *Biophys J*2013, 105:231–241. [PubMed: 23823243]
40. Delhase M, Hayakawa M, Chen Y, Karin M. Positive and negative regulation of I κ B kinase activity through IKK β subunit phosphorylation. *Science*1999, 284:309–313. [PubMed: 10195894]
41. Schröfelbauer B, Hoffmann A. How do pleiotropic kinase hubs mediate specific signaling by TNFR superfamily members? *Immunol Rev*2011, 244:29–43. [PubMed: 22017429]
42. Werner SL, Kearns JD, Zadorozhnaya V, Lynch C, O’Dea E, Boldin MP, Ma A, Baltimore D, Hoffmann A. Encoding NF- κ B temporal control in response to TNF: distinct roles for the negative regulators I κ B α and A20. *Genes Dev*2008, 22:2093–2101. [PubMed: 18676814]
43. Mitchell S, Tsui R, Hoffmann A. Studying NF- κ B signaling with mathematical models. *Methods Mol Biol*2015, 1280:647–661. [PubMed: 25736777]
44. Fagerlund R, Behar M, Fortmann KT, Lin YE, Vargas JD, Hoffmann A. Anatomy of a negative feedback loop: the case of I κ B α . *J R Soc Interface*2015, 12:0262. [PubMed: 26311312]
45. Sachdev S, Hoffmann A, Hannink M. Nuclear localization of I κ B α is mediated by the second ankyrin repeat: the I κ B α ankyrin repeats define a novel class of cis-acting nuclear import sequences. *Mol Cell Biol*1998, 18:2524–2534. [PubMed: 9566872]
46. Mathes E, O’Dea EL, Hoffmann A, Ghosh G. NF- κ B dictates the degradation pathway of I κ B α . *EMBO J*2008, 27:1357–1367. [PubMed: 18401342]
47. O’Dea EL, Barken D, Peralta RQ, Tran KT, Werner SL, Kearns JD, Levchenko A, Hoffmann A. A homeostatic model of I κ B metabolism to control constitutive NF- κ B activity. *Mol Syst Biol*2007, 3:111. [PubMed: 17486138]
48. Fortmann KT, Lewis RD, Ngo KA, Fagerlund R, Hoffmann A. A regulated, ubiquitin-independent degron in I κ B α . *J Mol Biol*2015, 427:2748–2756. [PubMed: 26191773]
49. Wuerzberger-Davis SM, Chen Y, Yang DT, Kearns JD, Bates PW, Lynch C, Ladell NC, Yu M, Podd A, Zeng H. Nuclear export of the NF- κ B inhibitor I κ B α is required for proper B cell and secondary lymphoid tissue formation. *Immunity*2011, 34:188–200. [PubMed: 21333553]
50. Mulero MC, Ferres-Marco D, Islam A, Margalef P, Pecoraro M, Toll A, Drechsel N, Charneco C, Davis S, Bellora N. Chromatin-bound I κ B α regulates a subset of polycomb target genes in differentiation and cancer. *Cancer Cell*2013, 24:151–166. [PubMed: 23850221]
51. Bergqvist S, Alverdi V, Mengel B, Hoffmann A, Ghosh G, Komives EA. Kinetic enhancement of NF- κ B[C1]DNA dissociation by I κ B α . *Proc Natl Acad Sci USA*2009, 106:19328–19333. [PubMed: 19887633]
52. Nelson D, Ihekwa A, Elliott M, Johnson J, Gibney C, Foreman B, Nelson G, See V, Horton C, Spiller D. Oscillations in NF- κ B signaling control the dynamics of gene expression. *Science*2004, 306:704–708. [PubMed: 15499023]

53. Regot S, Hughey JJ, Bajar BT, Carrasco S, Covert MW. High-sensitivity measurements of multiple kinase activities in live single cells. *Cell*2014, 157:1724–1734. [PubMed: 24949979]
54. Sung M-H, Bagain L, Chen Z, Karpova T, Yang X, Silvin C, Voss TC, McNally JG, Van Waes C, Hager GL. Dynamic effect of bortezomib on nuclear factor- κ B activity and gene expression in tumor cells. *Mol Pharmacol*2008, 74:1215–1222. [PubMed: 18684839]
55. Kellogg RA, Tay S. Noise facilitates transcriptional control under dynamic inputs. *Cell*2015, 160:381–392. [PubMed: 25635454]
56. Selimkhanov J, Taylor B, Yao J, Pilko A, Albeck J, Hoffmann A, Tsimring L, Wollman R. Accurate information transmission through dynamic biochemical signaling networks. *Science*2014, 346:1370–1373. [PubMed: 25504722]
57. Behar M, Hoffmann A. Understanding the temporal codes of intra-cellular signals. *Curr Opin Genet Dev*2010, 20:684–693. [PubMed: 20956081]
58. Shih VF-S, Kearns JD, Basak S, Savinova OV, Ghosh G, Hoffmann A. Kinetic control of negative feedback regulators of NF- κ B/RelA determines their pathogen-and cytokine-receptor signaling specificity. *Proc Natl Acad Sci USA*2009, 106:9619–9624. [PubMed: 19487661]
59. Savinova OV, Hoffmann A, Ghosh G. The Nfkb1 and Nfkb2 proteins p105 and p100 function as the core of high-molecular-weight heterogeneous complexes. *Mol Cell*2009, 34:591–602. [PubMed: 19524538]
60. Banoth B, Chatterjee B, Vijayaragavan B, Prasad MV, Roy P, Basak S. Stimulus-selective crosstalk via the NF- κ B signaling system reinforces innate immune response to alleviate gut infection. *Elife*2015, 4: eLife.05648.
61. Almaden JV, Tsui R, Liu YC, Birnbaum H, Shokhirev MN, Ngo KA, Davis-Turak JC, Otero D, Basak S, Rickert RC. A pathway switch directs BAFF signaling to distinct NF κ B transcription factors in maturing and proliferating B cells. *Cell Rep*2014, 9:2098–2111. [PubMed: 25497099]
62. Longo DM, Selimkhanov J, Kearns JD, Hasty J, Hoffmann A, Tsimring LS. Dual delayed feedback provides sensitivity and robustness to the NF- κ B signaling module. *PLoS Comput Biol*2013, 9:e1003112. [PubMed: 23825938]
63. Keams J, Basak S, Wemer S, Huang C, Hoffmann A. I κ Be provides negative feedback to control NF- κ B oscillations, signaling dynamics, and inflammatory gene expression. *J Cell Biol*2006, 173:659–664. [PubMed: 16735576]
64. Malek S, Chen Y, Huxford T, Ghosh G. I κ B β , but not I κ B α , functions as a classical cytoplasmic inhibitor of NF- κ B dimers by masking both NF- κ B nuclear localization sequences in resting cells. *J Biol Chem*2001, 276:45225–45235. [PubMed: 11571291]
65. Caldwell AB, Cheng Z, Vargas JD, Birnbaum HA, Hoffmann A. Network dynamics determine the autocrine and paracrine signaling functions of TNF. *Genes Dev*2014, 28:2120–2133. [PubMed: 25274725]
66. Davis-Turak JC, Allison K, Shokhirev MN, Ponomarenko P, Tsimring LS, Glass CK, Johnson TL, Hoffmann A. Considering the kinetics of mRNA synthesis in the analysis of the genome and epigenome reveals determinants of co-transcriptional splicing. *Nucleic Acids Res*2015, 43:699–707. [PubMed: 25541195]
67. Coope H, Atkinson P, Huhse B, Belich M, Janzen J, Holman M, Klaus G, Johnston L, Ley S. CD40 regulates the processing of NF- κ B2 p100 to p52. *EMBO J*2002, 21:5375–5385. [PubMed: 12374738]
68. Claudio E, Brown K, Park S, Wang H, Siebenlist U. BAFF-induced NEMO-independent processing of NF- κ B2 in maturing B cells. *Nat Immunol*2002, 3:958–965. [PubMed: 12352969]
69. Dejardin E, Droin NM, Delhase M, Haas E, Cao Y, Makris C, Li Z-W, Karin M, Ware CF, Green DR. The lymphotoxin- β receptor induces different patterns of gene expression via two NF- κ B pathways. *Immunity*2002, 17:525–535. [PubMed: 12387745]
70. Kayagaki N, Yan M, Seshasayee D, Wang H, Lee W, French DM, Grewal IS, Cochran AG, Gordon NC, Yin J. BAFF/BLyS receptor 3 binds the B cell survival factor BAFF ligand through a discrete surface loop and promotes processing of NF- κ B2. *Immunity*2002, 17:515–524. [PubMed: 12387744]

71. Novack DV, Yin L, Hagen-Stapleton A, Schreiber RD, Goeddel DV, Ross FP, Teitelbaum SL. The I κ B function of NF- κ B2 p100 controls stimulated osteoclastogenesis. *J Exp Med*2003, 198:771–781. [PubMed: 12939342]
72. Munroe ME, Bishop GA. Role of tumor necrosis factor (TNF) receptor-associated factor 2 (TRAF2) in distinct and overlapping CD40 and TNF receptor 2/CD120b-mediated B lymphocyte activation. *J Biol Chem*2004, 279:53222–53231. [PubMed: 15485859]
73. Saitoh T, Nakayama M, Nakano H, Yagita H, Yamamoto N, Yamaoka S. TWEAK induces NF- κ B2 p100 processing and long lasting NF- κ B activation. *J Biol Chem*2003, 278:36005–36012. [PubMed: 12840022]
74. Wicovsky A, Salzman S, Roos C, Ehrenschrwender M, Rosenthal T, Siegmund D, Henkler F, Gohlke F, Kneitz C, Wajant H. TNF-like weak inducer of apoptosis inhibits proinflammatory TNF receptor-1 signaling. *Cell Death Differ*2009, 16:1445–1459. [PubMed: 19557010]
75. DeJardin E The alternative NF- κ B pathway from biochemistry to biology: pitfalls and promises for future drug development. *Biochem Pharmacol*2006, 72:1161–1179. [PubMed: 16970925]
76. Bonizzi G, Bebién M, Otero DC, Johnson-Vroom KE, Cao Y, Vu D, Jegga AG, Aronow BJ, Ghosh G, Rickert RC, et al. Activation of IKK α target genes depends on recognition of specific κ B binding sites by RelB:p52 dimers. *EMBO J*2004, 23:4202–4210. [PubMed: 15470505]
77. Wang VY-F, Huang W, Asagiri M, Spann N, Hoffmann A, Glass C, Ghosh G. The transcriptional specificity of NF- κ B dimers is coded within the κ B DNA response elements. *Cell Rep*2012, 2:824–839. [PubMed: 23063365]
78. Sun S-C. Non-canonical NF- κ B signaling pathway. *Cell Res*2011, 21:71–85. [PubMed: 21173796]
79. Liao G, Zhang M, Harhaj EW, Sun S-C. Regulation of the NF- κ B-inducing kinase by tumor necrosis factor receptor-associated factor 3-induced degradation. *J Biol Chem*2004, 279:26243–26250. [PubMed: 15084608]
80. He JQ, Zarnegar B, Oganessian G, Saha SK, Yamazaki S, Doyle SE, Dempsey PW, Cheng G. Rescue of TRAF3-null mice by p100 NF- κ B deficiency. *J Exp Med*2006, 203:2413–2418. [PubMed: 17015635]
81. Vallabhapurapu S, Matsuzawa A, Zhang W, Tseng P-H, Keats JJ, Wang H, Vignali DA, Bergsagel PL, Karin M. Nonredundant and complementary functions of TRAF2 and TRAF3 in a ubiquitination cascade that activates NIK-dependent alternative NF- κ B signaling. *Nat Immunol*2008, 9:1364–1370. [PubMed: 18997792]
82. Xiao G, Harhaj EW, Sun S-C. NF- κ B-inducing kinase regulates the processing of NF- κ B2 p100. *Mol Cell*2001, 7:401–409. [PubMed: 11239468]
83. Senftleben U, Li Z-W, Baud V, Karin M. IKK β is essential for protecting T cells from TNF α -induced apoptosis. *Immunity*2001, 14:217–230. [PubMed: 11290332]
84. Shinkura R, Kitada K, Matsuda F, Tashiro K, Ikuta K, Suzuki M, Kogishi K, Serikawa T, Honjo T. Alympho-plasia is caused by a point mutation in the mouse gene encoding NF- κ B-inducing kinase. *Nat Genet*1999, 22:74–77. [PubMed: 10319865]
85. Ling L, Cao Z, Goeddel DV. NF- κ B-inducing kinase activates IKK- α by phosphorylation of Ser-176. *Proc Natl Acad Sci USA*1998, 95:3792–3797. [PubMed: 9520446]
86. Lallena M-J, Diaz-Meco MT, Bren G, Payá CV, Moscat J. Activation of I κ B kinase β by protein kinase C isoforms. *Mol Cell Biol*1999, 19:2180–2188. [PubMed: 10022904]
87. Lo JC, Basak S, James ES, Quiambo RS, Kinsella MC, Alegre ML, Weih F, Franzoso G, Hoffmann A, Fu YX. Coordination between NF- κ B family members p50 and p52 is essential for mediating LT β R signals in the development and organization of secondary lymphoid tissues. *Blood*2006, 107:1048–1055. [PubMed: 16195333]
88. Mordmüller B, Krappmann D, Esen M, Wegener E, Scheidereit C. Lymphotoxin and lipopolysaccharide induce NF- κ B-p52 generation by a co-translational mechanism. *EMBO Rep*2003, 4:82–87. [PubMed: 12524526]
89. Fusco AJ, Savinova OV, Talwar R, Kearns JD, Hoffmann A, Ghosh G. Stabilization of RelB requires multidomain interactions with p100/p52. *J Biol Chem*2008, 283:12324–12332. [PubMed: 18321863]
90. Basak S, Shih VF, Hoffmann A. Generation and activation of multiple dimeric transcription factors within the NF- κ B signaling system. *Mol Cell Biol*2008, 28:3139–3150. [PubMed: 18299388]

91. Solan NJ, Miyoshi H, Carmona EM, Bren GD, Paya CV. RelB cellular regulation and transcriptional activity are regulated by p100. *J Biol Chem*2002, 277:1405–1418. [PubMed: 11687592]
92. Liao G, Sun S-C. Regulation of NF- κ B2/p100 processing by its nuclear shuttling. *Oncogene*2003, 22:4868–4874. [PubMed: 12894228]
93. Busino L, Millman SE, Scotto L, Kyratsous CA, Basur V, O'Connor O, Hoffmann A, Elenitoba-Johnson KS, Pagano M. Fbxw7 α - and GSK3-mediated degradation of p100 is a pro-survival mechanism in multiple myeloma. *Nat Cell Biol*2012, 14:375–385. [PubMed: 22388891]
94. O'Dea EL, Kearns JD, Hoffmann A. UV as an amplifier rather than inducer of NF- κ B activity. *Mol Cell*2008, 30:632–641. [PubMed: 18538661]
95. Tam AB, Mercado EL, Hoffmann A, Niwa M. ER stress activates NF- κ B by integrating functions of basal IKK activity, IRE1 and PERK. *PLoS One*2012, 7: e45078. [PubMed: 23110043]
96. Mabb AM, Wuerzberger-Davis SM, Miyamoto S. PIASy mediates NEMO sumoylation and NF- κ B activation in response to genotoxic stress. *Nat Cell Biol*2006, 8:986–993. [PubMed: 16906147]
97. Ho JQ, Asagiri M, Hoffmann A, Ghosh G. NF- κ B potentiates caspase independent hydrogen peroxide induced cell death. *PLoS One*2011, 6:e16815. [PubMed: 21347231]
98. Huang TT, Wuerzberger-Davis SM, Wu Z-H, Miyamoto S. Sequential modification of NEMO/IKK γ by SUMO-1 and ubiquitin mediates NF- κ B activation by genotoxic stress. *Cell*2003, 115:565–576. [PubMed: 14651848]
99. Chen NX, Geist DJ, Genetos DC, Pavalko FM, Duncan RL. Fluid shear-induced NF κ B translocation in osteoblasts is mediated by intracellular calcium release. *Bone*2003, 33:399–410. [PubMed: 13678782]
100. Ganguli A, Persson L, Palmer IR, Evans I, Yang L, Smallwood R, Black R, Qwarnstrom EE. Distinct NF- κ B regulation by shear stress through Ras-dependent I κ B α oscillations: real-time analysis of flow-mediated activation in live cells. *Circ Res*2005, 96:626–634. [PubMed: 15731464]
101. Huang D-B, Vu D, Ghosh G. NF- κ B RelB forms an intertwined homodimer. *Structure*2005, 13:1365–1373. [PubMed: 16154093]
102. Guo S, Liu M, Gonzalez-Perez RR. Role of Notch and its oncogenic signaling crosstalk in breast cancer. *Biochim Biophys Acta*2011, 1815:197–213. [PubMed: 21193018]
103. Tsui R, Kearns JD, Lynch C, Vu D, Ngo KA, Basak S, Ghosh G, Hoffmann A. I κ B β enhances the generation of the low-affinity NF κ B/RelA homodimer. *Nat Commun*2015, 6:7068. [PubMed: 25946967]
104. Bren GD, Solan NJ, Miyoshi H, Pennington KN, Pobst LJ, Paya CV. Transcription of the RelB gene is regulated by NF- κ B. *Oncogene*2001, 20:7722–7733. [PubMed: 11753650]
105. Sung M-H, Li N, Lao Q, Gottschalk RA, Hager GL, Fraser ID. Switching of the relative dominance between feedback mechanisms in lipopolysaccharide-induced NF- κ B signaling. *Sci Signal*2014, 7:ra6. [PubMed: 24425788]
106. Beinke S, Ley S. Functions of NF- κ B1 and NF- κ B2 in immune cell biology. *Biochem J*2004, 382:393–409. [PubMed: 15214841]
107. Orián A, Gonen H, Bercovich B, Fajerman I, Eytan E, Israël A, Mercurio F, Iwai K, Schwartz AL, Ciechanover A. SCF β -TrCP ubiquitin ligase-mediated processing of NF- κ B p105 requires phosphorylation of its C-terminus by I κ B kinase. *EMBO J*2000, 19:2580–2591. [PubMed: 10835356]
108. Basak S, Hoffmann A. Crosstalk via the NF- κ B signaling system. *Cytokine Growth Factor Rev*2008, 19:187–197. [PubMed: 18515173]
109. Rao P, Hayden MS, Long M, Scott ML, West AP, Zhang D, Oeckinghaus A, Lynch C, Hoffmann A, Baltimore D, et al. I κ B β acts to inhibit and activate gene expression during the inflammatory response. *Nature*2010, 466:1115–1119. [PubMed: 20740013]
110. Saccani S, Marazzi I, Beg AA, Natoli G. Degradation of promoter-bound p65/RelA is essential for the prompt termination of the nuclear factor κ B response. *J Exp Med*2004, 200:107–113. [PubMed: 15226358]

111. Lawrence T, Bebién M, Liu GY, Nizet V, Karin M. IKK α limits macrophage NF- κ B activation and contributes to the resolution of inflammation. *Nature*2005, 434:1138–1143. [PubMed: 15858576]
112. Ryo A, Suizu F, Yoshida Y, Perrem K, Liou YC, Wulf G, Rottapel R, Yamaoka S, Lu KP. Regulation of NF- κ B signaling by Pin1-dependent prolyl isomerization and ubiquitin-mediated proteolysis of p65/RelA. *Mol Cell*2003, 12:1413–1426. [PubMed: 14690596]
113. Mao X, Gluck N, Li D, Maine GN, Li H, Zaidi IW, Repaka A, Mayo MW, Burstein E. GCN5 is a required cofactor for a ubiquitin ligase that targets NF- κ B/RelA. *Genes Dev*2009, 23:849–861. [PubMed: 19339690]
114. Mercurio F, DiDonato J, Rosette C, Karin M. p105 and p98 precursor proteins play an active role in NF- κ B-mediated signal transduction. *Genes Dev*1993, 7:705–718. [PubMed: 8458581]
115. Liptay S, Schmid RM, Nabel EG, Nabel GJ. Transcriptional regulation of NF- κ B2: evidence for κ B-mediated positive and negative autoregulation. *Mol Cell Biol*1994, 14:7695–7703. [PubMed: 7969113]
116. Huang B, Yang XD, Lamb A, Chen LF. Posttranslational modifications of NF- κ B: another layer of regulation for NF- κ B signaling pathway. *Cell Signal*2010, 22:1282–1290. [PubMed: 20363318]
117. Lu T, Stark GR. NF- κ B: regulation by methylation. *Cancer Res*2015, 75:3692–3695. [PubMed: 26337909]
118. Williams R, Timmis J, Qwarnstrom E. Computational models of the NF-KB signalling pathway. *Computation*2014, 2:131.
119. Shokhirev MN, Almaden J, Davis-Turak J, Birnbaum HA, Russell TM, Vargas JA, Hoffmann A. A multi-scale approach reveals that NF- κ B cRel enforces a B-cell decision to divide. *Mol Syst Biol*2015, 11:783. [PubMed: 25680807]
120. Fallahi-Sichani M, Kirschner DE, Linderman JJ. NF- κ B signaling dynamics play a key role in infection control in tuberculosis. *Front Physiol*2012, 3:170. [PubMed: 22685435]

FURTHER READING

- Hayden MS, Ghosh S. Signaling to NF- κ B. *Genes Dev*2004, 18:2195–2224. [PubMed: 15371334]
- Verstrepen L, Bekaert T, Chau TL, Tavernier J, Chariot A, Beyaert R. TLR-4, IL-1R and TNF-R signaling to NF- κ B: variations on a common theme. *Cell Mol Life Sci*2008, 65:2964–2978. [PubMed: 18535784]




FIGURE 1 |

The canonical nuclear factor κ B (NF κ B) activation pathway. (a) Schematic depiction of the canonical NF κ B signaling pathway. Multiple inflammatory signals activate the complex containing NEMO and IKK1/2. IKK1/2 phosphorylates NF κ B-bound I κ Bs, targeting them for ubiquitination and proteasomal degradation.^{24,25} Free I κ Bs also undergo constitutive degradation via a ubiquitin-independent proteasomal degradation pathway. As I κ Bs are degraded, free NF κ B is then able to translocate to the nucleus where it binds to κ B sites on DNA and activates gene expression.^{21,22} I κ B α , β , and ϵ are themselves NF κ B target genes, along with p100 that can form higher-molecular-weight complexes that inhibit NF κ B.^{25,26} (b) Diagram of the regulatory logic of the canonical NF κ B signaling network. Canonical signals activate NEMO/IKK, downregulating I κ Bs and reducing inhibition of NF κ B. Free NF κ B then translocates to the nucleus where it upregulates I κ Bs and p100 and in turn I κ B δ .




FIGURE 2 |

The noncanonical nuclear factor κ B (NFκB) activation pathway. (a) Schematic depiction of the noncanonical NFκB signaling pathway. Developmental signals activate the NIK/IKK1 complex that phosphorylates p100. Most p100 is found in a higher-molecular-weight inhibitory complex (IκBδ). Upon phosphorylation, p100 is processed into p52 and is then available to bind RelB, creating a dimer that localizes to the nucleus and binds DNA to activate transcription.^{82,83} Active NIK/IKK1 complex also phosphorylates the p100 within IκBδ, resulting in its partial degradation and releasing bound NFκB dimers for nuclear localization and gene activation.²⁶ (b) Diagram of the regulatory logic of the noncanonical NFκB signaling network. Noncanonical signals activate NIK/IKK1 that suppresses IκBδ and activates processing of p100 into p52. The suppression of IκBδ that was sequestering preexisting NFκB dimers results in nuclear localization of NFκB and early-phase gene

expression. NIK-dependent p100 processing results in p52 production and the formation of new RelB:p52 dimers that can activate a late-phase gene expression response.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

reduces the inhibition of NF κ B and increases nuclear NF κ B and target gene expression. In response to DNA damage, ATM is upregulated and activates NEMO through a complex with ELKS. Increased NEMO/IKK2 activation results in suppression of I κ Bs.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




FIGURE 4 |

Mechanisms regulating nuclear factor κ B (NF κ B) dimer generation. (a) Diagram of NF κ B monomer synthesis and processing.^{10,20} All NF κ B monomers and precursors are NF κ B target genes and induced, to varying extents, by NF κ B. RelA/RelB/cRel polypeptides are synthesized in a complete form, ready to dimerize into functional NF κ B dimers.^{61,104} p105 is a precursor to p50 that must be cleaved in a process thought to be dependent on IKK2.² p100 must be processed via a NIK/IKK1-dependent pathway into mature p52 before it can dimerize into NF κ B.⁸² (b) Schematic of the NF κ B dimerization process. Monomers must dimerize before they are transcriptionally active. The affinity of binding between monomers varies with two large, activation domain proteins having low affinity. I κ B β can act as a chaperone, enhancing the effective binding affinity of RelA to form homodimer by stabilizing this normally weak affinity dimer. (c) Table of the combinatorial composition of potential NF κ B dimers, indicating their capacities to bind DNA (indicated by horizontal

line) and to activate transcription (indicated by arrows). (d) Diagram of IKK's multiple points of control over NF κ B dimer formation. (1) The IKK kinases upregulates monomer expression by activating NF κ B-responsive promoters. (2) IKK1 and IKK2 activities promote processing of p100 to p52 and p105 to p50. (3) IKKs lead to the degradation of I κ Bs that may function as dimerization chaperones (as for example I κ B β for RelA homodimer) as well as inhibitors.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




FIGURE 5 |

Signaling crosstalk between canonical and noncanonical pathways. (a) Diagram of dual roles of Nfkb2/p100 and NIK within the noncanonical pathway that together with the inducible expression of Nfkb2/p100 mediate two crosstalk functions. (1) NIK/IKK1 processes p100 into p52, enabling the activity of RelB.⁸² (2) NIK degrades IκBδ, allowing for sustained RelA activity.⁶¹ (b) Canonical pathway activity may boost noncanonical pathway activation of RelB:p52.⁹⁰ Novel model simulations that illustrate how noncanonical pathway activation of RelB:p52 may be boosted by increasing constitutive canonical pathway activities. (c) A noncanonical pathway stimulus may prolong canonical pathway-induced NFκB activity. In B cells, BAFF may potentiate late IgM-induced cRel⁶¹ activity.