

# Carbapenem-Resistant *Klebsiella pneumoniae* in Southwest China: Molecular Characteristics and Risk Factors Caused by KPC and NDM Producers

Zhaoyinqian Li\*  
Zixuan Ding\*  
Jia Yang\*  
Yao Liu  
Xinrui Jin  
Jingling Xie  
Tingting Li  
Yinhuan Ding  
Zhangrui Zeng  
Jinbo Liu 

Department of Laboratory Medicine, The Affiliated Hospital of Southwest Medical University, Luzhou, 646000, Sichuan, People's Republic of China

\*These authors contributed equally to this work

**Background:** Carbapenem-resistant *Klebsiella pneumoniae* (CRKP) infection has attracted worldwide concern and became a serious challenge for clinical treatment. The aims of this study were to evaluate the molecular characteristics and risk factors for CRKP infection.

**Methods:** All the CRKP strains were screened for antimicrobial resistance genes, virulence genes, and integron by polymerase chain reaction (PCR). Plasmid typing was performed by plasmid conjugation assay and PCR-based replicon typing (PBRT). The genetic environments of *bla*<sub>KPC-2</sub> and *bla*<sub>NDM-1</sub> were analyzed by using overlapping PCR and molecular typing was performed by multi-locus sequence typing (MLST). Risk factors for CRKP infection were analyzed by logistic regression model.

**Results:** All the 66 CRKP isolates were multidrug-resistant, but all of them were susceptible to tetracycline and polymyxin B. Among the CRKP isolates, 42 *bla*<sub>KPC-2</sub>-positive strains were identified carrying IncFII plasmids. Meanwhile, 24 *bla*<sub>NDM</sub>-positive strains were found on IncX3 plasmids, including 20 *bla*<sub>NDM-1</sub> isolates and 4 *bla*<sub>NDM-5</sub> isolates. Most of CRKP isolates contained several virulence genes and the class I integron (*intI1*). The genetic environments of *bla*<sub>KPC-2</sub> and *bla*<sub>NDM-1</sub> revealed that the conserved regions (*tnpA-tnpR-IS kpn8-bla*<sub>KPC-2</sub>) and (*bla*<sub>NDM-1</sub>-*ble*<sub>MBL</sub>-*trpF-tat*) were associated with the dissemination of KPC-2 and NDM-1. ST11 was the most common type in this work. Hematological disease, tracheal cannula, and use of  $\beta$ -lactams and  $\beta$ -lactamase inhibitor combination were identified as independent risk factors for CRKP infection.

**Conclusion:** This study established the resistance pattern, molecular characteristics, clonal relatedness, and risk factors of CRKP infection. The findings of the novel strain that co-harboring *bla*<sub>NDM-5</sub> and *bla*<sub>IMP-4</sub>, and the novel ST4495 indicated that the brand-new types have spread in Southwest China, emphasizing the prevent and control the further dissemination of CRKP isolates are highly needed.

**Keywords:** carbapenem-resistant *Klebsiella pneumoniae*, molecular characteristics, genetic environments, plasmid, risk factors

## Introduction

*Klebsiella pneumoniae*, belonging to the family *Enterobacteriaceae*, is an important pathogen that causes opportunistic infections in hospitalized patients, and plays a primary role in pneumonia and neonatal sepsis.<sup>1</sup> With the increase of prevalence of multidrug-resistant (MDR) strains that producing extended-spectrum  $\beta$ -lactamase (ESBL) and/or carbapenemase, *K. pneumoniae* has emerged as a major threat in clinical and public health. Carbapenem-resistant *K. pneumoniae* (CRKP) has become

Correspondence: Jinbo Liu  
Department of Laboratory Medicine, The Affiliated Hospital of Southwest Medical University, Luzhou, 646000, Sichuan, People's Republic of China  
Tel/Fax +86 830 3165730  
Email Liulab2019@163.com



USA) and a blank disk with 1.5mg/mL EDTA were placed 10 mm between each disk on the plate. After 18 h incubation, an enlarged zone of inhibition was considered as EDTA-synergy test positive.<sup>18,19</sup>

The CIM was performed as described previously.<sup>18,20</sup> A single colony of CRKP isolates was inoculated into a tube containing 2 mL trypticase soy broth (TSB) (Haibo, Qingdao, China) and a tube containing 2 mL TSB with 5mM EDTA, respectively. A 10 µg meropenem disk was placed in each tube. After 4 h incubation, these disks were taken out and placed on a MH agar plate that was inoculated with a lawn of the meropenem-susceptible *Escherichia coli* ATCC25922 (0.5 McFarland standard). The results were interpreted according to CLSI.<sup>18</sup>

## The Detection of Resistance Genes, Virulence Genes, and Integrase-Associated Genes

All of template DNAs were extracted by bacterial DNA Kit (Tiangen, Beijing, China) and used to detect carbapenemase genes, ESBL genes, AmpC β-lactamase genes, virulence genes, and integrase-associated genes by polymerase chain reaction (PCR). The primers were shown in [Tables S1](#) and [S2](#). The PCR conditions were described previously.<sup>21–23</sup> Positive products were sequenced by Shanghai Jieli Biotechnology, and the variable region of the class I integron (*intI1*) and the insertion sequence common region I (ISCR I) were analyzed by BLAST (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>).

## Plasmid Conjugation and Analysis

In order to evaluate whether the carbapenemase genes are located on the plasmid and whether these genes can be horizontally transferred, a plasmid conjugation transfer experiment was performed. Four hundred µL donor strain (CRKP strains) and 200 µL receptor strain (sodium azide-resistant *E. coli* strain J53) at logarithmic phase were added into a glass tube containing 800 µL Luria-Bertani (LB) broth and cultured at 37°C for 18 h. Transconjugants were selected by using LB plates that contains sodium azide (180 µg/mL) and imipenem (0.5 µg/mL).<sup>21</sup> The transconjugants were verified by MALDI-TOF MS, and the conjugative carbapenemase genes were confirmed by PCR. In addition, the successful conjugative plasmids were analyzed by PCR-based replicon typing according to previous study.<sup>22</sup>

## Genetic Environments of KPC-2-Carrying Plasmids and NDM-1-Carrying Plasmids

The overlapping PCR was applied to investigate the genetic environments of the *bla*<sub>KPC-2</sub> and *bla*<sub>NDM-1</sub> ([Tables S3](#) and [S4](#)).<sup>23,24</sup> The PCR products were sequenced by Shanghai Jieli Biotechnology, and then analyzed by NCBI GenBank database.<sup>25</sup>

## Molecular Epidemiological Study

Multi-locus sequence typing (MLST) was performed to explore the genetic correlation of all the clinical CRKP isolates.<sup>26</sup> The positive products were sequenced by Shanghai Jieli Biotechnology, and the results were submitted to *K. pneumoniae* MLST database ([https://bigsdatabase.pasteur.fr/cgi-bin/bigsdatabase/bigsdatabase.pl?db=pubmlst\\_klebsiella\\_seqdef](https://bigsdatabase.pasteur.fr/cgi-bin/bigsdatabase/bigsdatabase.pl?db=pubmlst_klebsiella_seqdef)) for comparison.


## Statistical Analysis

All analyses and graphs were performed using SPSS v.24.0 software (SPSS Inc., Chicago, USA). The chi-square test or Fisher's exact test was used to analyze categorical variables. Continuous variables were presented as means ± standard deviation (SD), and were evaluated by Student's *t*-tests or Mann–Whitney *U*-test. Multivariable logistic regression analysis was operated to identify independent risk factors for CRKP infection. All biologically plausible variables with a value of *P* < 0.1 within univariate analysis were included in the following multiple logistic regression model. *P* < 0.05 was considered as statistically significant, and all probability values were two-tailed distribution.

## Results

### Distribution of Clinical CRKP Isolates

A total of 66 non-repetitive clinical CRKP isolates were collected from September 2016 to August 2019 in Southwest China. The clinical CRKP isolates were obtained from patients admitted to the 12 different departments, including the majority of neonatology department (27.3%, *n* = 18) and rehabilitation department (24.3%, *n* = 16) ([Figure 1A](#)). These isolates were mainly originated from sputum (34.8%, *n* = 23), followed by urine (25.8%, *n* = 17), blood (21.2%, *n* = 14), secretion (6.1%, *n* = 4), pleuroperitoneal fluids (6.1%, *n* = 4), catheter tip (3.0%, *n* = 2), and pus (3.0%, *n* = 2) ([Figure 1B](#)).



**Figure 1** Distribution of 66 clinical CRKP strains. **(A)** Department distribution of 66 CRKP strains; **(B)** sample types of 66 CRKP strains.

## Antimicrobial Susceptibility Profiles

All of 66 clinical CRKP isolates were multidrug-resistant and resistance rates to 12 antimicrobials reached 100%, which were cefepime, imipenem, piperacillin/tazobactam, ceftriaxone, ampicillin/sulbactam, cefotaxime, ceftazidime, meropenem, and ceftazidime. In addition, the resistance rates to sulperazon, aztreonam, tobramycin, gentamicin, levofloxacin, ciprofloxacin, cotrimoxazole, and amikacin were 95.5%, 87.9%, 83.3%, 83.3%, 81.9%, 80.3%, 65.2%, and 54.5%, respectively. Remarkably, all the CRKP strains in this study were sensitive to tigecycline and polymyxin B (Table 1).

## Carbapenem Resistance Phenotype and Molecular Characteristics

Both EDTA-synergy test and CIM test were shown that 24 (36.4%) clinical CRKP strains produced class B carbapenemases. The sequencing results confirmed that all of which isolates contained *bla*<sub>NDM</sub>, in which 20 isolates carried *bla*<sub>NDM-1</sub> and 4 isolates carried *bla*<sub>NDM-5</sub>, and 3 isolates possessed *bla*<sub>NDM</sub> and *bla*<sub>IMP</sub> simultaneously. The rest of 42 (63.6%) isolates contained *bla*<sub>KPC-2</sub>. Notably, in addition to the production of carbapenemase, 100% and 95.5% of the clinical CRKP isolates were positive for ESBL and AmpC genes, respectively. Furthermore, virulence genes were detected in all isolates, 54 (81.8%) isolates

detected three different virulence genes. Especially for strain Kpn497 and Kpn131, they carried 12 virulence genes and 11 virulence genes, respectively (Table 2).

## Plasmid and Integron-Associated Analysis

KPC-encoding plasmids were positive in 35 transconjugants and all of these plasmids belonged to IncFII. Meanwhile, NDM-encoding plasmids that belong to IncX3 were detected in 24 transconjugants (Table 2).

Nearly 85% (n = 56) strains were *intl1*-positive, of which 46 strains contained variable region. A total of 6 cassette arrays were found among them. The cassette arrays were revealed in this study included: *addA2* (82.6%), *ddfrA12* (6.5%), *dfrA12-addA2* (4.3%), *dfrA27-arr3* (2.2%), *gcuF-dfrA12* (2.2%), and *bla*<sub>OXA-10</sub>-*addA1-aacA4* (2.2%) (Table 2) In addition, ISCRI was detected in 25 strains.

## Genetic Environments of *bla*<sub>KPC-2</sub> and *bla*<sub>NDM-1</sub>

All of 42 KPC-2-producing strains could be divided into three different types (A, B, and C) based on the genetic structures that compared with pKP048 (GenBank Accession No.FJ628167). Type C was the most prevalent (n = 31), followed by type A (n = 8) and type B (n = 3). Type A exhibited the same structure as pKP048. The downstream of *ISKpn6-like* was deletion in type B, and three genetic elements downstream of *bla*<sub>KPC-2</sub> were

**Table 1** Antimicrobial Susceptibility Profiles of 66 Clinical CRKP Strains

Antimicrobial Agent	No.	%R	No.	%I	No.	%S
Amikacin	36	54.5	0	0	30	45.5
Cefepime	66	100	0	0	0	0
Imipenem	66	100	0	0	0	0
Piperacillin/tazobactam	66	100	0	0	0	0
Ceftriaxone	66	100	0	0	0	0
Ampicillin/sulbactam	66	100	0	0	0	0
Cefotaxime	66	100	0	0	0	0
Cefazolin	66	100	0	0	0	0
Cefuroxime	66	100	0	0	0	0
Ertapenem	66	100	0	0	0	0
Tobramycin	55	83.3	4	6.1	7	10.6
Sulperazon	63	95.5	2	3.0	1	1.5
Ceftazidime	66	100	0	0	0	0
Meropenem	66	100	0	0	0	0
Levofloxacin	54	81.9	0	0	12	18.1
Cefoxitin	66	100	0	0	0	0
Aztreonam	58	87.9	0	0	8	12.1
Ciprofloxacin	53	80.3	0	0	13	19.7
Gentamicin	55	83.3	0	0	11	16.7
Tigecycline	0	0	0	0	66	100
Compound sulfamethoxazole	43	65.2	0	0	23	34.8
Polymyxin B	0	0	0	0	66	100

**Abbreviations:** S, [susceptible](#); I, intermediate; R, resistant.

deletion in type C, including *ISKpn6-like*, *tnpR*, and *tnpA* (Figure 2A).

Furthermore, the genetic structure of pNDM-BJ01 (GenBank accession No. JQ001791) was used to compare with each NDM-1-producing strain, revealing A, B, and C types among these strains. As for type A (n = 12), the *ISAbal25* upstream of the *bla<sub>NDM-1</sub>* was truncated, all downstream genes of *tat* were complete deletion. *ISAbal25* upstream of the *bla<sub>NDM-1</sub>* was deletion in type B compared with type A. Type C presented as the *IS30* upstream of *bla<sub>NDM-1</sub>* and followed by *ble<sub>MBL</sub>*, *trpF*, *tat*, *cutA*, *groES*, *groEL* (Figure 2B).

### MLST of Clinical CRKP Isolates

A total of 13 MLSTs were identified among all the clinical CRKP isolates, and ST11 (56.2%, n = 37) was the most frequent ST type, followed by ST4495 (19.7%, n = 13), ST2407 (4.6%, n = 3), ST147 (3.0%, n = 2), ST307 (3.0%, n = 2), ST37 (3.0%, n = 2), ST15 (1.5%, n = 1), ST17 (1.5%, n = 1), ST152 (1.5%, n = 1), ST23 (1.5%, n = 1), ST405 (1.5%, n = 1), ST318 (1.5%, n = 1), and ST467 (1.5%, n = 1).

### Risk Factors and Multivariate Analysis of CRKP Infection

Statistically significant differences were observed for respiratory disease ( $P = 0.034$ ), renal disease ( $P = 0.039$ ), hematological disease ( $P = 0.02$ ), tracheal cannula ( $P < 0.001$ ), and use of  $\beta$ -lactams/ $\beta$ -lactamase inhibitor combination ( $P = 0.009$ ) between CRKP and CSKP groups (Table 3). Multivariate analysis revealed that hematological disease (odds ratio [OR], 2.568; 95% confidence interval [95% CI], 1.106 to 5.964;  $P = 0.028$ ), tracheal cannula (OR, 4.883; 95% CI, 1.797 to 13.265;  $P = 0.002$ ), and use of  $\beta$ -lactams/ $\beta$ -lactamase inhibitor combination (OR, 4.271; 95% CI, 1.760 to 10.365;  $P = 0.001$ ) were independent risk factors for CRKP infection.

### Discussion

The clinical CRKP strains were first reported in 1997,<sup>27,28</sup> and this type of resistance did not become common within that decade. However, with the increasing use of carbapenems in recent years, CRKP strains have been distributed around the world at a boosting rate, and it was listed as one of the critical-priority bacteria by World Health

**Table 2** Genotypes of 66 Clinical CRKP Strains

N	Carbapenemase	ESBLs	AmpC	Int	Virulence Gene	Gene Cassette	ISCR	MLST	Replicon Type
16	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>	ISCR I	ST11	IncFII
26	NDM-1	SHV+CTX-M	DHA		<i>entB+mrkD</i>		ISCR I	ST4495	IncX3
27	NDM-1	SHV+CTX-M	DHA		<i>entB+fimH+mrkD</i>		ISCR I	ST4495	IncX3
30	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>	ISCR I	ST11	IncFII
32	NDM-1	SHV+CTX-M	DHA		<i>entB+fimH+mrkD</i>		ISCR I	ST4495	IncX3
34	NDM-1	SHV+CTX-M	DHA		<i>entB+fimH+mrkD</i>		ISCR I	ST4495	IncX3
36	NDM-1	SHV+CTX-M	DHA		<i>entB+fimH+mrkD</i>		ISCR I	ST4495	IncX3
37	NDM-1	SHV+CTX-M	DHA		<i>entB+fimH+mrkD</i>		ISCR I	ST4495	IncX3
39	NDM-1	SHV	DHA		<i>entB+mrkD</i>		ISCR I	ST4495	IncX3
40	NDM-1	SHV	DHA+ACC		<i>entB+fimH</i>		ISCR I	ST37	IncX3
45	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	
46	NDM-1	SHV+CTX-M	DHA+ACC		<i>entB+fimH+mrkD</i>		ISCR I	ST4495	IncX3
49	NDM-1	SHV+CTX-M	DHA+ACC		<i>entB+fimH+mrkD</i>		ISCR I	ST4495	IncX3
51	KPC-2	SHV+CTX-M	ACC		<i>entB+fimH+mrkD</i>			ST11	
56	NDM-1	SHV+CTX-M	DHA+ACC		<i>entB+fimH+mrkD</i>		ISCR I	ST4495	IncX3
57	NDM-1	SHV+CTX-M	DHA+ACC		<i>entB+fimH+mrkD</i>		ISCR I	ST4495	IncX3
58	NDM-1+IMP-4	SHV+CTX-M	DHA+ACC		<i>entB+fimH</i>			ST4495	IncX3
101	NDM-1	SHV	ACC		<i>entB+fimH+mrkD</i>			ST37	IncX3
131	KPC-2	SHV+TEM	ACC	<i>IntI</i>	<i>entB+fimH+mrkD+rmpA/rmpA2+iucA+terB+aerobactin+HII B+iroN+iutA</i>	<i>addA2</i>	ISCR I	ST11	IncFII
210	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
211	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
214	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
215	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
221	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>	ISCR I	ST11	IncFII
223	KPC-2	SHV+TEM+CTX-M	ACC		<i>entB+fimH+mrkD</i>			ST147	
227	KPC-2	SHV+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
228	KPC-2	SHV+CTX-M	ACC		<i>entB+fimH+mrkD</i>			ST11	IncFII
230	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
232	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
233	KPC-2	SHV	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>dfrA12, addA2</i>		ST147	IncFII
234	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
241	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
244	KPC-2	SHV	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>dfrA12</i>		ST23	IncFII

(Continued)

Table 2 (Continued).

N	Carbapenemase	ESBLs	AmpC	Int	Virulence Gene	Gene Cassette	ISCR	MLST	Replicon Type
245	KPC-2	SHV+TEM+CTX-M	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
251	KPC-2	SHV+TEM+CTX-M	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
252	KPC-2	SHV+TEM+CTX-M		<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
257	KPC-2	SHV+TEM+CTX-M	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	
258	KPC-2	SHV+TEM+CTX-M	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
267	NDM-1	SHV	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>	ISCR 1	ST11	IncX3
271	KPC-2	SHV+TEM+CTX-M	ACC		<i>entB+fimH+mrkD</i>			ST11	IncFII
285	KPC-2	SHV+TEM+CTX-M	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
354	KPC-2	SHV+TEM+CTX-M	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
406	KPC-2	SHV+TEM	DHA	<i>Int1</i>	<i>fimH+mrkD</i>	<i>OXA-10, addA1, aacA4</i>	ISCR 1	ST318	IncFII
436	KPC-2	SHV+TEM	DHA	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST17	IncFII
440	NDM-1	SHV+TEM	ACC	<i>Int1</i>	<i>mrkD</i>	<i>addA2</i>		ST307	IncX3
445	NDM-1	SHV+TEM	ACC		<i>mrkD</i>			ST307	IncX3
473	NDM-1+IMP-4	SHV+TEM	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>	ISCR 1	ST152	IncX3
475	NDM-1	SHV+TEM	ACC	<i>Int1</i>	<i>fimH+mrkD</i>	<i>addA2</i>	ISCR 1	ST15	IncX3
478	KPC-2	SHV+TEM+CTX-M	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>	ISCR 1	ST11	IncFII
479	KPC-2	SHV+TEM+CTX-M	ACC	<i>Int1</i>	<i>entB+fimH</i>	<i>addA2</i>		ST11	IncFII
480	KPC-2	SHV+TEM+CTX-M	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
483	KPC-2	SHV+TEM+CTX-M	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
489	KPC-2	SHV+TEM+CTX-M	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>dfrA12, addA2</i>		ST11	
490	KPC-2	SHV	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>dfrA27, arr3</i>		ST467	IncFII
494	NDM-5	SHV+CTX-M	DHA	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>	ISCR 1	ST2407	IncX3
495	KPC-2	SHV+TEM+CTX-M		<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>dfrA12</i>	ISCR 1	ST11	IncFII
496	KPC-2	SHV+TEM	ACC	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>gcuF, dfrA12</i>		ST405	
497	KPC-2	SHV+TEM		<i>Int1</i>	<i>entB+fimH+mrkD+rmpA/rmpA2+iucA + terB+wcaG+ aerobactin+H11B+iroN+iutA</i>	<i>addA2</i>		ST11	IncFII
498	KPC-2	SHV+CTX-M	ACC	<i>Int1</i>	<i>entB+mrkD</i>	<i>dfrA12</i>		ST11	IncFII
499	NDM-5	SHV+CTX-M	DHA	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>	ISCR 1	ST2407	IncX3
500	KPC-2	SHV+TEM+CTX-M	ACC		<i>entB+fimH+mrkD</i>			ST11	
502	NDM-5	SHV+CTX-M	DHA	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST2407	IncX3
504	NDM-5+IMP-4	SHV+TEM+CTX-M	DHA	<i>Int1</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncX3

(Continued)

Table 2 (Continued).

N	Carbapenemase	ESBLs	AmpC	Int	Virulence Gene	Gene Cassette	ISCR	MLST	Replicon Type
505	KPC-2	SHV+TEM+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII
506	NDM-1	SHV+CTX-M	DHA	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST4495	IncX3
507	KPC-2	SHV+CTX-M	ACC	<i>IntI</i>	<i>entB+fimH+mrkD</i>	<i>addA2</i>		ST11	IncFII

**Abbreviations:** ESBL, extended-spectrum  $\beta$ -lactamase; Int, integron; ISCR, insertion sequence common region.

Organization (WHO).<sup>29</sup> In this work, the antimicrobial susceptibility profiles, molecular characteristics, plasmid and integron-associated analysis, genetic environments of *bla*<sub>KPC-2</sub> and *bla*<sub>NDM-1</sub>, and MLST were evaluated among 66 clinical CRKP isolates. Meanwhile, the risk factors for CRKP infection were also investigated. All the CRKP strains were MDR or even extensively drug-resistant (XDR), in this case, the tigecycline and polymyxin B were the last resort options for CRKP infection. However, the treatment was hardly successful because of the low blood concentrations when tigecycline was used in monotherapy.<sup>30</sup> Adverse events, such as hypofibrinogenemia, also resulted in treatment termination associated with tigecycline therapy.<sup>31</sup> In addition, nephrotoxicity and unclear dosing also limited the widespread clinical usage of polymyxin B.<sup>32,33</sup> Importantly, tigecycline- and polymyxin B-resistant CRKP strains have been reported in many regions,<sup>34,35</sup> therefore, the optimal therapeutic strategy for CRKP infection still needs to be explored.

Focusing on the phenotype and molecular analysis of CRKP isolates, the results revealed that the primary resistance mechanism of CRKP was carbapenemase production, which was consistent with other researches.<sup>36,37</sup> The *bla*<sub>KPC-2</sub> and *bla*<sub>NDM-1</sub> were two major carbapenemase genes in our study, accounting for 63.6% and 30.3%, respectively. KPC-2 is the most commonly identified enzyme among over 20 reported KPC variants so far,<sup>38</sup> and the majority of NDM-1-producers also harbored various other resistance genes but mostly were still susceptible to polymyxin B and tigecycline,<sup>39</sup> which were highly correlated with our results. Moreover, IMP is another important metallo- $\beta$ -lactamase, and *bla*<sub>IMP-1</sub> emerged in Japan in the late 1980s.<sup>40</sup> The *bla*<sub>IMP-4</sub> was one of the most prevalent *bla*<sub>IMP</sub> variants since it was first identified in *Acinetobacter* spp. in the mid-1990s,<sup>41</sup> particularly in China and Australia.<sup>42,43</sup> It is worth noting that *bla*<sub>IMP</sub> often co-exists with other resistance genes,<sup>44</sup> which is consistent with our results. In our work, 3 strains


were identified co-carrying *bla*<sub>NDM</sub> and *bla*<sub>IMP</sub> genes, including two isolates with *bla*<sub>NDM-1</sub> and *bla*<sub>IMP-4</sub> and one with *bla*<sub>NDM-5</sub> and *bla*<sub>IMP-4</sub>. Coexistence of *bla*<sub>NDM-1</sub> and *bla*<sub>IMP-4</sub> was still rarely distributed in China since first reported in 2012, but the strain co-harboring *bla*<sub>NDM-5</sub> and *bla*<sub>IMP-4</sub> has never been published before.<sup>45</sup>

In this work, *bla*<sub>KPC-2</sub> was only found on IncFII plasmids, and *bla*<sub>NDM</sub> was only detected on IncX3 plasmids. The IncFII plasmid family is commonly low copy number, not only carries multiple antimicrobial resistance and virulence genes, but also can replicate and disseminate in many different species of Enterobacteriaceae.<sup>46</sup> Furthermore, several studies have revealed that IncX3 plasmids were the predominant plasmid type that harboring a variety of *bla*<sub>NDM</sub> genes.<sup>47–49</sup> Of note, IncX3 plasmids can frequently disseminate in humans, animals, and environment, and more commonly spread in Asia, including China,<sup>49,50</sup> Myanmar,<sup>51</sup> South Korea,<sup>52</sup> and India.<sup>53</sup> In addition, 46 CRKP strains carried variable regions of *intI1*, and 7 resistance gene cassettes were detected among them. The *addA2* (40/46) and *addA1* (1/46) were the genes responsible for aminoglycoside resistance. The *dfrA12* (6/46), *dfrA27* (1/46) contributed to trimethoprim resistance. Furthermore, a  $\beta$ -lactams-resistant gene *bla*<sub>OXA-10</sub>, a gentamicin-resistant gene *arr3*, and a rifampicin-resistant gene *aacA4* were also detected.

A previous study has proved that the genetic environment of *bla*<sub>KPC-2</sub> in pKP048 from China was considered as a combination of the Tn3-based transposon and the partial Tn4401 structure.<sup>54</sup> In our study, KPC-2 producers presented a similar structure (*tnpA-tnpR-ISkpn8-bla*<sub>KPC-2</sub>), but the environment surrounding *bla*<sub>KPC-2</sub> was partly different from other reports in China due to the specific insertions and deletions.<sup>55,56</sup> Additionally, all the NDM-1 producers carried the highly conserved region (*bla*<sub>NDM-1-ble</sub><sub>MBL-trpF-tat) surrounding the *bla*<sub>NDM-1</sub> gene, which was</sub>




A



A: Kpn45, 233, 354, 478, 479, 480, 483, 490; B: Kpn406, 497, 498;

C: Kpn16, 30, 51, 131, 210, 211, 214, 215, 221, 223, 227, 228, 230, 232, 234, 241, 244, 245, 251, 252, 257, 258, 271, 285, 436, 489, 495, 496, 500, 505, 507.

B



A: Kpn26, 32, 34, 36, 37, 39, 56, 101, 267, 473, 475, 506; B: Kpn27, 46, 49, 57, 58; C: Kpn40, 440, 445.

**Figure 2** Comparison of the genetic elements surrounding the *bla*<sub>KPC-2</sub> and *bla*<sub>NDM-1</sub> genes identified in this study. (A) Comparison of the genetic environments of *bla*<sub>KPC-2</sub>, reference sequences: *K. pneumoniae* (pKP048, GenBank Accession No.FJ628167); (B) comparison of the genetic environments of *bla*<sub>NDM-1</sub>, reference sequences: *A. lwoffii* (pNDM-BJ01, GenBank accession NO. JQ001791).

identified not only in *K. pneumoniae*, but also in *Acinetobacter lwoffii* and *Enterobacter Cloacae*.<sup>25,57,58</sup>

Furthermore, the results showed that more than 80% clinical CRKP strains simultaneously carried virulence genes *entB*, *fimH*, and *mrkD*. The enterobactin that encoded by

*entB* is one of the ubiquitous siderophores in *K. pneumoniae*, which is an important factor involving in iron acquisition.<sup>59</sup>

The *fimH* and *mrkD* mediate bacterial adhesion through encoding type I and type III fimbriae of *K. pneumoniae* implied that these two genes may contribute to biofilm formation and

**Table 3** Clinical Characteristics of CRKP and CSKP Strains

Variable	CRKP (n=66)	CSKP (n=66)	P-value
Demographic, n (%) or IQR			
Age (years)	44.5 (0.60)	46.5 (0.61)	0.754
Sex (male)	47 (71.2%)	46 (69.7%)	0.894
Length of hospital stays	15 (7, 27)	25 (10, 38)	0.1
Admission to ICU	14 (21.1%)	7 (10.6%)	0.096
Co-morbidity, n (%)			
Malignant disease	7 (10.6%)	15 (22.7%)	0.062
Diabetes mellitus	11 (16.7%)	12 (18.2%)	0.819
Hypertension	16 (24.2%)	16 (24.2%)	1
Heart disease	9 (13.6%)	11 (16.7%)	0.627
Hepatobiliary disease	12 (18.2%)	15 (22.7%)	0.517
Respiratory disease	45 (68.2%)	33 (50%)	<b>0.034</b>
Renal disease	7 (10.6%)	16 (24.2%)	<b>0.039</b>
Urinary tract infection	10 (15.2%)	9 (13.6%)	0.804
Cranio-cerebral disease	15 (22.7%)	9 (13.6%)	0.176
Hematological disease	32 (48.5%)	19 (28.8%)	<b>0.02</b>
Gastrointestinal disease	8 (12.1%)	5 (7.6%)	0.381
Invasive procedures and devices			
Tracheal cannula	24 (36.4%)	7 (10.6%)	<b>&lt;0.001</b>
Central venous catheter	5 (7.6%)	12 (18.2%)	0.069
Foreign material in the body	14 (21.2%)	7 (10.6%)	0.096
Surgical operations after admission	9 (13.6%)	16 (24.2%)	0.12
Colostomy	3 (4.5%)	1 (1.5%)	0.612
Gastrostomy	2 (3%)	0	0.476
Antibiotic treatment, n (%)			
Penicillins	2 (3%)	5 (7.6%)	0.437
First, second-generation cephalosporins	7 (10.6%)	15 (22.7%)	0.21
Third, fourth-generation cephalosporins	17 (25.8%)	19 (28.8%)	0.696
Aminoglycosides	4 (6.1%)	1 (1.5%)	0.362
Quinolones	20 (30.3%)	21 (31.8%)	0.851
Metronidazole	0	1 (1.5%)	1
Carbapenems	30 (45.5%)	32 (48.5%)	0.727
β-lactams and β-lactamase inhibitor combination	40 (60.6%)	25 (37.9%)	<b>0.009</b>
Clinical outcomes, n (%)			
Patient outcome: mortality	5 (7.6%)	0	0.068

**Note:** Bold indicates  $P < 0.05$ .

**Abbreviations:** IQR, interquartile range; CR, carbapenem-resistant; CS, carbapenem-susceptible; ICU, intensive care unit.

development. Zhang et al have also observed that 95% tested *K. pneumoniae* isolates co-carrying *fimH* and *mrkD* genes in their study.<sup>60</sup> Additionally, the genes *peg-344*, *iroB*, *iucA*, *rmpA*, and *rmpA2* locate on the HVKP virulence plasmid and are regarded as regulators of hypervirulent phenotype.<sup>61</sup> Notably, two carbapenem-resistant HVKP (CR-HVKP) strains were found in this work, which contained *bla*<sub>KPC-2</sub> gene. There

are two hypotheses to explain the evolution of this novel phenotype. First, CRKP strains acquire HVKP-specific virulence plasmid; second, HVKP strains obtain the carbapenem-resistant genes by acquisition of resistance plasmid or by the insertion of resistance determinants into HVKP-specific virulence plasmid.<sup>59,62</sup> Compared with the second hypothesis, the first one was more likely to occur.<sup>63–65</sup>

According to the results of MLST, the 42 KPC-2-producing CRKP strains were grouped into 7 ST types, of which 35 strains belonged to ST11. It has been reported that most of KPC-2-producing strains belong to clone group (CG) 258; meanwhile, ST11 and ST258 are the dominant ST types.<sup>66</sup> ST258 has spread around the world since it emerged in the early 21st century, particularly in North America, Latin America, and several European countries.<sup>66,67</sup> However, ST11 is more prevalent in Asia, and usually accounts for more than half of all KPC-2-producing CRKP strains in China.<sup>47,68,69</sup> In addition, the 24 NDM-producing CRKP strains were classified into 8 ST types, and ST4495 was the most common ST type. Remarkably, ST4495 (4-1-99-1-9-5-5) has never been detected before, indicating a novel clone carrying *bla*<sub>NDM-1</sub> has spread in Southwest China.

Hematological disease, tracheal cannula, and exposure to  $\beta$ -lactams and  $\beta$ -lactamase inhibitor combination were independent risk factors for CRKP infection. Many studies have demonstrated that the history of tracheal cannula was independent risk factor for CRKP infection.<sup>70–73</sup> Meanwhile, exposure to antibiotic was also associated with CRKP infection, such as exposure to quinolones,<sup>74</sup>  $\beta$ -lactams and  $\beta$ -lactamase inhibitor combination, and carbapenems,<sup>75</sup> which was consistent with our study. Notably, hematological disease was rarely found to be a risk factor for CRKP infection in other studies, but it is reasonable due to patients with hematological disease were accompanied with severe immune function deficiency and were more prone to be infected. There are some limitations in our study, first, the number of patients included in this study is relatively small, although this is a common problem in studies assessing risk factors for multidrug-resistant microbial infections.<sup>74</sup> Secondly, the study was performed in a single-center setting, so that some important risk factors may be missed.

## Conclusion

The current study revealed the high prevalence of CRKP infection caused by NDM-1 and KPC-2 producers, and ST11 was the most common lineage. To the best of our knowledge, the novel ST4495 and co-exist of *bla*<sub>NDM-5</sub> and *bla*<sub>IMP-4</sub> were the first reported in this work. Moreover, plasmid, integron, ISCR, and other mobile genetic elements endowed bacteria a rapid adaptation ability in changing environments; meanwhile, the genetic environments of *bla*<sub>KPC-2</sub> and *bla*<sub>NDM-1</sub> showed

polymorphism. Logistic regression model indicated that hematological disease, tracheal cannula, and use of  $\beta$ -lactams and  $\beta$ -lactamase inhibitor combination were independent risk factors for CRKP infection, suggesting that appropriate clinical management and antimicrobials treatment were necessary for retarding the selection of CRKP strains. Effective measures to prevent and control the further dissemination of CRKP strains are highly needed.

## Data Sharing Statement

The data used and/or analyzed in this study are available from the corresponding author on reasonable request.

## Ethics Approval and Consent to Participate

This study was approved by the Institutional Review Board of affiliated hospital of southwest medical university (KY2020043). As for neonate, a parent or legal guardian provided informed consent for these patients, and that this study was conducted in accordance with the Declaration of Helsinki. Written informed consent was obtained from all participants.

## Acknowledgments

This work was supported by the grants from Sichuan Science and Technology Program (2021YFH001, 2021YFS0329), Luxian Government and Southwest Medical University Cooperation Program (2020LXXNYKD-04).

## Author Contributions

All authors contributed to data analysis, drafting or revising the article, have agreed on the journal to which the article will be submitted, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

## Disclosure

The authors declare that they have no conflict of interest.

## References

1. Wyres KL, Lam MMC, Holt KE. Population genomics of *Klebsiella pneumoniae*. *Nat Rev Microbiol*. 2020;18(6):344–359. doi:10.1038/s41579-019-0315-1
2. Li M, Wang X, Wang J, et al. Infection-prevention and control interventions to reduce colonisation and infection of intensive care unit-acquired carbapenem-resistant *Klebsiella pneumoniae*: a 4-year quasi-experimental before-and-after study. *Antimicrob Resist Infect Control*. 2019;8(1):8. doi:10.1186/s13756-018-0453-7

3. Hoxha A, Kärki T, Giambi C, et al. Attributable mortality of carbapenem-resistant *Klebsiella pneumoniae* infections in a prospective matched cohort study in Italy, 2012–2013. *J Hosp Infect.* 2016;92(1):61–66. doi:10.1016/j.jhin.2015.06.018
4. Nordmann P, Poirel L. Epidemiology and diagnostics of carbapenem resistance in gram-negative bacteria. *Clin Infect Dis.* 2019;69(Suppl 7):S521–s528. doi:10.1093/cid/ciz824
5. Paterson DL, Bonomo RA. Extended-spectrum beta-lactamases: a clinical update. *Clin Microbiol Rev.* 2005;18(4):657–686.
6. Jacoby GA. AmpC beta-lactamases. *Clin Microbiol Rev.* 2009;22(1):161–182. Table of Contents. doi:10.1128/CMR.00036-08
7. Bush K, Jacoby GA. Updated functional classification of beta-lactamases. *Antimicrob Agents Chemother.* 2010;54(3):969–976. doi:10.1128/AAC.01009-09
8. Patel G, Bonomo RA. “Stormy waters ahead”: global emergence of carbapenemases. *Front Microbiol.* 2013;4:48. doi:10.3389/fmicb.2013.00048
9. Zhu J, Sun L, Ding B, et al. Outbreak of NDM-1-producing *Klebsiella pneumoniae* ST76 and ST37 isolates in neonates. *Eur J Clin Microbiol Infect Dis.* 2016;35(4):611–618. doi:10.1007/s10096-016-2578-z
10. Zheng R, Zhang Q, Guo Y, et al. Outbreak of plasmid-mediated NDM-1-producing *Klebsiella pneumoniae* ST105 among neonatal patients in Yunnan, China. *Ann Clin Microbiol Antimicrob.* 2016;15(1):10. doi:10.1186/s12941-016-0124-6
11. Wang X, Xu X, Li Z, et al. An outbreak of a nosocomial NDM-1-producing *Klebsiella pneumoniae* ST147 at a teaching hospital in mainland China. *Microb Drug Resist.* 2014;20(2):144–149. doi:10.1089/mdr.2013.0100
12. Escobar Pérez JA, Olarte Escobar NM, Castro-Cardozo B, et al. Outbreak of NDM-1-producing *Klebsiella pneumoniae* in a neonatal unit in Colombia. *Antimicrob Agents Chemother.* 2013;57(4):1957–1960. doi:10.1128/AAC.01447-12
13. Monteiro J, Inoue FM, Lobo APT, Ibanes AS, Tufik S, Kiffer CRV. A major monoclonal hospital outbreak of NDM-1-producing *Klebsiella pneumoniae* ST340 and the first report of ST2570 in Brazil. *Infect Control Hosp Epidemiol.* 2019;40(4):492–494. doi:10.1017/ice.2018.333
14. Ripabelli G, Sammarco ML, Salzo A, Scutellà M, Felice V, Tamburro M. New Delhi metallo- $\beta$ -lactamase (NDM-1)-producing *Klebsiella pneumoniae* of sequence type ST11: first identification in a hospital of central Italy. *Lett Appl Microbiol.* 2020;71(6):652–659. doi:10.1111/lam.13384
15. Zhang Y, Zhao C, Wang Q, et al. High prevalence of hypervirulent *Klebsiella pneumoniae* infection in China: geographic distribution, clinical characteristics, and antimicrobial resistance. *Antimicrob Agents Chemother.* 2016;60(10):6115–6120. doi:10.1128/AAC.01127-16
16. Zhang Y, Jin L, Ouyang P, et al. Evolution of hypervirulence in carbapenem-resistant *Klebsiella pneumoniae* in China: a multicentre, molecular epidemiological analysis. *J Antimicrob Chemother.* 2020;75(2):327–336. doi:10.1093/jac/dkz446
17. Zhou C, Wu Q, He L, et al. Clinical and molecular characteristics of carbapenem-resistant hypervirulent *Klebsiella pneumoniae* isolates in a tertiary hospital in Shanghai, China. *Infect Drug Resist.* 2021;14:2697–2706. doi:10.2147/IDR.S321704
18. Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing.* 30th ed. CLSI M100. Wayne PA: CLSI; 2020.
19. Lee K, Chong Y, Shin HB, Kim YA, Yong D, Yum JH. Modified Hodge and EDTA-disk synergy tests to screen metallo-beta-lactamase-producing strains of *Pseudomonas* and *Acinetobacter* species. *Clin Microbiol Infect.* 2001;7(2):88–91. doi:10.1046/j.1469-0691.2001.00204.x
20. Sfeir MM, Hayden JA, Fauntleroy KA, et al. EDTA-modified carbapenem inactivation method: a phenotypic method for detecting metallo- $\beta$ -lactamase-producing enterobacteriaceae. *J Clin Microbiol.* 2019;57(5):e01757–18. doi:10.1128/JCM.01757-18
21. Xiang T, Chen C, Wen J, et al. Resistance of *Klebsiella pneumoniae* strains carrying bla (NDM-1) gene and the genetic environment of bla (NDM-1). *Front Microbiol.* 2020;11:700. doi:10.3389/fmicb.2020.00700
22. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods.* 2005;63(3):219–228. doi:10.1016/j.mimet.2005.03.018
23. Poirel L, Dortet L, Bernabeu S, Nordmann P. Genetic features of blaNDM-1-positive enterobacteriaceae. *Antimicrob Agents Chemother.* 2011;55(11):5403–5407. doi:10.1128/AAC.00585-11
24. Mishra S, Upadhyay S, Sen MR, Maurya AP, Choudhury D, Bhattacharjee A. Genetic acquisition of NDM gene offers sustainability among clinical isolates of *Pseudomonas aeruginosa* in clinical settings. *PLoS One.* 2015;10(1):e0116611. doi:10.1371/journal.pone.0116611
25. Jia X, Dai W, Ma W, et al. Carbapenem-resistant *E. cloacae* in Southwest China: molecular analysis of resistance and risk factors for infections caused by NDM-1-producers. *Front Microbiol.* 2018;9:658. doi:10.3389/fmicb.2018.00658
26. Diancourt L, Passet V, Verhoef J, Grimont PA, Brisse S. Multilocus sequence typing of *Klebsiella pneumoniae* nosocomial isolates. *J Clin Microbiol.* 2005;43(8):4178–4182. doi:10.1128/JCM.43.8.4178-4182.2005
27. Bradford PA, Urban C, Mariano N, Projan SJ, Rahal JJ, Bush K. Imipenem resistance in *Klebsiella pneumoniae* is associated with the combination of ACT-1, a plasmid-mediated AmpC beta-lactamase, and the fss of an outer membrane protein. *Antimicrob Agents Chemother.* 1997;41(3):563–569. doi:10.1128/AAC.41.3.563
28. MacKenzie FM, Forbes KJ, Dorai-John T, Amyes SG, Gould IM. Emergence of a carbapenem-resistant *Klebsiella pneumoniae*. *Lancet.* 1997;350(9080):783. doi:10.1016/S0140-6736(05)62567-6
29. Tacconelli E, Carrara E, Savoldi A, et al. Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. *Lancet Infect Dis.* 2018;18(3):318–327. doi:10.1016/S1473-3099(17)30753-3
30. Ruiz J, Ramirez P, Villarreal E, et al. Effect of pharmacokinetic/pharmacodynamic ratio on tigecycline clinical response and toxicity in critically ill patients with multidrug-resistant Gram-negative infections. *SAGE Open Med.* 2020;8:2050312120958897. doi:10.1177/2050312120958897
31. Zhang Q, Wang J, Liu H, Ma W, Zhou S, Zhou J. Risk factors for tigecycline-induced hypofibrinogenemia. *J Clin Pharm Ther.* 2020;45(6):1434–1441. doi:10.1111/jcpt.13250
32. Avedissian SN, Liu J, Rhodes NJ, et al. A review of the clinical pharmacokinetics of polymyxin B. *Antibiotics.* 2019;8(1):31.
33. Zavascki AP, Nation RL. Nephrotoxicity of polymyxins: is there any difference between colistimethate and polymyxin B? *Antimicrob Agents Chemother.* 2017;61(3):e02319–16. doi:10.1128/AAC.02319-16
34. Gao L, Lv Y, Li Y. Analysis of the drug resistance of carbapenem-resistant *Klebsiella pneumoniae* in the China antimicrobial resistance surveillance trial program, 2007–2018. *Microb Drug Resist.* 2020;26(8):944–950. doi:10.1089/mdr.2019.0299
35. Zhang Y, Wang Q, Yin Y, et al. Epidemiology of carbapenem-resistant enterobacteriaceae infections: report from the China CRE network. *Antimicrob Agents Chemother.* 2018;62(2):e01882–17. doi:10.1128/AAC.01882-17
36. Zheng B, Dai Y, Liu Y, et al. Molecular epidemiology and risk factors of carbapenem-resistant *Klebsiella pneumoniae* infections in Eastern China. *Front Microbiol.* 2017;8:1061. doi:10.3389/fmicb.2017.01061
37. Kumudaning WGM, Wijesooriya LI, Namalie-KD, Sunil-Chandra NP, Wijayasinghe YS. Epidemiology of multidrug-resistant enterobacteriaceae in Sri Lanka: first evidence of bla(KPC) harboring *Klebsiella pneumoniae*. *J Infect Public Health.* 2020;13(9):1330–1335. doi:10.1016/j.jiph.2020.04.010

38. Munoz-Price LS, Poirel L, Bonomo RA, et al. Clinical epidemiology of the global expansion of *Klebsiella pneumoniae* carbapenemases. *Lancet Infect Dis*. 2013;13(9):785–796. doi:10.1016/S1473-3099(13)70190-7
39. Lee CR, Lee JH, Park KS, Kim YB, Jeong BC, Lee SH. Global dissemination of carbapenemase-producing *klebsiella pneumoniae*: epidemiology, genetic context, treatment options, and detection methods. *Front Microbiol*. 2016;7:895.
40. Watanabe M, Iyobe S, Inoue M, Mitsuhashi S. Transferable imipenem resistance in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*. 1991;35(1):147–151. doi:10.1128/AAC.35.1.147
41. Chu YW, Afzal-Shah M, Houang ET, et al. IMP-4, a novel metallo-beta-lactamase from nosocomial *Acinetobacter* spp. collected in Hong Kong between 1994 and 1998. *Antimicrob Agents Chemother*. 2001;45(3):710–714. doi:10.1128/AAC.45.3.710-714.2001
42. Leung GH, Gray TJ, Cheong EY, Haertsch P, Gottlieb T. Persistence of related bla-IMP-4 metallo-beta-lactamase producing Enterobacteriaceae from clinical and environmental specimens within a burns unit in Australia - a six-year retrospective study. *Antimicrob Resist Infect Control*. 2013;2(1):35. doi:10.1186/2047-2994-2-35
43. Feng W, Zhou D, Wang Q, et al. Dissemination of IMP-4-encoding pIMP-HZ1-related plasmids among *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* in a Chinese teaching hospital. *Sci Rep*. 2016;6(1):33419. doi:10.1038/srep33419
44. Kizny Gordon A, Phan HTT, Lipworth SI, et al. Genomic dynamics of species and mobile genetic elements in a prolonged blaIMP-4-associated carbapenemase outbreak in an Australian hospital. *J Antimicrob Chemother*. 2020;75(4):873–882. doi:10.1093/jac/dkz526
45. Chen Z, Wang Y, Tian L, et al. First report in China of Enterobacteriaceae clinical isolates coharboring blaNDM-1 and blaIMP-4 drug resistance genes. *Microb Drug Resist*. 2015;21(2):167–170. doi:10.1089/mdr.2014.0087
46. Villa L, García-Fernández A, Fortini D, Carattoli A. Replicon sequence typing of IncF plasmids carrying virulence and resistance determinants. *J Antimicrob Chemother*. 2010;65(12):2518–2529. doi:10.1093/jac/dkq347
47. Zhang R, Liu L, Zhou H, et al. Nationwide surveillance of clinical carbapenem-resistant enterobacteriaceae (CRE) strains in China. *EBioMedicine*. 2017;19:98–106. doi:10.1016/j.ebiom.2017.04.032
48. Yoon EJ, Kang DY, Yang JW, et al. New Delhi Metallo-beta-lactamase-producing enterobacteriaceae in South Korea between 2010 and 2015. *Front Microbiol*. 2018;9:571. doi:10.3389/fmicb.2018.00571
49. Wang Y, Tong MK, Chow KH, et al. Occurrence of highly conjugative incx3 epidemic plasmid carrying bla (NDM) in enterobacteriaceae isolates in geographically widespread areas. *Front Microbiol*. 2018;9:2272. doi:10.3389/fmicb.2018.02272
50. Ho PL, Li Z, Lo WU, et al. Identification and characterization of a novel incompatibility group X3 plasmid carrying bla NDM-1 in enterobacteriaceae isolates with epidemiological links to multiple geographical areas in China. *Emerg Microb Infect*. 2012;1(11):e39. doi:10.1038/emi.2012.37
51. Sugawara Y, Akeda Y, Sakamoto N, et al. Genetic characterization of blaNDM-harboring plasmids in carbapenem-resistant *Escherichia coli* from Myanmar. *PLoS One*. 2017;12(9):e0184720. doi:10.1371/journal.pone.0184720
52. Kim JS, Jin YH, Park SH, et al. Horizontal transfer of bla(NDM-1)-carrying IncX3 plasmid between carbapenem-resistant enterobacteriaceae in a single patient. *J Infect*. 2020;81(5):816–846. doi:10.1016/j.jinf.2020.07.013
53. Choudhury NA, Paul D, Chakravarty A, Bhattacharjee A, Dhar Chanda D. IncX3 plasmid mediated occurrence of bla(NDM-4) within *Escherichia coli* ST448 from India. *J Infect Public Health*. 2018;11(1):111–114. doi:10.1016/j.jiph.2017.06.008
54. Shen P, Wei Z, Jiang Y, et al. Novel genetic environment of the carbapenem-hydrolyzing beta-lactamase KPC-2 among enterobacteriaceae in China. *Antimicrob Agents Chemother*. 2009;53(10):4333–4338. doi:10.1128/AAC.00260-09
55. Luo Y, Yang J, Ye L, et al. Characterization of KPC-2-producing *Escherichia coli*, *Citrobacter freundii*, *Enterobacter cloacae*, *Enterobacter aerogenes*, and *Klebsiella oxytoca* isolates from a Chinese hospital. *Microb Drug Resist*. 2014;20(4):264–269. doi:10.1089/mdr.2013.0150
56. Cai JC, Zhang R, Hu YY, Zhou HW, Chen GX. Emergence of *Escherichia coli* sequence type 131 isolates producing KPC-2 carbapenemase in China. *Antimicrob Agents Chemother*. 2014;58(2):1146–1152. doi:10.1128/AAC.00912-13
57. Wailan AM, Paterson DL, Kennedy K, Ingram PR, Bursle E, Sidjabat HE. Genomic characteristics of NDM-producing enterobacteriaceae isolates in Australia and their blaNDM genetic contexts. *Antimicrob Agents Chemother*. 2016;60(1):136–141. doi:10.1128/AAC.01243-15
58. Wailan AM, Sidjabat HE, Yam WK, et al. Mechanisms involved in acquisition of blaNDM genes by IncA/C2 and IncFIIY plasmids. *Antimicrob Agents Chemother*. 2016;60(7):4082–4088. doi:10.1128/AAC.00368-16
59. Russo TA, Marr CM. Hypervirulent *Klebsiella pneumoniae*. *Clin Microbiol Rev*. 2019;32(3):e00001–19. doi:10.1128/CMR.00001-19
60. Zhang S, Zhang X, Wu Q, et al. Clinical, microbiological, and molecular epidemiological characteristics of *Klebsiella pneumoniae*-induced pyogenic liver abscess in southeastern China. *Antimicrob Resist Infect Control*. 2019;8(1):166. doi:10.1186/s13756-019-0615-2
61. Russo TA, Olson R, Fang CT, et al. Identification of biomarkers for differentiation of hypervirulent *Klebsiella pneumoniae* from classical *K. pneumoniae*. *J Clin Microbiol*. 2018;56(9):e00776–18. doi:10.1128/JCM.00776-18
62. Wei DD, Wan LG, Deng Q, Liu Y. Emergence of KPC-producing *Klebsiella pneumoniae* hypervirulent clone of capsular serotype K1 that belongs to sequence type 11 in Mainland China. *Diagn Microbiol Infect Dis*. 2016;85(2):192–194. doi:10.1016/j.diagmicrobio.2015.03.012
63. Feng Y, Lu Y, Yao Z, Zong Z. Carbapenem-resistant hypervirulent *Klebsiella pneumoniae* of sequence type 36. *Antimicrob Agents Chemother*. 2018;62(7):e02644–17. doi:10.1128/AAC.02644-17
64. Zhang R, Lin D, Chan EW, Gu D, Chen GX, Chen S. Emergence of carbapenem-resistant serotype K1 hypervirulent *Klebsiella pneumoniae* strains in China. *Antimicrob Agents Chemother*. 2016;60(1):709–711. doi:10.1128/AAC.02173-15
65. Fu L, Tang L, Wang S, et al. Co-location of the bla(KPC-2), bla (CTX-M-65), rmtB and virulence relevant factors in an IncFII plasmid from a hypermucoviscous *Klebsiella pneumoniae* isolate. *Microb Pathog*. 2018;124:301–304. doi:10.1016/j.micpath.2018.08.055
66. Chen L, Mathema B, Chavda KD, DeLeo FR, Bonomo RA, Kreiswirth BN. Carbapenemase-producing *Klebsiella pneumoniae*: molecular and genetic decoding. *Trends Microbiol*. 2014;22(12):686–696. doi:10.1016/j.tim.2014.09.003
67. Di Tella D, Tamburro M, Guerrizio G, Fanelli I, Sammarco ML, Ripabelli G. Molecular epidemiological insights into colistin-resistant and carbapenemase-producing clinical *Klebsiella pneumoniae* isolates. *Infect Drug Resist*. 2019;12:3783–3795. doi:10.2147/IDR.S226416
68. Liu J, Yu J, Chen F, et al. Emergence and establishment of KPC-2-producing ST11 *Klebsiella pneumoniae* in a general hospital in Shanghai, China. *Eur J Clin Microbiol Infect Dis*. 2018;37(2):293–299. doi:10.1007/s10096-017-3131-4
69. Gao H, Liu Y, Wang R, Wang Q, Jin L, Wang H. The transferability and evolution of NDM-1 and KPC-2 co-producing *Klebsiella pneumoniae* from clinical settings. *EBioMedicine*. 2020;51:102599. doi:10.1016/j.ebiom.2019.102599

70. Akgul F, Bozkurt I, Sunbul M, Esen S, Leblebicioglu H. Risk factors and mortality in the Carbapenem-resistant *Klebsiella pneumoniae* infection: case control study. *Pathog Glob Health*. 2016;110(7–8):321–325. doi:10.1080/20477724.2016.1254976
71. Jiao Y, Qin Y, Liu J, et al. Risk factors for carbapenem-resistant *Klebsiella pneumoniae* infection/colonization and predictors of mortality: a retrospective study. *Pathog Glob Health*. 2015;109(2):68–74. doi:10.1179/2047773215Y.0000000004
72. Zhang Y, Guo LY, Song WQ, Wang Y, Dong F, Liu G. Risk factors for carbapenem-resistant *K. pneumoniae* bloodstream infection and predictors of mortality in Chinese paediatric patients. *BMC Infect Dis*. 2018;18(1):248. doi:10.1186/s12879-018-3160-3
73. Zheng X, Wang J-F, Xu W-L, Xu J, Hu J. Clinical and molecular characteristics, risk factors and outcomes of carbapenem-resistant *Klebsiella pneumoniae* bloodstream infections in the intensive care unit. *Antimicrob Resist Infect Control*. 2017;6(1):102. doi:10.1186/s13756-017-0256-2
74. Falagas ME, Rafailidis PI, Kofteridis D, et al. Risk factors of carbapenem-resistant *Klebsiella pneumoniae* infections: a matched case control study. *J Antimicrob Chemother*. 2007;60(5):1124–1130. doi:10.1093/jac/dkm356
75. Hu Y, Ping Y, Li L, Xu H, Yan X, Dai H. A retrospective study of risk factors for carbapenem-resistant *Klebsiella pneumoniae* acquisition among ICU patients. *J Infect Dev Ctries*. 2016;10(3):208–213. doi:10.3855/jidc.6697

## Infection and Drug Resistance

Dovepress

### Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>