



OPEN ACCESS

CLINICAL SCIENCE

Interaction between the *STAT4* rs11889341(T) risk allele and smoking confers increased risk of myocardial infarction and nephritis in patients with systemic lupus erythematosus

Sarah Reid ,¹ Niklas Hagberg ,¹ Johanna K Sandling ,¹ Andrei Alexsson,¹ Pascal Pucholt ,¹ Christopher Sjöwall ,² Karoline Lerang,³ Andreas Jönsen,⁴ Iva Gunnarsson,⁵ Ann-Christine Syvänen,¹ Anne Margrethe Troldborg,^{6,7} Anne Voss,⁸ Anders A Bengtsson,⁴ Øyvind Molberg,³ Søren Jacobsen,^{9,10} Elisabet Svenungsson ,⁵ Lars Rönnblom ,¹ Dag Leonard¹

Handling editor Josef S Smolen

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2020-219727>).

For numbered affiliations see end of article.

Correspondence to

Professor Lars Rönnblom, Medical Sciences and Science for Life Laboratory, Uppsala University, Uppsala 751 85, Sweden; Lars.Ronnblom@medsci.uu.se

Received 15 December 2020
Revised 10 February 2021
Accepted 6 March 2021
Published Online First
25 March 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Reid S, Hagberg N, Sandling JK, et al. *Ann Rheum Dis* 2021;**80**:1183–1189.

ABSTRACT

Objective To investigate how genetics influence the risk of smoking-related systemic lupus erythematosus (SLE) manifestations.

Methods Patients with SLE ($n_{\text{discovery cohort}}=776$, $n_{\text{replication cohort}}=836$) were genotyped using the 200K Immuchip single nucleotide polymorphisms (SNP) Array (Illumina) and a custom array. Sixty SNPs with SLE association ($p<5.0\times 10^{-8}$) were analysed. Signal transducer and activator of transcription 4 (*STAT4*) activation was assessed in *in vitro* stimulated peripheral blood mononuclear cells from healthy controls ($n=45$).

Results In the discovery cohort, smoking was associated with myocardial infarction (MI) (OR 1.96 (95% CI 1.09 to 3.55)), with a greater effect in patients carrying any rs11889341 *STAT4* risk allele (OR 2.72 (95% CI 1.24 to 6.00)) or two risk alleles (OR 8.27 (95% CI 1.48 to 46.27)).

Smokers carrying the risk allele also displayed an increased risk of nephritis (OR 1.47 (95% CI 1.06 to 2.03)). In the replication cohort, the high risk of MI in smokers carrying the risk allele and the association between the *STAT4* risk allele and nephritis in smokers were confirmed (OR 6.19 (95% CI 1.29 to 29.79) and 1.84 (95% CI 1.05 to 3.29), respectively).

The interaction between smoking and the *STAT4* risk allele resulted in further increase in the risk of MI (OR 2.14 (95% CI 1.01 to 4.62)) and nephritis (OR 1.53 (95% CI 1.08 to 2.17)), with 54% (MI) and 34% (nephritis) of the risk attributable to the interaction. Levels of interleukin-12-induced phosphorylation of *STAT4* in CD8+ T cells were higher in smokers than in non-smokers (mean geometric fluorescence intensity 1063 vs 565, $p=0.0063$).

Lastly, the *IL12A* rs564799 risk allele displayed association with MI in both cohorts (OR 1.53 (95% CI 1.01 to 2.31) and 2.15 (95% CI 1.08 to 4.26), respectively).

Conclusions Smoking in the presence of the *STAT4* risk gene variant appears to increase the risk of MI and nephritis in SLE. Our results also highlight the role of the IL12–*STAT4* pathway in SLE-cardiovascular morbidity.

Key messages**What is already known about this subject?**

► Neither traditional nor systemic lupus erythematosus (SLE)-related risk factors can fully account for the excess cardiovascular disease risk seen in patients with SLE, but interactions between traditional and SLE-specific risk factors have been scarcely investigated.

What does this study add?

► Our results show that the signal transducer and activator of transcription 4 (*STAT4*) risk allele rs11889341 enhances the effect of smoking on the risk of myocardial infarction and nephritis and that smoking is associated with increased interleukin (IL)-12-induced phosphorylation of *STAT4* in CD8+ T cells.
► We further demonstrate that the *IL12A* SLE risk variant rs564799 is associated with an increased risk of myocardial infarction, which further highlights the importance of the IL12–*STAT4* pathway in the aetiology of cardiovascular morbidity in SLE.

How might this impact on clinical practice or future developments?

► Our results suggest that genetic profiling of patients with SLE may be useful for predicting comorbidities of the disease, impact of environmental factors and for targeted smoking cessation interventions.

INTRODUCTION

Systemic lupus erythematosus (SLE) is a disease characterised by loss of tolerance to self-antigens, formation of immune complexes and an activated type I interferon (IFN) system.¹ A widely accepted view of the aetiology of SLE is that environmental factors trigger the disease in genetically susceptible individuals. The genetic background is complex, with more than 100 single nucleotide polymorphisms (SNPs) associated with risk

for SLE.² Exposure to certain environmental factors, including ultraviolet radiation and viral infections, is associated with SLE development and flare-ups of the disease.^{3,4} Several studies have evaluated smoking as a risk factor for SLE, with the largest meta-analysis to date showing a modest risk increase.⁵ While the results are not confirmed in prospective studies, both Cozier and Barbhuiya *et al* observed a trend of increased risk in smokers.^{6,7} The most extensive prospective study involving 286 cases with SLE demonstrated an association between smoking and development of SLE with increased anti-dsDNA, but no risk of overall SLE.⁷

Although death from active SLE has decreased since the 1950s,⁸ the mortality rate still exceeds that of the general population, with cardiovascular morbidity remaining considerably high and a strong risk factor for premature mortality.^{9–11} Both traditional and SLE-related risk factors, such as hypertension, nephritis and high disease activity have been identified as risk factors, but cannot fully account for the excess cardiovascular disease (CVD) risk seen in patients with SLE.^{12,13} To fully explain the aetiology of SLE or its comorbidities such as CVD, gene–gene or gene–environment interactions may be essential to consider. In rheumatoid arthritis, there is compelling evidence of a strong interaction between the HLA-DRB1*04 shared epitope and smoking on the development of anticitrullinated protein

autoantibodies¹⁴ and a high prevalence of cardiovascular events (CVE).¹⁵

In SLE, a few studies have investigated the interaction between genetic risk factors and smoking on the development of the disease.^{16,17} Recently, Cui *et al* demonstrated that an additive interaction between smoking and the cumulative genetic risk of SLE increases the risk of the disease.¹⁸ However, gene–smoking interactions on the development of specific manifestations or co-morbidities of SLE have been scarcely studied. This study, therefore, aims to investigate the effect of smoking on the development of specific manifestations of SLE, including CVE, end-stage renal disease (ESRD) and nephritis, and examine how the effect is modulated by the presence of genetic variants associated with an increased risk of SLE development.

METHODS

Patients of the discovery and replication cohort

The discovery cohort included 774 patients with SLE from Sweden. The replication cohort included 836 patients from Norway and Denmark. All subjects fulfilled ≥4 American College of Rheumatology (ACR)–82 and ACR-97 classification criteria for SLE and were of European descent. Clinical characteristics of the cohorts are described in table 1 and online supplemental

Table 1 Prevalence of clinical manifestations in smokers (n=371) and non-smokers (n=387) in the discovery cohort

	Smokers, n (%)	Non-smokers, n (%)	OR (95% CI)	P value
Age at last follow-up, mean (SD)	55 (15)	50 (17)		0.000025
Disease duration, mean (SD)	17 (11)	16 (12)		0.016
Male sex	49 (13)	48 (13)	0.94 (0.61 to 1.45)	0.79
Deceased at follow-up	55 (15)	39 (11)	1.19 (0.75 to 1.88)	0.45
ACR 1982 classification criteria ⁴⁰				
1 Malar rash	205 (55)	211 (57)	1.00 (0.74 to 1.34)	0.97
2 Discoid rash	83 (22)	80 (21)	0.96 (0.68 to 1.36)	0.81
3 Photosensitivity	262 (71)	246 (65)	1.27 (0.93 to 1.74)	0.13
4 Oral ulcer	103 (28)	96 (25)	1.20 (0.86 to 1.67)	0.27
5 Arthritis	306 (82)	301 (81)	1.11 (0.76 to 1.61)	0.59
6 Serositis	179 (48)	165 (45)	1.10 (0.82 to 1.47)	0.52
7 Renal disorder	129 (35)	132 (36)	1.10 (0.80 to 1.50)	0.56
8 Neurological disorder	33 (9)	40 (10)	0.86 (0.52 to 1.40)	0.54
9 Haematological disorder	213 (57)	258 (70)	0.62 (0.46 to 0.84)	0.0021
10 Immunological disorder	245 (66)	256 (69)	0.97 (0.71 to 1.33)	0.86
Anti-dsDNA	224 (61)	231 (63)	1.02 (0.76 to 1.38)	0.88
11 ANA	364 (98)	367 (99)	0.69 (0.19 to 2.49)	0.57
Renal variables				
WHO class I–II	12 (5)	20 (8)	0.71 (0.33 to 1.50)	0.37
WHO class III–IV	62 (21)	71 (22)	1.13 (0.75 to 1.69)	0.57
WHO class V	15 (6)	16 (6)	1.05 (0.50 to 2.20)	0.90
Other*	11 (4)	6 (2)	1.24 (0.50 to 3.06)	0.65
ESRD	14 (4)		1.41 (0.62 to 3.24)	0.42
Cardiovascular events				
MI	39 (11)	19 (5)	1.96 (1.09 to 3.55)	0.025
ICVD	45 (12)	30 (8)	1.37 (0.84 to 2.24)	0.21
VTE	61 (16)	52 (14)	1.14 (0.76 to 1.71)	0.52
Clinical APS	68 (20)	61 (18)	1.08 (0.73 to 1.60)	0.68
Anti-β2GP-I IgG	58 (19)	57 (18)	0.98 (0.69 to 1.39)	0.91
Anti-β2GP-I IgM	8 (11)	11 (12)	0.96 (0.36 to 2.55)	0.93
LA	62 (23)	57 (21)	1.19 (0.79 to 1.80)	0.40
aCL-IgG	86 (26)	90 (28)	1.05 (0.70 to 1.59)	0.080
aCL-IgM	34 (14)	33 (13)	0.96 (0.36 to 2.55)	0.93

Logistic regression models were used to assess differences between smokers and non-smokers. All analyses were adjusted for age at last follow-up and disease duration. p<0.05 (unadjusted for multiple comparisons) in bold.

*Patients with biopsies displaying signs of nephritis but not meeting the criteria for any of the above classes were classified as other.

ACR, American College of Rheumatology; ANA, antinuclear antibodies; Anti-β2GP-I, anti-β₂ Glycoprotein-I; APS, antiphospholipid syndrome; dsDNA, double-stranded DNA; ESRD, end-stage renal disease; ICVD, ischaemic cerebrovascular disease; LA, lupus anticoagulant; MI, myocardial infarction; VTE, venous thromboembolism.

table 1, respectively. Clinical data, including smoking status (ever-smoker, including current or a history of smoking, versus never-smoker), the ACR-82 classification criteria, antiphospholipid syndrome diagnosis, ESRD, renal biopsy data and CVE, defined as myocardial infarction (MI), ischaemic cerebrovascular disease (ICVD) or venous thromboembolism, was collected from medical records. For definitions, see online supplemental file. In the replication cohort, data regarding ESRD were not available, and data on smoking were available from 503 patients.

Genotyping and selection of SNPs

Genotyping of the discovery cohort was performed using the Illumina 200K ImmunoChip SNP array, for details, see online supplemental file. SNPs previously associated with SLE at genome-wide significance in the European population² were selected. For SNPs not included on the ImmunoChip, the SNP-proxy with the highest linkage disequilibrium (LD) ($r^2 \geq 0.96$) was selected. All SNPs were filtered for independent signals, removing the variant with the lowest SLE-OR for SNPs in LD ($r^2 > 0.2$). In total, 4 HLA and 56 non-HLA SNPs were investigated for associations with MI (online supplemental table 2). Individuals in the replication cohort were genotyped for three single nucleotide variants using a custom assay on the MassARRAY system (see online supplemental file).

Interleukin-12-induced phosphorylation of STAT4

Interleukin 12 (IL-12)-induced phosphorylation of signal transducer and activator of transcription 4 (pSTAT4) was previously determined in 72 healthy blood donors from Uppsala Bioresource using flow cytometry.¹⁹ Smoking data were available from 45 of these donors, of which 20 were past or current smokers and 25 were non-smokers.

Statistical analysis

To investigate associations between smoking and clinical manifestations, logistic regression models were used. As smoking was associated with longer disease duration and higher age at follow-up (table 1), these variables were included as covariates. In analysis of associations between genetic variants and MI, SNPs were first analysed separately. Next, all variants demonstrating a positive association with MI were included in a forwards conditional multiple regression model. All analyses were adjusted for age and disease duration. Results considered statistically significant (unadjusted $p < 0.05$) were reanalysed in the replication cohort using the same statistical model and covariates. Meta-analyses were performed on the two datasets and multiplicative and additive interactions between the *STAT4* risk allele and smoking were studied in a combined dataset through addition

of a *STAT4**smoking interaction term in the logistic models and by calculating the attributable proportion due to interaction, respectively.^{20,21} Differences in levels of pSTAT4 were assessed by Student's t-test and by a linear regression model allowing adjustment for age and the *STAT4* risk allele. R²² was used for all analyses except the meta-analyses which were performed in PLINK.²³

RESULTS

Smoking is modestly associated with MI

Initially, we assessed the association between smoking and clinical manifestations (table 1). We found no evidence of any associations between smoking and the ACR criteria, except the haematological criterion, which was less prevalent in smokers (table 1). Elevated levels of red and white blood cells in smokers is a well-known phenomenon.²⁴ Smoking was not associated with a history of DVT or ICVD, however, a significant association between smoking and MI was observed (OR 1.96 (95% CI 1.09 to 3.55), $p = 0.025$) (table 1).

Increased risk of MI in SLE-smokers with the *STAT4* risk allele

Next, we asked whether there are sub-groups of patients in which smoking plays a more prominent role in MI development. We initially examined 60 SNPs with established association with SLE ($p < 5.0 \times 10^{-8}$) for association with MI (online supplemental table 2). We found that the Neutrophil Cytosolic Factor 2 (*NCF2*), Interleukin-12A (*IL12A*) and *STAT4* risk alleles displayed independent, positive association with MI (table 2). In addition, patients carrying two alleles of both the *STAT4* and the *IL12A* risk variants ($n = 37$, 4.9% of the patients) displayed a substantially higher prevalence of MI compared with those with any other allele combination (27% vs 7%) (OR 5.88 (95% CI 2.44 to 14.17), $p = 7.9 \times 10^{-5}$) (figure 1A).

Next, we stratified patients by smoking status to determine whether each of the three SNPs displayed stronger association with MI in smokers. No significant associations were found for the *NCF2* or *IL12A* risk alleles (OR 1.58 (95% CI 0.89 to 2.78), $p = 0.12$ and OR 1.36 (95% CI 0.89 to 2.08), $p = 0.15$, respectively). However, the *STAT4* risk allele demonstrated a stronger association in smokers (OR 2.45 (95% CI 1.46 to 4.19), $p = 0.00086$) (figure 2A). Next, we assessed the association between smoking and MI in patients carrying the *STAT4* risk allele and observed an almost 3-fold increase in risk for the smokers compared with the non-smokers (OR 2.72 (95% CI 1.24 to 6.00), $p = 0.013$). In patients carrying two risk alleles, the risk was more than eightfold higher for smokers (OR 8.27 (95% CI 1.48 to 46.27), $p = 0.016$). In contrast, we could not demonstrate a significant association between smoking and MI

Table 2 Associations between SLE risk SNPs and myocardial infarction in the discovery and replication cohort

SNP	Gene name	Discovery cohort (n=763)				P value	Replication cohort (n=836)				
		HWE p	Risk allele frequency				HWE p	Risk allele frequency			P value
			MI+	MI-	OR			MI+	MI-	OR	
rs17849502	<i>NCF2</i>	0.013	0.12	0.08	2.00 (1.08 to 3.68)	0.027	8.80×10^{-7}	0.18	0.11	1.94 (0.94 to 4.04)	0.075
rs11889341	<i>STAT4</i>	0.81	0.44	0.34	1.76 (1.18 to 2.63)	0.0054	0.53	0.40	0.31	1.81 (0.94 to 3.47)	0.075
rs564799	<i>IL12A</i>	0.29	0.68	0.59	1.53 (1.01 to 2.31)	0.042	1.00	0.74	0.61	2.15 (1.08 to 4.26)	0.029

Using a forward conditional multiple logistic regression model, 60 genetic variants with previously established association with SLE ($p < 5 \times 10^{-8}$, online supplemental table 2) were analysed for associations with myocardial infarction in the discovery cohort. The table shows SNPs included in the final model. These SNPs were subsequently analysed in the replication cohort. Age at follow-up and disease duration were included as covariates. $P < 0.05$ (unadjusted for multiple comparisons) in bold. HWE was tested on all patients in the discovery and replication cohorts, respectively.

HWE, Hardy-Weinberg equilibrium; IL12a, Interleukin12A; MI, myocardial infarction; NCF2, neutrophil cytosolic factor 2; SLE, systemic lupus erythematosus; SNPs, single-nucleotide polymorphisms; STAT4, signal transducer and activator of transcription 4.




Figure 1 Prevalence of myocardial infarction (MI) in patients with different numbers of *STAT4* and *IL12A* risk alleles. The prevalence of MI for patients in the discovery (A) and the replication (B) cohorts, stratified by number of risk alleles of *STAT4* (rs11889341) and *IL12A* (rs564799). Patients carrying two risk alleles of both *STAT4* and *IL12A* ($n_{\text{discovery cohort}}=37$ (4.9%), $n_{\text{replication cohort}}=28$ (3.4%)) were compared with the remaining patients using logistic regression, adjusting for age at follow-up and disease duration. IL12, interleukin-12; *STAT4*, Signal Transducer and Activator of Transcription 4.

in patients without the risk allele (OR 1.20 (95% CI 0.49 to 2.96) $p=0.55$).

As patients with nephritis have previously been shown to have a higher prevalence of both MI and the *STAT4* risk allele.^{25–27} we hypothesised that the results would be similar if using nephritis, rather than MI, as the outcome variable. Without stratifying for smoking, the association between the *STAT4* risk allele and nephritis reached suggestive significance (OR 1.23 (95% CI 0.98 to 1.54), $p=0.072$). The effect was more pronounced in the smokers only (OR 1.47 (95% CI 1.06 to 2.03), $p=0.020$). In addition, we found moderate evidence that patients with nephritis carrying the *STAT4* risk allele were at a greater risk of developing ESRD (OR 1.85 (95% CI 0.96 to 3.59), $p=0.068$), and this risk was enhanced in smokers (OR 2.52 (95% CI 1.04 to 6.10), $p=0.040$) (figure 2B). Of note, despite the non-smoking group including more patients with nephritis ($n=140$ vs $n=129$), no evidence of an association between the *STAT4* risk allele and nephritis or ESRD could be demonstrated in this group (OR 1.07 (95% CI 0.77 to 1.46), $p=0.70$ and OR 1.10 (95% CI 0.38 to 3.16), $p=0.86$, respectively).

To validate our significant findings, we performed the same analyses in an independent cohort of patients with SLE (online supplemental table 1). Analysis of the genetic variants demonstrated that the *IL12A* risk allele was the only gene variant significantly associated with MI when not accounting for smoking

(table 2). Patients with two risk alleles of both *STAT4* and *IL12A* ($n=28$, 3.2% of the patients) were found to have a significantly higher prevalence of MI than patients without this combination of risk alleles (OR 7.21 (95% CI 1.36 to 38.27, $p=0.020$) (figure 1B). Similar to in the discovery cohort, we found a significant association between the *STAT4* risk allele and MI in smokers (OR 2.11 (95% CI 1.04 to 4.26), $p=0.038$). In addition, smoking was associated with MI in patients carrying the *STAT4* risk allele (OR 6.19 (95% CI 1.29 to 29.79), $p=0.023$). No evidence of these associations could be observed in the non-smoking group (OR 0.58 (95% CI 0.09 to 3.90), $p=0.58$ and OR 1.32 (95% CI 0.23 to 7.34), $p=0.75$, respectively). In patients carrying two risk alleles ($n=51$), the logistic regression could not be performed due to a ‘perfect separation’ between groups, with 9% of smokers having had a MI compared with 0% of never-smokers. As in the discovery cohort, we found an association between the *STAT4* risk allele and nephritis in smokers (OR 1.84 (95% CI 1.05 to 3.29), $p=0.035$), whereas the effect size was non-significant in non-smokers (OR 0.78 (95% CI 0.41 to 1.45), $p=0.44$).

Meta-analysis of the two cohorts showed the risk of MI to more than double with each additional *STAT4* risk allele in smokers (OR 2.28, $p=0.00010$). In contrast, no association could be detected in the never-smoker group (OR 0.80, $p=0.52$). Similarly, in meta-analysis of patients with 2 *STAT4* risk alleles ($n=282$, 22%), smoking was found to be a strong risk factor for MI (OR 8.27, $p=0.016$). For nephritis, each *STAT4* risk allele increased the risk by ~50% in smokers (OR 1.52, $p=0.00051$), whereas no increase in risk was found in the non-smoker group (OR 0.82, $p=0.33$).

Interaction between *STAT4* risk allele and smoking results in a higher risk of MI and nephritis

We subsequently performed interaction analyses on all patients and found a significant multiplicative interaction between the *STAT4* risk allele and smoking on the development of MI (OR 2.14 (95% CI 1.01 to 4.62), $p=0.049$) as well as nephritis (OR 1.53 (95% CI 1.08 to 2.17), $p=0.020$) (online supplemental table 3). Next, we examined additive interaction and observed an attributable proportion due to interaction of 0.54 (95% CI 0.24 to 0.83, $p=0.00019$) and 0.34 (95% CI 0.080 to 0.61, $p=0.0051$) for MI and nephritis, respectively.

To determine whether the effect of the *STAT4* risk allele on nephritis in smokers could explain the association with MI, we performed stratification of the combined dataset and investigated the association between the *STAT4* risk allele and MI in smokers and non-smokers without nephritis. In the smokers, the association between the *STAT4* risk allele and MI remained significant ($n=428$, OR 2.43 (95% CI 1.40 to 4.27), $p=0.0017$) (figure 3). Similarly, the association between the *STAT4* risk allele and nephritis in smokers remained significant after excluding patients with MI from the analysis ($n=623$, OR 1.54 (95% CI 1.20 to 1.98), $p=0.00076$).

Lastly, as both SLE risk alleles in *STAT4* and smoking have shown association with the development of aPL in previous studies,^{28 29} we assessed the association between the *STAT4* risk allele and aPL in smokers, however, it was not significant (OR 1.56 (95% CI 0.71 to 3.72, $p=0.26$). Next, we performed a multiple regression model in the smoking group including the *STAT4* risk allele, any aPL, nephritis, age at follow-up, and disease duration as covariates. We found the association between the *STAT4* risk allele and MI to remain significant (OR 3.26 (95% CI 1.15 to 9.20), $p=0.026$), whereas neither aPL nor




Figure 2 Prevalence of myocardial infarction (MI) and end-stage renal disease (ESRD) in the discovery cohort. Prevalence of MI (A) and ESRD (B) in the discovery cohort in all patients ($n=776$), smokers only ($n=371$) and non-smokers only ($n=405$), stratified by the number of *STAT4* (rs11889341) risk alleles. Patients with 0, 1 or 2 risk alleles in each group were compared using logistic regression, adjusting for age at follow-up and disease duration. *STAT4*, Signal Transducer and Activator of Transcription 4.




Figure 3 The prevalence of myocardial infarction (MI) in patients without nephritis. To investigate whether the association between the rs11889341 (*STAT4*) allele and MI was dependent on the association between the *STAT4*–nephritis association, all patients with nephritis were excluded and the prevalence of MI was subsequently plotted for smoking (A) and non-smoking (B) patients with 0, 1 or 2 of the *STAT4* risk allele. Patients with 0, 1 or 2 risk alleles in each group were compared using logistic regression, adjusting for age at follow-up and disease duration. *STAT4*, signal transducer and activator of transcription 4.

nephritis displayed significant association with MI ($p=0.87$ and $p=0.50$, respectively), (online supplemental table 4).

Levels of activated *STAT4* are increased in smokers

The *STAT4* risk allele is associated with higher levels of p*STAT4* in CD8+ T cells from SLE patients on stimulation with IL-12.¹⁹ Furthermore, higher levels of *STAT4* expression have demonstrated association with increased cardiovascular damage in patients with SLE.³⁰ To investigate whether smoking also increases the levels of p*STAT4*, we analysed IL-12 stimulated CD8+ T cells from healthy blood donors who were smokers ($n=20$) or never-smokers ($n=25$). We found the levels of p*STAT4* to be higher in smokers ($p=0.0063$), with a mean value of 1063 compared with 565 in non-smokers (figure 4). When adjusting for age and the *STAT4* risk allele, which is in LD ($r^2=1.00$) with a *STAT4* risk variant previously shown to influence levels of p*STAT4* in these individuals,^{19,31} the association between smoking and levels of p*STAT4* remained significant ($\beta=396$, $p=0.023$).




Figure 4 Levels of p*STAT4* in CD8+ T cells after stimulation with IL-12. The levels of IL-12–induced p*STAT4* were compared between healthy blood donors who were smokers (current or past) ($n=20$) and never-smokers ($n=25$) by Student's t-test. IL-12, interleukin-12; p*STAT4*, phosphorylated Signal Transducer and Activator of Transcription 4.

DISCUSSION

In the present study, we demonstrate that smoking substantially increases the risk of MI in a subset of patients with SLE carrying a variant of the *STAT4* SLE-risk gene. In both the discovery and replication cohorts, the effect size increased with an increasing number of *STAT4* risk alleles, with smoking giving rise to a more than 8-fold risk of MI in homozygous individuals. We believe that our results add important knowledge in the understanding of how SLE-risk alleles can modulate the effect of traditional risk factors.

The prevalence of MI is higher in SLE patients with nephritis than patients without renal manifestations^{26,32} and SLE-risk alleles in *STAT4* have previously been linked to both nephritis and severe renal insufficiency.^{25,27} We, therefore, speculated that the smoking-*STAT4* risk allele interaction did not directly affect MI, but rather, was a consequence of an interaction between the *STAT4* risk allele and smoking on the development of nephritis. Indeed, we found that this gene-environment combination also results in a higher risk of nephritis, as well as ESRD. Interestingly, however, the *STAT4* risk allele/smoking effect on MI did not decrease when adjusting the model for nephritis or when completely removing the patients with nephritis from the analysis. Similarly, the effect of the *STAT4* risk allele/smoking on nephritis remained significant after excluding all patients with MI from the analysis, indicating that the associations were independent.

Based on these results, we hypothesised that the increased risk in individuals who smoke and carry the risk allele is connected with the levels of activated *STAT4* in these individuals. Hagberg *et al* have shown that the rs7574865 *STAT4* risk allele—which is in perfect LD ($r^2=1.00$) with the SNP used in this study³¹—is associated with increased levels of p*STAT4* in activated CD8+T cells of SLE patients.¹⁹ Therefore, we assessed whether smoking elevates p*STAT4* in this cell type and found that the levels were almost twofold higher in smokers. This observation is in line with previous findings by Di Stefano *et al*, who demonstrated higher levels of p*STAT4* in bronchial T cells from healthy smokers compared with non-smokers.³³ When *STAT4* is activated and phosphorylated, it homodimerises and translocates to the nucleus where it induces expression of hundreds of genes, resulting in production of IFN- γ , T-helper type 1 and 17 differentiation and activation of monocytes.³⁴ Increased *STAT4* mRNA expression is associated with increased cardiovascular damage in patients with SLE,³⁰ and several studies on animal models indicate a link between *STAT4* and the development of atherosclerosis.³⁵ The mechanism of how smoking leads to increased levels of activated *STAT4* is unclear, however, we speculate that epigenetics may constitute the bridge between smoking, genetics and SLE. It is well known that smoking affects both overall DNA methylation and specific gene promoters.^{36,37} Epigenetic regulation is further believed to play an important role both in cardiovascular biology and in SLE development.^{36,38} Whether smoking is associated with epigenetic changes in SLE-specific genes, and if such changes are associated with specific manifestations of SLE, deserves further studies.

The analyses of individual SLE risk alleles identified the SLE-risk SNP *IL12A* to be associated with an increased risk of MI, and that patients in the discovery and replication cohort carrying two alleles of both the *IL12A* and *STAT4* risk SNPs had a more than fivefold and eightfold risk of MI, respectively. The *IL12A* SNP is located within the fourth intron of the *IL12A* gene, which encodes the p35 subunit of the IL-12 protein. On binding to its receptor, IL-12 induces phosphorylation of *STAT4*.³⁴ We

believe that the association of the *IL12A* risk allele, in addition to the *STAT4* risk allele, with MI points to the importance of this pathway in the development of the comorbidity. Previous work has demonstrated that JAK-inhibitors efficiently block the increase in pSTAT4 levels, and ameliorate murine lupus as well as its associated vascular dysfunction.^{19 39} Due to the potential therapeutic strategy of JAK-inhibitors for patients with SLE displaying an altered activity in this particular pathway, we believe that further studies of the effect of this pathway on development of CVE are warranted.

This study's strength is the large, well-characterised discovery cohort, that results were validated in a second large cohort, and the analysis of healthy control cells, which confirmed that pSTAT4 levels are higher in smokers. In addition, the quality control of genetic data was rigorous, and the patients' long mean follow-up time of 17 years allowed for many outcome variables such as MI to be recorded. There are, however, some limitations. First, our study is based on retrospective data and we lacked data on the year of smoking cessations. As patients who were past smokers at the last follow-up may have been active smokers at the time of their CVE, we could not analyse previous and current smoking separately. Second, we did not have data on number of pack-years, which may have generated more precise results. Third, the study includes only Scandinavian patients with SLE, and whether the associations are generalisable to patients of other ethnicities needs further investigation.

CONCLUSION

We demonstrate that smokers carrying the *STAT4* risk allele are at an increased risk of MI and nephritis and that the *IL12-STAT4* pathway may be important for the development of MI. Our results stress the importance of smoking cessation in SLE and particularly among those carrying this risk allele.

Author affiliations

¹Department of Medical Sciences, Rheumatology and Science for Life Laboratory, Uppsala University, Uppsala, Sweden

²Department of Biomedical and Clinical Sciences, Division of Inflammation and Infection, Linköping University, Linköping, Sweden

³Department of Rheumatology, Oslo University Hospital, Oslo, Norway

⁴Department of Clinical Sciences Lund, Rheumatology, Lund University, Skane University Hospital, Lund, Sweden

⁵Division of Rheumatology, Department of Medicine, Karolinska Institutet and Karolinska University Hospital, Stockholm, Sweden

⁶Department of Rheumatology, Aarhus University Hospital, Aarhus, Denmark

⁷Department of Biomedicine, Aarhus University, Aarhus, Denmark

⁸Department of Rheumatology, Odense University Hospital, Odense, Denmark

⁹Center for Rheumatology and Spine Diseases, Copenhagen University Hospital Rigshospitalet, Copenhagen, Denmark

¹⁰Institute of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark

Acknowledgements We thank the patients and healthy volunteers for participating, clinicians and researchers contributing to the replication cohort and Maija-Leena Eloranta, Cane Yaka, Rezvan Kiani and Karolina Tandre.

Contributors SR, DL and LR designed the study. SR, NH, JKS, AA, PP, CS, AJ, IG, A-CS, AMT, AV, AAB, ES, LR and DL collected the data for the discovery cohort. KL, AMT, AV, ØM and SJ collected the data for the replication cohort. SR, DL and LR analysed the data. SR, DL and LR wrote the manuscript. All authors revised the manuscript critically for important intellectual content and approved the final version of the manuscript.

Funding This work was supported by grants from the Swedish Research Council for Medicine and Health (D0283001), the Swedish Rheumatism Association, King Gustav V's 80-year Foundation, the Swedish Society of Medicine and the Ingegerd Johansson donation, Åke Wiberg's foundation, Gustafsson's foundation, Selander's foundation, Norsk Revmatikerforbunds Forskningsfond, Pahles legat og Stortuens legat and the Swedish Society for Medical Research (S20-0127).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The study protocol was approved by the Regional Ethical Review Board, Uppsala (DNR 2020–05065) and the local ethics committees.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Deidentified patient data and statistical code are available on request with project plan.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Sarah Reid <http://orcid.org/0000-0003-4065-6875>

Niklas Hagberg <http://orcid.org/0000-0003-2064-2716>

Johanna K Sandling <http://orcid.org/0000-0003-1382-2321>

Pascal Pucholt <http://orcid.org/0000-0003-3342-1373>

Christopher Sjöwall <http://orcid.org/0000-0003-0900-2048>

Elisabet Svenungsson <http://orcid.org/0000-0003-3396-3244>

Lars Rönnblom <http://orcid.org/0000-0001-9403-6503>

REFERENCES

- Rönnblom L, Leonard D. Interferon pathway in SLE: one key to unlocking the mystery of the disease. *Lupus Sci Med* 2019;6:e000270.
- Chen L, Morris DL, Vyse TJ. Genetic advances in systemic lupus erythematosus: an update. *Curr Opin Rheumatol* 2017;29:423–33.
- Barbhaiya M, Costenbader KH. Ultraviolet radiation and systemic lupus erythematosus. *Lupus* 2014;23:588–95.
- James JA, Neas BR, Moser KL, *et al*. Systemic lupus erythematosus in adults is associated with previous Epstein-Barr virus exposure. *Arthritis Rheum* 2001;44:1122–6.
- Jiang F, Li S, Jia C. Smoking and the risk of systemic lupus erythematosus: an updated systematic review and cumulative meta-analysis. *Clin Rheumatol* 2015;34:1885–92.
- Cozier YC, Barbhaiya M, Castro-Webb N, *et al*. Relationship of cigarette smoking and alcohol consumption to incidence of systemic lupus erythematosus in a prospective cohort study of black women. *Arthritis Care Res* 2019;71:671–7.
- Barbhaiya M, Tedeschi SK, Lu B, *et al*. Cigarette smoking and the risk of systemic lupus erythematosus, overall and by anti-double stranded DNA antibody subtype, in the nurses' health study cohorts. *Ann Rheum Dis* 2018;77:196–202.
- Tektonidou MG, Lewandowski LB, Hu J, *et al*. Survival in adults and children with systemic lupus erythematosus: a systematic review and Bayesian meta-analysis of studies from 1950 to 2016. *Ann Rheum Dis* 2017;76:2009–16.
- Björnådal L, Yin L, Granath F, *et al*. Cardiovascular disease a hazard despite improved prognosis in patients with systemic lupus erythematosus: results from a Swedish population based study 1964–95. *J Rheumatol* 2004;31:713–9.
- Gustafsson JT, Simard JF, Gunnarsson I, *et al*. Risk factors for cardiovascular mortality in patients with systemic lupus erythematosus, a prospective cohort study. *Arthritis Res Ther* 2012;14:R46.
- Ingvarsson RF, Landgren AJ, Bengtsson AA, *et al*. Good survival rates in systemic lupus erythematosus in southern Sweden, while the mortality rate remains increased compared with the population. *Lupus* 2019;28:1488–94.
- Esdaile JM, Abrahamowicz M, Grodzicky T, *et al*. Traditional Framingham risk factors fail to fully account for accelerated atherosclerosis in systemic lupus erythematosus. *Arthritis Rheum* 2001;44:2331–7.
- Boulos D, Koelmeyer RL, Morand EF, *et al*. Cardiovascular risk profiles in a lupus cohort: what do different calculators tell us? *Lupus Sci Med* 2017;4:e000212.
- Padyukov L, Silva C, Stolt P, *et al*. A gene-environment interaction between smoking and shared epitope genes in HLA-DR provides a high risk of seropositive rheumatoid arthritis. *Arthritis Rheum* 2004;50:3085–92.
- López-Longo FJ, Oliver-Miñarro D, de la Torre I, *et al*. Association between anti-cyclic citrullinated peptide antibodies and ischemic heart disease in patients with rheumatoid arthritis. *Arthritis Rheum* 2009;61:419–24.

- 16 Kiyohara C, Washio M, Horiuchi T, *et al.* Risk modification by CYP1A1 and GSTM1 polymorphisms in the association of cigarette smoking and systemic lupus erythematosus in a Japanese population. *Scand J Rheumatol* 2012;41:103–9.
- 17 Kiyohara C, Washio M, Horiuchi T, *et al.* Cigarette smoking, STAT4 and TNFRSF1B polymorphisms, and systemic lupus erythematosus in a Japanese population. *J Rheumatol* 2009;36:2195–203.
- 18 Cui J, Raychaudhuri S, Karlson EW, *et al.* Interactions between genome-wide genetic factors and smoking influencing risk of systemic lupus erythematosus. *Arthritis Rheumatol* 2020;72:1863–71.
- 19 Hagberg N, Joelsson M, Leonard D, *et al.* The STAT4 SLE risk allele rs7574865[T] is associated with increased IL-12-induced IFN- γ production in T cells from patients with SLE. *Ann Rheum Dis* 2018;77:1070–7.
- 20 Andersson T, Alfredsson L, Källberg H, *et al.* Calculating measures of biological interaction. *Eur J Epidemiol* 2005;20:575–9.
- 21 Mathur MB, VanderWeele TJ. R function for additive interaction measures. *Epidemiology* 2018;29:e5–6.
- 22 Team RC. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing, 2017.
- 23 Purcell S, Neale B, Todd-Brown K, *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;81:559–75.
- 24 Pedersen KM, Çolak Y, Ellervik C, *et al.* Smoking and increased white and red blood cells. *Arterioscler Thromb Vasc Biol* 2019;39:965–77.
- 25 Bolin K, Sandling JK, Zickert A, *et al.* Association of STAT4 polymorphism with severe renal insufficiency in lupus nephritis. *PLoS One* 2013;8:e84450.
- 26 Hermansen M-L, Lindhardtsen J, Torp-Pedersen C, *et al.* The risk of cardiovascular morbidity and cardiovascular mortality in systemic lupus erythematosus and lupus nephritis: a Danish nationwide population-based cohort study. *Rheumatology* 2017;56:kew475–15.
- 27 Taylor KE, Remmers EF, Lee AT, *et al.* Specificity of the STAT4 genetic association for severe disease manifestations of systemic lupus erythematosus. *PLoS Genet* 2008;4:e1000084.
- 28 Gustafsson JT, Gunnarsson I, Källberg H, *et al.* Cigarette smoking, antiphospholipid antibodies and vascular events in systemic lupus erythematosus. *Ann Rheum Dis* 2015;74:1537–43.
- 29 Svenungsson E, Gustafsson J, Leonard D, *et al.* A STAT4 risk allele is associated with ischaemic cerebrovascular events and anti-phospholipid antibodies in systemic lupus erythematosus. *Ann Rheum Dis* 2010;69:834–40.
- 30 Uzrail AH, Assaf AM, Abdalla SS. Correlations of Expression Levels of a Panel of Genes (*IRF5*, *STAT4*, *TNFSF4*, *MECP2*, and *TLR7*) and Cytokine Levels (IL-2, IL-6, IL-10, IL-12, IFN- γ , and TNF- α) with Systemic Lupus Erythematosus Outcomes in Jordanian Patients. *Biomed Res Int* 2019;2019:1–11.
- 31 Machiela MJ, Chanock SJ. LDlink: a web-based application for exploring population-specific haplotype structure and linking correlated alleles of possible functional variants. *Bioinformatics* 2015;31:3555–7.
- 32 Gustafsson JT, Herlitz Lindberg M, Gunnarsson I, *et al.* Excess atherosclerosis in systemic lupus erythematosus, -A matter of renal involvement: Case control study of 281 SLE patients and 281 individually matched population controls. *PLoS One* 2017;12:e0174572.
- 33 Di Stefano A, Caramori G, Capelli A, *et al.* Stat4 activation in smokers and patients with chronic obstructive pulmonary disease. *Eur Respir J* 2004;24:78–85.
- 34 Yang C, Mai H, Peng J, *et al.* Stat4: an immunoregulator contributing to diverse human diseases. *Int J Biol Sci* 2020;16:1575–85.
- 35 Dobrian AD, Hatcher MA, Brotman JJ, *et al.* Stat4 contributes to adipose tissue inflammation and atherosclerosis. *J Endocrinol* 2015;227:13–24.
- 36 Breitling LP. Current genetics and epigenetics of smoking/tobacco-related cardiovascular disease. *Arterioscler Thromb Vasc Biol* 2013;33:1468–72.
- 37 Zong D, Liu X, Li J, *et al.* The role of cigarette smoke-induced epigenetic alterations in inflammation. *Epigenetics Chromatin* 2019;12:65.
- 38 Miceli-Richard C. Epigenetics and lupus. *Joint Bone Spine* 2015;82:90–3.
- 39 Furumoto Y, Smith CK, Blanco L, *et al.* Tofacitinib ameliorates murine lupus and its associated vascular dysfunction. *Arthritis Rheumatol* 2017;69:148–60.
- 40 Tan EM, Cohen AS, Fries JF, *et al.* The 1982 revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1982;25:1271–7.