

# AnCF, the CCAAT Binding Complex of *Aspergillus nidulans*, Contains Products of the *hapB*, *hapC*, and *hapE* Genes and Is Required for Activation by the Pathway-Specific Regulatory Gene *amdR*

STEFAN STEIDL,<sup>1,2†</sup> PETER PAPAGIANNOPOULOS,<sup>1</sup> OLIVIER LITZKA,<sup>2</sup> ALEX ANDRIANOPOULOS,<sup>1</sup> MERYL A. DAVIS,<sup>1</sup> AXEL A. BRAKHAGE,<sup>2†</sup> AND MICHAEL J. HYNES<sup>1\*</sup>

*Department of Genetics, University of Melbourne, Parkville, Victoria 3052, Australia,<sup>1</sup> and Institut fuer Genetik und Mikrobiologie, Ludwig-Maximilians-Universität, Munich 80638, Germany<sup>2</sup>*

Received 2 July 1998/Returned for modification 25 August 1998/Accepted 29 September 1998

CCAAT binding factors (CBFs) positively regulating the expression of the *amdS* gene (encoding acetamidase) and two penicillin biosynthesis genes (*ipnA* and *aatA*) have been previously found in *Aspergillus nidulans*. The factors were called AnCF and PENR1, respectively. Deletion of the *hapC* gene, encoding a protein with significant similarity to Hap3p of *Saccharomyces cerevisiae*, eliminated both AnCF and PENR1 binding activities. We now report the isolation of the genes *hapB* and *hapE*, which encode proteins with central regions of high similarity to Hap2p and Hap5p of *S. cerevisiae* and to the CBF-B and CBF-C proteins of mammals. An additional fungus-specific domain present in HapE was revealed by comparisons with the homologs from *S. cerevisiae*, *Neurospora crassa*, and *Schizosaccharomyces pombe*. The HapB, HapC, and HapE proteins have been shown to be necessary and sufficient for the formation of a CCAAT binding complex in vitro. Strains with deletions of each of the *hapB*, *hapC*, and *hapE* genes have identical phenotypes of slow growth, poor conidiation, and reduced expression of *amdS*. Furthermore, induction of *amdS* by omega amino acids, which is mediated by the AmdR pathway-specific activator, is abolished in the *hap* deletion mutants, as is growth on  $\gamma$ -aminobutyric acid as a sole nitrogen or carbon source. AmdR and AnCF bind to overlapping sites in the promoters of the *amdS* and *gatA* genes. It is known that AnCF can bind independently of AmdR. We suggest that AnCF binding is required for AmdR binding in vivo.

The sequence CCAAT is found in the 5' region of approximately 30% of eukaryotic genes between 50 and 200 bp from the start point of transcription and can be present in either orientation (8). There is evidence for a conserved multimeric protein activator that recognizes this sequence in a wide variety of organisms (reviewed in references 31 and 32). The first complex identified was the Hap complex of *Saccharomyces cerevisiae*, which consists of four subunits, Hap2p, Hap3p, Hap4p, and Hap5p (33, 39, 42). Hap2p, Hap3p, and Hap5p are essential for the formation of a DNA binding complex, while Hap4p is not directly involved in DNA binding but carries a functional activation domain (18, 53). The Hap complex is required for the expression of genes involved in oxidative phosphorylation in response to growth on nonfermentable carbon sources (42). In addition, genes not involved in respiration have been found to be subject to regulation by the Hap complex (13, 14). Conserved homologs of HAP2, HAP3, and HAP5 have been found in a variety of organisms (reviewed in reference 32). The mammalian CCAAT binding factor (CBF; also known as NF-Y) consists of three subunits, CBF-A, -B, and -C; in this case, CBF-B and CBF-C have been found to carry transcriptional activation domains. Evidence for a fourth subunit equivalent to Hap4p is lacking (11; reviewed in reference 31).

In the filamentous fungus *Aspergillus nidulans*, CCAAT sequences in the promoters of the *amdS* gene (encoding the catabolic enzyme acetamidase) and the *acvA*, *ipnA*, and *aatA* genes (encoding enzymes for biosynthesis of the secondary metabolite penicillin) have been shown to affect the expression of these genes (27, 29, 49). Proteins binding to these sequences have been detected—AnCF in the case of *amdS* (51) and PENR1 in the case of the penicillin biosynthesis genes (29, 49). Similarly, a protein designated AnCP from *A. nidulans* has been found to bind to a CCAAT sequence present in the Taka-amylase gene promoter of *Aspergillus oryzae* (24, 36).

The *hapC* gene of *A. nidulans* was identified as a homolog of the HAP3 gene of *S. cerevisiae* (40). Deletion of this gene resulted in low levels of expression of *amdS*, *aatA*, and *ipnA* reporter gene fusions (28, 40). In addition, deletion of *hapC* resulted in loss of binding of AnCF and PENR1 to CCAAT sequences derived from *amdS*, the penicillin biosynthesis genes, and the Taka-amylase gene, indicating that these complexes contain a *hapC*-encoded component (25, 28, 40). This was confirmed directly in assays using anti-HapC antiserum (25, 28). A CCAAT binding complex was reconstituted by using recombinant HapC together with expressed Hap2p and Hap5p from *S. cerevisiae* (25).

We have now cloned genes encoding two additional components of the AnCF complex. These genes, *hapB* and *hapE*, encode proteins with a central core with a high similarity to Hap2p and Hap5p of *S. cerevisiae* as well as to the corresponding mammalian proteins. We have shown that the HapB, HapC, and HapE subunits are necessary and sufficient for DNA binding by reconstitution of the complex in vitro. Furthermore, deletions of the *hapB* and *hapE* genes result in loss

\* Corresponding author. Mailing address: Department of Genetics, University of Melbourne, Parkville, Victoria 3052, Australia. Phone: (61 3) 9344 6246. Fax: (61 3) 9344 5139. E-mail: hynes.lab@genetics.unimelb.edu.au.

† Present address: Institut für Mikrobiologie und Genetik, Technische Universität Darmstadt, 64287 Darmstadt, Germany.

TABLE 1. *A. nidulans* strains used in this study

Strain	Genotype <sup>a</sup>
MH54	..... <i>biA1; niiA4</i>
MH3018	..... <i>yA1, pabaA1; amdS368 argB2; amdA7</i>
MH3408	..... <i>biA1; amdS::lacZ; niiA4</i>
MH9092	..... <i>yA1, pabaA1; amdS368 argB2; amdA7; hapEΔ::argB</i>
MH9099	..... <i>yA1, pabaA1; amdS368 argB2; amdA7; hapBΔ::argB</i>
MH9094	.....MH9092 transformed with pPTR2
MH9210	.....MH9099 transformed with pBShapB4.0
MH9206	..... <i>yA1, pabaA1; amdS::lacZ; hapEΔ::argB</i>
MH9207	..... <i>yA1, pabaA; amdS::lacZ; hapBΔ::argB</i>
MH8194	..... <i>hapCΔ::riboB, amdS::lacZ; pyroA4, niiA4</i>

<sup>a</sup> Gene symbols are as given by Clutterbuck (10). Construction of the original *amdS::lacZ* fusion strain is described by Davis et al. (16). The *hapBΔ* and *hapEΔ* deletions are described in this paper. Generation of the *hapCΔ* deletion has been described elsewhere (40).

of AnCF binding activity and in phenotypes similar to those observed with the *hapC* deletion. However, resolution of the binding complex into two bands by using electrophoretic mobility shift assays together with the observation that HapE shares a fungus-specific domain with the *S. cerevisiae* and *Schizosaccharomyces pombe* homologs has raised the possibility that one or more additional components may associate with the core HapB/C/D complex and may carry activation domains.

We also have found for the first time evidence that AnCF also affects a pathway-specific regulatory circuit. The *amdR* gene encodes a Zn(II)<sub>2</sub>Cys<sub>6</sub> binuclear cluster DNA binding protein that is required for omega-amino acid induction of the *amdS* gene and the genes for omega-amino acid utilization (3). Deletion of the *hapB* and *hapE* genes results in loss of omega-amino acid induction of *amdS* expression. In addition, the *hapB* and *hapE* deletion strains are unable to use  $\gamma$ -aminobutyric acid (GABA) as a sole nitrogen or carbon source.

#### MATERIALS AND METHODS

**Strains, media, genetic techniques, and transformation.** *A. nidulans* strains used in this study are described in Table 1. Standard *A. nidulans* growth media and conditions were as described by Cove (12). Nitrogen sources were added to a final concentration of 10 mM. In plate growth tests, sodium acetate (50 mM), ethanol (1%, vol/vol), glycerol (1%, vol/vol), GABA (10 mM), and proline (10 mM) were used as carbon sources. Genetic manipulation of haploid *A. nidulans* strains was carried out as described by Clutterbuck (10). Protoplast preparation and DNA transformation were performed as described by Andrianopoulos and Hynes (2). *Escherichia coli* NM522 (New England Biolabs) was used for propagation of plasmids and DNA manipulation; strain BL21(DE3) (Novagen) was used for expression of His<sub>6</sub>-HapBct encoded by pT7-hapBct (see below). Molecular cloning techniques were performed as specified by Sambrook et al. (45).

**Isolation of *hapB* and *hapE* genes.** For amplification of *hapB* cDNA fragments, degenerate oligonucleotides HAPB1 (5' CARCCITTYTAYGTIAAYGC 3') and HAPBB1 (5' GCRITTIACRTARAAGGYTG 3') were used together with the T7 or T3 oligonucleotide or primer HAPB5 (5' CAGACATCATAACCAGC ATAAC 3') and an *A. nidulans* cDNA library in  $\lambda$  ZAPII (constructed by R. Aramayo; available from the Fungal Genetics Stock Center, Kansas City, Kans.) previously amplified by using the universal and reverse primers (52) as the template for PCR. With the primer combination HAPB1A and T7, a single 700-bp fragment was generated and upon sequencing found to have a high level of similarity to Hap2p of *S. cerevisiae*. The primer combination HAPBB1 and T3 yielded a fragment of 800 bp with no similarity to Hap2p but which subsequently proved to contain the 5' region of *hapB*. To isolate a genomic clone, a 520-bp PCR fragment, amplified with the primers HAPB1 and HAPB5, was used to screen a genomic *A. nidulans* cosmid library (7) by colony hybridization. The clone pCOSHAB was isolated and a 4.0-kb *Bam*HI fragment that hybridized to the probe in Southern blot analysis was subcloned into pBluescriptII SK<sup>+</sup>. To obtain the corresponding cDNA, a T3 or T7 primer together with internal primers derived from this open reading frame (ORF) were used to amplify PCR fragments from the *A. nidulans* cDNA library.

Plasmid pCAABI-1.5 (kindly provided by J. A. Kinsey, Kansas City, Kans.) contains a 1.5-kb *Eco*RI *aab-1* cDNA clone (9). An *A. nidulans* cDNA library was probed with the 1.5-kb *Eco*RI fragment derived from this clone. A hybridizing clone (pHAPE-9) containing a cDNA insert of 1.2 kb was sequenced, and the insert showed 63% sequence identity to *aab-1* at the DNA level. The gene was

designated *hapE*. The 1.2-kb insert of pHAPE-9 was found to hybridize to a 6.0-kb *Bgl*II-*Pst*I genomic fragment in Southern blot analysis. A plasmid partial library was created by cloning genomic *Bgl*II-*Pst*I fragments of approximately 6.0 kb into *Bam*HI-*Pst*I-digested pBluescriptII SK<sup>+</sup> and a *hapE*-containing clone was identified by colony hybridization with the pHAPE-9 cDNA insert. Southern blot analysis showed that the *hapE* gene was contained within a 2.8-kb *Bam*HI-*Pst*I fragment, which was subcloned into pBluescriptII SK<sup>+</sup> and sequenced.

**Oligonucleotides and plasmids.** Plasmid pBRhapB carries a genomic 4.0-kb *Bam*HI *hapB* fragment in the *Bam*HI site of pBR322. pKOhapB was constructed by excising a 1.6-kb *Sma*I-*Xho*I *hapB* fragment from pBRhapB and replacing it with a 1.8-kb *Sma*I-*Xho*I *argB* fragment derived from plasmid pDC1 (4). pPTR18 carries a genomic 2.5-kb *Bam*HI-*Sal*I *hapE* fragment in pBluescriptII SK<sup>+</sup>. pKOhapE was constructed by replacing an 0.8-kb *Eco*RV-*Sph*I fragment of pPTR18 with a *Sma*I-*Sph*I *argB* fragment derived from plasmid pDC1, thereby eliminating the evolutionary conserved domain of *hapE*. pMalE-HapC and pMalE-HapE, were kindly provided by M. Kato and N. Tsukagoshi (Nagoya, Japan).

pT7-hapBct, used for expression of a N-terminal truncated HapB protein, was generated as follows. A full-length *hapB* PCR product was cloned in frame into the *Xmn*I/*Bam*HI sites of pMal-c2 (New England Biolabs) to give pMalE-hapB. The integrity of the cloned PCR product was verified by DNA sequencing. A *Sac*I-*Hind*III digest was performed to transfer the insert of pMalE-hapB into pQE31 (Qiagen), resulting in pQE31-hapB. An *Eco*RI-*Hind*III fragment carrying the full-length *hapB*, an *E. coli* ribosome binding site, and a His<sub>6</sub> tag was then cloned into pT7-5 (48) to give pT7-hapB. The expression constructs pMalE-hapB, pQE31-hapB, and pT7-hapB did not allow high-level expression of full-length HapB in *E. coli*. However, an N-terminal part of HapB (amino acid [aa] 166 to 349) could be highly expressed. After an inverse PCR using primers Bct1 (TGGATAGGTACCGGTCATCCTTC) and Bct2 (CCATGTCCGGTACCCC GCACG), with pT7-hapB as the template, a PCR fragment missing the *hapB* N-terminal part coding for aa 3 to 165 was created. Digestion with *Kpn*I and ligation of the PCR fragment resulted in plasmid pT7-hapBct, which allows expression of the His<sub>6</sub>-tagged C-terminal 183 aa of HapB (His<sub>6</sub>-HapBct). The *hapB* sequence was checked by DNA sequencing.

**DNA sequencing.** Double-stranded plasmids were sequenced by the dideoxy-chain termination method of Sanger et al. (46) on an ABI Prism (Perkin-Elmer) automated sequencer.

**Expression and purification of recombinant proteins.** MalE-HapC and MalE-HapE fusion proteins were prepared as described by Kato et al. (25). Cultures (200 ml) of *E. coli* NM522 harboring plasmid pMalE-HapC or pMalE-HapE were grown to an optical density at 600 nm of ~0.5, and induction was achieved by the addition of 1 mM isopropyl- $\beta$ -D-thiogalactopyranoside. After induction for 3 h at 37°C, cells were harvested, taken up in 15 ml of buffer A (20 mM Tris-Cl, 1 mM EDTA, 1 mM dithiothreitol [DTT], 200 mM NaCl [pH 7.5]), lysed by sonication, and centrifuged at 12,100  $\times$  g for 20 min. The crude cellular extracts were applied to amylose columns (New England Biolabs) and washed with buffer A until the flowthrough optical density at 280 nm was <0.01. The fusion proteins were eluted with buffer A containing 10 mM maltose. His<sub>6</sub>-HapBct-containing crude cellular *E. coli* extracts were obtained by the same protocol. The crude cellular extracts were applied to a nitrilotriacetic acid-agarose column (Qiagen). Washing steps were performed according to the supplier. The His<sub>6</sub>-tagged protein was eluted by a 50 to 500 mM gradient of imidazole in washing buffer. The purity of all recombinant proteins was examined by sodium dodecyl sulfate-polyacrylamide gel electrophoresis and Coomassie blue staining.

**Preparation of *A. nidulans* nuclear extracts and electrophoretic mobility shift assays.** *A. nidulans* nuclear extracts were prepared by the method of Papagiannopoulos et al. (40). Oligonucleotides MH10 (GATCGCCAGCCAATCACCA GCTAGCCACCAGCTAAACCC) and MH11 (GATCGGGTTTAGCTGGTG CCTAGCTGGTATTGGCTGGC) (referred to as oligonucleotides 10 and 11) are complementary oligonucleotides derived from the CCAAT box region of the *amdS* promoter which have 5' GATC overhangs after annealing (51). Labeling of the probe and electrophoretic mobility shift assays were performed as described by Papagiannopoulos et al. (40) except that 20  $\mu$ g of nuclear protein extract was incubated at 25°C for 30 min together with the probe before the mixtures were loaded on 6% polyacrylamide gels. For reconstitution experiments, the DNA binding reaction mixture contained DTT at a concentration of 5 mM and 10 ng of each recombinant purified protein.

**$\beta$ -Galactosidase assays.** Reporter gene studies used strains containing a single copy of the *amdS::lacZ* fusion lacking vector sequences in place of the native *amdS* gene (16).  $\beta$ -Galactosidase assays were carried out by the method of Davis et al. (16).

**Nucleotide sequence accession numbers.** The complete *hapB* DNA and protein sequences have been submitted to EMBL/GenBank databases under accession no. Y13768; *hapE* DNA and protein sequences are available under accession no. U96847.

## RESULTS

**Cloning and sequencing of *hapB*.** The DNA binding and subunit interaction domains of Hap2p and CBF-B reside in an








FIG. 2. Deletion strategy for *hapB* and *hapE*. (A) Partial restriction maps of the genomic regions of *hapE* and *hapB*. ORFs and direction of transcription are indicated by rightward arrows. (B) Schematic representation of the deletions generated by gene replacement of *hapB* (strain MH9099) and *hapE* (strain MH9092) by the selectable marker *argB* after a homologous double-recombination event with the linearized deletion constructs pKO*hapB* and pKO*hapE*, respectively. The direction of *argB* transcription is shown (leftward arrows).

mobilities (Fig. 4, lane 2), whereas the nuclear extracts prepared from the *hapB* and *hapE* deletion strains showed loss of both complexes (Fig. 4, lanes 3 and 4), clearly indicating that HapB and HapE are also required for AnCF binding, as has been shown for HapC (40). In contrast to previous results where the resolution of two complexes was not clearly seen (40), two CCAAT binding complexes were consistently resolved with different preparations of wild-type nuclear extracts and different CCAAT-containing probes (not shown), although the intensity of the two complexes showed some variation. Two complexes binding to a CCAAT-containing sequence have also been observed in *N. crassa* (9).

**Expression of *amdS* is reduced in *hapB* $\Delta$  and *hapE* $\Delta$  backgrounds.** To assess *amdS* expression in *hapB* $\Delta$  and *hapE* $\Delta$  backgrounds, strains MH9094 and MH9210 were crossed with strain MH3408, carrying a single-copy *amdS::lacZ* reporter gene at the *amdS* locus (16).  $\beta$ -Galactosidase expression in the resulting *hapB* $\Delta$  and *hapE* $\Delta$  backgrounds was compared to that in the parent strain MH3408. The two deletions reduced *amdS* expression in similar manners (Table 2). As noted for a *hapC* deletion strain (40), the reduction of *amdS* expression was more significant under carbon-limiting conditions (0.1% glucose) than with 1% glucose. However, the *amdS::lacZ* fusion was sensitive to nitrogen metabolite repression in the deletion strains, as shown by derepression by growth on the limiting nitrogen source L-alanine compared with ammonium-grown mycelia (Tables 2 and 3).

**In vitro reconstitution of a CCAAT binding complex.** HapB, HapC, and HapE were expressed in *E. coli* as fusion proteins and purified by affinity chromatography (Materials and Methods). HapC (aa 1 to 186) and HapE (aa 1 to 265) were purified as MalE fusion proteins. HapB was purified as a His<sub>6</sub>-tagged truncated protein (HapBct) containing the C-terminal 183 aa of HapB (aa 168 to 349), which includes the predicted conserved DNA binding and subunit interaction domains (Fig. 1A) characterized in *S. cerevisiae* Hap2p (39) and rat CBF-B (30). When combinations of two of the *A. nidulans* Hap proteins were mixed and incubated with the DNA probe 10-11, no DNA binding was detected in mobility gel shifts (Fig. 5, lanes

3 to 5). Only when all three proteins were combined was binding observed (Fig. 5, lane 6). The subunits HapC and HapE were each fused to the 42.7-kDa MalE protein; therefore, the newly formed DNA binding protein complex had a mobility lower than that of the wild-type Hap complex(es) (lane 2). The MalE-HapE fusion protein was cleaved into the two components MalE and HapE, using the protease factor Xa cleavage site in the linker region. When this HapE-MalE mixture was used in a mobility shift together with His<sub>6</sub>-HapBct and MalE-HapC, a complex with increased mobility was observed, indicating that the presence of HapE instead of the larger MalE-HapE fusion protein reduced the complex size. DNA binding could be reconstituted only in the presence of 5 mM DTT. Subunit association and DNA binding of the mammalian CBF have previously been shown to be dependent on reduction of the CBF subunits (37).

**The *A. nidulans* Hap complex is required for AmdR-dependent induction.** The binding sites for the transcriptional activator AmdR and the AnCF complex are contained within a 19-bp sequence in the *amdS* promoter (51). Furthermore, other functional AmdR binding sites are found in the *lamA/lamB* and *gatA* promoters, directly flanked by a GCAAT or a CCAAT sequence within the 19-bp element (44). The *hap* deletion mutants allowed us to examine whether a functional AnCF complex is a prerequisite for AmdR-mediated activation of *amdS*. Induction by omega amino acid GABA or  $\beta$ -alanine (five- or eightfold, respectively, in a wild-type strain) mediated by AmdR was eliminated in *hapB* and *hapE* deletion backgrounds (Table 3). This result suggested that AmdR is dependent on a functional AnCF complex for activation of *amdS* expression.

Utilization of GABA as carbon or nitrogen source requires the expression of *gaba* (GABA permease) and *gatA* (GABA transaminase), which are positively regulated by AmdR (6, 43). The three *hap* deletion strains show greatly reduced levels of growth on GABA as a nitrogen source compared to the wild-type strain MH54 and do not form any aerial mycelia (Fig. 1B). Similarly, the *hap* deletion strains grew extremely poorly on GABA as a sole carbon and nitrogen source or as a sole carbon




FIG. 3. Effects of *hap* deletions on growth and morphology. Strains MH9099 (*hapB* $\Delta$ ), MH8194 (*hapC* $\Delta$ ), and MH9092 (*hapE* $\Delta$ ) are compared to a wild-type (WT) strain (MH54). (A) Growth on complete medium and sodium acetate (50 mM), ethanol (1%, vol/vol) and glycerol (1%, vol/vol) as sole carbon sources in the presence of 10 mM  $\text{NH}_4\text{Cl}$ . (B) Growth on proline (10 mM) or GABA (10 mM) as the sole nitrogen source in the presence of 1% glucose or as the sole carbon and nitrogen source. The strains were incubated at 37°C for 48 h, except for the growth tests on ethanol and on proline or GABA as the sole carbon and nitrogen source, in which cases incubation was for 78 h at 37°C.

source in the presence of ammonium. Strain MH8194 (*hapC* $\Delta$ ) carries an additional deletion of *amdR* (40), whereas strains MH9092 (*hapE* $\Delta$ ) and MH9099 (*hapB* $\Delta$ ) are *amdR*<sup>+</sup>. Although *amdR* is intact in these two strains, growth levels were comparable to those observed for the *hapC* $\Delta$  *amdR* $\Delta$  double mutant MH8194. This finding indicated that AmdR-mediated activation of GABA utilization does not occur in the absence of the AnCF complex. Proline and GABA are metabolized via independent pathways leading to glutamate. Growth of the *hap* deletion mutants on proline either as a sole nitrogen source or as a sole carbon and nitrogen source was comparable to growth on glucose and ammonium medium (Fig. 3). In addition, growth on L-proline, L-alanine, and L-glutamate as sole carbon sources in the presence of ammonium was not affected by the *hap* deletion mutations (results not shown). Therefore, the observed weak growth on GABA reflects a pathway-specific effect on the genes responsible for the utilization of GABA.

## DISCUSSION

The finding that deletion of each of the three genes *hapB*, *hapC*, and *hapE* results in identical slowly growing and weakly




FIG. 4. Loss of CCAAT binding activity in *hapB* $\Delta$  and *hapE* $\Delta$  strains. For mobility shift analysis, the end-labeled oligonucleotide pair 10-11 was incubated with 20  $\mu\text{g}$  of crude nuclear extract prepared from a *hap*<sup>+</sup> strain (lane 2), a *hapB* $\Delta$  strain (MH9099; lane 3), or a *hapE* $\Delta$  strain (MH9092; lane 4), and the mixtures were fractionated on a 6% polyacrylamide gel. Free probe and AnCF complexes are marked by arrows.

condensing phenotypes is consistent with all three gene products interacting to form a functional complex in which each subunit is essential. Loss of AnCF binding in each of the three *hap* deletion strains together with *in vitro* reconstitution of a binding complex with expressed subunits provides further compelling evidence for this.

The *in vitro* reconstitution of a binding complex shows that the absence of the N-terminal 182 HapB amino acids in His<sub>6</sub>-HapBct does not impair subunit interaction or DNA binding. These residues are therefore not essential for functional DNA binding complex formation *in vitro*. Furthermore, this experiment demonstrates that a heteromeric complex consisting only of the subunits HapB, HapC, and HapE creates a protein surface able to bind to DNA containing CCAAT sequences, as has been found for the equivalent three subunits in *S. cerevisiae* (33) and mammals (47).

Although the DNA binding and subunit association domains of HapB and HapE have been highly conserved among fungi and mammals (Fig. 1), the proteins have diverged significantly in overall structure. The functional domains are not conserved with respect to their positions within the proteins (Fig. 6), and when regions apart from the conserved domains are compared, no significant homology between yeast, mammals, and *A. nidulans* can be found. Therefore, while the ability of the subunits of the HAP-CBF complexes to interact with each other and with DNA has been maintained, the specific functions of the complexes have been altered by the divergence of the flanking

TABLE 2. Expression of *amdS::lacZ* fusions in the *hapB* $\Delta$  and *hapE* $\Delta$  backgrounds

Strain	Relevant genotype	$\beta$ -Galactosidase activity <sup>a</sup> (U)	
		1% Glucose + 10 mM $\text{NH}_4$ -tartrate	0.1% Glucose + 10 mM $\text{NH}_4$ -tartrate
MH3408	<i>amdS::lacZ</i>	1.94 (0.12)	20.18 (2.49)
MH9207	<i>amdS::lacZ; hapB</i> $\Delta$	0.38 (0.12)	1.07 (0.24)
MH9206	<i>amdS::lacZ; hapE</i> $\Delta$	0.31 (0.11)	0.42 (0.06)

<sup>a</sup> Mycelia were grown in 100 ml of medium at 37°C for 16 h (1% glucose) and 20 h (0.1% glucose). Data represent the results of four separate experiments; standard errors are given in parentheses. Assays of endogenous  $\beta$ -galactosidase in a wild-type strain and in *hapB* $\Delta$  and *hapE* $\Delta$  strains lacking the inserted *amdS::lacZ* fusion gene gave activities of 0.1 to 0.4 and were not affected by the growth conditions used for the experiments reported here or in Table 3.

TABLE 3. Expression of *amdS::lacZ* fusions in the *hapBΔ* and *hapEΔ* backgrounds under induction with omega amino acids

Strain	Relevant genotype	β-Galactosidase activity <sup>a</sup> (U)		
		10 mM L-alanine	10 mM L-alanine + 10 mM GABA	10 mM L-alanine + 10 mM β-alanine
MH3408	<i>amdS::lacZ</i>	19.14 (2.02)	94.24 (7.16)	148.70 (3.56)
MH9207	<i>amdS::lacZ; hapBΔ</i>	1.95 (0.34)	2.26 (0.55)	1.10 (0.07)
MH9206	<i>amdS::lacZ; hapEΔ</i>	1.14 (0.08)	1.21 (0.07)	0.83 (0.07)

<sup>a</sup> Mycelia were grown in 100 ml of medium at 37°C for 16 h with 1% glucose. Nitrogen sources were added as indicated. Data represent the results of four separate experiments; standard errors are given in parentheses.

sequences during evolution. HapE and the *N. crassa* Aab-1 protein show considerable similarity within their central conserved domains and also in the flanking sequences. Although these species are the closest relatives in this group, they represent distinct taxa within the fungi, suggesting that the Hap complexes in these two filamentous fungi may be involved in the regulation of similar genes and processes.

McNabb et al. (34) identified a second domain in *S. pombe* that is conserved in *S. cerevisiae* and showed that this domain is involved in recruiting the activator subunit Hap4p to the Hap complex in *S. cerevisiae*. This domain is also present in HapE and in Aab-1 but not in CBF-C (Fig. 1B and 6) and therefore may be specific to the fungi. As suggested by McNabb et al. (34), this is a strong indication that fungal CCAAT binding complexes may have a fourth Hap4p-like subunit. Our results provide evidence that this may be the case for *A. nidulans*. When wild-type extracts are used in electrophoretic mobility shifts, two distinct DNA-protein complexes are formed, but both are lost when nuclear extracts derived from *hapΔ* mutants are used (Fig. 4). The complex with higher mobility may be composed of the DNA binding core complex HapB/C/E, whereas the complex with lower mobility contains a fourth, unknown component. The predicted molecular mass of a HapB/C/E complex is 87 kDa. Gel filtration experiments determining the molecular mass of the AnCF and PENR1 complexes have given estimates of 120 to 130 kDa (25, 28). These results provide further support for the idea that one (or more) additional subunit is associated with the HapB/C/E




FIG. 5. In vitro reconstitution of a CCAAT binding complex. The recombinant proteins His<sub>6</sub>-HapBct, MalE-HapC, and MalE-HapE were expressed in *E. coli* and purified by affinity chromatography (Materials and Methods). The recombinant proteins were tested for DNA binding in electrophoretic mobility shift analysis using the oligonucleotide 10-11 probe. Binding reaction mixtures contained 10 ng of each recombinant protein, indicated by + (lanes 3 to 6), or 20 μg of crude nuclear extract prepared from a wild-type *A. nidulans* strain (lane 2). Free probe, AnCF complexes, and the reconstituted CCAAT binding complex are marked by arrows.




FIG. 6. Positions of the functional domains of the HapB and HapE homologs are not conserved. Each protein is depicted with the N terminus on the left and aligned with respect to the conserved domains. Highlighted boxes indicate the evolutionarily conserved domains that are depicted in Fig. 1. Black boxes represent the conserved domains found in all of the HapB or HapE homologs, whereas grey boxes represent the homology in the Hap4p interaction domain characterized for Hap5p (33, 34) present in all of the fungal homologs.

core. It is possible that different Hap4p-like activator subunits can be associated with the core complex and provide specificity for particular sets of genes. In *S. cerevisiae*, there is evidence for the regulation of genes by the Hap complex that is not absolutely dependent on Hap4p (13, 14).

In *S. cerevisiae*, the Hap complex plays a major role in the control of genes involved in respiration in response to the carbon source, and the activator subunit Hap4p is strongly up-regulated by a shift from glucose to nonfermentable carbon sources (18). The *A. nidulans* Hap complex, in contrast, appears to be not absolutely required for activation of genes involved in oxidative phosphorylation. The ability of *hapB/C/E* deletion strains to grow on carbon sources such as acetate and glycerol (Fig. 3) clearly distinguishes *A. nidulans*, an obligate aerobe, from *S. cerevisiae hap* mutants, which are not viable on nonfermentable carbon sources (33, 42).

The CBF complex in mammals has a wide range of target genes (32). In filamentous fungi, the complex has been implicated in the regulation of genes involved in the catabolism of sole carbon sources (e.g., Taka-amylase [24, 36]), sole nitrogen sources (formamide hydrolysis [19]), and sources of both carbon and nitrogen (acetamide [reference 40 and references therein]). In addition, the complex affects the expression of genes involved in secondary metabolism (penicillin biosynthesis [28, 29, 49]) and in ammonium assimilation (NADP-dependent glutamate dehydrogenase [9, 15]). A protein binding to a CCAAT sequence in the 5' region of the conidiation-specific *yA* gene of *A. nidulans* has been detected, but it has not been established that the protein is AnCF (1, 5).

The viability of the *A. nidulans hapB*, *hapC*, and *hapE* deletion strains indicates that the *hap* complex is not absolutely required for the transcription of any essential genes. It is likely that the phenotype of slow growth results from suboptimal expression of many genes which are subject to additional regulatory controls. This is exemplified by the *amdS* gene: although the CCAAT sequence and the *hap* complex have a major effect on expression, the gene can still respond to other control mechanisms. In addition, penicillin production is reduced but not abolished in a *hapC* deletion strain (28).

The *amdS* gene is subject to multiple wide domain- and pathway-specific regulatory controls, one of which is mediated by the *amdR* gene, which is necessary for omega-amino acid induction (reviewed in reference 23). The *amdI93* deletion, which removes both the CCAAT site and the binding site for



AmdR, does not affect regulation by the *amdA*, *areA*, *facB*, or *creA* gene (21, 22, 50). Mutation of the *amdS* 5' CCAATCA 3' sequence to 5' CAAGTCT 3' reduces *amdS* expression to levels seen in the *hap* deletion strains and eliminates *amdR*-mediated induction (20). However, the effects of the other regulatory circuits, including responses to the gain-of-function mutations *amdA7* and *areA102* are still observed (20). We have shown here that deletion of the *hapB* and *hapE* genes eliminates omega-amino acid induction of the *amdS::lacZ* reporter gene and also specifically affects GABA utilization, which is dependent on AmdR activity. Therefore, we conclude that activation of gene expression by the pathway-specific regulator AmdR is dependent on AnCF function.

AmdR contains an N-terminal Zn(II)<sub>2</sub>Cys<sub>6</sub> binuclear cluster DNA binding domain which has been shown to confer DNA binding specificity in vivo (3, 41). Studies on transformants overexpressing AmdR due to multiple copies (2, 3) together with domain swapping experiments with the FacB activator and the analysis of *amdR<sup>c</sup>* constitutive mutations (41) lead to the suggestion that omega-amino acid inducers do not affect AmdR DNA binding but rather increase the ability of bound protein to activate transcription. Both AmdR and AnCF have been found to bind in vitro to conserved sequences found within a 19-bp element in the *amdS*, *gatA*, and *lamA/B* 5' regions (44, 51). Therefore, in each gene controlled by AmdR, there is a close association with AnCF binding sites.

The dependence of AmdR function on the AnCF complex could result from effects on DNA binding and/or activation. AnCF activity is not dependent on AmdR, since all genes regulated by AnCF are not also regulated by AmdR and the basal levels of *amdS* expression can be restored by the insertion of the CCAAT-containing oligonucleotide pair 10-11 without restoring AmdR control. Fusion of the activation domains of AmdR to the DNA binding domain of the acetate-specific regulator FacB results in activation of acetate utilization genes and in omega-amino acid induction of *amdS* when tested in *amdR* and *facB* mutant strains (41). Since AnCF is not required for acetate utilization (Fig. 3) and does not affect FacB-mediated regulation of *amdS*, this suggests that it is less likely that AmdR dependence on AnCF results from cooperative activation of transcription. It is unlikely that AnCF is required for AmdR expression, since no CCAAT sites are present in the 5' region of *amdR* and the *amdR* transcript is present at a low constitutive level (2, 3). It is therefore more likely that AnCF facilitates AmdR binding to DNA in vivo. Previous studies of in vitro binding of nuclear extracts showed that AmdR could bind to an oligonucleotide pair derived from the *amdS* 5' region in which the CCAAT motif was changed to CCTTT, while AnCF could not bind to this sequence. This demonstrates the independent nature of the binding sites but does not necessarily reflect the in vivo binding capacity of AmdR. We suggest that AnCF binding to its target sequence is a prerequisite for a change in chromatin structure necessary for AmdR binding. Support for this view comes from preliminary data from this laboratory which indicate that a CCAAT sequence and the integrity of the AnCF complex are necessary for a change in DNase I hypersensitivity (38).

Interactions of HAP-like CCAAT binding complexes with other transcription factors have been observed in both mammals and fungi. The *Kluyveromyces lactis* Hap complex stabilizes the binding of Abf1p, and activation of transcription is achieved synergistically (35). Mutations in the CCAAT boxes of the invariant chain promoter have been shown to abolish the binding of two other transcription factors (Sp1 and RFX) to their target sites in in vivo footprinting studies (26). In the 3-hydroxy-3-methylglutaryl coenzyme A synthase gene, a bind-

ing site for the sterol regulatory element binding proteins is separated by 17 bp from an inverted CCAAT box. It has been shown that binding of the CBF complex to the CCAAT box is required for sterol-mediated regulation of 3-hydroxy-3-methylglutaryl coenzyme A synthase. Furthermore, a direct interaction between CBF and the basic helix-loop-helix sterol regulatory element binding proteins occurs (17). Other examples of CBF interactions with regulatory circuits are reviewed by Maity and de Crombrughe (31).

Complex protein-protein interactions between transcription factors have been discovered with increasing frequency in recent years. Characterization of interactions between the HapB/C/E complex and Hap4p homologs as well as gene-specific regulatory proteins such as AmdR are required to determine how the specificity of Hap complex-mediated regulation of gene expression in filamentous fungi is modulated.

#### ACKNOWLEDGMENTS

We gratefully acknowledge Norihiro Tsukagoshi and Masashi Kato for providing the expression plasmids pMalE-HapC and pMalE-HapE and Jack Kinsey for plasmid pCAAB1-1.5.

This work was supported by the Australian Research Council, by the Deutsche Forschungsgemeinschaft (Sonderforschungsbereich 369), and by a DAAD Doktorandenstipendium im Rahmen des gemeinsamen Hochschulniederprogramms III von Bund und Laendern granted to S.S.

#### REFERENCES

1. Andrianopoulos, A. Unpublished data.
2. Andrianopoulos, A., and M. J. Hynes. 1988. Cloning and analysis of the positively acting regulatory gene *amdR* from *Aspergillus nidulans*. *Mol. Cell. Biol.* **8**:3532-3541.
3. Andrianopoulos, A., and M. J. Hynes. 1990. Sequence and functional analysis of the positively acting regulatory gene *amdR* from *Aspergillus nidulans*. *Mol. Cell. Biol.* **10**:3194-3203.
4. Aramayo, R., T. H. Adams, W. E. Timberlake. 1989. A large cluster of highly expressed genes is dispensable for growth and development in *Aspergillus nidulans*. *Genetics* **122**:65-71.
5. Aramayo, R., and W. E. Timberlake. 1993. The *Aspergillus nidulans* *yaA* gene is regulated by *abaA*. *EMBO J.* **12**:1039-2048.
6. Arst, H. N. 1976. Integrator gene in *Aspergillus nidulans*. *Nature* **262**:231-234.
7. Borges-Walmsley, M. I., G. Turner, A. M. Bailey, J. Brown, J. Lehmbeck, and I. G. Clausen. 1995. Isolation and characterisation of genes for sulphate activation and reduction in *Aspergillus nidulans*: implications for evolution of an allosteric control region by gene duplication. *Mol. Gen. Genet.* **247**:423-429.
8. Bucher, P. 1990. Weight matrix descriptions of four eukaryotic RNA polymerase II promoter elements derived from 502 unrelated promoter sequences. *J. Mol. Biol.* **212**:563-578.
9. Chen, H., J. W. Crabb, and J. A. Kinsey. 1998. The *Neurospora aab-1* gene encodes a CCAAT binding protein homologous to yeast HAP5. *Genetics* **148**:123-130.
10. Clutterbuck, A. J. 1984. Loci and linkage map of the filamentous fungus *Aspergillus nidulans*, p. 265-273. In S. J. O'Brien (ed.), *Genetic maps*, vol. 3. Cold Spring Harbor Laboratory Press, New York, N.Y.
11. Coustry, F., S. Sinha, S. N. Maity, and B. de Crombrughe. 1998. The two activation domains of the CCAAT-binding factor CBF interact with the dTAFII110 component of the Drosophila TFIID complex. *Biochem. J.* **331**:291-297.
12. Cove, D. J. 1966. The induction and repression of nitrate reductase in the fungus *Aspergillus nidulans*. *Biochim. Biophys. Acta* **113**:51-56.
13. Dang, V. D., C. Bohn, M. Bolotin-Fukuhara, and B. Daignan-Fornier. 1996. The CCAAT box-binding factor stimulates ammonium assimilation in *Saccharomyces cerevisiae*, defining a new cross-pathway regulation between nitrogen and carbon metabolisms. *J. Bacteriol.* **178**:1842-1849.
14. Dang, V. D., M. Valens, M. Bolotin-Fukuhara, and B. Daignan-Fornier. 1996. Cloning of the ASN1 and ASN2 genes encoding asparagine synthetases in *Saccharomyces cerevisiae*: differential regulation by the CCAAT-box-binding factor. *Mol. Microbiol.* **22**:681-692.
15. Davis, M. A. Unpublished data.
16. Davis, M. A., C. S. Cobbet, and M. J. Hynes. 1988. An *amdS-lacZ* fusion for studying gene regulation in *Aspergillus*. *Gene* **63**:199-212.
17. Dooley, K. A., S. Millinder, and T. F. Osborne. 1998. Sterol regulation of 3-hydroxy-3-methylglutaryl-coenzyme A synthase gene through a direct interaction between sterol regulatory element binding protein and the trimeric

- CCAAT-binding factor/nuclear factor Y. *J. Biol. Chem.* **273**:1349–1356.
18. **Forsburg, S. L., and L. Guarente.** 1989. Identification and characterization of HAP4: a third component of the CCAAT-bound HAP2/HAP3 heteromer. *Genes Dev.* **3**:1166–1178.
  19. **Fraser, J. A.** Unpublished data.
  20. **Hynes, M. J.** Unpublished data.
  21. **Hynes, M. J.** 1980. A mutation, adjacent to the gene *amdS*, defining the site of action of positive-control gene *amdR* in *Aspergillus nidulans*. *J. Bacteriol.* **142**:400–406.
  22. **Hynes, M. J., C. M. Corrick, J. M. Kelly, and T. G. Littlejohn.** 1988. Identification of the sites of action for regulatory genes controlling the *amdS* gene of *Aspergillus nidulans*. *Mol. Cell. Biol.* **8**:2589–2596.
  23. **Hynes, M. J., and M. A. Davis.** 1996. Regulation of acetamide catabolism, p. 381–393. In R. Bramble and G. A. Marzluf (ed.), *The Mycota, biochemistry and molecular biology*, vol. III. Springer-Verlag, Heidelberg, Germany.
  24. **Kato, M., A. Aoyama, F. Naruse, T. Kobayashi, and N. Tsukagoshi.** 1997. An *Aspergillus nidulans* nuclear protein, AnCP, involved in enhancement of Taka-amylase A gene expression, binds to the CCAAT-containing *taaG2*, *amdS*, and *gatA* promoters. *Mol. Gen. Genet.* **254**:119–126.
  25. **Kato, M., A. Aoyama, F. Naruse, Y. Tateyama, K. Hayashi, M. Miyazaki, P. Papagiannopoulos, M. A. Davis, M. J. Hynes, T. Kobayashi, and N. Tsukagoshi.** 1998. The *Aspergillus nidulans* CCAAT-binding factor AnCP/AnCF is a heteromeric protein analogous to the HAP complex of *Saccharomyces cerevisiae*. *Mol. Gen. Genet.* **257**:404–411.
  26. **Linhoff, M. W., K. L. Wright, and J. P. Ting.** 1997. CCAAT-binding factor NF-Y and RFX are required for in vivo assembly of a nucleoprotein complex that spans 250 base pairs: the invariant chain promoter as a model. *Mol. Cell. Biol.* **17**:4589–4596.
  27. **Littlejohn, T. G., and M. J. Hynes.** 1992. Analysis of the site of action of the *amdR* product for the regulation of the *amdS* gene of *Aspergillus nidulans*. *Mol. Gen. Genet.* **235**:81–88.
  28. **Litzka, O., P. Papagiannopoulos, M. A. Davis, M. J. Hynes, and A. A. Brakhage.** 1998. The penicillin regulator PENR1 of *Aspergillus nidulans* is a HAP-like transcriptional complex. *Eur. J. Biochem.* **251**:758–767.
  29. **Litzka, O., K. Then Bergh, and A. A. Brakhage.** 1996. The *Aspergillus nidulans* penicillin-biosynthesis gene *aat(penDE)* is controlled by a CCAAT-containing DNA element. *Eur. J. Biochem.* **238**:675–682.
  30. **Maity, S. N., and B. de Crombrughe.** 1992. Biochemical analysis of the B subunit of the heteromeric CCAAT-binding factor. A DNA-binding domain and a subunit interaction domain are specified by two separate segments. *J. Biol. Chem.* **267**:8286–8292.
  31. **Maity, S. N., and B. de Crombrughe.** 1998. Role of the CCAAT-binding protein CBF/NF-Y in transcription. *Trends Biochem. Sci.* **23**:174–178.
  32. **Mantovani, R.** 1998. A survey of 178 NF-Y binding CCAAT boxes. *Nucleic Acids Res.* **26**:1135–1143.
  33. **McNabb, D. S., Y. Xing, and L. Guarente.** 1995. Cloning of yeast HAP5: a novel subunit of a heteromeric complex required for CCAAT binding. *Genes Dev.* **9**:47–58.
  34. **McNabb, D. S., K. A. Tseng, and L. Guarente.** 1997. The *Saccharomyces cerevisiae* Hap5p homolog from fission yeast reveals two conserved domains that are essential for assembly of heterotetrameric CCAAT-binding factor. *Mol. Cell. Biol.* **17**:7008–7018.
  35. **Mulder, W., I. H. Scholten, and L. A. Grivell.** 1995. Distinct transcriptional regulation of a gene coding for a mitochondrial protein in the yeasts *Saccharomyces cerevisiae* and *Kluyveromyces lactis* despite similar promoter structures. *Mol. Microbiol.* **17**:813–824.
  36. **Nagata, O., T. Takashima, M. Tanaka, and N. Tsukagoshi.** 1993. *Aspergillus nidulans* nuclear proteins bind to a CCAAT element and the adjacent upstream sequence in the promoter region of the starch-inducible Taka-amylase A gene. *Mol. Gen. Genet.* **237**:251–260.
  37. **Nakshatri, H., P. Bhat-Nakshatri, and R. A. Currie.** 1996. Subunit association and DNA binding activity of the heterotrimeric transcription factor NF-Y is regulated by cellular redox. *J. Biol. Chem.* **271**:28784–28791.
  38. **Narendja, F.** Unpublished data.
  39. **Olesen, J. T., and L. Guarente.** 1990. The HAP2 subunit of yeast CCAAT transcriptional activator contains adjacent domains for subunit association and DNA recognition: model for the HAP2/3/4 complex. *Genes Dev.* **4**:1714–1729.
  40. **Papagiannopoulos, P., A. Andrianopoulos, J. A. Sharp, M. A. Davis, M. A., and M. J. Hynes.** 1996. The *hapC* gene of *Aspergillus nidulans* is involved in the expression of CCAAT-containing promoters. *Mol. Gen. Genet.* **251**:412–421.
  41. **Parsons, L. M., M. A. Davis, and M. J. Hynes.** 1992. Identification of functional regions of the positively acting regulatory gene *amdR* from *Aspergillus nidulans*. *Mol. Microbiol.* **6**:2999–3007.
  42. **Pinkham, J., and L. Guarente.** 1985. Cloning and molecular analysis of the HAP2 locus: a global regulator of respiratory genes in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **5**:3410–3416.
  43. **Richardson, I. B., S. K. Hurley, and M. J. Hynes.** 1989. Cloning and molecular characterisation of the *amdR* controlled *gatA* gene of *Aspergillus nidulans*. *Mol. Gen. Genet.* **217**:118–125.
  44. **Richardson, I. B., M. E. Katz, and M. J. Hynes.** 1992. Molecular characterisation of the *lam* locus and sequences involved in regulation by the *AmdR* protein of *Aspergillus nidulans*. *Mol. Cell. Biol.* **12**:337–346.
  45. **Sambrook, J., E. F. Fritsch, and T. Maniatis.** 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Press, Cold Spring Harbor, N.Y.
  46. **Sanger, F., S. Nicklen, and A. R. Coulson.** 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
  47. **Sinha, S., S. N. Maity, J. Lu, and B. de Crombrughe.** 1995. Recombinant rat CBF-C, the third subunit of CBF/NFY, allows formation of a protein-DNA complex with CBF-A and CBF-B and with yeast HAP2 and HAP3. *Proc. Natl. Acad. Sci. USA* **92**:1624–1628.
  48. **Tabor, S., and C. Richardson.** 1985. A bacteriophage T7 RNA polymerase/promoter system for controlled exclusive expression of specific genes. *Proc. Natl. Acad. Sci. USA* **82**:1074–1078.
  49. **Then Bergh, K., O. Litzka, and A. A. Brakhage.** 1996. Identification of a major *cis*-acting DNA element controlling the bidirectionally transcribed penicillin biosynthesis genes *acvA (pcbAB)* and *ipnA (pcbC)* of *Aspergillus nidulans*. *J. Bacteriol.* **178**:3908–3916.
  50. **Todd, R. B., A. Andrianopoulos, M. A. Davis, and M. J. Hynes.** 1998. FacB, the *Aspergillus nidulans* activator of acetate utilization genes, binds dissimilar DNA sequences. *EMBO J.* **17**:2042–2054.
  51. **van Heeswijk, R., and M. J. Hynes.** 1991. The *amdR* product and a CCAAT-binding factor bind to adjacent, possibly overlapping DNA sequences in the promoter region of the *Aspergillus nidulans amdS* gene. *Nucleic Acids Res.* **19**:2655–2660.
  52. **Weidner, G., B. Steffan, and A. A. Brakhage.** 1997. The *Aspergillus nidulans lysF* gene encodes homoaconitase, an enzyme involved in the fungus-specific lysine biosynthesis pathway. *Mol. Gen. Genet.* **255**:237–247.
  53. **Xing, Y., J. D. Fikes, and L. Guarente.** 1993. Mutations in yeast HAP2/HAP3 define a hybrid CCAAT box binding domain. *EMBO J.* **12**:4647–4655.