



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.e-jmii.com

Correspondence

SARS-CoV-2 infection in vulnerable population in Goiania, Central Brazil



KEYWORDS

COVID-19;
SARS-CoV-2;
Vulnerable populations

Dear Editor,

Brazil faces a severe economic, social and politic crisis, and consequently is one of the leading countries with the greatest number of cases and deaths due to COVID-19 caused by SARS-CoV-2.¹ Despite infectious diseases affect mainly socially and economically vulnerable people,² there is few data about the frequency of SARS-CoV-2 infection among them. Therefore, between July and October 2020, we investigated the prevalence of SARS-CoV-2 among recyclable waste collectors, homeless people, immigrants and refugees, LGBTQ people, sex workers, people using illicit drugs, and patients with HIV in Goiânia, a large city in Midwest Brazil.

All participants were interviewed, and samples of oropharynx and nasopharynx were collected for the detection of RNA-SARS-CoV-2. RNA extraction from SARS-CoV-2 was performed using the QIAamp Viral RNA Mini Kit (Qiagen, Maryland, USA), and real-time quantitative polymerase chain reaction (RT-PCR) was performed using AgPath-IDTM One-Step RT-PCR Reagents (Thermo Fisher Scientific, Massachusetts, USA). The assays were performed using probes from the 2019-nCoV kit (Integrated DNA Technologies, San Diego, USA).

This project was approved by the Research Ethics Committee (process no. 4145237).

A total of 517 individuals were recruited and 510 agreed to participate in the study (Table 1). The median age was 33 years (interquartile range [IQR]: 22 years), and 57.8% were

male. The majority of individuals (79.2%) were non-white. The median number of years of formal education was 10 (IQR: 6), and the monthly income was approximately 222 US\$ (IQR: 198).

In 17 samples, the results of RT-PCR were inconclusive. RNA-SARS-CoV-2 was positive in 95/493 (19.3%; 95% CI: 16.0–23.0) samples. The frequency of SARS-CoV-2 positivity was higher among immigrants/refugees (47.1%), followed by recycled waste collectors (15.9%) and LGBT (15.2%) subgroups (Table 1).

Currently, Brazil is the epicenter of the COVID-19 pandemic in Latin America¹ and, as observed in other contagious diseases, socially and economically vulnerable populations are disproportionately affected.² We detected RNA-SARS-CoV-2 in samples of 19.3% (95% CI: 16.0–23.0) of the individuals, and this increased to 37.3% (95% CI: 27.7–48.1) among those who reported with at least one COVID-19 sign and symptom. This is close to that observed among symptomatic Brazilian healthcare workers in São Paulo (42.4%; 95% CI: 36.9–48.1),³ a regional COVID-19 epicenter. It is noteworthy that healthcare workers are frontline professionals in coping with the COVID-19 pandemic.

In this study, immigrants were the groups most affected, following by LGBT and recycled waste collectors. Therefore, the COVID-19 pandemic represents an additional challenge to be overcome by them, considering that unemployment, precarious housing, and food conditions have intensified their social, economic, and health difficulties, mirroring the existing social inequality. Thus, face masks, water and soap/sanitizer for hand hygiene, social distancing, and more recently the access to COVID-19 vaccine are challenges for these individuals.

The high frequency of COVID-19 reinforces the priority of this population for vaccination against SARS-CoV-2 in Brazil, as well as the necessity to improve the tracking of new cases, such as expanded testing and the creation of useful public policies aimed at vulnerable populations.

<https://doi.org/10.1016/j.jmii.2021.08.005>

1684-1182/Copyright © 2021, Taiwan Society of Microbiology. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1 RNA-severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection among 493 vulnerable individuals in Midwest region.

Characteristic	n	RNA-SARS-CoV-2 (+)	%
Age (median; IQR)	493	29 (15)	
Education (years) (median; IQR)	493	12 (5)	
Monthly income (US\$) (median; IQR)	493	213 (202)	
Population subgroup			
LGBT ^a	79	12	15.2
Recycled waste collectors	145	23	15.9
Immigrants/refugees	121	57	47.1
Homeless	131	3	2.3
Sex workers	4	0	
Illicit drug users	10	0	
PLWHIV ^b	6	0	
Gender			
Female	208	41	19.7
Male	285	54	18.9
Color			
White	104	22	21.2
Non-white	389	73	18.8

^a LGBT: lesbian, gay, bisexual, transexual; PLWHIV: people living with HIV

^b IQR: interquartile range.

among symptomatic healthcare workers in a large university tertiary hospital in São Paulo, Brazil [Internet] *BMC Infect Dis* 2020;20(1):917 [cited 2021 mar 15]. Available from: <https://bmcinfectdis.biomedcentral.com/articles/10.1186/s12879-020-05662>.

Kamila Cardoso dos Santos
Grazielle Rosa da Costa e Silva
Winy Éveny Alves Moura
Larissa Silva Magalhães
Universidade Federal de Goiás, Faculdade de Enfermagem, Brazil

Brunna Rodrigues de Oliveira
Universidade Federal de Goiás, Instituto de Patologia Tropical e Saúde Pública, Brazil

Paulie Marcelly R. dos Santos Carvalho
Pontifícia Universidade Católica de Goiás, Escola de Ciências Sociais e da Saúde, Brazil

Karlla Antonieta Amorim Caetano
Universidade Federal de Goiás, Faculdade de Enfermagem, Brazil

Megmar Aparecida dos Santos Carneiro
Universidade Federal de Goiás, Instituto de Patologia Tropical e Saúde Pública, Brazil

Leonora Rezende Pacheco
Universidade Federal de Goiás, Faculdade de Enfermagem, Brazil

Clayton Luiz Borges
Juliana Alves Parente-Rocha
Gabriela Silvério Bazílio
Universidade Federal de Goiás, Instituto de Ciências Biológicas, Laboratório de Análises Clínicas e Ensino em Saúde, Brazil

Robert L. Cook
Krishna Vaddiparti
University of Florida, Department of Epidemiology, United States of America, Brazil

Claci Fátima Weirich Rosso
Sheila Araújo Teles*
Universidade Federal de Goiás, Faculdade de Enfermagem, Brazil

*Corresponding author. Universidade Federal de Goiás, Rua 227, Viela Q. 68, S/N - Setor Leste Universitário, Goiânia, GO, 74605-080, Brazil.
E-mail addresses: sateles@ufg.br, sheila.fen@ufg.br (S.A. Teles)

11 May 2021

Available online 27 August 2021

Funding/support statement

This study was funded by the National Council on Scientific and Technological Development/Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

And Foundation for Research Support in Goiás/Fundação de Amparo a Pesquisa de Goiás.

Declaration of competing interest

None.

References

- World Health Organization. *WHO coronavirus disease (COVID-19) dashboard* [Internet]. Data last updated: 5:52pm CET. 8 July 2021 [cited 2021 Jul 8]. Available from: <https://covid19.who.int/>.
- Engels D, Zhou X. Neglected tropical diseases: an effective global response to local poverty-related disease priorities. *Infect Dis Poverty* 2020;9(1):10 [Internet].
- Buonafine CP, Paiatto BNM, Leal FB, de Matos SF, de Moraes CO, Guerra GG, et al. High prevalence of SARS-CoV-2 infection