




# Accelerated Atherosclerosis in Systemic Lupus Erythematosus: Role of Fibroblast Growth Factor 23- Phosphate Axis

Yaser Ammar <sup>1</sup>  
Amira Mohamed <sup>1</sup>  
Gihane Khalil <sup>2</sup>  
Dalia Maharem <sup>1</sup>

<sup>1</sup>Internal Medicine Department, Medical Research Institute, Alexandria University, Alexandria, Egypt; <sup>2</sup>Chemical Pathology Department, Medical Research Institute, Alexandria University, Alexandria, Egypt

**Purpose:** Despite management advances, accelerated atherosclerotic cardiovascular disease (ACVD) remains a major cause of morbimortality in systemic lupus erythematosus (SLE) patients; that is not fully explained by traditional risk factors. Fibroblast growth factor-23 (FGF23) is a bone-derived phosphaturic hormone with multiple klotho-dependent and independent effects, including promotion of atherosclerosis and vascular calcification, particularly in the context of chronic kidney disease. Increased circulating FGF23 was reported in SLE patients, particularly with lupus nephritis (LN); but its atherogenic role in these disorders was not explored.

**Subjects and Methods:** Three study groups of predominantly middle-aged females were categorized by the 2012 SLE International Collaborating Clinics (SLICC) criteria as SLE (without LN), LN, or controls matching for traditional CVD risk profile. Measures of SLE activity, damage, steroid therapy, and glomerular filtration rate were calculated. Fasting blood samples were checked for serum lipid profile, anti-DNA, urea, creatinine, uric acid, proteins, albumin, calcium, phosphorus, C3, C4, CRP, vitamin-D3, intact parathyroid hormone and FGF23 (iFGF23). By carotid ultrasonography, mean common carotid artery intima-media thickness (CC-IMT), plaque score (PS) and internal carotid resistive index (ICRI) were recorded.

**Results:** CC-IMT, ICRI and serum iFGF23 differed along the study groups (LN>SLE>controls). In both SLE and LN patients, serum iFGF23 had a significant positive correlation with serum phosphorus, CC-IMT and PS. On multivariate analysis, the strongest predictor of increased CC-IMT was cumulative steroid dose in SLE and serum iFGF23 in LN patients. Most significant independent predictors of increased serum iFGF23 were hyperphosphatemia in SLE and proteinuria in LN patients.

**Conclusion:** FGF23-phosphate axis has a key role in accelerated ACVD in SLE patients. Serum phosphorus and iFGF23 should be included in ACVD risk profile assessment of these patients. Prospective studies shall define the role of dietary and/or pharmacologic control of hyperphosphatemia and proteinuria in reducing circulating iFGF23 and ACVD in them.

**Keywords:** atherosclerosis, FGF23, hyperphosphataemia, lupus, nephritis, ultrasonography

## Introduction

Systemic lupus erythematosus (SLE) is a prototype autoimmune disease predominantly affecting middle-aged women, typically with multiple organ-system involvement, and a protracted course with remissions, exacerbation and cumulative tissue damage. Despite marked geographical disparities, a review of recent epidemiologic studies denoted a trend for increasing global prevalence.<sup>1</sup> Although patients' life

Correspondence: Yaser Ammar  
Internal Medicine Department, Medical Research Institute, Alexandria University, Egypt  
Tel +2 1142389128  
Fax +203 – 4283719  
Email yaser\_ammara@yahoo.com

expectancy has improved in line with advances in diagnosis and treatment, it remains considerably lower compared with the general population.<sup>2</sup> Accelerated atherosclerotic cardiovascular disease (ACVD) constitutes the major comorbidity and the leading cause of death, with a frightening risk of myocardial infarction and stroke disproportionately striking young female patients, who are otherwise known to have low ACVD risk.<sup>3</sup> It has been long-recognized that this amplified risk cannot be fully accounted for by traditional (Framingham) risk factors, like hypertension, obesity and dyslipidaemia.<sup>4</sup> The immunologically geared chronic inflammatory status that typify the disease is a major inducer of ACVD in itself.<sup>3</sup> Corticosteroids, that have long constituted the backbone of effective anti-inflammatory therapy, are also atherogenic in the long-term.<sup>5</sup> The inclusion of SLE diagnosis and corticosteroid use in the newly developed QRISK3 score has significantly improved its CVD risk prediction in SLE patients.<sup>6</sup> Looking for novel, non-traditional, ACVD risk factors, with a focus on recently identified molecular biomarkers, is an evolving area of SLE research.<sup>7,8</sup>

There is now ample evidence for a link between bone disease and ACVD, both in the general population,<sup>9</sup> and in patients with SLE and other autoimmune diseases.<sup>10,11</sup> This disturbed bone-vascular axis may result from perturbations of vitamin D,<sup>12</sup> parathyroid hormone (PTH),<sup>13</sup> or fibroblast growth factor-23 (FGF23).<sup>14</sup> Principally secreted from osteocytes and osteoblasts in response to phosphorus loading, FGF23 has been identified as the major phosphaturic hormone (phosphatonin), that decreases proximal renal tubular phosphate reabsorption, intestinal phosphorus absorption, and vitamin D activation.<sup>15</sup> It was soon realized that FGF23 is a pleiotropic molecule, with a host of klotho-dependent and independent autocrine, paracrine and endocrine effects on almost all body tissues.<sup>16</sup> Increased circulating FGF23 is now recognized as the earliest biochemical abnormality in chronic kidney disease – mineral bone disorder (CKD-MBD).<sup>17</sup> Progressive renal impairment is paralleled with an exponential increase in circulating FGF23, approaching several thousands of normal in patients with end-stage renal disease.<sup>18</sup>

Epidemiologic studies have inked increased circulating FGF23 in subjects with normal renal function with both early functional alterations (impaired forearm flow-mediated dilatation)<sup>14</sup> and established carotid ultrasonographic (US) changes (increased intima-media thickness [IMT] and plaque presence)<sup>19,20</sup> of atherosclerosis in the

community. Similar associations were confirmed in CKD patients.<sup>21,22</sup> Therefore, serum FGF23 could improve the power of Framingham risk score to predict increased carotid IMT.<sup>23</sup> FGF23 may induce/augment some traditional atherosclerosis risk factors, such as hypertension,<sup>24</sup> dyslipidaemia,<sup>25</sup> insulin resistance and obesity.<sup>26</sup> Of more importance, however, is the association of FGF23 with a host of non-traditional ACVD risk factors such as chronic inflammation,<sup>27</sup> hypovitaminosis D,<sup>28</sup> and vascular calcification.<sup>29</sup> These latter factors become particularly prominent in presence of CKD. Increased FGF23 is also a significant predictor of CKD occurrence<sup>30</sup> and progression.<sup>31</sup>


Two small studies reported significantly higher serum FGF23 in SLE patients compared with controls, with a still higher level in patients with lupus nephritis (LN).<sup>32,33</sup> The inflammatory reaction of active lupus upregulates FGF23 production by osteocytes.<sup>34</sup> The development of LN is conducive for a further rise of circulating FGF23, possibly induced by hyperparathyroidism,<sup>35</sup> hyperphosphataemia,<sup>36</sup> impaired renal clearance,<sup>37</sup> and klotho deficiency.<sup>38</sup> Therefore, LN patients typically have in parallel significantly higher level of FGF23 and higher burden of ACVD, compared with SLE patients without nephritis.<sup>33</sup> FGF23 may thus prove to be an important mediator of accelerated ACVD in SLE patients, with a particularly more prominent role in patients with LN. Controlling the circulating level of this key molecule may then become a novel therapeutic approach to reduce cardiovascular morbimortality in SLE patients. Of note, however, studies exploring the cardiovascular risk of circulating FGF23 in SLE patients are lacking.<sup>39</sup>

## Materials and Methods

### Study Design and Participants

This was a cross-sectional, case control study, conducted in accordance with the Declaration of Helsinki and approved by the institutional medical ethics committee. Between March and November 2018, eligible participants providing written informed consent were recruited from the outpatient clinics of the Medical Research Institute, Alexandria University, Egypt, and subjected to full clinical evaluation and medical records review, followed by laboratory and imaging studies (Figure 1).

Diagnosis of SLE was based on the 2012 SLE International Collaborating Clinics (SLICC) criteria.<sup>40</sup>



**Figure 1** Study flowchart.

**Abbreviation:** SLICC, SLE International Collaborating Clinics.

LN was defined by persistent proteinuria  $> 0.5\text{gm/day}$  in the context of SLE.<sup>41</sup> SLE disease activity was assessed by SLE Disease Activity Index (SLEDAI),<sup>42</sup> whereas the extent of established damage was assessed by SLICC Damage Index.<sup>43</sup> Average steroid dose (Av\_S) was calculated as the average equivalent prednisone dose in mg/day. Cumulative steroid dose (Cum\_S) was obtained by multiplying Av\_S by disease duration in years. Steroid pulses were defined as short-term (3–5 days) intensification of basic steroid dose, mostly given parenterally during in-hospital admission. Traditional atherosclerosis risk factors (age decade, male sex, hypertension, dyslipidaemia, obesity, treatment for these conditions, and presence of ACVD in different vascular beds) were defined by standard criteria and given one point each. The total score (termed Second Manifestations of ARterial disease, or SMART score) provided a validated semiquantitative estimate of the burden of traditional ACVD risk factors.<sup>44,45</sup>

We excluded patients with diabetes mellitus, morbid obesity, smoking (current or past), pregnancy, estimated glomerular filtration rate (eGFR)  $< 30\text{ mL/min/1.73m}^2$ , age  $> 50$  years or insufficient data. Finally, 68 subjects (age 20–50 years, 9 males) were triaged into 3 study groups:

- G1 (Controls) (N = 20): selected on individual basis to be matching with the other groups in age, sex and SMART score.
- G2 (SLE) (N = 20): having SLE without LN.
- G3 (LN) (N = 28): having LN.

### Laboratory Studies<sup>46</sup>

After 12 hours fast, blood samples were drawn into EDTA tubes (for complete blood count) and serum separator tubes that were immediately transported and centrifuged, keeping serum at  $-80^{\circ}\text{C}$  until batch analysis was made for levels of serum low-density lipoproteins (LDL), high-density

lipoproteins (HDL), triglycerides (TGs), anti-double-stranded DNA (anti-DNA), complement components C3 and C4, C-reactive protein (CRP), urea, creatinine, uric acid, total proteins, albumin, calcium, phosphorus, vitamin D3, intact parathyroid hormone (iPTH) (3<sup>rd</sup> generation assay), and intact FGF23 (iFGF23) using commercially available kits according to manufacturers' instructions. Morning void urine was used for complete urinalysis. Twenty-four hours urine collections were used for quantitation of proteinuria. eGFR was calculated from stable serum creatinine using CKD-epidemiology collaboration (CKD-EPI) equation.<sup>47</sup>

## Carotid Duplex Ultrasonography<sup>48</sup>

It was done by an experienced sonographer blinded to the patients' data, using Acuson X300 US System, Siemens Healthineers (formerly Siemens Healthcare), USA. The patient was placed in a supine position with a slight head tilt away from the examined side. A linear-array transducer (VF10-5) with a frequency of 8 MHz was used to scan the carotid system on both sides and calculate the following:

1. Plaque Score (PS):<sup>49</sup> A carotid plaque was defined as either a focal protrusion of the intima-media layer encroaching into the arterial lumen (protuberant plaque), or a diffuse thickening of the intima-media layer measuring > 1.5 mm (diffuse plaque). These were looked for thoroughly in three distinct segments (distal common carotid, bulb, and proximal internal carotid arteries). The total number of plaque-bearing segments on both sides represented the PS, which ranged 0 to 6.
2. Mean Common Carotid Artery Intima-Media Thickness (CC-IMT):<sup>50</sup> The distal segment of the common carotid artery on each side was scanned with the probe in 3 directions (anterior, lateral, and posterior). Each time, a short clip was saved in which the intima-lumen and the media-adventitia interfaces on the arterial far wall were clearly delineated. Some clearly captured image frames were then analysed by the Syngo Arterial Health Package (AHP), an automatic edge detection software, to calculate the mean intima-media thickness along a one-centimetre plaque-free arterial segment. The composite mean CC-IMT was calculated by averaging the 6 readings (3 from each side).
3. Internal Carotid Artery Resistive Index (ICRI):<sup>51</sup> The insonation angle (between transducer beam

and axis of blood flow) was kept <60° and the pulse repetition frequency was adjusted to prevent aliasing while maximizing sensitivity and waveform size. Aided by colour flow mapping, the sampling gate was placed completely within the internal carotid flow. The spectral velocity-time curve was recorded while the patient holds breath. The ICRI was calculated as: [(peak-systolic flow velocity – end-diastolic flow velocity) divided by (peak-systolic flow velocity)] (Pourcelot's equation, measured from the same waveform). Two readings were obtained on each side and the average of the four readings was recorded.

Test-retest variabilities of the ultrasonographic measurements were generally < 5%.

## Statistical Methods

Data were analysed using SPSS software package version 20 (SPSS Inc., Chicago, Illinois, USA). Categorical data were expressed as absolute numbers (percentages) and compared by Chi-square or Fisher exact test. Continuous data were tested for normality using Shapiro Wilk test. Parametric data were presented as mean ± SD and compared by analysis of variance (ANOVA). Non-parametric data were presented as median (interquartile range) and compared by Mann-Whitney U or Kruskal-Wallis *H*-test. Post-hoc analysis was performed for pairwise comparisons. Correlations were tested by the Spearman's rank correlation coefficient. Models for multivariate linear regression analysis were constructed separately in G2 and G3, introducing clinically relevant predictors with unadjusted *P* < 0.08. Significance was judged at the 5% level.

## Results


The three study groups were comparable regarding sex, age, frequency of hypertension and dyslipidaemia, SMART score, white blood cells, serum TGs, HDL, uric acid, and PS (Table 1). Compared with the other two groups, controls had significantly higher haemoglobin, serum C4 and vitamin D3, and significantly lower serum anti-DNA, CRP, and iPTH. SLE (G2) patients had significantly higher serum proteins and significantly lower platelets than the other two groups. LN patients had significantly higher proteinuria, serum phosphorus and ICRI and significantly lower serum calcium than the other two groups (Figure 2). Compared with controls, LN patients had significantly higher serum

**Table 1** Comparison Between the 3 Study Groups

Parameter	Controls (Cont) (N = 20)	Lupus Only (SLE) (N = 20)	Lupus Nephritis (LN) (N = 28)	Statistical Test	P value (Post- hoc Differences)
Male Female	3 (15%) 17 (85%)	2 (10%) 18 (90%)	4 (14.3%) 24 (85.7%)	Fisher exact test	0.57
Age (Years)	35.56 ± 7.5	31.15 ± 7.8	36.11 ± 9.3	ANOVA	0.127
HTN No HTN	6 (30%) 14 (70%)	3 (15%) 17 (85%)	11 (39.3%) 17 (60.7%)	Chi square	0.19
DL No DL	8 (40%) 12 (60%)	6 (30%) 14 (70%)	15 (53.6%) 13 (46.4%)	Chi square	0.524
SMART Score	5 (4–6)	4.5 (4–5)	4 (4–6)	KWH	0.743
S. LDL (mg/dL)	121.89 ± 21	108.6 ± 15.2	130.96 ± 33.1	ANOVA	0.016* (LN > SLE)
S. TGs (mg/dL)	170.5 (151–188.3)	160.5 (136–171.3)	155 (120–198.3)	KWH	0.311
S. HDL (mg/dL)	44 (41.3–49.8)	44.5 (42.8–47)	45 (41.8–49.3)	KWH	0.828
Haemoglobin (gm/dL)	13.54 ± 0.9	9.87 ± 1.9	10.18 ± 1.4	ANOVA	< 0.001** (Cont > Others)
WBCs (k/uL)	7 (4.5–7.5)	7.8 (3.7–9.5)	6.9 (5.5–9.2)	KWH	0.627
Platelets (k/uL)	288.5 (263.5–309)	217 (192.3–273.5)	297 (245–320.5)	KWH	0.018* (SLE < Others)
S. Anti-DNA (IU/mL)	37 (28.3–43.5)	71 (48.8–102.8)	135 (69.5–210.3)	KWH	< 0.001** (Cont < Others)
S. C3 (mg/dL)	132 ± 29.7	93.85 ± 24.4	56.83 ± 26.4	ANOVA	0.001** (LN < SLE < Cont)
S. C4 (mg/dL)	48.67 ± 9.3	21.33 ± 9.1	18.89 ± 9.9	ANOVA	< 0.001** (Cont > Others)
S. CRP (mg/L)	3.2 (2.4–4)	5.5 (4–7)	7 (5.8–8.3)	KWH	< 0.001** (Cont < Others)
S. Urea (mg/dL)	23.5 (22–30)	31 (27.8–44.5)	48 (29.8–80.3)	KWH	< 0.001** (LN > Cont)
S. Creatinine (mg/dL)	0.9 (0.8–1.1)	0.8 (0.7–1)	1.1 (0.9–1.6)	KWH	0.011* (LN > SLE)
eGFR (mL/min/1.73)	89.2 (66.9–103.3)	99.6 (75.2–120.7)	70.2 (44.7–95.8)	KWH	0.006** (LN < SLE)
S. Uric Acid (mg/dL)	5.2 (4.2–6.3)	4.7 (4.3–5.4)	5.7 (4.8–7.2)	KWH	0.237
Proteinuria (gm/day)	0.24 (0.2–0.3)	0.15 (0.1–0.3)	2.5 (0.86–4.78)	KWH	< 0.001** (LN > Others)
S. Proteins (gm/dL)	6.4 (5.8–7.2)	7.3 (6.6–7.6)	6.6 (6.2–7)	KWH	0.007** (SLE > Others)
S. Albumin (gm/dL)	3.4 (3.2–3.7)	4 (3.4–4.2)	3.2 (2.7–3.7)	KWH	< 0.001** (LN < SLE)
S. Calcium (mg/dL)	10 (9.5–10.5)	10.1 (9–10.6)	8.8 (8.3–8.9)	KWH	< 0.001** (LN < Others)
S. Phosphorus (mg/dL)	3.73 ± 0.6	3.74 ± 0.6	4.53 ± 0.8	ANOVA	0.001** (LN > Others)
S. Vitamin D3 (ng/mL)	34.5 (29.6–40.8)	12.5 (11–15.8)	12 (10.9–13.1)	KWH	< 0.001** (Cont > Others)
S. iPTH (pg/mL)	46.5 (32.5–51.8)	107.5 (84–121.3)	105 (92.3–120)	KWH	< 0.001** (Cont < Others)
S. iFGF23 (pg/mL)	108.5 (92.4–152.2)	409.5 (313–483.2)	771.7 (579.2–900)	KWH	< 0.001** (LN > SLE > Cont)
PS	0 (0–0)	0 (0–1)	0 (0–1)	KWH	0.427
CC-IMT (mm)	0.47 (0.4–0.5)	0.517 (0.464–0.577)	0.541 (0.425–0.661)	KWH	0.01* (LN > Cont)
ICRI	0.61 (0.6–0.6)	0.62 (0.58–0.63)	0.66 (0.61–0.71)	KWH	0.001** (LN > Others)

**Notes:** Data are expressed as mean + SD or median (interquartile range), \*Significant (P < 0.05), \*\*Highly significant (P < 0.01).

**Abbreviations:** HTN, hypertension; DL, dyslipidemia; S, serum; LDL, low-density lipoproteins; TGs, triglycerides; HDL, high-density lipoproteins; WBCs, white blood cells; k, X1000; CRP, C-reactive protein; eGFR, estimated glomerular filtration rate; iPTH, intact parathyroid hormone; iFGF23, intact fibroblast growth factor-23; PS, Plaque score; CC-IMT, common carotid intima-media thickness; ICRI, internal carotid resistive index; ANOVA, analysis of variance; KWH, Kruskal–Wallis H-Test.



**Figure 2** Comparison of serum phosphorus in the 3 study groups. \*\*Highly significant ( $P < 0.01$ ).


urea and CC-IMT (Figure 3). Compared with SLE (G2), LN patients had significantly higher serum LDL and creatinine and significantly lower eGFR and serum albumin. Serum C3 and iFGF23 showed significant changes between the 3 groups (Figure 4). Compared with SLE (G2), LN patients had significantly higher Cum\_S and SLEDAI (Table 2).

In both G2 and G3, serum iFGF23 had a statistically significant positive correlation with serum phosphorus, CC-IMT and PS; and these two US parameters significantly correlated with each other (Table 3, Figure 5). Serum iFGF23 had a statistically significant negative correlation with eGFR (in G2), and a statistically significant positive correlation with proteinuria (in G3). Serum phosphorus had a statistically significant positive correlation with CC-IMT and PS (in G2), and a statistically significant negative correlation with eGFR (in G3). Cum\_S had a statistically significant positive correlation with CC-IMT in G2; this correlation was reversed and of weaker significance in G3.

In the regression analysis, the strongest predictor of increased CC-IMT was Cum\_S in G2 and serum iFGF23 in G3. The strongest predictor of increased serum iFGF23 was hyperphosphatemia in G2 and proteinuria in G3 (Table 4).

## Discussion

This study confirms previous reports that serum FGF23 is elevated in SLE patients (more so in LN),<sup>32,33</sup> and generates a conceptual framework for its atherogenic role in these patients. We excluded subjects with exaggerated ACVD risk profile because of diabetes mellitus, obesity, advanced renal disease or age. Controls (G1) were individually selected to match with lupus patients (G2,3) in the traditional ACVD risk profile, so that inter-group differences are largely determined by the lupus disease and related factors. We used CKD-EPI equation for eGFR calculation as it performed best in SLE patients, compared with other creatinine-based formulas.<sup>52</sup> Subclinical atherosclerosis was evaluated in the carotid arteries by 3 integrative US measures, of proven benefit for quantitation of atherosclerosis extent,<sup>53</sup> and progression,<sup>54,55</sup> as well as prediction of future CVD events in SLE patients.<sup>56</sup> IMT is a sensitive early marker of generalised atherosclerosis.<sup>57</sup> Its measurement at the distal common carotid artery offers easier accessibility and higher accuracy and reproducibility, over other carotid segments.<sup>58</sup> However, IMT may increase because of media thickening as a function of age and increased blood pressure, rather than




**Figure 3** Comparison of common carotid intima-media thickness in the 3 study groups. \*Significant ( $P < 0.05$ ).

atherosclerosis.<sup>59</sup> Therefore, the measurement of established focal atherosclerotic protuberances by PS was intended to further improve the quantitation of subclinical atherosclerosis and CVD risk.<sup>60</sup> The simplified PS score we adopted may be less prone to assessment errors, and has repeatedly proved useful for quantification of carotid atherosclerosis in SLE patients and assessment of its progression.<sup>54,61</sup> Carotid US included a measurement of ICRI, a potentially more sensitive measure for subtle early functional vascular alterations,<sup>62</sup> with comparable cardiovascular morbimortality predictor power to the standard IMT.<sup>51</sup>

Among these predominantly middle-aged females with relatively low traditional ACVD risk profile, both CC-IMT and ICRI showed a graded rise along the 3 study groups (LN > SLE > Controls). LN patients had significantly higher CC-IMT compared with controls, and significantly higher ICRI compared with the other 2 groups. These results conform with the previous reports that SLE patients suffer an increased burden of subclinical atherosclerosis<sup>3,5,7,13</sup> and that this burden is much significantly amplified and to a large extent determined, by the presence of LN.<sup>63–66</sup> Contrary to some previous

studies,<sup>67,68</sup> we did not find significant differences between the study groups regarding PS, which may be explained by the relatively young age, low overall ACVD risk and low plaque occurrence, in the study subjects. Serum iFGF23 varied the same way as CC-IMT and ICRI (LN > SLE > Controls), with significant differences between the 3 study groups, also in line with the few previous reports.<sup>32,33</sup>

Our study revealed multiple possible correlates for the increased ACVD risk in SLE patients. Some correlates encompassed all SLE patients (G 2,3); and some others were particularly distinguished among LN patients. Correlates common for all SLE patients were mainly indicative of the overwhelming immunologically driven, chronic inflammatory nature of the disease; and have been reported by previous studies, like increased serum anti-DNA,<sup>69</sup> and CRP,<sup>70</sup> and depressed serum C3 and C4.<sup>63</sup> Clearly, the lupus-related parameters included in Table 2 also tackle SLE patients as whole; and may constitute an indirect measure of the immune-inflammatory disease pathogenesis, which is thought to progress in parallel with atherogenesis.<sup>71</sup> Hypovitaminosis D was also common to all SLE (G2,3) patients, a nearly universal finding in previous studies, that is mainly ascribed to



**Figure 4** Comparison of serum iFGF23 in the 3 study groups. \*Significant ( $P < 0.05$ ), \*\*Highly significant ( $P < 0.01$ ).

insufficient sun exposure and drug-induced accelerated vitamin D catabolism.<sup>12,72–74</sup> Somewhat unexpectedly, PTH was significantly increased in all SLE patients (including those without LN). Increased circulating PTH and its association with ACVD in SLE patients was only recently reported,<sup>13</sup> and may be the result of concomitant hypovitaminosis D.<sup>75,76</sup>

As shown in Table 1, LN patients have, in comparison with other SLE patients, the added risks of proteinuria, renal impairment, hyperphosphatemia and a higher FGF23 elevation. In community studies, both proteinuria,<sup>77–80</sup> and

GFR decline,<sup>81–84</sup> have significant, additive, dose-dependent associations with subclinical atherosclerosis burden and CVD events. These associations were firmly reproduced in studies of LN patients.<sup>63–66</sup> Clearly, the development of LN engenders a more aggressive lupus phenotype, evidenced in the present cohort by having significantly lower serum C3, and significantly higher LDL, SLEDAI, and Cum\_S, compared with SLE (G2) patients.

An intriguing finding is that Cum\_S had a statistically significant positive correlation with CC-IMT in G2 (SLE

**Table 2** Comparison of Lupus-Related Parameters in SLE Patients with and without Lupus Nephritis (LN)

Parameter	SLE (without LN) (N = 20)	LN (N = 28)	P (Mann Whitney U-Test)
Lupus Duration (Y)	3 (2–4.3)	3.5 (2.4–6)	0.194
Average Steroid Dose	22.5 (18.8–30)	30 (20–30)	0.272
Cumulative Steroid Dose	60 (40–105)	90 (60–165)	0.049*
Steroid Pulses	1 (0–1.3)	1 (0–2)	0.217
SLEDAI Activity Index	6.5 (4–9.3)	10 (6–12)	0.046*
SLICC Damage Index	0 (0–1.3)	1 (0–2)	0.628

**Notes:** Average Steroid Dose: Expressed as average equivalent prednisone dose in mg/day. Cumulative Steroid Dose: Average steroid dose multiplied by disease duration in years. Steroid Pulses: Total number of pulses given throughout the disease history; a pulse is defined as an exceptionally high steroid dose given for 3–5 days. \*Significant ( $P < 0.05$ ).

**Abbreviations:** SLEDAI, systemic lupus erythematosus disease activity index; SLICC, SLE International Collaborating Clinics damage index.



**Table 3** Statistical Correlations Between Some Parameters in SLE Patients with and without Lupus Nephritis (LN)

		eGFR	Proteinuria	S. Phosphorus	S. iFGF23	Cum_S	PS	CC-IMT
eGFR	r	<b>SLE LN</b>	- 0.334	- 0.153	- 0.446	- 0.56	- 0.245	- 0.297
	P		0.15	0.519	0.049*	0.01*	0.299	0.203
Proteinuria	r	- 0.139	<b>SLE LN</b>	- 0.184	0.244	0.479	0.245	0.152
	P	0.48		0.437	0.301	0.033*	0.297	0.523
S. Phosphorus	r	-0.458	0.34	<b>SLE LN</b>	0.513	0.295	0.552	0.539
	P	0.014*	0.076		0.021*	0.207	0.012*	0.014*
S. iFGF23	r	- 0.31	0.479	0.383	<b>SLE LN</b>	0.414	0.675	0.568
	P	0.108	0.01*	0.044*		0.069	0.001**	0.009**
Cum_S	r	- 0.234	0.087	0.115	- 0.12	<b>SLE LN</b>	0.263	0.689
	P	0.23	0.659	0.56	0.543		0.263	0.001**
PS	r	0.315	0.295	- 0.054	0.431	- 0.281	<b>SLE LN</b>	0.485
	P	0.102	0.128	0.784	0.022*	0.147		0.03*
CC-IMT	r	0.057	0.338	- 0.06	0.422	- 0.396	0.478	<b>SLE LN</b>
	P	0.774	0.078	0.76	0.025*	0.037*	0.01*	


**Notes:** \*Significant ( $P < 0.05$ ), \*\*Highly significant ( $P < 0.01$ ). Correlations in G2 (SLE without LN,  $N = 20$ ) and G3 (LN,  $N = 28$ ) are shown in right upper and left lower halves of the table, respectively.

**Abbreviations:** S, serum; eGFR, estimated glomerular filtration rate; iFGF23, intact fibroblast growth factor 23; Cum\_S, cumulative steroid dose; PS, plaque score; CC-IMT, common carotid intima-media thickness.

only) that remained marginally significant ( $P = 0.05$ ) in the adjusted model, whereas this correlation was reversed (still significant but somewhat weaker) in G3 (LN patients). Corticosteroids in SLE represent a double-edged sword.<sup>85</sup> It is possible that their pro-atherogenic effects (insulin resistance, dyslipidaemia, obesity, and hypertension) dominated in the absence of renal disease.<sup>5</sup> LN, as a hallmark of more aggressive disease, determined a greater need for the immunosuppressive and anti-inflammatory steroid actions, imparting a better benefit/risk ratio and a net anti-atherogenic action.<sup>86</sup> Possibly also the steroid dose was not precisely tailored for the different patient needs, being relatively overdosed in absence, and underdosed in presence, of LN; or there may be endogenous individual variations that determine the patient's response to these medications.<sup>85</sup>

The FGF23/klotho axis is the principal regulator of phosphorus metabolism.<sup>87</sup> Tight regulation of serum phosphorus is essential since hyperphosphatemia, and even minor elevations of serum phosphorus within the normal range, have been increasingly linked in community studies with impaired vasoreactivity,<sup>88</sup> subclinical atherosclerosis,<sup>89,90</sup> vascular calcification,<sup>91</sup> cardiovascular events,<sup>92,93</sup> and mortality.<sup>94</sup> Similar strong associations were also found in CKD

patients.<sup>95–97</sup> Growing evidence for cardiovascular phosphorus toxicity led some authors suggest it the “new cholesterol”.<sup>98</sup> Surprisingly, serum phosphorus was rarely reported in SLE patients. Similar to the present study, one study reported serum phosphorus to be significantly higher in LN patients compared with lupus patients without nephritis and controls.<sup>99</sup> There is evidence that increased serum phosphorus in LN patients would augment further systemic inflammation, vascular calcification and faster ACVD.<sup>100–103</sup> Another study found serum phosphorus to be marginally elevated in SLE patients with atheromatous plaques or increased arterial IMT compared with patients lacking these signs, but the differences were insignificant.<sup>13</sup> In the present study, serum phosphorus in G2 (SLE only) had a significant correlation with CC-IMT and PS, although it was within normal range ( $< 4.5$  mg/dL) in all but one patient. A Finnish population study found a significant direct correlation between dietary phosphorus intake and CC-IMT.<sup>90</sup> Unawareness about excessive dietary phosphorus consumption is a recently recognised global health problem,<sup>104</sup> particularly afflicting less-privileged societies habitually ingesting highly absorbable phosphorus-rich food additives.<sup>105</sup>



**Figure 5** Correlation of serum iFGF23 with common carotid intima-media thickness (upper) and plaque score (lower) in SLE patients (groups 2 and 3 together). \*\*Highly significant ( $P < 0.01$ ).

Normally, increased secretion of the counter-regulatory phosphaturic hormone, FGF23, occurs in a robust response to phosphorus loading.<sup>15,106</sup> This adaptive FGF23 rise (typically not exceeding 200 pg/mL), utilizing klotho-dependent pathways, generates a controlled phosphaturic response destined to prevent hyperphosphatemia and subsequent perturbations of the bone-vascular axis (bone disease, atherosclerosis, and vascular calcification). A maladaptive FGF23 surge, largely signalling through klotho-independent pathways, approaches much higher values but fails to prevent hyperphosphatemia and rather produces off target effects like hypovitaminosis D, bone rarefaction, ACVD, and

myocardial hypertrophy.<sup>107,108</sup> A full-blown picture of this maladaptive FGF23 increase occurs in CKD, when klotho deficiency, low GFR, and hypovitaminosis D generate a state of progressive FGF23 resistance;<sup>109</sup> leading in advanced CKD to a marked rise of both serum phosphorus and FGF23 in a characteristic strong direct correlation.<sup>110,111</sup> A hallmark for the disturbed FGF23-phosphorus axis in SLE patients (G2 and 3) in the present study was the rise of both biomarkers and the presence of a significant direct correlation between them. The maladaptive FGF23 rise was not restricted to LN patients; it involved SLE patients without evidence of renal disease as well. Indeed, G2 patients (SLE


**Table 4** Multivariate Linear Regression Analysis for Predictors of Increased CC-IMT and S. iFGF23 in SLE Patients with and without Lupus Nephritis (LN)

<b>A: Outcome: Increased CC-IMT</b>					
<b>G2: SLE Patients (N = 20)</b>			<b>G3: LN Patients (N = 28)</b>		
<b>Model R<sup>2</sup></b>		<b>0.614</b>	<b>Model R<sup>2</sup></b>		<b>0.339</b>
<b>Model Significance</b>		<b>0.029*</b>	<b>Model Significance</b>		<b>0.042*</b>
<b>Predictors</b>	<b>β Coefficient</b>	<b>Adjusted P</b>	<b>Predictors</b>	<b>β Coefficient</b>	<b>Adjusted P</b>
<b>Cum_S</b>	0.833	0.05*	<b>S. iFGF23</b>	0.442	0.09
<b>S. iFGF23</b>	0.195	0.475	<b>Cum_S</b>	- 0.053	0.792
<b>S. Phosphorus</b>	0.043	0.86	<b>Dyslipidaemia</b>	0.163	0.43
<b>Haemoglobin</b>	- 0.420	0.089	<b>Proteinuria</b>	0.099	0.64
<b>Dyslipidaemia</b>	- 0.364	0.214			
<b>Age</b>	- 0.308	0.292			
*Significant (P < 0.05).					
<b>B: Outcome: Increased S. iFGF23</b>					
<b>G2: SLE Patients (N = 20)</b>			<b>G3: LN Patients (N = 28)</b>		
<b>Model R<sup>2</sup></b>		<b>0.611</b>	<b>Model R<sup>2</sup></b>		<b>0.463</b>
<b>Model Significance</b>		<b>0.005**</b>	<b>Model Significance</b>		<b>0.028*</b>
<b>Predictors</b>	<b>β Coefficient</b>	<b>Adjusted P</b>	<b>Predictors</b>	<b>β Coefficient</b>	<b>Adjusted P</b>
<b>S. LDL</b>	0.072	0.73	<b>SMART Score</b>	0.233	0.277
<b>S. Phosphorus</b>	0.397	0.049*	<b>Dyslipidaemia</b>	0.231	0.209
<b>S. Creatinine</b>	0.276	0.197	<b>Proteinuria</b>	0.466	0.026*
<b>Cum_S</b>	0.314	0.121	<b>S. Creatinine</b>	- 0.046	0.828
			<b>Av_S</b>	0.059	0.762
			<b>S. Phosphorus</b>	0.011	0.954
*Significant (P < 0.05), **Highly significant (P < 0.01).					

**Abbreviations:** CC-IMT, common carotid intima-media thickness; Cum\_S, cumulative steroid dose; S, serum; iFGF23, intact fibroblast growth factor 23; LDL, low-density lipoproteins; Av\_S, average steroid dose.

only) had a significantly increased serum iFGF23 over controls (exceeding the typical level of adaptive increase and matching with a previous study),<sup>32</sup> and an even stronger direct correlation between serum iFGF23 and phosphorus compared to this correlation in LN patients. In multivariate analysis, serum phosphorus persisted in G2 as the most (and only) significant independent predictor of increased serum iFGF23. We thus infer that the maladaptive FGF23 hypersecretion can occur in SLE patients as an effect of the lupus phenotype itself, independently of renal disease. At least two

salient, renal function independent, features of SLE can synergistically reduce klotho gene expression and induce significant FGF23 resistance and hypersecretion, namely, chronic inflammation,<sup>27,34,112,113</sup> and hypovitaminosis D.<sup>114,115</sup> The latter might also result in increased serum PTH which provides another stimulus for FGF23 hypersecretion,<sup>116</sup> a possibility supported by the finding of increased serum PTH in SLE patients in the present study and a previous one.<sup>13</sup> The issue of FGF23 resistance in SLE requires a separate study.



**Figure 6** Simplified schema for the role of FGF23 in atherogenesis in SLE patients. Factors in the upper (yellow) part are typical early findings in all (inflammation) or a large proportion of (hypovitaminosis D and dyslipidaemia) SLE patients, whereas the other factors (lower gold part) are either dependent on (proteinuria and renal impairment), or largely determined by, the presence and severity of LN.

Previous studies built a consensus that ACVD in SLE mainly (but by no means exclusively) develops in conjunction with LN, with or without renal impairment.<sup>63–66</sup> Consistent with this, we found the median CC-IMT in SLE patients, with or without LN, increased over controls; but the difference was statistically significant only with LN patients. Furthermore, we found a significantly increased ICRI in LN patients over the other two groups, denoting a more generalized vascular pathology conducive for atherogenesis in LN patients.<sup>62</sup> Serum iFGF23 had a statistically significant positive correlation with CC-IMT and PS, both in G2 and G3 (Table 3), and a similar correlation with ICRI in G2 (not shown). Regression analysis in LN patients revealed that the strongest independent predictor of increased CC-IMT was the increased serum iFGF23; and that proteinuria was the strongest (and the only significant) independent predictor of increased serum iFGF23 in these patients.

Therefore, increased FGF23 may, at least in part, explain the strong association between proteinuria and ACVD observed in LN patients,<sup>63–66</sup> an association that has been reported as well in type 2 diabetic patients,<sup>117</sup> and even in the healthy population.<sup>118</sup> The explanation for FGF23 rise in proteinuric nephropathies was based on results of the KNOW-CKD<sup>119</sup> and subsequent studies.<sup>120,121</sup> Proteinuria decreases the biologic activity of FGF23 on renal tubules, independent of renal function.<sup>120</sup> Albuminuria induces a state endoplasmic reticulum stress in renal tubular cells, leading to downregulation of klotho production.<sup>122</sup> This leads to FGF23 hypersecretion that fails to correct hyperphosphatemia (more so if renal function is impaired as well). In these cases, an enormous ACVD risk burden would result from the combined effects of increased FGF23, hyperphosphatemia, hypovitaminosis D, and renal impairment.<sup>123</sup> Identification of this multifactorial model,

with the focal role of the proteinuria-hyperphosphatemia axis, is an important addition to our understanding of the ACVD in SLE patients.

Based on the present work and current knowledge, we constructed a simplified conceptual framework (Figure 6) for the role of FGF23 in the pathogenesis of ACVD in SLE patients. Although not all inclusive, the factors numbered one through ten may be regarded as the key players in this paradigm. Factors in the upper (yellow) pane are nearly always present in all SLE patients since the disease beginning. Inflammation is meant to denote the SLE disease process itself, as well as markers of disease activity and treatment; which continue to have a significant impact on ACVD risk and long-term outcome.<sup>3,5,6</sup> Factors in the lower (gold) pane occur mainly or exclusively in patients with LN, when all risk factors become significantly amplified. FGF23 lies at the focal point of intersection of several self-perpetuating (vicious) cycles. For example, increased circulating FGF23 further exaggerates hypovitaminosis D, which is present in most SLE patients.<sup>28,72–74</sup> Hypovitaminosis D exaggerates inflammation<sup>12,124</sup> and hyperparathyroidism<sup>75,76</sup> which independently complete two vicious circles for FGF23 hypersecretion. We can likewise describe another vicious circle in which proteinuria causes hyperphosphatemia and FGF23 hypersecretion;<sup>120,121</sup> then FGF23 fails to correct hyperphosphatemia which even continues to progress causing further renal damage,<sup>14,30,31</sup> and resistance to the renoprotective effects of angiotensin converting enzyme inhibitors,<sup>125</sup> and low protein diet.<sup>126</sup> Unless one or more of these vicious circles is interrupted, FGF23 levels would continue to rise inexorably with increased burden of ACVD and further CKD progression. We propose that breaking these cycles to control circulating FGF23 level (for example by controlling hyperphosphatemia and proteinuria) might provide novel approaches for reduction of ACVD risk in SLE patients. It should be noticed, however, that directly targeting the FGF23 itself by neutralizing antibodies in a rat model of CKD-MBD led to a dose-dependent increase in serum phosphorus, aortic calcification, and mortality.<sup>127</sup> Therefore, well-designed prospective studies shall define the optimal circulating FGF23 range offering the best compromise between adaptive and maladaptive effects and resulting in the best measures of long-term outcome.

To the best of our knowledge, this is the first study to explore the role of FGF23 and related parameters in subclinical atherosclerosis in SLE patients with or without LN. We acknowledge the limitations of the relatively

small sample size and the cross-sectional design of the study; so that a cause-effect relationship between the study parameters cannot be readily inferred. The simplified PS we adopted does not account for the multiplicity of plaques in one arterial segment or variations in their area and size, as offered by more elaborate scores.<sup>49</sup> Subclinical atherosclerosis assessed in only one vascular bed might not sufficiently reflect its generalized burden. Better exploration of FGF23 actions would have required assessment of dietary phosphorus intake, urinary phosphorus excretion and circulating klotho level.

## Conclusion

The FGF23-phosphate axis has a key role in accelerated ACVD in SLE patients. A maladaptive FGF23 hypersecretion coupled with renal tubular resistance to its phosphaturic action operates several self-perpetuating cycles leading to progressive hyperphosphatemia, hyperparathyroidism, hypovitaminosis D and ACVD. Most significant independent predictors of FGF23 hypersecretion were hyperphosphatemia and proteinuria, in SLE patients without and with LN, respectively. Serum phosphorus and iFGF23 should be included in the ACVD risk profile assessment of SLE patients. Dietary and/or pharmacologic control of hyperphosphatemia and proteinuria can be feasible therapeutic targets to reduce circulating FGF23 and ACVD burden in SLE patients. Large-scale prospective studies are needed to define the circulating FGF23 target range in SLE/LN patients providing the best compromise between its adaptive and maladaptive effects, and to assess the effects of reduction of serum phosphorus, iFGF23 and proteinuria on ACVD progression and CVD events in SLE/LN patients. Dietary vigilance for phosphorus-rich food additives may be a readily feasible intervention that can be broadly promulgated in less-privileged communities.

## Acknowledgments

This study has been presented in an abstract form in the 56th ERA-EDTA Congress (June 13–16, 2019, Budapest, Hungary) and published in *Nephrology Dialysis Transplantation*, Volume 34, Issue Supplement\_1, June 2019, gfz106.FP220, <https://doi.org/10.1093/ndt/gfz106.FP220>.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Rees F, Doherty M, Grainge MJ, Lanyon P, Zhang W. The world-wide incidence and prevalence of systemic lupus erythematosus: a systematic review of epidemiological studies. *Rheumatology*. 2017;56(11):1945–1961. doi:10.1093/rheumatology/kex260
- Urowitz MB, Gladman DD, Tom BD, Ibanez D, Farewell VT. Changing patterns in mortality and disease outcomes for patients with systemic lupus erythematosus. *J Rheumatol*. 2008;35(11):2152–2158. doi:10.3899/jrheum.080214
- Appleton BD, Major AS. The latest in systemic lupus erythematosus-accelerated atherosclerosis: related mechanisms inform assessment and therapy. *Curr Opin Rheumatol*. 2021;33(2):211–218. doi:10.1097/BOR.0000000000000773
- Esdaile JM, Abrahamowicz M, Grodzicky T, et al. Traditional Framingham risk factors fail to fully account for accelerated atherosclerosis in systemic lupus erythematosus. *Arthritis Rheum*. 2001;44(10):2331–2337. doi:10.1002/1529-0131(200110)44:10<2331::AID-ART395>3.0.CO;2-I
- Doria A, Shoenfeld Y, Wu R, et al. Risk factors for subclinical atherosclerosis in a prospective cohort of patients with systemic lupus erythematosus. *Ann Rheum Dis*. 2003;62(11):1071–1077. doi:10.1136/ard.62.11.1071
- Edwards N, Langford-Smith AW, Parker BJ, et al. QRISK3 improves detection of cardiovascular disease risk in patients with systemic lupus erythematosus. *Lupus Sci Med*. 2018;5(1):e000272. doi:10.1136/lupus-2018-000272
- Teixeira V, Tam L-S. Novel insights in systemic lupus erythematosus and atherosclerosis. *Front Med*. 2018;4:262. doi:10.3389/fmed.2017.00262
- Rho YH, Chung CP, Oeser A, et al. Novel cardiovascular risk factors in premature coronary atherosclerosis associated with systemic lupus erythematosus. *J Rheumatol*. 2008;35(9):1789–1794.
- den Uyl D, Nurmohamed MT, van Tuyl LH, Raterman HG, Lems WF. (Sub) clinical cardiovascular disease is associated with increased bone loss and fracture risk; a systematic review of the association between cardiovascular disease and osteoporosis. *Arthritis Res Ther*. 2011;13(1):1–19. doi:10.1186/ar3224
- Ajeganova S, Gustafsson T, Jogestrand T, Frostegård J, Hafström I. Bone mineral density and carotid atherosclerosis in systemic lupus erythematosus: a controlled cross-sectional study. *Arthritis Res Ther*. 2015;17(1):1–13. doi:10.1186/s13075-015-0595-4
- Bultink IE. Bone disease in connective tissue disease/systemic lupus erythematosus. *Calcif Tissue Int*. 2018;102(5):575–591. doi:10.1007/s00223-017-0322-z
- Hiemstra TF, Lim K, Thadhani R, Manson JE. Vitamin D and atherosclerotic cardiovascular disease. *J Clin Endocrinol Metab*. 2019;104(9):4033–4050. doi:10.1210/je.2019-00194
- Giannelou M, Skarlis C, Stamouli A, Antypa E, Moutsopoulos HM, Mavragani CP. Atherosclerosis in SLE: a potential role for serum parathormone levels. *Lupus Sci Med*. 2020;7(1):e000393. doi:10.1136/lupus-2020-000393
- Mirza MA, Larsson A, Lind L, Larsson TE. Circulating fibroblast growth factor-23 is associated with vascular dysfunction in the community. *Atherosclerosis*. 2009;205(2):385–390. doi:10.1016/j.atherosclerosis.2009.01.001
- Kocelak P, Olszanecka-Glinianowicz M, Chudek J. Fibroblast growth factor 23—structure, function and role in kidney diseases. *Adv Clin Exp Med*. 2012;21(3):391–401.
- Erben RG. Update on FGF23 and Klotho signaling. *Mol Cell Endocrinol*. 2016;432:56–65. doi:10.1016/j.mce.2016.05.008
- Isakova T, Wahl P, Vargas GS, et al. Fibroblast growth factor 23 is elevated before parathyroid hormone and phosphate in chronic kidney disease. *Kidney Int*. 2011;79(12):1370–1378. doi:10.1038/ki.2011.47
- Chathoth S, Al-Mueilo S, Cyrus C, et al. Elevated fibroblast growth factor 23 concentration: prediction of mortality among chronic kidney disease patients. *Cardiorenal Med*. 2016;6(1):73–82. doi:10.1159/000440984
- Rodríguez-Ortiz ME, Alcalá-Díaz JF, Canalejo A, et al. Fibroblast growth factor 23 predicts carotid atherosclerosis in individuals without kidney disease. The CORDIOPREV study. *Europ J Int Med*. 2020;74:79–85. doi:10.1016/j.ejim.2019.12.008
- Shah NH, Dong C, Elkind MS, et al. Fibroblast growth factor 23 is associated with carotid plaque presence and area: the Northern Manhattan Study. *Arterioscler Thromb Vasc Biol*. 2015;35(9):2048–2053. doi:10.1161/ATVBAHA.115.305945
- Yilmaz MI, Sonmez A, Saglam M, et al. FGF-23 and vascular dysfunction in patients with stage 3 and 4 chronic kidney disease. *Kidney Int*. 2010;78(7):679–685. doi:10.1038/ki.2010.194
- Yilmaz G, Ustundag S, Temizoz O, et al. Fibroblast growth factor-23 and carotid artery intima media thickness in chronic kidney disease. *Clin Lab*. 2015;61(8):1061–1070.
- Hu X, Ma X, Luo Y, et al. Contribution of fibroblast growth factor 23 to Framingham risk score for identifying subclinical atherosclerosis in Chinese men. *Nutr Metab Cardiovasc Dis*. 2017;27(2):147–153. doi:10.1016/j.numecd.2016.11.009
- Fyfe-johnson AL, Alonso A, Selvin E, et al. Serum fibroblast growth factor-23 and incident hypertension: the atherosclerosis risk in communities study. *J Hypertens*. 2016;34(7):1266–1272. doi:10.1097/HJH.0000000000000936
- Mirza MA, Alsiö J, Hammarstedt A, et al. Circulating fibroblast growth factor-23 is associated with fat mass and dyslipidemia in two independent cohorts of elderly individuals. *Arterioscler Thromb Vasc Biol*. 2011;31(1):219–227. doi:10.1161/ATVBAHA.110.214619
- Hanks LJ, Casazza K, Judd SE, Jenny NS, Gutiérrez OM. Associations of fibroblast growth factor-23 with markers of inflammation, insulin resistance and obesity in adults. *PLoS One*. 2015;10(3):e0122885. doi:10.1371/journal.pone.0122885
- Sharaf El Din U, Salem MM, Abdulazim DO. FGF23 and inflammation. *World J Nephrol*. 2017;6(1):57–58. doi:10.5527/wjn.v6.i1.57
- Prié D, Friedlander G. Reciprocal control of 1, 25-dihydroxyvitamin D and FGF23 formation involving the FGF23/Klotho system. *Clin J Am Soc Nephrol*. 2010;5(9):1717–1722. doi:10.2215/CJN.02680310
- Donate-Correa J, Martín-Núñez E, Hernández-Carballo C, et al. Fibroblast growth factor 23 expression in human calcified vascular tissues. *Aging*. 2019;11(18):7899–7913. doi:10.18632/aging.102297
- De Jong MA, Eisenga MF, van Ballegooijen AJ, et al. Fibroblast growth factor 23 and new-onset chronic kidney disease in the general population: the Prevention of Renal and Vascular Endstage Disease (PREVEND) study. *Nephrol Dial Transplant*. 2021;36(1):121–128. doi:10.1093/ndt/gfz266
- Fliser D, Kollerits B, Neyer U, et al. Fibroblast growth factor 23 (FGF23) predicts progression of chronic kidney disease: the Mild to Moderate Kidney Disease (MMKD) Study. *J Am Soc Nephrol*. 2007;18(9):2600–2608. doi:10.1681/ASN.2006080936
- Falcini F, Masi L, Capannini S, et al. Fibroblast growth factor (FGF23) serum levels in juvenile onset systemic lupus erythematosus (JSLE): a possible link with renal involvement. *Arthritis Rheum*. 2010;2:62.
- Resende AL, Elias RM, Wolf M, et al. Serum levels of fibroblast growth factor 23 are elevated in patients with active lupus nephritis. *Cytokine*. 2017;91:124–127. doi:10.1016/j.cyt.2016.12.022
- Zhou M, Li S, Pathak JL. Pro-inflammatory cytokines and osteocytes. *Curr Osteoporos Rep*. 2019;17(3):97–104. doi:10.1007/s11914-019-00507-z

35. Rhee Y, Bivi N, Farrow E, et al. Parathyroid hormone receptor signaling in osteocytes increases the expression of fibroblast growth factor-23 in vitro and in vivo. *Bone*. 2011;49(4):636–643. doi:10.1016/j.bone.2011.06.025
36. Burnett SAM, Gunawardene SC, Bringham FR, Juppner H, Lee H, Finkelstein JS. Regulation of C-terminal and intact FGF-23 by dietary phosphate in men and women. *J Bone Miner Res*. 2006;21(8):1187–1196. doi:10.1359/jbmr.060507
37. van Ballegooijen AJ, Rhee EP, Elmariah S, de Boer IH, Kestenbaum B. Renal clearance of mineral metabolism biomarkers. *J Am Soc Nephrol*. 2016;27(2):392–397. doi:10.1681/ASN.2014121253
38. Lu X, Hu MC. Klotho/FGF23 axis in chronic kidney disease and cardiovascular disease. *Kidney Dis*. 2017;3(1):15–23. doi:10.1159/000452880
39. Kirchl C, Husar-Memmer E, Rappersberger K, Thaler K, Fritsch-Stork R. Type I Interferon as cardiovascular risk factor in systemic and cutaneous lupus erythematosus: a systematic review. *Autoimmun Rev*. 2021;102794. doi:10.1016/j.autrev.2021.102794
40. Petri M, Orbai AM, Alarcón GS, et al. Derivation and validation of the Systemic Lupus International Collaborating Clinics classification criteria for systemic lupus erythematosus. *Arthritis Rheum*. 2012;64(8):2677–2686. doi:10.1002/art.34473
41. Hahn BH, McMahon MA, Wilkinson A, et al. American College of Rheumatology guidelines for screening, treatment, and management of lupus nephritis. *Arthritis Care Res*. 2012;64(6):797–808. doi:10.1002/acr.21664
42. Bombardier C, Gladman DD, Urowitz MB, et al. Derivation of the SLEDAI. A disease activity index for lupus patients. *Arthritis Rheum*. 1992;35(6):630–640. doi:10.1002/art.1780350606
43. Gladman D, Ginzler E, Goldsmith C, et al. The development and initial validation of the Systemic Lupus International Collaborating Clinics/American College of Rheumatology damage index for systemic lupus erythematosus. *Arthritis Rheum*. 1996;39(3):363–369. doi:10.1002/art.1780390303
44. Simons PC, Algra A, Bots ML, Grobbee DE, van der Graaf Y. Common carotid intima-media thickness and arterial stiffness: indicators of cardiovascular risk in high-risk patients the SMART study (Second Manifestations of ARterial Disease). *Circulation*. 1999;100(9):951–957. doi:10.1161/01.CIR.100.9.951
45. Uthoff H, Staub D, Socrates T, et al. PROCAM-, FRAMINGHAM-, SCORE-and SMART-risk score for predicting cardiovascular morbidity and mortality in patients with overt atherosclerosis. *Vasa*. 2010;39(4):325–333. doi:10.1024/0301-1526/a000057
46. Burtis CA, Ashwood ER, Bruns DE. *Tietz Textbook of Clinical Chemistry and Molecular Diagnostics-e-Book*. Elsevier Health Sciences; 2012.
47. Levey AS, Stevens LA. Estimating GFR using the CKD epidemiology collaboration (CKD-EPI) creatinine equation: more accurate GFR estimates, lower CKD prevalence estimates, and better risk predictions. *Am J Kidney Dis*. 2010;55(4):622. doi:10.1053/j.ajkd.2010.02.337
48. Stein JH, Korcarz CE, Hurst RT, et al. Use of carotid ultrasound to identify subclinical vascular disease and evaluate cardiovascular disease risk: a consensus statement from the American Society of Echocardiography Carotid Intima-Media Thickness Task Force endorsed by the Society for Vascular Medicine. *J Am Soc Echocardiogr*. 2008;21(2):93–111.
49. Johri AM, Nambi V, Naqvi TZ, et al. Recommendations for the assessment of carotid arterial plaque by ultrasound for the characterization of atherosclerosis and evaluation of cardiovascular risk: from the American Society of Echocardiography. *J Am Soc Echocardiogr*. 2020;33(8):917–933. doi:10.1016/j.echo.2020.04.021
50. Rivera LMS, Ríos LV, Díaz CH, Villaseñor CP. Usefulness of measuring common carotid intima-media thickness: ultrasound diagnosis of sub-clinical atherosclerosis in rheumatic diseases. A literature review. *Rev Colomb Reumatol*. 2016;23(2):92–101.
51. Staub D, Meyerhans A, Bundi B, Schmid HP, Frauchiger B. Prediction of cardiovascular morbidity and mortality: comparison of the internal carotid artery resistive index with the common carotid artery intima-media thickness. *Stroke*. 2006;37(3):800–805. doi:10.1161/01.STR.0000202589.47401.c6
52. Martínez-Martínez MU, Martínez-Martínez MU, Mandeville P, Llamazares-Azuara L, Abud-Mendoza C. CKD-EPI is the most reliable equation to estimate renal function in patients with systemic lupus erythematosus. *Nefrología*. 2013;33(1):99–106.
53. Belibou C, Ancuța C, Filoș C, Chiriac R. Carotid intima-media thickness and plaque as surrogate biomarkers of atherosclerosis among consecutive women with systemic lupus erythematosus. *Rom J Morphol Embryol*. 2012;53(1):29–34.
54. Telles RW, Lanna CC, Sousa AJ, et al. Progression of carotid atherosclerosis in patients with systemic lupus erythematosus. *Clin Rheumatol*. 2013;32(9):1293–1300. doi:10.1007/s10067-013-2264-9
55. Ajeganova S, Gustafsson T, Lindberg L, Hafström I, Frostegård J. Similar progression of carotid intima-media thickness in 7-year surveillance of patients with mild SLE and controls, but this progression is still promoted by dyslipidaemia, lower HDL levels, hypertension, history of lupus nephritis and a higher prednisolone usage in patients. *Lupus Sci Med*. 2020;7(1):e000362.
56. Kao AH, Lertratanakul A, Elliott JR, et al. Relation of carotid intima-media thickness and plaque with incident cardiovascular events in women with systemic lupus erythematosus. *Am J Cardiol*. 2013;112(7):1025–1032. doi:10.1016/j.amjcard.2013.05.040
57. Salonen JT, Salonen R. Ultrasound B-mode imaging in observational studies of atherosclerotic progression. *Circulation*. 1993;87(3 Suppl):II56–65.
58. Touboul P, Hennerici M, Meairs S, Adams H, Amarencu P, Bornstein N. Mannheim carotid intima-media thickness consensus (2004–2006). *Cerebrovasc Dis*. 2007;23(1):75–80. doi:10.1159/000097034
59. Finn AV, Kolodgie FD, Virmani R. Correlation between carotid intimal/medial thickness and atherosclerosis: a point of view from pathology. *Arterioscler Thromb Vasc Biol*. 2010;30(2):177–181. doi:10.1161/ATVBAHA.108.173609
60. Inaba Y, Chen JA, Bergmann SR. Carotid plaque, compared with carotid intima-media thickness, more accurately predicts coronary artery disease events: a meta-analysis. *Atherosclerosis*. 2012;220(1):128–133. doi:10.1016/j.atherosclerosis.2011.06.044
61. Roman MJ, Crow MK, Lockshin MD, et al. Rate and determinants of progression of atherosclerosis in systemic lupus erythematosus. *Arthritis Rheum*. 2007;56(10):3412–3419. doi:10.1002/art.22924
62. Frauchiger B, Schmid HP, Roedel C, Moosmann P, Staub D. Comparison of carotid arterial resistive indices with intima-media thickness as sonographic markers of atherosclerosis. *Stroke*. 2001;32(4):836–841. doi:10.1161/01.STR.32.4.836
63. Gustafsson JT, Herlitz Lindberg M, Gunnarsson I, et al. Excess atherosclerosis in systemic lupus erythematosus,—a matter of renal involvement: case control study of 281 SLE patients and 281 individually matched population controls. *PLoS One*. 2017;12(4):e0174572. doi:10.1371/journal.pone.0174572
64. Hermansen M-L, Lindhardsen J, Torp-Pedersen C, Faurschou M, Jacobsen S. The risk of cardiovascular morbidity and cardiovascular mortality in systemic lupus erythematosus and lupus nephritis: a Danish nationwide population-based cohort study. *Rheumatology*. 2017;56(5):709–715.

65. Hermansen M-L, Sandholt B, Fuchs A, et al. Atherosclerosis and renal disease involvement in patients with systemic lupus erythematosus: a cross-sectional cohort study. *Rheumatology*. 2018;57(11):1964–1971. doi:10.1093/rheumatology/key201
66. Falaschi F, Ravelli A, Martignoni A, et al. Nephrotic-range proteinuria, the major risk factor for early atherosclerosis in juvenile-onset systemic lupus erythematosus. *Arthritis Rheum*. 2000;43(6):1405–1409. doi:10.1002/1529-0131(200006)43:6<1405::AID-ANR26>3.0.CO;2-V
67. Souza A, Hatta FS, Miranda JF, Sato EI. Atherosclerotic plaque in carotid arteries in systemic lupus erythematosus: frequency and associated risk factors. *Sao Paulo Med J*. 2005;123(3):137–142. doi:10.1590/S1516-31802005000300010
68. Ahmad Y, Shelmerdine J, Bodill H, et al. Subclinical atherosclerosis in systemic lupus erythematosus (SLE): the relative contribution of classic risk factors and the lupus phenotype. *Rheumatology*. 2007;46(6):983–988. doi:10.1093/rheumatology/kem002
69. Skaggs BJ, Hahn BH, McMahon M. Accelerated atherosclerosis in patients with SLE—mechanisms and management. *Nat Rev Rheumatol*. 2012;8(4):214–223. doi:10.1038/nrrheum.2012.14
70. O'Neill SG, Isenberg DA, Rahman A. Could antibodies to C-reactive protein link inflammation and cardiovascular disease in patients with systemic lupus erythematosus? *Ann Rheum Dis*. 2007;66(8):989–991. doi:10.1136/ard.2007.073312
71. Manzi S. Systemic lupus erythematosus: a model for atherogenesis? *Rheumatology*. 2000;39(4):353–359. doi:10.1093/rheumatology/39.4.353
72. Kamen DL, Cooper GS, Bouali H, Shaftman SR, Hollis BW, Gilkeson GS. Vitamin D deficiency in systemic lupus erythematosus. *Autoimmun Rev*. 2006;5(2):114–117. doi:10.1016/j.autrev.2005.05.009
73. Kamen DL. Vitamin D in lupus: new kid on the block? *Bull NYU Hosp Jt Dis*. 2010;68(3):218–222.
74. Ruiz-Irastorza G, Egurbide M, Olivares N, Martinez-Berriotxo A, Aguirre C. Vitamin D deficiency in systemic lupus erythematosus: prevalence, predictors and clinical consequences. *Rheumatology*. 2008;47(6):920–923. doi:10.1093/rheumatology/ken121
75. Oh R. Vitamin D insufficiency as a cause of hyperparathyroidism. *Am Fam Physician*. 2005;71(1):46–49.
76. Lips P. Vitamin D deficiency and secondary hyperparathyroidism in the elderly: consequences for bone loss and fractures and therapeutic implications. *Endocrine Rev*. 2001;22(4):477–501. doi:10.1210/edrv.22.4.0437
77. Currie G, Delles C. Proteinuria and its relation to cardiovascular disease. *Int J Nephrol Renovasc Dis*. 2013;7:13–24.
78. Perkovic V, Verdon C, Ninomiya T, et al. The relationship between proteinuria and coronary risk: a systematic review and meta-analysis. *PLoS Med*. 2008;5(10):e207. doi:10.1371/journal.pmed.0050207
79. Sung JK, Kim J-Y, Youn YJ, et al. Urine albumin creatinine ratio is associated with carotid atherosclerosis in a community based cohort: atherosclerosis risk of rural area in Korean general population study. *J Cardiovasc Ultrasound*. 2010;18(4):134–138. doi:10.4250/jcu.2010.18.4.134
80. Jorgensen L, Jenssen T, Johnsen SH, et al. Albuminuria as risk factor for initiation and progression of carotid atherosclerosis in non-diabetic persons: the Tromso Study. *Eur Heart J*. 2007;28(3):363. doi:10.1093/eurheartj/ehl394
81. Xu R, Cai H, Fan Z, Wan Y, Gao X. The change in kidney function was associated with carotid artery plaque in a community-based population: a cohort study. *Nutr Metab Cardiovasc Dis*. 2021;31(1):119–126. doi:10.1016/j.numecd.2020.08.016
82. Kajitani N, Uchida HA, Suminoe I, et al. Chronic kidney disease is associated with carotid atherosclerosis and symptomatic ischaemic stroke. *J Int Med Res*. 2018;46(9):3873–3883. doi:10.1177/0300060518781619
83. Ohara T, Kokubo Y, Toyoda K, et al. Impact of chronic kidney disease on carotid atherosclerosis according to blood pressure category: the Suita study. *Stroke*. 2013;44(12):3537–3539. doi:10.1161/STROKEAHA.113.002957
84. Jankowski J, Floege J, Fliser D, Böhm M, Marx N. Cardiovascular disease in chronic kidney disease: pathophysiological insights and therapeutic options. *Circulation*. 2021;143(11):1157–1172. doi:10.1161/CIRCULATIONAHA.120.050686
85. Bruce I. 'Not only ... but also': factors that contribute to accelerated atherosclerosis and premature coronary heart disease in systemic lupus erythematosus. *Rheumatology*. 2005;44(12):1492–1502. doi:10.1093/rheumatology/kei142
86. Ng M, Celermajer D. *Glucocorticoid Treatment and Cardiovascular Disease*. BMJ Publishing Group Ltd; 2004.
87. Donate-Correa J, Muros-de-fuentes M, Mora-Fernández C, Navarro-González JF. FGF23/Klotho axis: phosphorus, mineral metabolism and beyond. *Cytokine Growth Factor Rev*. 2012;23(1–2):37–46. doi:10.1016/j.cytogfr.2012.01.004
88. Shuto E, Taketani Y, Tanaka R, et al. Dietary phosphorus acutely impairs endothelial function. *J Am Soc Nephrol*. 2009;20(7):1504–1512. doi:10.1681/ASN.2008101106
89. Onufrak SJ, Bellasi A, Shaw LJ, et al. Phosphorus levels are associated with subclinical atherosclerosis in the general population. *Atherosclerosis*. 2008;199(2):424–431. doi:10.1016/j.atherosclerosis.2007.11.004
90. Itkonen ST, Karp HJ, Kemi VE, et al. Associations among total and food additive phosphorus intake and carotid intima-media thickness—a cross-sectional study in a middle-aged population in Southern Finland. *Nutr J*. 2013;12(1):1–10. doi:10.1186/1475-2891-12-94
91. Sheridan K, Logomarsino JV. Effects of serum phosphorus on vascular calcification in a healthy, adult population: a systematic review. *J Vasc Nurs*. 2017;35(3):157–169. doi:10.1016/j.jvn.2017.01.003
92. Foley RN. Phosphate levels and cardiovascular disease in the general population. *Clin J Am Soc Nephrol*. 2009;4(6):1136–1139. doi:10.2215/CJN.01660309
93. Osuka S, Razaque MS. Can features of phosphate toxicity appear in normophosphatemia? *J Bone Miner Metab*. 2012;30(1):10–18. doi:10.1007/s00774-011-0343-z
94. Bai W, Li J, Liu J. Serum phosphorus, cardiovascular and all-cause mortality in the general population: a meta-analysis. *Clinica Chimica Acta*. 2016;461:76–82. doi:10.1016/j.cca.2016.07.020
95. Disthabanchong S. Phosphate and cardiovascular disease beyond chronic kidney disease and vascular calcification. *Int J Nephrol*. 2018;1:3162806.
96. Zhou C, Shi Z, Ouyang N, Ruan X. Hyperphosphatemia and cardiovascular disease. *Front Cell Dev Biol*. 2021;9:644363. doi:10.3389/fcell.2021.644363
97. Ritter CS, Slatopolsky E. Phosphate toxicity in CKD: the killer among us. *Clin J Am Soc Nephrol*. 2016;11(6):1088–1100. doi:10.2215/CJN.11901115
98. Ellam TJ, Chico TJ. Phosphate: the new cholesterol? The role of the phosphate axis in non-uremic vascular disease. *Atherosclerosis*. 2012;220(2):310–318. doi:10.1016/j.atherosclerosis.2011.09.002
99. Keshk W, Soliman N, Esheba N. Study of some biomarkers changes in patients with lupus nephritis and their correlation with disease activity and progression. *Bull Egypt Soc Physiol Sci*. 2013;33(1):17–34.



100. Cunningham SE, Czaya BE, Faul C. Elevated phosphate levels induce markers of systemic inflammation and anemia in murine hepatocytes. *THE FASEB J.* 2020;34(S1):1.
101. Yamada S, Tokumoto M, Tatsumoto N, et al. Phosphate overload directly induces systemic inflammation and malnutrition as well as vascular calcification in uremia. *Am J Physiol Renal Physiol.* 2014;306(12):F1418–F1428. doi:10.1152/ajprenal.00633.2013
102. Czaya B, Richter B, Yanucil C, Campos I, Heitman K, Faul C. Hyperphosphatemia contributes to inflammation and iron dysregulation in models of normal and impaired renal function. *Blood.* 2019;134(Suppl 1):2238. doi:10.1182/blood-2019-122144
103. Voelkl J, Egli-Spichtig D, Alesutan I, Wagner CA. Inflammation: a putative link between phosphate metabolism and cardiovascular disease. *Clin Sci.* 2021;135(1):201–227.
104. Erem S, Razzaque MS. Dietary phosphate toxicity: an emerging global health concern. *Histochem Cell Biol.* 2018;150(6):711–719. doi:10.1007/s00418-018-1711-8
105. Gutiérrez OM, Anderson C, Isakova T, et al. Low socioeconomic status associates with higher serum phosphate irrespective of race. *J Am Soc Nephrol.* 2010;21(11):1953–1960. doi:10.1681/ASN.2010020221
106. Wöhrle S, Bonny O, Beluch N, et al. FGF receptors control vitamin D and phosphate homeostasis by mediating renal FGF-23 signaling and regulating FGF-23 expression in bone. *J Bone Miner Res.* 2011;26(10):2486–2497. doi:10.1002/jbmr.478
107. Quarles LD. FGF-23 and  $\alpha$ -klotho co-dependent and independent functions. *Curr Opin Nephrol Hypertens.* 2019;28(1):16–25. doi:10.1097/MNH.0000000000000467
108. Gutierrez O, Isakova T, Rhee E, et al. Fibroblast growth factor-23 mitigates hyperphosphatemia but accentuates calcitriol deficiency in chronic kidney disease. *J Am Soc Nephrol.* 2005;16(7):2205–2215. doi:10.1681/ASN.2005010052
109. Koh N, Fujimori T, Nishiguchi S, et al. Severely reduced production of klotho in human chronic renal failure kidney. *Biochem Biophys Res Commun.* 2001;280(4):1015–1020. doi:10.1006/bbrc.2000.4226
110. Kritmetapak K, Losbanos L, Berent TE, et al. Hyperphosphatemia with elevated serum PTH and FGF23, reduced 1, 25 (OH) 2 D and normal FGF7 concentrations characterize patients with CKD. *BMC Nephrol.* 2021;22(1):1–8. doi:10.1186/s12882-021-02311-3
111. Vogt I, Haffner D, Leifheit-Nestler M. FGF23 and phosphate-cardiovascular toxins in CKD. *Toxins.* 2019;11(11):647. doi:10.3390/toxins11110647
112. Martín-Núñez E, Donate-Correa J, López-Castillo A, et al. Soluble levels and endogenous vascular gene expression of KLOTHO are related to inflammation in human atherosclerotic disease. *Clin Sci.* 2017;131(21):2601–2609. doi:10.1042/CS20171242
113. Rodríguez-Ortiz ME, Díaz-Tocados JM, Muñoz-Castañeda JR, et al. Inflammation both increases and causes resistance to FGF23 in normal and uremic rats. *Clin Sci.* 2020;134(1):15–32. doi:10.1042/CS20190779
114. Forster RE, Jurutka PW, Hsieh J-C, et al. Vitamin D receptor controls expression of the anti-aging klotho gene in mouse and human renal cells. *Biochem Biophys Res Commun.* 2011;414(3):557–562. doi:10.1016/j.bbrc.2011.09.117
115. Haussler MR, Whitfield GK, Haussler CA, et al. 1, 25-Dihydroxyvitamin D and Klotho: a tale of two renal hormones coming of age. *Vitam Horm.* 2016;100:165–230.
116. Lavi-Moshayoff V, Wasserman G, Meir T, Silver J, Naveh-Many T. PTH increases FGF23 gene expression and mediates the high-FGF23 levels of experimental kidney failure: a bone parathyroid feedback loop. *Am J Physiol Renal Physiol.* 2010;299(4):F882–F889. doi:10.1152/ajprenal.00360.2010
117. Bae J, Lee Y-H, Kang ES, Cha B-S, Lee B-W. Proteinuria is associated with carotid artery atherosclerosis in non-albuminuric type 2 diabetes: a cross-sectional study. *J Clin Med.* 2020;9(1):136. doi:10.3390/jcm9010136
118. Furtner M, Kiechl S, Mair A, et al. Urinary albumin excretion is independently associated with carotid and femoral artery atherosclerosis in the general population. *Europ Heart J.* 2005;26(3):279–287. doi:10.1093/eurheartj/ehi014
119. Oh K-H, Park SK, Park HC, et al. KNOW-CKD (KoreaN cohort study for Outcome in patients With Chronic Kidney Disease): design and methods. *BMC Nephrol.* 2014;15(1):1–9. doi:10.1186/1471-2369-15-80
120. De Seigneux S, Courbebaisse M, Rutkowski JM, et al. Proteinuria increases plasma phosphate by altering its tubular handling. *J Am Soc Nephrol.* 2015;26(7):1608–1618. doi:10.1681/ASN.2014010104
121. Jung JY, Ro H, Chang JH, et al. Mediation of the relationship between proteinuria and serum phosphate: insight from the KNOW-CKD study. *PLoS One.* 2020;15(6):e0235077. doi:10.1371/journal.pone.0235077
122. Delitsikou V, Jarad G, Rajaram RD, et al. Klotho regulation by albuminuria is dependent on ATF3 and endoplasmic reticulum stress. *FASEB J.* 2020;34(2):2087–2104. doi:10.1096/fj.201900893R
123. de Seigneux S, Delitsikou V, Martin P-Y. The KNOW-CKD study: evidence for a link between proteinuria and alterations of mineral metabolism. *Nephrol Dial Transplant.* 2020;35(3):382–385. doi:10.1093/ndt/gfz083
124. Leal LKAM, Lima LA, de Aquino PEA, et al. Vitamin D (VD3) antioxidative and anti-inflammatory activities: peripheral and central effects. *Europ J Pharmacol.* 2020;879:173099. doi:10.1016/j.ejphar.2020.173099
125. Zoccali C, Ruggenti P, Perna A, et al. Phosphate may promote CKD progression and attenuate renoprotective effect of ACE inhibition. *J Am Soc Nephrol.* 2011;22(10):1923–1930. doi:10.1681/ASN.2011020175
126. Di Iorio BR, Bellizzi V, Bellasi A, et al. Phosphate attenuates the anti-proteinuric effect of very low-protein diet in CKD patients. *Nephrol Dial Transplant.* 2013;28(3):632–640. doi:10.1093/ndt/gfs477
127. Shalhoub V, Shatzen EM, Ward SC, et al. FGF23 neutralization improves chronic kidney disease-associated hyperparathyroidism yet increases mortality. *J Clin Invest.* 2012;122(7):2543–2553. doi:10.1172/JCI61405