



Published in final edited form as:

*Dig Dis Sci.* 2022 April ; 67(4): 1337–1344. doi:10.1007/s10620-021-06949-9.

## History of Diverticulitis and Risk of Incident Cardiovascular Disease in Men: A cohort study

Idy Tam<sup>1,\*</sup>, Po-Hong Liu<sup>2,3,\*</sup>, Wenjie Ma<sup>2,4</sup>, Yin Cao<sup>2,4,5</sup>, Manol Jovani<sup>2,4</sup>, Kana Wu<sup>6</sup>, Eric B. Rimm<sup>6,7,8</sup>, Lisa L. Strate<sup>9,†</sup>, Edward L. Giovannucci<sup>6,7,8,†</sup>, Andrew T. Chan<sup>2,4,8,10,†</sup>

<sup>1</sup>Tufts University School of Medicine, Boston, MA

<sup>2</sup>Clinical and Translational Epidemiology Unit, Massachusetts General Hospital and Harvard Medical School, Boston, MA

<sup>3</sup>Digestive and Liver Diseases Division, UT Southwestern Medical Center, Dallas, TX

<sup>4</sup>Division of Gastroenterology, Massachusetts General Hospital and Harvard Medical School, Boston, MA

<sup>5</sup>Division of Public Health Sciences, Department of Surgery, Washington University School of Medicine, St. Louis, MO

<sup>6</sup>Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA

<sup>7</sup>Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA

<sup>8</sup>Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA

<sup>9</sup>Division of Gastroenterology, University of Washington School of Medicine, Harborview Medical Center, Seattle, WA

<sup>10</sup>Broad Institute of MIT and Harvard, Cambridge, MA

### Abstract

**Background:** Diverticulitis and cardiovascular disease (CVD) are two highly prevalent disorders sharing common risk factors which are hypothesized to have an inflammatory basis.

**Aims:** To examine the association between history of diverticulitis and risk of incident CVD.

**Methods:** We conducted a prospective cohort study of 43,904 men aged 40 to 75 years without a history of CVD (fatal or nonfatal myocardial infarction and stroke) at enrollment who were followed up from 1986 to 2012 in the Health Professionals Follow-Up Study. Lifestyle factors, dietary intake, and disease information were self-reported biennially or quadrennially. Incident diverticulitis and CVD were confirmed by review of medical records. We used Cox proportional

---

**Correspondence:** Dr. Andrew T. Chan, Clinical and Translational Epidemiology Unit, Massachusetts General Hospital, 55 Fruit Street, Boston, MA 02114; Email: achan@mgh.harvard.edu; Tel: 617-726-7802; Fax: 617-726-3673.

\*Drs. Tam and Liu contributed equally.

†Drs. Strate, Giovannucci, and Chan contributed equally.

**Disclosures:** A.T.C. previously served as a consultant for Bayer Pharma AG, Janssen Pharmaceuticals and Pfizer Inc. for work unrelated to the topic of this manuscript. This study was not funded by Bayer Pharma AG, Janssen Pharmaceuticals or Pfizer Inc.

hazard models to calculate age- and multivariable-adjusted hazard ratios (HR) and 95% confidence intervals (CI) of incident CVD. We conducted a stratified analysis according to the presence or absence of CVD risk factors (smoking, hypertension, hyperlipidemia, and diabetes).

**Results:** We identified 3,848 incident cases of CVD during 856,319 person-years of follow-up. Men with diverticulitis had higher incidence of CVD (727 cases per 100,000 person-years) compared to men without diverticulitis [446 cases per 100,000 person-years, multivariate HR of 1.35 (95% CI: 1.07-1.70)]. The association of diverticulitis and subsequent CVD appeared more evident among men without known CVD risk factors (HR: 4.06, 95% CI: 2.04-8.08) compared to those with 1 or more CVD risk factors (HR: 1.27, 95% CI: 0.98-.63).

**Conclusions:** Diverticulitis may be an independent risk factor of incident CVD, suggesting possible common etiopathogenic mechanisms. Diagnosis of diverticulitis underscores the importance of preventive measures to reduce future CVD.

### Keywords

Diverticulitis; cardiovascular disease prevention; cardiovascular disease risk factors

---

## INTRODUCTION

Diverticular disease is a highly prevalent disease in developed countries. It is the third most common gastrointestinal diagnosis on hospital discharge in the US, accounting for \$3.6 billion in health care costs each year.<sup>1, 2</sup> The prevalence of diverticulosis increases with age, affecting approximately 60% of individuals 70 years of age.<sup>3</sup> Among patients with diverticulosis, an estimated 4% will develop acute diverticulitis.<sup>4</sup> Diverticulitis is inflammation and micro-perforation of one or more diverticula in the colon that may progress to complications including abscess, fistula, or peritonitis. Risk factors for diverticulitis include low fiber intake, obesity, hypertension, smoking, and physical inactivity.<sup>5</sup> These risk factors also increase the risk of cardiovascular disease (CVD) contributing to the hypothesis of shared or exacerbated risk between these two diseases.<sup>6-11</sup> In addition, diverticulitis may arise in a background of low-grade, chronic inflammation potentially mediated by changes in the microbiota composition which could predispose to the development of CVD.<sup>12-15</sup>

Several studies have demonstrated an association between diverticular disease and risk of CVD.<sup>16-18</sup> One large retrospective cohort study reported a 25% increased risk for acute coronary syndrome in patients with a history of diverticulitis.<sup>18</sup> However, existing studies are based on medical or insurance registries which utilize ICD codes for identification of exposures and outcomes, and therefore, were not able to differentiate diverticulitis, diverticular bleeding, and uncomplicated diverticulosis or account for important potential risk factors such as lifestyle and dietary factors. Therefore, we performed this study to analyze the association between a history of diverticulitis and risk of incident CVD in a large prospective cohort of US men in the Health Professional Follow-up Study (HPFS) in which we collected detailed information on risk factors for both CVD and diverticulitis and confirmed incident cases of CVD and diverticulitis through medical record review.

## METHODS

### Study population

The HPFS is a large ongoing prospective cohort study of 51,529 US male dentists, veterinarians, pharmacists, optometrists, osteopathic physicians, and podiatrists aged 40-75 years at baseline in 1986. The cohort members report on demographics, lifestyle and medical history biennially and dietary information every 4 years via self-administered questionnaires by mail. The questionnaires for each cycle can be found in the following website: <https://sites.sph.harvard.edu/hpfs/hpfs-questionnaires/>. The biennial follow-up rate averages 94%.<sup>19</sup> This study was approved by the institutional review board at the Harvard T.H. Chan School of Public Health.

We excluded participants who reported CVD, malignancy (except for non-melanoma skin cancer), or non-plausible calorie intake (less than 800 or greater than 4,200 kcal daily) at baseline. The final analysis included 43,904 men.

### Assessment of diverticulitis

History of diverticulitis was based on responses from supplemental diverticular disease questionnaires, which were sent to men who reported newly diagnosed diverticulitis or diverticulosis on biennial questionnaires since 1990. Diverticulitis was defined as abdominal pain attributed to diverticular disease and one of the following criteria: (1) complicated by perforation, abscess, fistula or obstruction; (2) requiring hospitalization, antibiotics or surgery; or (3) pain categorized as severe or acute or abdominal pain presenting with fever, requiring antibiotics or confirmed using abdominal CT. In 2006, supplemental questionnaires were updated to further assess diverticular complications including diverticular bleeding; uncomplicated diverticulitis; diverticular abscess, obstruction, perforation, and fistula. We have previously validated these methods and used these case definitions in prior analyses.<sup>6, 10</sup>

### Assessment of cardiovascular disease

The primary endpoint of this study was incident CVD, defined as fatal myocardial infarction, nonfatal myocardial infarction, and stroke. For participants who reported newly diagnosed CVD or stroke, permission was requested for medical records. All medical records were reviewed by physicians who were blind to the questionnaire data to confirm the diagnosis. Nonfatal myocardial infarction is defined according to World Health Organization criteria, which required typical symptoms plus either diagnostic electrocardiographic findings or elevated cardiac enzyme concentrations. We also considered myocardial infarctions requiring hospital admission and corroborated by phone interview or letter only as probable cases. Stroke was confirmed by use of the National Survey of Stroke criteria, requiring a constellation of neurological deficits, sudden or rapid onset, and duration of  $\geq$  24 hours or until death. When medical records were not available, interviews or letters confirmed coronary and stroke events, which were designated as probable cases. Deaths were identified from the state vital statistics records and the National Death Index or were reported by families and the postal system.<sup>20-24</sup> The cause listed on the death certificate was not considered sufficient to confirm a coronary or stroke death. Underlying

and contributing causes of deaths were confirmed by study physicians via review of medical records or autopsy reports with permission of family members.

### Assessment of covariables

Medical history and lifestyle factors were assessed biennially. Covariables that were considered included age, body mass index (BMI), smoking status, alcohol intake, physical activity, and regular use of aspirin, non-steroidal anti-inflammatory drugs (NSAID), acetaminophen, cholesterol-lowering medication, and anti-hypertensive medication (e.g. thiazide diuretics, calcium-channel blockers, beta-blockers, other antihypertensives). Self-reported hypertension, hypercholesterolemia, diabetes mellitus, family history of myocardial infarction and diabetes mellitus were also assessed biennially. Dietary factors including dietary fiber, total energy intake, unprocessed and processed red meat intake, and the Alternate Healthy Eating Index (AHEI) were derived from validated food frequency questionnaires administered every four years.<sup>25</sup>

### Statistical analysis

In our primary analysis, we examined the association between history of diverticulitis and risk of incident CVD. We used biennially updated reports of diverticulitis to define the exposure. We also defined recent history of diverticulitis as having diverticulitis within 4 years of incident CVD disease. Participants with diverticulitis at least 4 years prior to questionnaire cycle were defined as having remote histories of diverticulitis. Person-years were calculated from baseline (1986) until the development of CVD disease, death, or end of follow up (January 2012), whichever came first. Participants who developed CVD and cancer during the follow-up were censored. Age-stratified cox proportional hazard models with time-varying covariables were used to estimate age-adjusted and multivariable-adjusted hazard ratios (HRs) and 95% confidence intervals (CIs) for risk of incident CVD. The following were included in multivariable models: age, BMI (6 categories, <18.5, 18.5-22.9, 23-24.9, 25-29.9, 30-34.9, and ≥35 kg/m<sup>2</sup>), physical activity (metabolic equivalents per week in quintiles), smoking status (pack-years in categories: never smokers, 1-4.9, 5-19.9, 20-39.9, ≥40), alcohol intake (nondrinker, 0.1-4.9, 5-14.9, 15-29.9, ≥30 g/d), total energy intake (in quintiles), AHEI (in quintiles), total red meat (in quintiles), dietary fiber (in quintiles), history of hypertension, hypercholesterolemia, and diabetes (all yes or no), family history of myocardial infarction and diabetes (both yes or no), and regular use of aspirin, NSAIDs, acetaminophen, lipid-lowering medication, and anti-hypertensive medication (yes or no). We updated history of diverticulitis and other covariables at 2- or 4-year intervals.

We also considered the possibility that participants who did not report history of diverticulitis and CVD or who reported fewer cardiovascular risk factors may have less interaction with health care. We therefore conducted a sensitivity analysis restricting to participants who had routine physical exams for health screening in the same questionnaire cycle. In addition, we conducted stratified analyses in patients with history of diverticulitis according to the presence or absence of the following CVD risk factors: hypertension, hypercholesterolemia, diabetes, or smoking. Statistical analyses were performed using SAS version 9.4 (SAS Institute Inc.). Two-sided *p*-values less than 0.05 were considered statistically significant.

## Participants involvement

Participants were not involved in the design of research questions or outcome measures. They were not involved in the design and implantation of the study. There are no plans to involve participants in the dissemination of results.

## RESULTS

The mean ages at baseline were 53.8 in men without diverticulitis and 52.4 in men with diverticulitis. During 856,319 person-years of follow-up, we documented 3,848 incident cases of CVD. Compared to men without a history of diverticulitis, men with history of diverticulitis tended to have CVD risk factors, such as higher total red meat intake, greater pack-years of smoking, less physical activity, and a higher prevalence of hypercholesterolemia. In addition, men with a history of diverticulitis used aspirin, NSAID, and acetaminophen more often (Table 1).

Participants with a history of diverticulitis had higher incidence of CVD (727 cases per 100,000 person-years) compared to men without diverticulitis (446 cases per 100,000 person-years). The multivariate HR after adjustment for age, BMI, smoking status, alcohol intake, physical activity, red meat intake, dietary fiber, and AHEI were 1.40 (95% CI: 1.11-1.76). Further adjustment for personal history of hypertension, hyperlipidemia, diabetes, family history of myocardial infarction and diabetes, and use of aspirin, non-aspirin NSAIDs, acetaminophen, lipid-lowering medication, and anti-hypertensive medication showed similar results (HR: 1.35, 95% CI: 1.07-1.70, Table 2).

As an exploratory analysis, we evaluated the influence of a recent (within 4 years) or remote (greater than 4 years) history of diverticulitis and risk of incident CVD. We observed trends toward increases in risk of CVD with both recent (HR: 1.45, 95% CI: 0.98-2.14) and remote (HR: 1.30, 95% CI: 0.97-1.73) history of diverticulitis, even though these did not reach statistical significance. Our main results remained unchanged in a sensitivity analysis limiting to participants who had received screening physical examinations (Table 3).

In the stratified analyses, we found that history of diverticulitis was associated with higher risk of incident CVD (HR: 4.06, 95% CI: 2.04-8.08) among participants without CVD risk factors (hypertension, hypercholesterolemia, diabetes, and smoking, Table 4), but not in those with CVD risk factors (HR: 1.27, 95% CI: 0.98-1.63,  $p$  for interaction = 0.12).

## DISCUSSION

In this large prospective cohort of men, we found that a history of diverticulitis was associated with a significant 34% increased risk of incident CVD, independent of conventional lifestyle and dietary risk factors. The trend towards an increased risk of incident CVD was observed with both recent history and remote history of diverticulitis, although this association did not reach statistical significance. Results were similar when we limited the exposure to those with regular contact with the health care system through regular physical examinations. In addition, the association between diverticulitis and risk of CVD was more evident in participants without traditional CVD risk factors.

Our findings are consistent with evidence from previous limited studies.<sup>14–18</sup> In particular, a large prospective cohort study of more than 77 thousand patients with incident diverticular disease identified in the Danish medical registries found that patients with diverticular disease had a modestly increased risk of acute myocardial infarction and venous thromboembolic events when compared to matched controls, with adjusted incidence rate ratios 1.11 (95% CI: 1.07-1.14) and 1.35 (95% CI: 1.30-1.43), respectively. These studies were limited by the use of diagnostic coding from medical and insurance registries, and therefore, were unable to identify and differentiate among diverticulitis, diverticular bleeding and uncomplicated diverticulosis. In addition, previous studies could not adjust for important covariables such as lifestyle factors and dietary intake. In contrast, in our study, we utilized a prospective cohort-based design with updated assessments of lifestyle and dietary factors for over 26 years of follow-up, which minimized the probability of selection and recall bias. In addition, we provided validated assessments of diverticulitis and CVD cases in comparison to existing studies and comprehensively controlled for multiple potential confounding factors.

Our results have biological plausibility. Chronic low-grade systemic inflammation due to gut microbiome dysbiosis may play a role in explaining the association between cardiovascular and diverticular disease. Studies have postulated that gut microbiota is a contributor to atherosclerosis, a precursor of CVD through modulation of host metabolism and inflammation, as well as various CVD risk factors such as metabolic syndrome, obesity, and diabetes.<sup>13, 14, 26–29</sup> Previous studies have shown that gut microbiota metabolizes choline, phosphatidylcholine, and L-carnitine to trimethylamine which is oxidized to trimethylamine-N-oxide, a proatherogenic metabolite.<sup>30, 31</sup> Gut microbiota composition of patients with atherosclerotic CVD have an abundance of species that have been also associated with inflammatory bowel disease.<sup>32</sup> Interestingly, studies that have examined gut microbiota composition in patients with diverticular disease found changes, such as abundance of *Enterobacteriaceae* and depletion of *Faecalibacterium prausnitzii*, that are similar to those in patients with atherosclerotic CVD.<sup>12</sup> Initial data suggest that patients with diverticular disease could harbor changes in gut microbiota composition which alter host immune defenses and cause dysfunction of mucosal barrier, leading to cross-reactivity and release of proinflammatory cytokines leading to systemic inflammation.<sup>13, 33, 34</sup> These changes contribute to the development of chronic low-grade systemic inflammation which can then contribute to atherosclerotic plaque formation.<sup>13, 31, 32</sup> Hence, dysbiosis in gut microbiota and an attendant chronic inflammatory state may be the link for the increased risk of CVD in patients with history of diverticulitis observed in this study. Indeed, a recent prospective analysis found that patients with a history of diverticulitis had a higher inflammatory potential of diet as well as higher circulating levels of C-reactive protein (CRP), interleukin 6 (IL6), markers of chronic inflammation.<sup>35, 36</sup> Therefore, patients with a history of diverticulitis appear to be in a state of chronic low-grade inflammation and may harbor an increased risk of CVD through this mechanism. The observed increased risk of CVD in men with either a recent or remote history of diverticulitis may be explained by this chronic inflammatory state rather than any acute events occurring at the time of a diverticulitis episode.

In addition, we found that stratification by several main components of the Framingham Risk Score<sup>37</sup> CVD risk factors modified the association between diverticulitis and CVD, although formal testing was not statistically significant ( $p$  for interaction = 0.12). Remarkably, risks of CVD associated with history of diverticulitis appeared to be more prominent in presumably healthier men, *i.e.*, those without hypertension, hypercholesterolemia, diabetes, and smoking history. Since CVD risk factors such as hypertension, hypercholesterolemia and diabetes are often asymptomatic and may go undiagnosed, it is possible we did not fully account for their potential confounding in our multivariate models. However, our results were largely unchanged when we limited our analysis to those participants with routine physical examinations in which these conditions are likely to be diagnosed through asymptomatic screening. An alternative explanation is that men with hypertension, hypercholesterolemia, diabetes, and smoking history had a higher absolute risk to CVD; therefore, the increased risk attributable to diverticulitis might be less evident compared to those men without CVD risk factors. A history of diverticulitis may be associated with gut microbial changes that also predispose to the development of obesity, diabetes, and hypertension, which in turn is associated with CVD. Thus, adjusting for these factors may attenuate a potential association. Finally, it is possible that patients who are diagnosed with hypertension, hypercholesterolemia, and diabetes might make healthier lifestyle modifications compared to men without these diagnoses, which could mitigate an association between diverticulitis and risk of CVD among patients with CVD risk factors. Further studies are warranted to explore these underlying mechanisms.

The main strengths of this study include large sample size and a prospective population-based design with update assessment of lifestyle and dietary factors for over 26 years of follow-up. The comprehensive lifestyle information available allowed us to finely adjust for potential confounders, anti-hypertensive, and lipid-lowering medications, and risk factors shared between diverticulitis and CVD. In addition, we were able to demonstrate a similar effect among participants with regular contact with health care, minimizing possible under-reporting of CVD risk factors. Finally, we were also able to differentiate diagnoses of diverticulitis from uncomplicated diverticulosis and diverticular bleeding as these diseases have distinct clinical manifestations and risk factors.

Nevertheless, our study has some potential limitations that need to be considered. First, it is possible that the association between history of diverticulitis and incident CVD can be confounded by a higher prevalence of CVD risk factors, such as low fiber intake, hypertension, smoking, and physical inactivity.<sup>6-11</sup> However, even after adjusting for these risk factors, the risk of developing CVD remained higher in individuals with history of diverticulitis, which suggests that shared risk factors may not fully account for the association that we observed. In addition, although we considered a comprehensive list of potential confounders, there is possibility of residual, unmeasured confounding. However, in our secondary analysis, we limited the analysis to those who have routine health physicals to minimize unmeasured confounding of asymptomatic diseases. Second, diagnoses of diverticulitis were self-reported and could potentially be misclassified. However, the validity of self-reported diverticulitis has been confirmed in this cohort and health professionals are more likely to provide an accurate account of medical information.<sup>6, 10</sup> Third, our study may lack generalizability due to the homogeneity of our population, particularly for women and

other racial or ethnic groups. Finally, due to the observational nature of this study, a causal relationship between diverticulitis and CVD could not be proven.

## CONCLUSION

Our results indicate that a history of diverticulitis is associated with an increased risk of incident CVD. These findings highlight the possibility that there are biological pathways which underlie both conditions. Clinically, this suggests that a diagnosis of diverticulitis may be an early marker of an underlying predisposition to CVD and underscores the importance of frequent monitoring of lifestyle factors that could influence not only risk of recurrent diverticulitis, but also other health conditions associated with chronic inflammation.

## Acknowledgement:

We would like to thank the participants and staff of the Health Professionals Follow-Up Study for their valuable contributions.

## Sources of Funding:

This work was supported by grants R01 DK101495, R01 DK084157, K24 DK098311, and R01 HL35464 and UM1 CA167552 from the National Institutes of Health.

## REFERENCES

1. Peery AF, Crockett SD, Barritt AS, Dellon ES, Eluri S, Gangarosa LM, Jensen ET, Lund JL, Pasricha S, Runge T, Schmidt M, Shaheen NJ and Sandler RS. Burden of Gastrointestinal, Liver, and Pancreatic Diseases in the United States. *Gastroenterology*. 2015;149:1731–1741 e3. [PubMed: 26327134]
2. Peery AF, Dellon ES, Lund J, Crockett SD, McGowan CE, Bulsiewicz WJ, Gangarosa LM, Thiny MT, Stizenberg K, Morgan DR, Ringel Y, Kim HP, DiBonaventura MD, Carroll CF, Allen JK, Cook SF, Sandler RS, Kappelman MD and Shaheen NJ. Burden of gastrointestinal disease in the United States: 2012 update. *Gastroenterology*. 2012;143:1179–1187 e3. [PubMed: 22885331]
3. Peery AF, Barrett PR, Park D, Rogers AJ, Galanko JA, Martin CF and Sandler RS. A high-fiber diet does not protect against asymptomatic diverticulosis. *Gastroenterology*. 2012;142:266–72 e1. [PubMed: 22062360]
4. Shahedi K, Fuller G, Bolus R, Cohen E, Vu M, Shah R, Agarwal N, Kaneshiro M, Atia M, Sheen V, Kurzbard N, van Oijen MGH, Yen L, Hodgkins P, Erder MH and Spiegel B. Long-term Risk of Acute Diverticulitis Among Patients With Incidental Diverticulosis Found During Colonoscopy. *Clin Gastroenterol Hepatol*. 2013;11:1609–1613. [PubMed: 23856358]
5. Liu PH, Cao Y, Keeley BR, Tam I, Wu K, Strate LL, Giovannucci EL and Chan AT. Adherence to a Healthy Lifestyle is Associated With a Lower Risk of Diverticulitis among Men. *Am J Gastroenterol*. 2017;112:1868–1876. [PubMed: 29112202]
6. Strate LL, Liu YL, Aldoori WH, Syngal S and Giovannucci EL. Obesity increases the risks of diverticulitis and diverticular bleeding. *Gastroenterology*. 2009;136:115–122 e1. [PubMed: 18996378]
7. Aldoori WH, Giovannucci EL, Rockett HR, Sampson L, Rimm EB and Willett WC. A prospective study of dietary fiber types and symptomatic diverticular disease in men. *J Nutr*. 1998;128:714–9. [PubMed: 9521633]
8. Rosemar A, Angeras U and Rosengren A. Body mass index and diverticular disease: a 28-year follow-up study in men. *Dis Colon Rectum*. 2008;51:450–5. [PubMed: 18157570]
9. Hjern F, Wolk A and Hakansson N. Smoking and the risk of diverticular disease in women. *Br J Surg*. 2011;98:997–1002. [PubMed: 21462366]



10. Strate LL, Liu YL, Aldoori WH and Giovannucci EL. Physical activity decreases diverticular complications. *Am J Gastroenterol.* 2009;104:1221–30. [PubMed: 19367267]
11. Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, de Ferranti SD, Floyd J, Fornage M, Gillespie C, Isasi CR, Jimenez MC, Jordan LC, Judd SE, Lackland D, Lichtman JH, Lisabeth L, Liu S, Longenecker CT, Mackey RH, Matsushita K, Mozaffarian D, Mussolino ME, Nasir K, Neumar RW, Palaniappan L, Pandey DK, Thiagarajan RR, Reeves MJ, Ritchey M, Rodriguez CJ, Roth GA, Rosamond WD, Sasson C, Towfighi A, Tsao CW, Turner MB, Virani SS, Voeks JH, Willey JZ, Wilkins JT, Wu JH, Alger HM, Wong SS, Muntner P, American Heart Association Statistics C and Stroke Statistics S. Heart Disease and Stroke Statistics-2017 Update: A Report From the American Heart Association. *Circulation.* 2017;135:e146–e603. [PubMed: 28122885]
12. Barbara G, Scaioli E, Barbaro MR, Biagi E, Laghi L, Cremon C, Marasco G, Colecchia A, Picone G, Salfi N, Capozzi F, Brigidi P and Festi D. Gut microbiota, metabolome and immune signatures in patients with uncomplicated diverticular disease. *Gut.* 2017;66:1252–1261. [PubMed: 27618836]
13. Chistiakov DA, Bobryshev YV, Kozarov E, Sobenin IA and Orekhov AN. Role of gut microbiota in the modulation of atherosclerosis-associated immune response. *Front Microbiol.* 2015;6:671. [PubMed: 26175728]
14. Boulange CL, Neves AL, Chilloux J, Nicholson JK and Dumas ME. Impact of the gut microbiota on inflammation, obesity, and metabolic disease. *Genome Med.* 2016;8:42. [PubMed: 27098727]
15. Baothman OA, Zamzami MA, Taher I, Abubaker J and Abu-Farha M. The role of Gut Microbiota in the development of obesity and Diabetes. *Lipids Health Dis.* 2016;15:108. [PubMed: 27317359]
16. Foster KJ, Holdstock G, Whorwell PJ, Guyer P and Wright R. Prevalence of diverticular disease of the colon in patients with ischaemic heart disease. *Gut.* 1978;19:1054–6. [PubMed: 730073]
17. Strate LL, Erichsen R, Horvath-Puho E, Pedersen L, Baron JA and Sorensen HT. Diverticular disease is associated with increased risk of subsequent arterial and venous thromboembolic events. *Clin Gastroenterol Hepatol.* 2014;12:1695–701 e1. [PubMed: 24316104]
18. Lin JN, Lin CL, Yang CH, Lin MC, Lai CH, Lin HH and Kao CH. Increased Risk of Acute Coronary Syndrome in Patients With Diverticular Disease: A Nationwide Population-Based Study. *Medicine (Baltimore).* 2015;94:e2020. [PubMed: 26559302]
19. Rimm EB, Stampfer MJ, Colditz GA, Giovannucci E and Willett WC. Effectiveness of various mailing strategies among nonrespondents in a prospective cohort study. *Am J Epidemiol.* 1990;131:1068–71. [PubMed: 2343859]
20. Rimm EB, Giovannucci EL, Willett WC, Colditz GA, Ascherio A, Rosner B and Stampfer MJ. Prospective study of alcohol consumption and risk of coronary disease in men. *Lancet.* 1991;338:464–8. [PubMed: 1678444]
21. Walker AE, Robins M and Weinfeld FD. The National Survey of Stroke. Clinical findings. *Stroke.* 1981;12:113–44. [PubMed: 7222164]
22. Colditz GA, Martin P, Stampfer MJ, Willett WC, Sampson L, Rosner B, Hennekens CH and Speizer FE. Validation of questionnaire information on risk factors and disease outcomes in a prospective cohort study of women. *Am J Epidemiol.* 1986;123:894–900. [PubMed: 3962971]
23. Sotos-Prieto M, Bhupathiraju SN, Mattei J, Fung TT, Li Y, Pan A, Willett WC, Rimm EB and Hu FB. Changes in Diet Quality Scores and Risk of Cardiovascular Disease Among US Men and Women. *Circulation.* 2015;132:2212–9. [PubMed: 26644246]
24. Fung TT, Chiuve SE, McCullough ML, Rexrode KM, Logroscino G and Hu FB. Adherence to a DASH-style diet and risk of coronary heart disease and stroke in women. *Arch Intern Med.* 2008;168:713–20. [PubMed: 18413553]
25. Feskanich D, Rimm EB, Giovannucci EL, Colditz GA, Stampfer MJ, Litin LB and Willett WC. Reproducibility and validity of food intake measurements from a semiquantitative food frequency questionnaire. *J Am Diet Assoc.* 1993;93:790–6. [PubMed: 8320406]
26. Karlsson FH, Fak F, Nookaew I, Tremaroli V, Fagerberg B, Petranovic D, Backhed F and Nielsen J. Symptomatic atherosclerosis is associated with an altered gut metagenome. *Nature communications.* 2012;3:1245.

27. Forslund K, Hildebrand F, Nielsen T, Falony G, Le Chatelier E, Sunagawa S, Prifti E, Vieira-Silva S, Gudmundsdottir V, Pedersen HK, Arumugam M, Kristiansen K, Voigt AY, Vestergaard H, Hercog R, Costea PI, Kultima JR, Li J, Jorgensen T, Levenez F, Dore J, Meta HITc, Nielsen HB, Brunak S, Raes J, Hansen T, Wang J, Ehrlich SD, Bork P and Pedersen O. Disentangling type 2 diabetes and metformin treatment signatures in the human gut microbiota. *Nature*. 2015;528:262–266. [PubMed: 26633628]
28. Li DY and Tang WHW. Gut Microbiota and Atherosclerosis. *Current atherosclerosis reports*. 2017;19:39. [PubMed: 28842845]
29. Musso G, Gambino R and Cassader M. Obesity, diabetes, and gut microbiota: the hygiene hypothesis expanded? *Diabetes Care*. 2010;33:2277–84. [PubMed: 20876708]
30. Wang Z, Klipfell E, Bennett BJ, Koeth R, Levison BS, Dugar B, Feldstein AE, Britt EB, Fu X, Chung YM, Wu Y, Schauer P, Smith JD, Allayee H, Tang WH, DiDonato JA, Lusis AJ and Hazen SL. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature*. 2011;472:57–63. [PubMed: 21475195]
31. Tang WH, Wang Z, Levison BS, Koeth RA, Britt EB, Fu X, Wu Y and Hazen SL. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *N Engl J Med*. 2013;368:1575–84. [PubMed: 23614584]
32. Jie Z, Xia H, Zhong SL, Feng Q, Li S, Liang S, Zhong H, Liu Z, Gao Y, Zhao H, Zhang D, Su Z, Fang Z, Lan Z, Li J, Xiao L, Li J, Li R, Li X, Li F, Ren H, Huang Y, Peng Y, Li G, Wen B, Dong B, Chen JY, Geng QS, Zhang ZW, Yang H, Wang J, Wang J, Zhang X, Madsen L, Brix S, Ning G, Xu X, Liu X, Hou Y, Jia H, He K and Kristiansen K. The gut microbiome in atherosclerotic cardiovascular disease. *Nature communications*. 2017;8:845.
33. Daniels L, Budding AE, de Korte N, Eck A, Bogaards JA, Stockmann HB, Consten EC, Savelkoul PH and Boermeester MA. Fecal microbiome analysis as a diagnostic test for diverticulitis. *Eur J Clin Microbiol Infect Dis*. 2014;33:1927–36. [PubMed: 24894339]
34. Guarner F and Malagelada JR. Gut flora in health and disease. *Lancet*. 2003;361:512–9. [PubMed: 12583961]
35. Ma W, Jovani M, Nguyen LH, Tabung FK, Song M, Liu PH, Cao Y, Tam I, Wu K, Giovannucci EL, Strate LL and Chan AT. Association Between Inflammatory Diets, Circulating Markers of Inflammation, and Risk of Diverticulitis. *Clin Gastroenterol Hepatol*. 2020;18:2279–2286 e3. [PubMed: 31712072]
36. Ruiz-Canela M, Bes-Rastrollo M and Martinez-Gonzalez MA. The Role of Dietary Inflammatory Index in Cardiovascular Disease, Metabolic Syndrome and Mortality. *International journal of molecular sciences*. 2016;17.
37. D'Agostino RB Sr., Vasan RS, Pencina MJ, Wolf PA, Cobain M, Massaro JM and Kannel WB. General cardiovascular risk profile for use in primary care: the Framingham Heart Study. *Circulation*. 2008;117:743–53. [PubMed: 18212285]

**Table 1.** Characteristics of men in the Health Professionals Follow-Up Study at baseline (1986) and midpoint (1998) according to presence or absence of diverticulitis

| Characteristics*  | Baseline 1986                   |                             | Midpoint 1998                   |                             |
|---|---------------------------------|-----------------------------|---------------------------------|-----------------------------|
|   | No Diverticulitis<br>n = 42,744 | Diverticulitis<br>n = 1,160 | No Diverticulitis<br>n = 33,545 | Diverticulitis<br>n = 1,013 |
| Age, years  | 53.8 (9.6)                      | 52.4 (8.7)                  | 63.9 (8.9)                      | 63.4 (8.5)                  |
| Total red meat intake, g/day                                  | 76.2 (55.0)                     | 81.7 (54.5)                 | 70.3 (53.2)                     | 74.9 (52.8)                 |
| Dietary fiber, g/day  | 20.9 (7.0)                      | 19.8 (6.4)                  | 23.6 (7.5)                      | 22.7 (6.9)                  |
| All physical activity, MET-hours/week                         | 19.0 (26.3)                     | 16.7 (22.1)                 | 29.1 (29.2)                     | 28.2 (28.2)                 |
| Vigorous physical activity, MET-hours/week                    | 13.2 (26.3)                     | 10.5 (21.1)                 | 13.6 (25.4)                     | 11.7 (25.8)                 |
| Body mass index, kg/m <sup>2</sup>                            | 25.5 (3.3)                      | 25.6 (3.0)                  | 26.2 (3.7)                      | 26.5 (3.6)                  |
| Ever smokers, %   | 51.2                            | 56.1                        | 48.8                            | 55.0                        |
| Pack-years among ever smokers                                 | 24.3 (18.6)                     | 25.9 (18.8)                 | 23.2 (18.3)                     | 25.8 (19.3)                 |
| Alcohol intake, g/day   | 11.3 (15.4)                     | 11.4 (14.6)                 | 11.1 (14.2)                     | 10.6 (13.2)                 |
| Regular aspirin use, %  | 27.0                            | 28.9                        | 51.9                            | 56.1                        |
| Regular NSAID use, %  | 5.4                             | 7.2                         | 17.5                            | 22.9                        |
| Regular acetaminophen use, %                                  | 5.7                             | 8.4                         | 7.7                             | 9.9                         |
| Hypertension, %   | 20.3                            | 19.9                        | 34.8                            | 38.3                        |
| Hyperlipidemia, %   | 10.7                            | 11.9                        | 45.2                            | 50.2                        |
| Diabetes, %   | 0.4                             | 0.4                         | 4.3                             | 4.7                         |
| Any hypertension, hyperlipidemia, or diabetes, %              | 27.3                            | 27.9                        | 61.3                            | 67.0                        |
| Any hypertension, hyperlipidemia, diabetes, or ever smoker, % | 61.4                            | 64.1                        | 77.8                            | 81.8                        |

MET, metabolic equivalent of tasks; NSAID, non-steroid anti-inflammatory drugs

\* Mean (standard deviation) is presented for continuous variables.

**Table 2.** History of diverticulitis and risk of incident cardiovascular disease among all participants

|                         | Any history of diverticulitis |                  | Recent history of diverticulitis within past 4 years |                  | Remote history of diverticulitis beyond 4 years |                  |
|-------------------------|-------------------------------|------------------|--|------------------|---|------------------|
|                         | No                            | Yes              | No   | Yes              | No  | Yes              |
| Number of CVD events    | 3,772                         | 76               | 3,772  | 27               | 3,772   | 49               |
| Person-years            | 845,861                       | 10,458           | 845,861  | 3,432            | 845,861   | 7,036            |
| Age-adjusted            | 1 (Ref)                       | 1.49 (1.18-1.88) | 1 (Ref)  | 1.61 (1.09-2.36) | 1 (Ref)   | 1.43 (1.07-1.90) |
| *Multivariable-adjusted | 1 (Ref)                       | 1.40 (1.11-1.76) | 1 (Ref)  | 1.52 (1.03-2.23) | 1 (Ref)   | 1.34 (1.00-1.79) |
| <sup>†</sup> Full model | 1 (Ref)                       | 1.35 (1.07-1.70) | 1 (Ref)  | 1.45 (0.98-2.14) | 1 (Ref)   | 1.30 (0.97-1.73) |

  

| History of diverticulitis and risk of incident cardiovascular disease among participants below age of 65 years |                               |                  |  |                  |   |                  |
|--|-------------------------------|------------------|--|------------------|---|------------------|
|  | Any history of diverticulitis |                  | Recent history of diverticulitis within past 4 years |                  | Remote history of diverticulitis beyond 4 years |                  |
|  | No                            | Yes              | No   | Yes              | No  | Yes              |
| Number of CVD events   | 1,466                         | 21               | 1,466  | 8                | 3,772   | 13               |
| Person-years   | 512,650                       | 4,392            | 512,650  | 1,829            | 845,861   | 2,563            |
| Age-adjusted   | 1 (Ref)                       | 1.50 (0.97-2.32) | 1 (Ref)  | 1.34 (0.67-2.71) | 1 (Ref)   | 1.62 (0.93-2.81) |
| *Multivariable-adjusted  | 1 (Ref)                       | 1.36 (0.88-2.11) | 1 (Ref)  | 1.27 (0.63-2.56) | 1 (Ref)   | 1.43 (0.82-2.48) |
| <sup>†</sup> Full model  | 1 (Ref)                       | 1.28 (0.83-1.98) | 1 (Ref)  | 1.20 (0.59-2.44) | 1 (Ref)   | 1.33 (0.76-2.32) |

  

| History of diverticulitis and risk of incident cardiovascular disease among participants above age of 65 years |                               |                  |  |                  |   |                  |
|--|-------------------------------|------------------|--|------------------|---|------------------|
|  | Any history of diverticulitis |                  | Recent history of diverticulitis within past 4 years |                  | Remote history of diverticulitis beyond 4 years |                  |
|  | No                            | Yes              | No   | Yes              | No  | Yes              |
| Number of CVD events   | 2,306                         | 55               | 2,306  | 19               | 2,306   | 36               |
| Person-years   | 333,211                       | 6,066            | 333,211  | 1,603            | 333,211   | 4,463            |
| Age-adjusted   | 1 (Ref)                       | 1.48 (1.12-1.94) | 1 (Ref)  | 1.75 (1.11-2.78) | 1 (Ref)   | 1.36 (0.97-1.90) |
| *Multivariable-adjusted  | 1 (Ref)                       | 1.41 (1.07-1.85) | 1 (Ref)  | 1.67 (1.05-2.65) | 1 (Ref)   | 1.30 (0.93-1.81) |
| <sup>†</sup> Full model  | 1 (Ref)                       | 1.36 (1.04-1.80) | 1 (Ref)  | 1.60 (1.01-2.54) | 1 (Ref)   | 1.27 (0.90-1.77) |

Note: Hazard ratio and 95% confidence interval are shown.

\* Adjusted for lifestyle factors (red meat intake, fiber intake, body mass index, calorie intake, smoking, Alternate Healthy Eating Index, alcohol consumption, and physical activity).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

† Additionally adjusted for family history (myocardial infarction or diabetes), medication (aspirin, NSAID, acetaminophen, lipid-lowering, and anti-hypertensive), hypertension, diabetes, and hyperlipidemia.

**Table 3.**

History of diverticulitis and risk of incident cardiovascular disease among participants with regular screening practices.

|                         | Any history of diverticulitis |                  | Recent history of diverticulitis within past 4 years |                  | Remote history of diverticulitis beyond 4 years |                  |
|-------------------------|-------------------------------|------------------|--|------------------|---|------------------|
|                         | No                            | Yes              | No   | Yes              | No  | Yes              |
| Number of CVD events    | 2,001                         | 49               | 2,001  | 14               | 2,001   | 35               |
| Person-years            | 445,383                       | 6,294            | 445,383  | 1,792            | 445,383   | 4,502            |
| Age-adjusted            | 1 (Ref)                       | 1.69 (1.26-2.27) | 1 (Ref)  | 1.53 (0.89-2.62) | 1 (Ref)   | 1.77 (1.25-2.49) |
| *Multivariable-adjusted | 1 (Ref)                       | 1.56 (1.17-2.09) | 1 (Ref)  | 1.40 (0.82-2.41) | 1 (Ref)   | 1.64 (1.16-2.32) |
| †Full model             | 1 (Ref)                       | 1.51 (1.13-2.03) | 1 (Ref)  | 1.40 (0.82-2.41) | 1 (Ref)   | 1.56 (1.10-2.20) |

Note: Hazard ratio and 95% confidence interval are shown.

\* Adjusted for lifestyle factors (red meat intake, fiber intake, body mass index, calorie intake, smoking, Alternate Healthy Eating Index, alcohol consumption, and physical activity).

† Additionally adjusted for family history (myocardial infarction or diabetes), medication (aspirin, NSAID, acetaminophen, lipid-lowering, and anti-hypertensive), hypertension, diabetes, and hyperlipidemia.

**Table 4.**

History of diverticulitis and risk of incident cardiovascular disease in participants with and without cardiovascular risk factors

|                         | Without hypertension, diabetes, hyperlipidemia, and smoking |                  | With either hypertension, diabetes, hyperlipidemia, or smoking |                  |
|-------------------------|---|------------------|--|------------------|
|                         | History Diverticulitis                                      |                  | History Diverticulitis   |                  |
|                         | No  | Yes              | No   | Yes              |
| Number of CVD events    | 466   | 11               | 3,306  | 65               |
| Person-years            | 207,933   | 1,601            | 637,928  | 8,857            |
| Age-adjusted            | 1 (Ref)   | 3.72 (1.88-7.35) | 1 (Ref)  | 1.39 (1.08-1.79) |
| *Multivariable-adjusted | 1 (Ref)   | 3.89 (1.96-7.73) | 1 (Ref)  | 1.32 (1.03-1.70) |
| †Full model             | 1 (Ref)   | 4.06 (2.04-8.08) | 1 (Ref)  | 1.27 (0.98-1.63) |

Note: Hazard ratio and 95% confidence interval are shown.

\* Adjusted for lifestyle factors (red meat intake, fiber intake, body mass index, calorie intake, smoking, Alternate Healthy Eating Index, alcohol consumption and physical activity).

† Additionally adjusted for family history (myocardial infarction or diabetes), medication (aspirin, NSAID, acetaminophen, lipid-lowering, and anti-hypertensive), hypertension, diabetes, and hyperlipidemia.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript