

Catalog of 5' Fusion Partners in ALK-positive NSCLC Circa 2020



Sai-Hong Ignatius Ou, MD, PhD,^{a,*} Viola W. Zhu, MD, PhD,^a Misako Nagasaka, MD^{b,c}

^aDepartment of Medicine, Division of Hematology-Oncology, Chao Family Comprehensive Cancer Center, University of California Irvine School of Medicine, Orange, California

^bDepartment of Oncology, Karmanos Cancer Institute/Wayne State University School of Medicine, Detroit, Michigan

^cDepartment of Advanced Medical Innovations, St. Marianna University Graduate School of Medicine, Kawasaki, Kanagawa, Japan

Received 4 February 2020; accepted 5 February 2020

Available online - 19 February 2020

ABSTRACT

Since the discovery of anaplastic lymphoma kinase fusion-positive (*ALK+*) NSCLC in 2007, the methods to detect *ALK+* NSCLC have evolved and expanded from fluorescence in situ hybridization and immunohistochemistry to next-generation DNA sequencing, targeted RNA sequencing, and whole transcriptome sequencing. As such, the deep sequencing methods have resulted in the expansion of distinct fusion partners identified in *ALK+* NSCLC to 90 (one variant *PLEKHM2-ALK* is found in small cell lung cancer but included in this catalog) by the end of January 2020; about 65 of them (since 2018) and most of the recent novel fusion partners were reported from China. Thirty-four of the distinct fusion partners are located on the short arm of chromosome 2; 28 of these 34 fusion partners are located on 2p21-25, in which *ALK* is located on 2p23.2-p23.1. Many of these new *ALK+* NSCLC fusion variants have responded to *ALK* tyrosine kinase inhibitors (TKIs). Several of these novel *ALK* fusion variants were identified as being resistant to EGFR TKIs or as dual 3'*ALK* fusions. In addition, at least 28 intergenic *ALK* rearrangements have also been reported, with three of them reported as responding to crizotinib. This review aims to serve as a central source of reference of fusion partners in *ALK+* NSCLC for clinicians and scientists. We aim to update and improve the list going forward.

© 2020 The Authors. Published by Elsevier Inc. on behalf of the International Association for the Study of Lung Cancer. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: *ALK* fusion partners; Next-generation sequencing; *ALK+* NSCLC; Whole transcriptome sequencing

Introduction

Since the discovery of anaplastic lymphoma kinase fusion-positive (*ALK+*) NSCLC (*EML4-ALK*, *TPF-ALK*) in 2007,^{1,2} there has been a rapid development of *ALK* tyrosine kinase inhibitors (TKIs) to treat *ALK+* NSCLC with five *ALK* TKIs approved in the United States (crizotinib, ceritinib, alectinib, brigatinib, and lorlatinib) by 2018. At the same time, the detection of *ALK+* NSCLC has expanded and shifted from the original methods of fluorescence in situ hybridization and immunohistochemistry (IHC) to next-generation sequencing (NGS), targeted RNA sequencing, and even whole transcriptome sequencing being offered by commercial sequencing companies. Targeted RNA sequencing and whole transcriptome sequencing have been used to supplement

*Corresponding author.

Disclosure: Dr. Ou has stock ownership and was on the scientific advisory board of Turning Point Therapeutics, Inc. (until Feb 28, 2019). He has received speaker honorarium from Merck, Roche/Genentech, Astra Zeneca, Takeda/ARIAD, and Pfizer and has received advisory fees from Roche/Genentech, Astra Zeneca, Takeda/ARIAD, Pfizer, Foundation Medicine Inc., and Spectrum. Dr. Zhu has received honoraria from AstraZeneca, Biocept, Roche-Foundation Medicine, Roche/Genentech, and Takeda, and has stock ownership of Turning Point Therapeutics, Inc. Dr. Nagasaka has received honorarium from Astra Zeneca and Tempus.

Address for correspondence: Sai-Hong Ignatius Ou, MD, PhD, Chao Family Comprehensive Cancer Center, University of California Irvine School of Medicine, 200 South Manchester Avenue, Suite 407, Orange, CA 92868-3298. E-mail: siou@uci.edu

Cite this article as: Ou S-HI, et al. Catalog of 5' Fusion Partners in *ALK*-positive NSCLC Circa 2020. *JTO Clin Res Rep* 1:100015

© 2020 The Authors. Published by Elsevier Inc. on behalf of the International Association for the Study of Lung Cancer. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

ISSN: 2666-3643

<https://doi.org/10.1016/j.jtocr.2020.100015>

Table 1. Catalog of Fusion Partners in ALK+ NSCLC

No.	Fusion Partner	Year Published in Print/Presented	Chromosomal Location	Fusion Breakpoint	Response to ALK TKI at the Time of Publication	Tumor Source	Method of Detection	Variant Frequency in Tumor	FISH/IHC	References ^a
1	<i>EML4</i>	2007	2p21	(E13, A21)	Not treated with ALK TKI	Tumor	PCR/Sanger sequencing	NR	ND/ND	Soda, 2007 ¹
		2007	2p21	(E13, A20)	Not treated with ALK TKI	Cell line/ Tumor	5'RACE PCR DNA sequencing	NR	ND/ND	Rikova, 2007 ²
2	<i>TFG</i>	2007	3q12.2	(T3, A20)	Not treated with ALK TKI	Tumor	5'RACE PCR DNA sequencing	NR	ND/ND	Rikova, 2007 ²
		2007	3q12.2	NR	Not treated with ALK TKI	Tumor	PCR/Sanger sequencing	NR	ND/ND	Soda, 2007 ¹
3	<i>KIF5B</i>	2009	10p11.22	(K24, A20)	Not treated with ALK TKI	Tumor	RT-PCR	NR	+/+	Takeuchi, 2009 ⁶
		2011	10p11.22	(K15, A20)	Not treated with ALK TKI	Tumor	RT-PCR	NR	+/+	Won, 2011 ⁷
		2012	10p11.22	(K17, A20)	Not treated with ALK TKI	Tumor	RT-PCR	NR	+/+	Takeuchi, 2012 ⁸
4	<i>KLC1</i>	2012	14q32.33	(K9, A 20)	Not treated with ALK TKI	Tumor	RT-PCR	NR	+/ND	Togashi, 2012 ⁹
5	<i>STRN</i>	2013	2p22.2	(S3, A20)	NR	Tumor	RT-PCR	NR	+/+	Majewski, 2013 ¹⁰
		2017	2p22.2	(S3, A20)	PR to crizotinib	Plasma	DNA NGS	1%	ND/ND	Yang, 2017 ¹¹
		2017	2p22.2	(S3, A20)	PR to alectinib	Tumor	RNA sequencing	NR	+/+	Nakanishi, 2017 ¹²
	<i>STRN^b</i>	2019	2p22.2	(S3, A20)	Not treated with ALK TKI	Tumor	DNA NGS	NR	NR/NR	Xu, 2019 ¹³
6	<i>HIP1</i>	2014	7q11.23	(H2, A20)	Not treated with ALK TKI	Tumor	RNA sequencing	NR	ND/ND	Fang, 2014 ¹⁴
		2014	7q11.23	(H21, A 20)	PR to crizotinib	Tumor	RT-PCR	NR	+/+	Hong, 2014 ¹⁵
		2014	7q11.23	(H30, A20)	PR to crizotinib and alectinib	Tumor	DNA NGS	NR	+/ND	Ou, 2014 ¹⁶
7	<i>TPR</i>	2014	1q31.1	(T15, A20)	Not treated with ALK TKI	Tumor	PCR	NR	+/+	Choi, 2014 ¹⁷
8	<i>BIRC6</i>	2015	2p22.3	NR	PR to crizotinib	Tumor	DNA NGS	NR	-/+	Shan 2015 ¹⁸
9	<i>DCTN1</i>	2015	2p13.1	(D26, A20)	NR	Tumor	DNA NGS	NR	+/ND	Iyevleva, 2015 ¹⁹
10	<i>SQSTM1</i>	2015	5q35.3	(S5, A20)	NR	Tumor	DNA NGS	NR	+/ND	Iyevleva, 2015 ¹⁹
11	<i>SOCS5</i>	2015	2p21	NR	NR	Tumor	NGS	NR	-/ND	Drilon, 2015 ²⁰
12	<i>SEC31A</i>	2016	4q21.22	(S21, A20)	Adjuvant setting, not treated with ALK TKI	Tumor	NGS	NR	+/+	Kim, 2016 ²¹
13	<i>CLTC</i>	2016	17q23.1	(C31, A20)	Unknown	Tumor	NGS	NR	NR/NR	Ali, 2016 ²²
14	<i>PRKAR1A</i>	2016	17q24.2	(P5, A20)	PR to crizotinib	Tumor	NGS	NR	+/+	Ali, 2016 ²²
15	<i>PPM1B</i>	2016	2p21	(P1, A20)	PR to crizotinib	Tumor	NGS	NR	NR/NR	Ali, 2016 ²²
16	<i>EIF2AK3</i>	2016	2p11.2	(E2, A20)	PR to crizotinib	Tumor	NGS	NR	-/-	Ali, 2016 ²²
17	<i>CRIM1</i>	2016	2p22.2	NR	NR	Tumor	NGS	NR	NR/NR	Tan, 2016 ²³
18	<i>CEBPZ</i>	2017	2p22.2	(C2, A20)	Not treated with ALK TKI	Tumor	NGS	25.3%	+/+	Li, 2017 ²⁴
	<i>CEBPZ^b</i>	2019	2p22.2	NR	Crizotinib, unknown results	Tumor	NGS	NR	NR/NR	Xu, 2019 ¹³
19	<i>PICALM</i>	2017	11q14.2	(P19, A20)	Not treated with ALK TKI	Tumor	NGS	10.2%	-/+	Li, 2017 ²⁴
20	<i>CLIP1</i>	2017	12q24.31	(C22, A20)	PR to crizotinib	Tumor	Targeted RNA sequencing	NR	+/+	Vendrelli, 2017 ²⁵

(continued)

Table 1. Continued

No.	Fusion Partner	Year Published in Print/Presented	Chromosomal Location	Fusion Breakpoint	Response to ALK TKI at the Time of Publication	Tumor Source	Method of Detection	Variant Frequency in Tumor	FISH/IHC	References ^a
21	<i>BCL11A</i> <i>BCL11A</i> ^c	2017	2p16.1	(B4, A20)	PR to crizotinib	Tumor	DNA and RNA NGS	NR	ND/ND	Tian, 2017 ²⁶
		2019	2p16.1	(B2, A18)	PR to crizotinib	Tumor and plasma	DNA NGS	54.2% (PPFE) 14.9% (plasma)	ND/ND	Qin 2019 ²⁷
22	<i>GCC2</i>	2017	2q12.3	(G12, A20)	NR	Tumor	RT-PCR, NGS	NR	+/+	Noh, 2017 ²⁸
		2017	2q12.3	(G19, A20)	Adjuvant setting, not treated with ALK TKI	Tumor	Targeted RNA sequencing	NR	+/+	Vendrell, 2017 ²⁵
		2018	2q12.3	(G18, A20)	PR to crizotinib and then ceritinib	Tumor	RT-PCR, sanger sequencing	NR	NR/NR	Jiang, 2018 ²⁹
23	<i>LMO7</i>	2017	13q22.2	(L15, A20)	NR	Tumor	RT-PCR, NGS	NR	+/+	Noh, 2017 ²⁸
24	<i>PHACTR1</i>	2017	6p24.1	(P7, A20)	NR	Tumor	RT-PCR, NGS	NR	+/+	Noh, 2017 ²⁸
25	<i>CMTR1</i>	2018	6p21.2	(C2, A20)	No with crizotinib, SD with pemetrexed	Tumor	NGS	~7.5%	-/-	Du, 2018 ³⁰
26	<i>VIT</i>	2018	2p22.2	(V7, A20)	PR to alectinib	Tumor	NGS	NR	+/+	Hu, 2018 ³¹
27	<i>DYSF</i>	2018	2p13.2	NR	Extracranial PR but intracranial progression to crizotinib	Pleural effusion	DNA NGS	23.7%	ND/+	Yin 2018 ³²
28	<i>ITGAV</i>	2018	2q32.1	NR	Extracranial PR but intracranial progression to crizotinib	Pleural effusion	DNA NGS	15.2%	ND/+	Yin, 2018 ³²
29	<i>PLEKHA7</i> ^b	2018	11p15.2-p15.1	(P26, A19)	PR to alectinib + osimertinib	Plasma	DNA NGS	NR	ND/ND	Schrock, 2018 ³³
30	<i>CUX1</i>	2018	7q22.1	(C8, A20)	PR to crizotinib	Tumor	NGS	11%	NR/NR	Zhang 2018 ³⁴
31	<i>VKORC1L1</i>	2018	7q11.21	(V1, A20)	PR with crizotinib and alectinib	Plasma	NGS	NR	+/ND	Zhu, 2018 ³⁵
32	<i>FBXO36</i>	2018	2q36.3	NR	PR to crizotinib	Tumor	NGS	NR	ND/+	Xu, 2018 ³⁶
33	<i>SPTBN1</i> ^c	2018	2p16.2	NR	NR	Plasma	NGS	NR	NR/NR	Ramalingam, 2018 ³⁷
34	<i>EML6</i> ^d	2018	2p16.1	(E1, A20)	PR to crizotinib	Tumor	NGS	NR	ND/+	Lin, 2018 ³⁸
35	<i>FBXO11</i> ^d	2018	2p16.3	(F1, A20)	PR to crizotinib	Tumor	NGS	NR	ND/+	Lin, 2018 ³⁸
36	<i>CLIP4</i>	2018	2p23.2	(C7, A20)	NR	Tumor	NGS	NR	ND/+	Zhao, 2018 ³⁹
37	<i>CAMKMT</i>	2019	2p21	(C3, A 20)	Not treated with ALK TKI	Tumor	NGS	NR	+/+	Hu, 2019 ⁴⁰
38	<i>NCOA1</i>	2019	2p23.3	(N12, A20)	PR to crizotinib, PFS > 18 months	Tumor	NGS	NR	ND/+	Cao, 2019 ⁴¹
39	<i>MYT1L</i>	2019	2p25.3	(M14, A20)	PR on crizotinib, PD on ceritinib and alectinib	Tumor	NGS	NR	-/ND	Tsou, 2019 ⁴²
40	<i>SRBD1</i>	2019	2p21	(S20, A20)	Not treated with ALK TKI	Tumor	NGS	2.6%	ND/+	Hou, 2019 ⁴³
41	<i>SRD5A2</i>	2019	2p23.1	(S1, A20)	NR	Tumor	NGS	NR	ND/+	Zhao, 2019 ⁴⁴

(continued)

Table 1. Continued

No.	Fusion Partner	Year Published in Print/Presented	Chromosomal Location	Fusion Breakpoint	Response to ALK TKI at the Time of Publication	Tumor Source	Method of Detection	Variant Frequency in Tumor	FISH/IHC	References ^a
42	<i>NYAP2</i> (<i>KIAA 1486</i>)	2019	2q36.3	(N3, A20)	NR	Tumor	NGS	NR	ND/-	Zhao, 2019 ⁴⁴
43	<i>MPRIP</i>	2019	17p11.2	(M21, A20)	PR to crizotinib	Tumor	RNA sequencing	NR	+/+	Fan, 2019 ⁴⁵
44	<i>ADAM17</i>	2019	2p25.1	(A4, A20)	PR to alectinib	Plasma	DNA NGS	3.68%	NR/NR	Supplee, 2019 ⁴⁶
45	<i>ALK</i>	2019	2p23.2-p23.1	(A6, A20)	NR	Plasma	DNA NGS	26.63%	NR/NR	Supplee, 2019 ⁴⁶
46	<i>LPIN1^b</i>	2019	2p25.1	NR	Response to crizotinib + erlotinib	Tumor	NR	NR	NR/NR	Supplee 2019 ⁴⁶
47	<i>WDPCP</i>	2019	2p15	(W17, A20)	PR to crizotinib	Tumor	DNA NGS	52.6%	+/+	He, 2019 ⁴⁷
48	<i>CEP55</i>	2019	10q23.33	(C3, A20)	NR	Tumor	DNA NGS	NR	NR/NR	Couëtoux du Tertr, 2019 ⁴⁸
49	<i>ERC1^e</i>	2019	12p13.33	(E15, A20)	NR	Tumor	DNA NGS	NR	NR/NR	Couëtoux du Tertr, 2019 ⁴⁸
		2019	12p13.33	NR	NR	Tumor/ plasma	DNA NGS	NR	NR/NR	Zhou, 2019 ⁴⁹
50	<i>SLC16A7^e</i>	2019	12q14.1	(S1, A 20)	PR to crizotinib prolonged PFS	Tumor	DNA NGS	NR	NR/NR	Couëtoux du Tertr, 2019 ⁴⁸
51	<i>TNIP2</i>	2019	4p16.3	(T5, A20)	PR to crizotinib	Tumor/ plasma	DNA NGS	0.1% (plasma) 3.3% (tumor)	ND/+	Feng, 2019 ⁵⁰
52	<i>ATAD2B</i>	2019	2p24.1-p23.3	(A1, A20)	Treated with crizotinib	Tumor	DNA NGS	NR	ND/+	Bai, 2019 ⁵¹
53	<i>SLMAP</i>	2019	3p14.3	(S12, A20) (S13, A20)	Unknown, adjuvant treatment with crizotinib	Tumor	Anchored Multiplex RNA sequencing	NR	+/+	Paga, 2019 ⁵²
54	<i>FBN1</i>	2019	15q21.1	NR	NR	Tumor/ plasma	DNA NGS	NR	NR/NR	Zhou, 2019 ⁴⁹
55	<i>SWAP70</i>	2019	11p15.4	NR	NR	Tumor/ plasma	DNA NGS	NR	NR/NR	Zhou, 2019 ⁴⁹
56	<i>TCF12</i>	2019	15q21.3	NR	NR	Tumor/ plasma	DNA NGS	NR	NR/NR	Zhou, 2019 ⁴⁹
57	<i>TRIM66</i>	2019	11p15.4	NR	NR	Tumor/ plasma	DNA NGS	NR	NR/NR	Zhou, 2019 ⁴⁹
58	<i>WNK3</i>	2019	Xp11.22	NR	NR	Tumor/ plasma	DNA NGS	NR	NR/NR	Zhou, 2019 ⁴⁹
59	<i>AKAP8L</i>	2019	19p13.12	NR	ensartinib	plasma	DNA NGS	NR	NR/NR	Horn, 2019 ⁵³
60	<i>SPECC1L</i>	2019	22q11.23	(S9, A20)	Not treated with ALK TKI	Tumor	DNA NGS	NR	NR/NR	Pan, 2019 ⁵⁴
61	<i>PRKCB^f</i>	2019	16p12.2-p12.1	(P2, A19)	PR to crizotinib, disappearance of PRKCB-ALK fusion variant	Tumor and plasma	NGS	2.6% (tumor) 0.8% (plasma)	NR/NR	Luo, 2019 ⁵⁵

(continued)

Table 1. Continued

No.	Fusion Partner	Year Published in Print/Presented	Chromosomal Location	Fusion Breakpoint	Response to ALK TKI at the Time of Publication	Tumor Source	Method of Detection	Variant Frequency in Tumor	FISH/IHC	References ^a
62	<i>CDK15</i> ^l	2019	2q33.1	(C10, A19)	NR	Tumor	DNA NGS	NR	NR/NR	Wen, 2019 ⁵⁶
63	<i>LCLAT1</i>	2019	2p23.1	NR	NR	Tumor	DNA NGS	NR	NR/NR	Wen, 2019 ⁵⁶
64	<i>YAP1</i>	2019	11q22.1	NR	NR	Tumor	DNANGS	NR	NR/NR	Wen, 2019 ⁵⁶
65	<i>PLEKHM2</i> (<i>SCLC</i>)	2020	1p36.21	(P7, A20)	SD to crizotinib and brigatinib	Tumor	NGS	NR	ND/+	Li, 2020 ⁵⁷
66	<i>DCHS1</i>	2020	11p15.4	NR	PR or SD to ensartinib	Tumor	NGS	NR	NR/NR	Yang, 2020 ⁵⁸
67	<i>PPF1BP1</i>	2020	12p11.23- p11.22	NR	PR or SD to ensartinib	Tumor	NGS	NR	NR/NR	Yang, 2020 ⁵⁸
68	<i>ATP13A4</i>	2020	3q29	(A9, A19)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
69	<i>C12orf75</i>	2020	12q23.3	(C1, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
70	<i>EPAS1</i>	2020	2p21	(E1, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
71	<i>FAM179A</i> (<i>TOGARAM2</i>)	2020	2p23.2	(F1, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
		2020	2p23.2	(F13, A20)	NR	Plasma	NGS	NR	ND/NR	Zhang, 2020 ⁶⁰
72	<i>FUT8</i>	2020	14q23.3	(F3, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
73	<i>LIMD1</i>	2020	3p21.31	(L2, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
74	<i>LINC00327</i>	2020	13q12.12	(L2, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
75	<i>LOC349160</i>	2020	7q33	(L1, A20)	SD to crizotinib	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
76	<i>LYPD1</i>	2020	2q21.2	(L3, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
77	<i>RBM20</i>	2020	10q25.2	(R1, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
78	<i>TACR1</i>	2020	2p12	(T1, A20)	PR to crizotinib	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
79	<i>TANC1</i>	2020	2q24.2	(T3, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
80	<i>TTC27</i>	2020	2p22.3	(T12, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
81	<i>TUBBB</i>	2020	6p21.33	(T3, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
82	<i>SMPD4</i>	2020	2q21.1	(S1, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
83	<i>SORCS1</i>	2020	10q25.1	(S10, A20)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
84	<i>LINC00211</i>	2020	2p22.2	(L?, A20)	PR with crizotinib and alectinib, SD with lorlatinib	CSF	NGS	33.2%	NR/+	Li, 2020 ⁶¹
85	<i>SOS1</i>	2020	2p22.1	(S2, A20)	PR to crizotinib	FFPE	NGS	NR	ND/ND	Chen, 2020 ⁶²
86	<i>C9orf3</i>	2020	9q22.32	(C12, A20)	NR	FFPE	NGS	22.6%	ND/+	Zhang, 2020 ⁶⁰
87	<i>CYBRD1</i>	2020	2q31.1	(C21, A20)	NR	FFPE	NGS	12.5%	ND/NR	Zhang, 2020 ⁶⁰
88	<i>MTA3</i> ^s	2020	2p21	(M6, A 20)	SD with crizotinib, no response to alectinib	FFPE	NGS	15.3%	ND/NR	Zhang, 2020 ⁶⁰
89	<i>THADA</i>	2020	2p21	(T25, A20)	SD to crizotinib, PR to ceritinib	Plasma	NGS	0.3%	ND/NR	Zhang, 2020 ⁶⁰

(continued)

Table 1. Continued

No.	Fusion Partner	Year Published in Print/Presented	Chromosomal Location	Fusion Breakpoint	Response to ALK TKI at the Time of Publication	Tumor Source	Method of Detection	Variant Frequency in Tumor	FISH/IHC	References ^a
90	<i>TSPYL6</i> ^f	2020	2p16.2	(T6, A20)	PR to crizotinib, SD to alectinib	FFPE	NGS	8.5%	ND/NR	Zhang, 2020 ⁶⁰
91	<i>WDR37</i>	2020	10p15.3	(W6, A20)	PR to crizotinib	FFPE	NGS	30.2%	ND/NR	Zhang, 2020 ⁶⁰
92	<i>PLEKHH2</i>	2020	2p21	(P6, A20)	PR to alectinib	FFPE	Targeted RNA sequencing	NR	+/+	M. Nagasaka, written communication, 2020

^hThe earlier detected ALK fusion partners were not treated with crizotinib at the time of publication; but all of them have been shown to respond to ALK TKIs. The column entry is for the later discovery of ALK fusion partners.

^gThe first report(s) are cited except when response information from ALK TKIs are from later reports on some of the rare fusion partners, or if the fusion is identified as a resistance mechanism to EGFR TKI.

^bALK fusions identified as resistance to EGFR TKIs.

^cDual fusion with *EML4-ALK* (E18, A20).

^dDual fusions (*EML6* and *FBXO11*) together.

^eDual fusion (*ERC1* and *SLC16A7*) together.

^fDual fusion with *EML4-ALK* (E6, A20)

^gDual fusion with *EML4-ALK* (E7; A18)

+, positive; -, negative; ALK, anaplastic lymphoma kinase; CSF, cerebrospinal fluid; FISH, fluorescence in situ hybridization; FFPE, formalin-fixed paraffin embedded; FNA, fine-needle aspiration; IHC, immunohistochemistry; ND, not done; NGS, next-generation sequencing; NR, not reported; PFS, progression-free survival; PR, partial response; SD, stable disease; TKI, tyrosine kinase inhibitor; *ADAM17*, ADAM metallopeptidase domain 17; *AKAP8L*, A-kinase anchoring protein 8 like; *ATAD2B*, ATPase family AAA domain containing 2B; *ATP13A4*, ATPase 13A4; *BCL11A*, BAF chromatin remodeling complex subunit; *BIRC6*, baculoviral IAP repeat containing 6; *C12orf75*, chromosome 12 open reading frame 75; *CAMKMT*, calmodulin-lysine N-methyltransferase; *CDK15*, cyclin dependent kinase 15; *CEBPZ*, CCAAT enhancer binding protein zeta; *CLIP1*, CAP-Gly domain containing linker protein family member 1; *CLIP4*, CAP-Gly domain containing linker protein family member 4; *CMTR1*, cap methyltransferase 1; *CRIM1*, cysteine rich transmembrane BMP regulator 1; *CUX1*, cut like homeobox 1; *CYBRD1*, cytochrome b reductase 1; *DCHS1*, dachshous cadherin-related 1; *DCTN1*, dynactin subunit 1; *DYSF*, dysferlin; *EIF2AK3*, eukaryotic translation initiation factor 2 alpha kinase 3; *EML4*, echinoderm microtubule-associated protein-like 4; *EML6*, EMAP like 6; *EPAS1*, endothelial PAS domain protein 1; *ERC1*, ELKS/RAB6-interacting/CAST family member 1; *FAM179A*, family with sequence similarity 179 member A; *FBN1*, fibrillin 1; *FBXO11*, F-box protein 11; *FBXO36*, F-box protein 36; *FUT8*, fucosyltransferase 8; *GCC2*, GRIP and coiled-coil domain containing 2; *HIP1*, huntingtin interacting protein 1; *ITGAV*, integrin subunit alpha V; *KLC1*, kinesin light chain 1; *KIF5B*, kinesin family member 5B; *LCLAT1*, lysocardiolipin acyltransferase 1; *LIMD1*, LIM domains containing 1; *LINC00211*, long intergenic non-protein coding RNA 211; *LINC00327*, long intergenic non-protein coding RNA 327; *LMO7*, LIM domain 7; *LOC349160*, uncharacterized LOC349160; *LPIN1*, lipin 1; *LYPD1*, LY6/PLAUR domain containing 1; *MPRIP*, myosin phosphatase Rho interacting protein; *MTA3*, metastasis associated 1 family member 3; *MYT1L*, myelin transcription factor 1 like; *NCOA1*, nuclear receptor coactivator 1; *NYAP2*, neuronal tyrosine-phosphorylated phosphoinositide-3-kinase adaptor 2; *PHACTR1*, phosphatase and actin regulator 1; *PICALM*, phosphatidylinositol binding clathrin assembly protein; *PLEKHA7*, pleckstrin homology domain containing A7; *PLEKHH2*, pleckstrin homology, MyTH4 and FERM domain containing H2; *PLEKHM2*, pleckstrin homology and RUN domain containing M2; *PPFIBP1*, Liprin-beta-1/PPF1A. binding protein 1; *PPM1B*, protein phosphatase, Mg²⁺/Mn²⁺ dependent 1B; *PRKAR1A*, protein kinase cAMP-dependent type I regulatory subunit alpha; *PRKCB*, protein kinase C beta; *RBM20*, RNA binding motif protein 20; *SEC31A*, SEC31 homolog A, COPII coat complex component; *SLC16A7*, solute carrier family 16 member 7; *SLMAP*, sarcolemma associated protein; *SMPD4*, sphingomyelin phosphodiesterase 4; *SOCS5*, suppressor of cytokine signaling 5; *SORCS1*, sortilin related VPS10 domain containing receptor 1; *SOS1*, Son of sevenless Ras/Rac guanine nucleotide exchange factor 1; *SPECC1L*, sperm antigen with calponin homology and coiled-coil domains 1 like; *SRBD1*, S1 RNA binding domain 1; *SRD5A2*, steroid 5 alpha-reductase 2; *SPTBN1*, spectrin beta, non-erythrocytic 1; *SQSTM1*, sequestosome 1; *STRN*, Striatin; *SWAP70*, switching B cell complex subunit SWAP70; *TACR1*, tachykinin receptor 1; *TANC1*, tetratricopeptide repeat, ankyrin repeat and coiled-coil containing 1; *TCF12*, transcription factor 12; *TFG*, trafficking from ER to golgi regulator; *THADA*, THADA (thyroid adenoma associated) armadillo repeat containing; *TNIP2*, TNFAIP3 interacting protein 2; *TOGARAM2*, TOG array regulator of axonemal microtubules 2; *TPR*, translocated promoter region, nuclear basket protein; *TRIM66*, tripartite motif containing 66; *TSPYL6*, TSPY like 6; *TTC27*, tetratricopeptide repeat domain 27; *TUBB*, tubulin beta class I; *VIT*, vitrin; *VKORC1L1*, vitamin K epoxide reductase complex subunit 1 like 1; *WDR37*, WD repeat domain 37; *WDPCP*, WD repeat containing planar cell polarity effector; *WNK3*, WNK lysine deficient protein kinase 3; *YAP1*, Yes1 associated transcriptional regulator.

Table 2. List of Chromosomal Locations of Intergenic Translocations With Potential Fusion Partners

No.	Year Published in Print/Presented	Chromosomal Location	Potential Fusion Partner Gene	Response to ALK TKI at the Time of Publication	Tumor Source	Method of Detection	Variant Frequency in Tumor	FISH/IHC	References
1	2019	12q23.3	<i>RIC8B</i>	NR	Tumor	NGS	NR	ND/NR	Zhao, 2019 ⁴⁴
2	2019	2p21	<i>LOC388942</i> (<i>LINC01913</i>)	NR	Tumor	NGS	NR	ND/NR	Zhao, 2019 ⁴⁴
	2020	2p21	<i>LOC388942</i> (<i>LINC01913</i>)	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
3	2019	2q22.1-q22.2	<i>LRP1B</i>	NR	Tumor	NGS	NR	ND/NR	Zhao, 2019 ⁴⁴
4	2019	2p16.2	<i>MIR4431</i>	NR	Tumor	NGS	NR	ND/NR	Zhao, 2019 ⁴⁴
5	2019	2p23.3	<i>CENPA/DPYSL5</i>	PR to crizotinib	Tumor	NGS	NR	+/+	Fei, 2019 ⁶³
6	2020	18q12.1	<i>CDH2</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
7	2020	18q12.2	<i>CELF4</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
8	2020	2p23.3	<i>CENPA</i>	PR to crizotinib	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
9	2020	15q13.3	<i>CHRNA7</i>	PR to crizotinib	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
10	2020	2q14.3	<i>CNTNAP5</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
11	2020	2p21	<i>COX7A2L</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
12	2020	2p13.2	<i>DYSF</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
13	2020	2p16.3	<i>FSHR</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
14	2020	13q12.11	<i>GJB6</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
15	2020	3q22.3	<i>LINC01210</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
16	2020	2p22.3	<i>MEMO1</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
17	2020	2p22.3	<i>MIR548AD</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
18	2020	4q31.1	<i>MGST2</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
19	2020	2q11.2	<i>PDCL3</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
20	2020	2p22.2	<i>QPCT</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
21	2020	2p23.3	<i>RAB10</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
22	2020	2p22.1	<i>SLC8A1</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
23	2020	2q32.3	<i>STK17B</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
24	2020	6q24.1-q24.2	<i>VTA1</i>	NR	Tumor	NGS	NR	NR/NR	Tian, 2020 ⁵⁹
25	2020	2p22.2	<i>CDC42EP3</i> ^a	No response to crizotinib and alectinib	Plasma	NGS	13.0%	ND/+	Zhang, 2020 ⁶⁰
26	2020	19q13.42	<i>PR11-433C9.2</i> (<i>PRPF31</i>)	NR	Tumor	NGS	18.6%	ND/NR	Zhang, 2020 ⁶⁰
27	2020	3p22.1	<i>RPSA</i>	NR	Tumor	NGS	7.9%	ND/+	Zhang, 2020 ⁶⁰
28	2020	2p23.3	<i>UBXN2A</i>	NR	Tumor	NGS	25.4%	ND/NR	Zhang, 2020 ⁶⁰

^aTogether with *EML4-ALK* (E6, A20) and breakpoint is 3'UTR of *CDC43EP3* to exon 20 of *ALK*. +, positive.

ALK, anaplastic lymphoma kinase; FISH, fluorescence in situ hybridization; IHC, immunohistochemistry; ND, not done; NGS, next-generation sequencing; NR, not reported; PR, partial response; SD, stable disease; *CENPA*, centromere protein A; *CDC42EP3*, *CDC42* effector protein 3; *CDH2*, cadherin 2; *CELF4*, CUGBP Elav-like family member 4; *CNTNAP5*, contactin associated protein family member 5; *COX7A2L*, cytochrome c oxidase subunit 7A2 like; *DPYSL5*, dihydropyrimidinase like 5; *DYSF*, dysferlin; *FSHR*, follicle stimulating hormone receptor; *GJB6*, gap junction protein beta 6; *LINC01210*, long intergenic non-protein coding RNA 1210; *LINC01913*, long intergenic non-protein coding RNA 1913; *LRP1B*, LDL receptor related protein 1B; *MEMO1*, mediator of cell motility 1; *MIR4431*, microRNA 4431; *MIR548AD*, microRNA 548ad; *MGST2*, microsomal glutathione S-transferase 2; *PDCL3*, phosducin like 3; *PRPF31*, pre-mRNA processing factor 31; *QPCT*, glutaminyl-peptide cyclotransferase; *RAB10*, RAB10, member RAS oncogene family; *RIC8B*, RIC8 guanine nucleotide exchange factor B; *RPSA*, ribosomal protein SA; *SLC8A1*, solute carrier family 8 member A1; *STK17B*, serine/threonine kinase 17b; *UBXN2A*, UBX domain protein 2A; *VTA1*, vesicle trafficking 1.

DNA NGS to detect even rare actionable driver mutations such as *NTRK* and *NRG1*.^{3,4} Although *EML4-ALK* (with multiple fusion breakpoints in *EML4*) remains the major fusion variant in *ALK+* NSCLC (accounting for approximately 95% of *ALK* fusion variants⁵), multiple case reports have reported novel *ALK* fusion partners in *ALK+* NSCLC. In this article, we have compiled a list of the *ALK* fusion partners including intergenic rearrangements identified in the literature for easy reference.

Methods and Results

We searched PubMed publications, conference/congress abstracts, and presentations extensively to identify novel *ALK* fusion partners (including noncoding RNAs). We included only those fusion partners that retained the 3'*ALK* kinase domain. Reciprocal/nonreciprocal *ALK* translocations involving 5'-*ALK* gene rearrangements (most frequently *ALK* exons 1-19 fused to a 3'-truncated gene [*ALK-XXX*]) were not listed although these nonfunctional 5'-*ALK* fusion variants are usually listed as *ALK* fusion variants in the literature. Overall, a total of 90 distinct *ALK* fusion partners (including noncoding RNAs) have been identified in the literature (by the end of January 2020) (Table 1). Many of these novel *ALK* fusion variants have been reported to respond to *ALK* TKIs or shown to be *ALK* IHC positive. Twenty-five intergenic rearrangements to exon 20 of *ALK* have also been identified and listed separately in Table 2. Three of these intergenic *ALK* rearrangements have been shown to respond to crizotinib, but the significance of these intergenic rearrangements remains to be determined, including whether functional fusion RNAs are translated from these intergenic rearrangements.

Discussion

With the increasing adoption of NGS for molecular profiling of NSCLC, especially in China, the pace at which new fusion partners are being identified and reported has rapidly increased since 2018. In particular, from 2018 onwards, approximately 65 of the 90 fusion partners reported in the literature (calculated at the time page numbers were assigned for this publication) were almost exclusively identified from China, indicating the widespread use of NGS there. Dual in-frame 3'-*ALK* fusion variants with different 5' fusion partners are now being recognized; however, whether the relative contribution of each of the dual *ALK* fusion variant to oncogenesis depends on the allele frequency of each fusion variant remains to be elucidated. We identified at least 28 intergenic 3'-*ALK* rearrangements. Whether these translate to a functional (and truncated)? *ALK* RNA fusion transcript and whether these intergenic

rearrangements are related to the isolated 3'-*ALK* fusion signals remain to be determined.

The concluding perspectives are as follows:

1. *ALK+* NSCLC is a heterogeneous disease with at least 90 distinct fusion partners identified in the literature by January 2020;
2. It is likely that many more fusion partners and intergenic rearrangements will continue to be identified with the ever-increasing adoption of targeted RNA sequencing and whole transcriptome sequencing owing to the need to identify rare actionable fusions such as *NTRK* and *NRG1* fusions;
3. The role of individual 3'-*ALK* fusion variant in a dual 3'-*ALK* fusion variants will need to be elucidated; and
4. The functional significance of intergenic rearrangements remains to be determined.

We recommend that clinicians from around the world to continue to report these novel fusions or intergenic rearrangements with information on the exon or fusion breakpoints, response to *ALK* TKIs, allele frequency, and if possible, whether the tumor is *ALK* fluorescence in situ hybridization and IHC positive.

References

1. Soda M, Choi YL, Enomoto M, et al. Identification of the transforming *EML4-ALK* fusion gene in non-small-cell lung cancer. *Nature*. 2007;448:561-566.
2. Rikova K, Guo A, Zeng Q, et al. Global survey of phosphotyrosine signaling identifies oncogenic kinases in lung cancer. *Cell*. 2007;131:1190-1203.
3. Benayed R, Offin M, Mullaney K, et al. High yield of RNA sequencing for targetable kinase fusions in lung adenocarcinomas with no mitogenic driver alteration detected by DNA sequencing and low tumor mutation burden. *Clin Cancer Res*. 2019;25:4712-4722.
4. Jonna S, Feldman RA, Swensen J, et al. Detection of *NRG1* gene fusions in solid tumors. *Clin Cancer Res*. 2019;25:4966-4972.
5. Ross JS, Ali SM, Fasan O, et al. *ALK*. Fusions in a Wide Variety of Tumor Types Respond to Anti-*ALK* Targeted Therapy. *Oncologist*. 2017;22:1444-1450.
6. Takeuchi K, Choi YL, Togashi Y, et al. KIF5B-*ALK*, a novel fusion onco-kinase identified by an immunohistochemistry-based diagnostic system for *ALK*-positive lung cancer. *Clin Cancer Res*. 2009;15:3143-3149.
7. Wong DW, Leung EL, Wong SK, et al. A novel KIF5B-*ALK* variant in nonsmall cell lung cancer. *Cancer*. 2011;117:2709-2718.
8. Takeuchi K, Soda M, Togashi Y, et al. RET, ROS1 and *ALK* fusions in lung cancer. *Nat Med*. 2012;18:378-381.
9. Togashi Y, Soda M, Sakata S, et al. KLC1-*ALK*: a novel fusion in lung cancer identified using a formalin-fixed paraffin-embedded tissue only. *PLoS One*. 2012;7:e31323.

10. Majewski IJ, Mittenpergher L, Davidson NM, et al. Identification of recurrent FGFR3 fusion genes in lung cancer through kinome-centred RNA sequencing. *J Pathol.* 2013;230:270-276.
11. Yang Y, Qin SK, Zhu J, et al. A rare STRN-ALK fusion in lung adenocarcinoma identified using next-generation sequencing-based circulating tumor DNA profiling exhibits excellent response to crizotinib. *Mayo Clin Proc Innov Qual Outcomes.* 2017;1:111-116.
12. Nakanishi Y, Masuda S, Iida Y, Takahashi N, Hashimoto S. Case report of non-small cell lung cancer with STRN-ALK translocation: a nonresponder to alectinib. *J Thorac Oncol.* 2017;12:e202-e204.
13. Xu H, Shen J, Xiang J, et al. Characterization of acquired receptor tyrosine-kinase fusions as mechanisms of resistance to EGFR tyrosine-kinase inhibitors. *Cancer Manag Res.* 2019;11:6343-6351.
14. Fang DD, Zhang B, Gu Q, et al. HIP1-ALK, a novel ALK fusion variant that responds to crizotinib. *J Thorac Oncol.* 2014;9:285-294.
15. Hong M, Kim RN, Song JY, et al. HIP1-ALK, a novel fusion protein identified in lung adenocarcinoma. *J Thorac Oncol.* 2014;9:419-422.
16. Ou SH, Klempner SJ, Greenbowe JR, et al. Identification of a novel HIP1-ALK fusion variant in non-small-cell lung cancer (NSCLC) and discovery of ALK I1171 (I1171N/S) mutations in two ALK-rearranged NSCLC patients with resistance to alectinib. *J Thorac Oncol.* 2014;9:1821-1825.
17. Choi YL, Lira ME, Hong M, et al. A novel fusion of TPR and ALK in lung adenocarcinoma. *J Thorac Oncol.* 2014;9:563-566.
18. Shan L, Jiang P, Xu F, et al. BIRC6-ALK, a novel fusion gene in ALK break-apart FISH-negative lung adenocarcinoma, responds to crizotinib. *J Thorac Oncol.* 2015;10:e37-e39.
19. Iyevleva AG, Raskin GA, Tiurin VI, et al. Novel ALK fusion partners in lung cancer. *Cancer Lett.* 2015;362:116-121.
20. Drilon A, Wang L, Arcila ME, et al. Broad, hybrid capture-based next-generation sequencing identifies actionable genomic alterations in lung adenocarcinomas otherwise negative for such alterations by other genomic testing approaches. *Clin Cancer Res.* 2015;21:3631-3639.
21. Kim RN, Choi YL, Lee MS, et al. SEC31A-ALK fusion gene in lung adenocarcinoma. *Cancer Res Treat.* 2016;48:398-402.
22. Ali SM, Hensing T, Schrock AB, et al. Comprehensive genomic profiling identifies a subset of crizotinib-responsive ALK-rearranged non-small cell lung cancer not detected by fluorescence in situ hybridization. *Oncologist.* 2016;21:762-770.
23. Tan DSW, Kim DW, Thomas M, et al. Genetic landscape of ALK+ non-small cell lung cancer (NSCLC) patients (pts) and response to ceritinib in ASCEND-1. *J Clin Oncol.* 2016;34(suppl 15):9064.
24. Li W, Zhang J, Guo L, Chuai S, Shan L, Ying J. Combinational analysis of FISH and immunohistochemistry reveals rare genomic events in ALK fusion patterns in NSCLC that responds to crizotinib treatment. *J Thorac Oncol.* 2017;12:94-101.
25. Vendrell JA, Taviaux S, Béganton B, et al. Detection of known and novel ALK fusion transcripts in lung cancer patients using next-generation sequencing approaches. *Sci Rep.* 2017;7:12510.
26. Tian Q, Deng WJ, Li ZW. Identification of a novel crizotinib-sensitive BCL11A-ALK gene fusion in a non-small cell lung cancer patient. *Eur Respir J.* 2017;49.
27. Qin BD, Jiao XD, Liu K, Wu Y, Zang YS. Identification of a novel EML4-ALK, BCL11A-ALK double-fusion variant in lung adenocarcinoma using next-generation sequencing and response to crizotinib. *J Thorac Oncol.* 2019;14:e115-e117.
28. Noh KW, Lee MS, Lee SE, et al. Molecular breakdown: a comprehensive view of anaplastic lymphoma kinase (ALK)-rearranged non-small cell lung cancer. *J Pathol.* 2017;243:307-319.
29. Jiang J, Wu X, Tong X, et al. GCC2-ALK as a targetable fusion in lung adenocarcinoma and its enduring clinical responses to ALK inhibitors. *Lung Cancer.* 2018;115:5-11.
30. Du X, Shao Y, Gao H, et al. *CMTR1-ALK*: an ALK fusion in a patient with no response to ALK inhibitor crizotinib. *Cancer Biol Ther.* 2018;19:962-966.
31. Hu S, Li Q, Peng W, Feng C, Zhang S, Li C. VIT-ALK, a novel alectinib-sensitive fusion gene in lung adenocarcinoma. *J Thorac Oncol.* 2018;13:e72-e74.
32. Yin J, Zhang Y, Zhang Y, Peng F, Lu Y. Reporting on two novel fusions, *DYSF-ALK* and *ITGAV-ALK*, coexisting in one patient with adenocarcinoma of lung, sensitive to crizotinib. *J Thorac Oncol.* 2018;13:e43-e45.
33. Schrock AB, Zhu VW, Hsieh WS, et al. Receptor tyrosine kinase fusions and BRAF kinase fusions are rare but actionable resistance mechanisms to EGFR tyrosine kinase inhibitors. *J Thorac Oncol.* 2018;13:1312-1323.
34. Zhang M, Wang Q, Ding Y, et al. *CUX1-ALK*, a novel ALK rearrangement that responds to crizotinib in non-small cell lung cancer. *J Thorac Oncol.* 2018;13:1792-1797.
35. Zhu VW, Schrock AB, Bosemani T, Benn BS, Ali SM, Ou SI. Dramatic response to alectinib in a lung cancer patient with a novel *VKORC1L1-ALK* fusion and an acquired ALK T1151K mutation. *Lung Cancer (Auckl).* 2018;9:111-116.
36. Xu CW, Wang WX, Chen YP, et al. Simultaneous VENTANA IHC and RT-PCR testing of ALK status in Chinese non-small cell lung cancer patients and response to crizotinib. *J Transl Med.* 2018;16:93.
37. Ramalingam SS, Cheng Y, Zhou C, et al. Mechanisms of acquired resistance to first-line osimertinib: preliminary data from the phase III FLAURA study. *Ann Oncol.* 2018;29(suppl 8):viii740.
38. Lin H, Ren G, Liang X. A novel EML6-ALK FBXO11-ALK double fusion variant in lung adenocarcinoma and response to crizotinib. *J Thorac Oncol.* 2018;13:e234-e236.
39. Zhao J, Li Q, Lin G, et al. Distribution, differences in clinical characteristics and resistance mechanism of ALK variants in Chinese lung cancer patients. *J Thorac Oncol.* 2018;13:5584.
40. Hu X, Cui Q, Wang M. A novel CAMKMT exon3-ALK exon20 fusion variant was identified in a primary pulmonary mucinous adenocarcinoma. *J Thorac Oncol.* 2019;14:e11-e12.

41. Cao Q, Liu Z, Huang Y, Qi C, Yin X. NCOA1-ALK: a novel ALK rearrangement in one lung adenocarcinoma patient responding to crizotinib treatment. *OncoTargets Ther.* 2019;12:1071-1074.
42. Tsou TC, Gowen K, Ali SM, et al. Variable response to ALK inhibitors in NSCLC with a novel MYT1L-ALK fusion. *J Thorac Oncol.* 2019;14:e29-e30.
43. Hou X, Xu H, Chen L. SRBD1-ALK, a novel ALK fusion gene identified in an adenocarcinoma patient by next-generation sequencing. *J Thorac Oncol.* 2019;14:e72-e73.
44. Zhao R, Zhang J, Han Y, et al. Clinicopathological features of ALK expression in 9889 cases of non-small-cell lung cancer and genomic rearrangements identified by capture-based next-generation sequencing: A Chinese retrospective analysis. *Mol Diagn Ther.* 2019;23:395-405.
45. Fang W, Gan J, Hong S, Lu F, Zhang L. MPRIP-ALK, a novel ALK rearrangement that responds to ALK inhibition in NSCLC. *J Thorac Oncol.* 2019;14:e148-e151.
46. Supplee JG, Milan MSD, Lim LP, et al. Sensitivity of next-generation sequencing assays detecting oncogenic fusions in plasma cell-free DNA. *Lung Cancer.* 2019;134:96-99.
47. He Z, Wu X, Ma S, et al. Next-generation sequencing identified a novel WDPCP-ALK fusion sensitive to crizotinib in lung adenocarcinoma. *Clin Lung Cancer.* 2019;20:e548-e551.
48. Couëtoux du Tertre M, Marques M, Tremblay L, et al. Analysis of the genomic landscape in ALK+ NSCLC patients identifies novel aberrations associated with clinical outcomes. *Mol Cancer Ther.* 2019;18:1628-1636.
49. Zhou X, Shou J, Sheng J, et al. Molecular and clinical analysis of Chinese patients with anaplastic lymphoma kinase (ALK)-rearranged non-small cell lung cancer. *Cancer Sci.* 2019;110:3382-3390.
50. Feng T, Chen Z, Gu J, Wang Y, Zhang J, Min L. The clinical responses of TNIP2-ALK fusion variants to crizotinib in ALK-rearranged lung adenocarcinoma. *Lung Cancer.* 2019;137:19-22.
51. Bai H, Jia X, Jin X, et al. ATAD2B-ALK, a novel fusion in lung adenocarcinoma identified using next-generation sequencing (NGS). *J Thorac Oncol.* 2019;14:S844.
52. Pagan C, Barua S, Hsiao SJ, et al. Targeting SLMAP-ALK-a novel gene fusion in lung adenocarcinoma. *Cold Spring Harb Mol Case Stud.* 2019;5:a003939.
53. Horn L, Whisenant JG, Wakelee H, et al. Monitoring therapeutic response and resistance: analysis of circulating tumor DNA in patients with ALK+ lung cancer. *J Thorac Oncol.* 2019;14:1901-1911.
54. Pan Y, Zhang Y, Ye T, et al. Detection of novel NRG1, EGFR, and MET Fusions in lung adenocarcinomas in the Chinese population. *J Thorac Oncol.* 2019;14:2003-2008.
55. Luo J, Gu D, Lu H, Liu S, Kong J. Coexistence of a novel PRKCB-ALK, EML4-ALK double-fusion in a lung adenocarcinoma patient and response to crizotinib. *J Thorac Oncol.* 2019;14:e266-e268.
56. Wen S, Dai L, Wang L, et al. Genomic signature of driver genes identified by target next-generation sequencing in Chinese non-small cell lung cancer. *Oncologist.* 2019;24:e1070-e1081.
57. Li T, Zhang F, Wu Z, et al. PLEKHM2-ALK: A novel fusion in small-cell lung cancer and durable response to ALK inhibitors. *Lung Cancer.* 2020;139:146-150.
58. Yang Y, Zhou J, Zhou J, et al. Efficacy, safety, and biomarker analysis of ensartinib in crizotinib-resistant, ALK-positive non-small-cell lung cancer: a multicentre, phase 2 trial. *Lancet Respir Med.* 2020;8:45-53.
59. Tian P, Liu Y, Zeng H, et al: Unique molecular features and clinical outcomes in young patients with non-small cell lung cancer harboring ALK fusion genes [epub ahead of print] *J Cancer Res Clin Oncol.* <https://doi.org/10.1007/s00432-019-03116-6>, accessed January 21, 2020.
60. Zhang Y, Zeng L, Zhou C, et al. Detection of non-reciprocal/reciprocal ALK translocation as poor predictive marker in first-line crizotinib-treated ALK-rearranged non-small cell lung cancer patients. *J Thorac Oncol.* 2020. <https://doi.org/10.1016/j.jtho.2020.02.007>.
61. Li Z, Li P, Yan B, et al. Sequential ALK inhibitor treatment benefits patient with leptomeningeal metastasis harboring non-EML4-ALK rearrangements detected from cerebrospinal fluid: A case report. *Thorac Cancer.* 2020;11:176-180.
62. Chen HF, Wang WX, Xu CW, et al. A novel SOS1-ALK fusion variant in a metastatic lung adenocarcinoma patient with remarkable response to crizotinib. *Lung Cancer.* 2020. in press.
63. Fei X, Zhu L, Zhou H, Qi C, Wang C. A novel intergenic region between CENPA and DPYSL5-ALK exon 20 fusion variant responding to crizotinib treatment in a patient with lung adenocarcinoma. *J Thorac Oncol.* 2019;14:e191-e193.